Accepted Manuscript

Title: The extra-intestinal avian pathogenic *Escherichia coli* strain BEN2908 invades avian and human epithelial cells and survives intracellularly

Authors: Nathalie Katy Chanteloup, Gaëlle Porcheron, Bernadette Delaleu, Pierre Germon, Catherine Schouler, Maryvonne Moulin-Schouleur, Philippe Gilot

PII: S0378-1135(10)00348-2
DOI: doi:10.1016/j.vetmic.2010.07.013
Reference: VETMIC 4969

To appear in: *VETMIC*

Received date: 7-5-2010
Revised date: 9-7-2010
Accepted date: 14-7-2010

Please cite this article as: Chanteloup, N.K., Porcheron, G., Delaleu, B., Germon, P., Schouler, C., Moulin-Schouleur, M., Gilot, P., The extra-intestinal avian pathogenic *Escherichia coli* strain BEN2908 invades avian and human epithelial cells and survives intracellularly, *Veterinary Microbiology* (2010), doi:10.1016/j.vetmic.2010.07.013

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Short Communication

The extra-intestinal avian pathogenic *Escherichia coli* strain BEN2908 invades avian and human epithelial cells and survives intracellularly

Nathalie Katy Chanteloup^{1*}, Gaëlle Porcheron¹, Bernadette Delaleu², Pierre Germon¹, Catherine Schouler¹, Maryvonne Moulin-Schouleur¹, Philippe Gilot¹

¹ INRA, UR1282 Infectiologie Animale et Santé Publique, Laboratoire de Pathogénie Bactérienne, F-37380 Nouzilly, France

² INRA, UMR85 Physiologie de la Reproduction et des Comportements, CNRS, Université de Tours, Haras Nationaux, F-37380 Nouzilly, France

* Corresponding author. Tel.: +33 2 47 42 76 79; Fax : +33 2 47 42 77 74.

E-mail address: Nathalie.Chanteloup@tours.inra.fr

Abstract

Extra-intestinal pathogenic *Escherichia coli* (ExPEC) are responsible for a wide range of diseases in humans and animals. By using *in vitro* invasion assays and transmission electron microscopy, we showed that BEN2908, an ExPEC strain of avian origin (also termed APEC for Avian Pathogenic E. coli), is able to usurp cellular endocytic pathways to invade A549 human type II pneumocytes and LMH avian hepatocytes where it is able to survive over several days. Although type 1 fimbriae are the major adhesin of BEN2908, proportions of adherent fimbriated or afimbriated bacteria that entered cells were comparable. Internalization of BEN2908 into human pneumocytes reinforces previous studies indicating that APEC strains could represent a zoonotic risk.

ExPEC / APEC/ intracellular/ epithelial cells / type 1 fimbriae/ zoonotic risk

1. Introduction

Extra-intestinal pathogenic *Escherichia coli* (ExPEC) strains are responsible for a wide range of diseases in humans and animals, including urinary tract infections, neonatal meningitis, septicaemia and pneumonia (Russo and Johnson, 2003). In avian species, ExPEC strains, also termed APEC for avian pathogenic *E. coli*, are responsible for a systemic disease that starts with a respiratory tract infection and evolves into septicaemia and colonization of internal organs (heart, liver, spleen etc.). Phylogenetic studies and virulence-factor pattern analysis have shown close relationships between ExPEC strains of human and avian origin, suggesting that APEC strains could represent a zoonotic risk (Rodriguez-Siek et al., 2005; Moulin-Schouleur et al., 2007).

In the present study, to gather new knowledge on the possible zoonotic aspect of APEC, we examined the ability of the APEC strain BEN2908 to enter and survive intracellularly in human type II pneumocytes, an epithelial cell line originating from a target (lung) of early phase of ExPEC infection. As no avian pneumocyte cell lines are available, we then evaluated the ability of BEN2908 to interact with avian hepatocytes, a cell line originating from a target (liver) of late phase of APEC infection.

2. Materials and methods

2.1. Cell lines, bacterial strains and growth conditions

A549 human type II pneumocytes (Giard et al., 1973) and LMH avian hepatocytes (Kawaguchi et al., 1987) were grown with 5% CO₂ at 37 °C in DMEM and at 40 °C in DMEM/Ham's F12 (1:1), respectively. Media were supplemented with 2 mM L-glutamine and 10% heat-inactivated fetal bovine serum. The APEC strain BEN2908, O2:K1:H5 (*fim*⁺, *pap*⁻) is a nalidixic acid-resistant derivative of strain MT78 which was isolated from the trachea of a chicken with respiratory infection (Dho and Lafont,

1982). Strain DM34 is an isogenic mutant of BEN2908 lacking the *fim* operon encoding type 1 fimbriae. Electron microscopic examination of DM34 revealed only flagella as cell surface appendages (Marc et al., 1998). Bacteria were routinely grown in Luria-Bertani (LB) broth at 37 °C with shaking. Growth of strains BEN2908 and DM34 in LB and cell culture media was found to be similar, under the conditions as described above.

2.2. Invasion, association and survival assays

For invasion assays, A549 and LMH cells were seeded in 24-well plates to reach approximately 2.5×10^5 and 5×10^5 cells per well after 24 h of culture, respectively. Cell monolayers were infected with mid-log-phase bacteria ($OD_{600} \approx 0.350$) at a multiplicity of infection (MOI) of 10 and incubated for 2 h. Cells were washed and remaining extracellular bacteria were killed by incubating for an additional 1h 30 min in medium containing gentamicin (100 µg/ml). Cells were then washed and lysed with sterile water for 30 min at 4 °C. For association assays, the additional incubation with gentamicin was omitted. Inocula, cell-associated (adherent and intracellular) and intracellular bacteria were enumerated by viable counts on LB agar. To provide a true adhesion value, the number of intracellular bacteria was subtracted from the number of cell-associated bacteria. Adhesion and invasion were respectively calculated as the number of adherent or intracellular bacteria, divided by the number of bacteria in the inoculum, and expressed as a percentage. Invasion efficiency was calculated as the number of intracellular bacteria divided by the number of adherent bacteria, and expressed as a percentage.

For invasion-blocking experiments, inhibitors were added to cell monolayers 30 min prior to inoculation with bacteria and maintained throughout the invasion assay. Chlorpromazine, cytochalasin D, filipin III and nystatine were all purchased from

Sigma-Aldrich and stock solutions were prepared in DMSO with the exception of chlorpromazine, which was solubilized in water. The chosen concentration of each inhibitor (0.8 μ M cytochalasin D, 14 μ M chlorpromazine, 25 μ M nystatin and 6 μ M filipin III) was the lowest concentration necessary to obtain the maximal inhibitory effect on bacterial uptake. Invasion efficiency was calculated and expressed in relation to that of the untreated controls. Inhibitors and DMSO, at the final concentrations used, were controlled to have no notable effects on the viability and growth of bacteria (as checked by viable counts) or eukaryotic cells [as checked by trypan blue exclusion assay (Longo-Sorbello et al., 2006)]. DMSO at the used concentrations (0.05% for cytochalasin D and nystatin, 0.1% for filipin III) was proven not to interfere with the adhesion of bacteria with eukaryotic cells. Adhesion, invasion and invasion-blocking experiments were conducted in triplicate and were independently repeated at least three times.

For intracellular survival assays, A549 and LMH cells were seeded in 6-well plates to reach 1×10^6 and 2.5×10^5 cells per well after 24 h of culture, respectively. The invasion assay was performed as described above. To delay cell overgrowth after the initial killing of extracellular bacteria with 100 μ g/ml of gentamicin, the infected cells were cultured in medium containing 2.5% of serum only. At that time, the gentamicin concentration was reduced to 10 μ g/ml (a concentration shown to kill BEN2908) to avoid putative accumulation within cells during prolonged incubation. The medium was changed daily. Intracellular and extracellular bacteria were enumerated by viable counts on LB agar every 24 h until 192 h post-inoculation. Intracellular survival assay was conducted in duplicate and was independently repeated at least two times.

2.3. Transmission electron microscopy (TEM)

At 2 h post-infection, infected A549 and LMH monolayers were fixed with 2% glutaraldehyde in 0.1 M cacodylate buffer (pH 7.4) for 24 h at 4°C. Infected LMH monolayers were also fixed similarly at 72 h post-infection. They were then postfixed in 2% OsO₄ – 1.5% potassium ferrocyanide – 0.1 M cacodylate buffer for 1h 30 min at room temperature. After dehydration using increasing ethanol concentrations, the monolayers were embedded in Epoxy Resin. Ultrathin sections (70 nm) were collected on copper grids, contrasted with uranyl acetate and lead citrate, and observed with a CM10 electron microscope (Philips, Eindhoven, Netherlands). Analysis software (Soft Imaging System, Germany) was used for image acquisition.

2.4. Statistical analysis

Data were analysed using Student's *t*-test with $P < 0.05$ considered as significant.

3. Results

3.1. Interaction of strain BEN2908 with epithelial cells

The adhesive and invasive abilities of strain BEN2908 were tested on A549 human type II pneumocytes and LMH avian hepatocytes. Figure 1A (black bars) shows that BEN2908 adhered to LMH and A549 cells to the same extent (366% and 317% of the inoculum, respectively). Bacterial uptake occurred in both cell lines although the internalization of BEN2908 was significantly higher ($P = 0.004$) in LMH cells (23.10% of the inoculum) than in A549 cells (0.756% of the inoculum) (Fig. 1B, black bars). The low internalization level of BEN2908 in A549 cells could not be increased by increasing the MOI from 10 to 100 (data not shown).

The ability of BEN2908 to invade LMH and A549 cells was also examined by transmission electron microscopy. Figure 2 illustrates evidence of invasion of both cell lines by BEN2908. Intracellular bacteria were observed both singly and in groups and were clearly enclosed in membrane-bound vacuoles (Figs. 2A to 2D). *E. coli* were observed bound to the cell surface of LMH cells and engulfed by lamellipodia suggesting a zipper-like mechanism of internalization (Fig. 2B, arrowhead). Moreover, some images of synchronized dividing intracellular bacteria suggest that intravacuolar multiplication may occur during infection of LMH cells (Fig. 2C, arrowheads). In infected A549 monolayers, *E. coli* were observed bound to the cell surface and surrounded by microvilli (Figs. 2D and 2F, arrowheads). In some cases, bacteria were associated with endocytic pits at the cell membrane (Fig. 2E, arrowheads), suggesting that lipid rafts/caveolae- or clathrin- mediated endocytic pathways could be involved in BEN2908 internalization.

3.2. Role of type I fimbriae in the interaction of the APEC strain BEN2908 with avian hepatocytes and human type II pneumocytes

As type I fimbriae have been shown to be involved in the interaction of BEN2908 with avian primary pharyngeal or tracheal cells (Marc et al., 1998), we quantified their impact on the interactions of BEN2908 with LMH and A549 cells. To that end, we monitored the adhesive and invasive abilities of strain DM34, a *fim* operon deletion mutant of BEN2908 that does not express type I fimbriae (Marc et al., 1998). As shown in Figure 1A, the adhesion of DM34 with LMH and A549 cells was drastically reduced compared to that of BEN2908 (11.55% versus 366% of the inoculum for LMH cells and 1.805% versus 317% for A549 cells). Similarly, the internalization of DM34 in LMH and A549 cells was significantly lower than that of BEN2908 (0.923% versus 23.10%

of the inoculum and 0.003% versus 0.756%, respectively) (Fig. 1B). Although bacterial uptake occurred in both cell lines, the adhesive and invasive abilities of DM34 towards LMH cells were significantly higher than towards A549 cells ($P = 0.031$ for the capacity of adhesion, $P = 0.022$ for the capacity of invasion). Electron microscopy examination of LMH cells infected with DM34 confirmed the presence of cell-surface adherent and intravacuolar bacteria (data not shown). Such investigations were not performed on A549 cells regarding the very low level of DM34 internalization in this cell type.

3.3. Influence of endocytosis inhibitors on the internalization of BEN2908 into eukaryotic cells

As bacteria were frequently observed near endocytic pits at the cell membrane, we investigated the involvement of the lipid rafts/caveolae and of the clathrin endocytic pathways in BEN2908 uptake. To that end, LMH and A549 cells were treated with inhibitors known to affect these endocytic processes. Nystatin and filipin III, two inhibitors of the lipid rafts/caveolae endocytic pathway, have no effect on bacterial uptake by LMH and A549 cells (Fig. 3A and 3B, black bars). However, cytochalasin D, an inhibitor of actin polymerization, and chlorpromazine, an inhibitor of clathrin-coated pit formation, significantly reduced entry of strain BEN2908 into LMH (96% and 95% inhibition, respectively) and A549 cells (82% and 97% inhibition, respectively) in a dose-dependent manner, giving good indication of the involvement of clathrin in the internalization process (Fig. 3A and 3B, black bars; data not shown). As BEN2908 is able to invade cells independently of type 1 fimbriae, we tested the role of actin and clathrin in this process and we found that cytochalasin D and chlorpromazine also significantly reduced (50% and 76% inhibition, respectively) the entry of strain DM34

into LMH cells (Fig. 3A, white bars). Such tests could not be done on A549 cells due to the very low invasion rate of these cells by DM34.

3.4. Fate of internalized BEN2908 bacteria

We investigated the ability of strain BEN2908 to survive inside LMH and A549 cells (Fig. 4). In LMH cells, the number of intracellular bacteria decreased slowly during the first 48 h post-infection and then remained stable until 192 h post-infection. Electron microscopy examination of LMH cells infected since 72 h confirmed the presence of intracellular bacteria (Fig. 2G). In contrast, in A549 cells the number of intracellular bacteria decreased steadily, reaching 0.6% of the initial number of intracellular bacteria at 120 h post-infection. These data demonstrate that BEN2908 is able to survive inside epithelial cells to a greater or lesser extent depending on the cell lines or host species.

4. Discussion

In this study, we found that BEN2908, an ExPEC strain of avian origin is able to adhere to and invade human pneumocytes and avian hepatocytes. Intracellular bacteria were observed enclosed in membrane-bound vacuoles (Fig. 2). Internalization of this ExPEC strain of avian origin into human pneumocytes indicates that APEC do not have an absolute host specificity for at least one of their entry site. This reinforces previous studies showing the close phylogenetic proximity of ExPEC strains of avian and human origin and indicating that APEC could represent a zoonotic risk (Rodriguez-Siek et al., 2005; Moulin-Schouleur et al., 2007). BEN2908 invades LMH hepatocytes with greater efficiency than A549 pneumocytes, although it adheres similarly to both cell lines (Fig. 1). This difference in invasiveness could be due to cellular type per se and/or species origin of the cells.

Previous *in vivo* studies have shown that BEN2908 is still present in the lungs and liver of chicken 144 h post-inoculation (Dozois et al., 1994). This could be explained in part by the ability of strain BEN2908 to survive intracellularly as shown by intracellular persistence of BEN2908 in the cell lines used in this study (Fig. 4). Although synchronized dividing intravacuolar bacteria were observed in the first time of invasion of LMH cells (Fig. 2C), the number of intracellular bacteria did not increase post-infection (Fig. 4), suggesting that BEN2908 does not replicate within cells. This apparently stable number of intracellular bacteria observed from 48 h to 192 h in LMH cells could have resulted from a balance between intracellular replication and intracellular degradation. However, as the antibiotic was maintained in extracellular media throughout the survival assay, we cannot exclude the possibility of some extracellular release of bacteria which were subsequently killed by the antibiotic.

Our data (Fig. 1) confirm that type 1 fimbriae are the major adhesin of strain BEN2908 but suggest that other minor components are also involved in adhesion and invasion. These putative BEN2908 components have not been identified to date. Unpublished PCR experiments from our laboratory indicated that BEN2908 is devoided of the *pap* genes coding for P fimbriae but factors such as curli, Tsh and Yqi were described to be involved in adhesion and/or invasion of other APEC strains (Provence and Curtiss, 1994; Gophna et al., 2001; Antao et al., 2009). Although type 1 fimbriae are the major adhesin of BEN2908 no significant difference was nevertheless observed between the invasive efficiency of fimbriated and afimbriated strains. Indeed, 7.83% of adherent DM34 bacteria were internalized in LMH cells versus 6.32% of BEN2908 bacteria. In A549 cells, these percentages were 0.232% and 0.207% for DM34 and BEN2908, respectively. This suggests either that the host cell receptors for type 1 fimbriae and for

the other minor adhesin(s) of BEN2908 allow similar rates of bacterial internalization, or that type 1 fimbriae are not directly involved in the BEN2908 internalization process. Bacterial pathogens have developed various strategies to usurp host endocytic pathways and enter epithelial cells (Pizarro-Cerda and Cossart, 2006). Studies of ExPEC entry into epithelial cells have mainly focused on strains responsible for human urinary tract infections. Interaction between type 1 fimbriae of uropathogenic ExPEC and host cells has been shown to trigger localized actin rearrangements (Martinez et al., 2000; Eto et al., 2008) leading to bacterial internalization via endocytosis pathway. Type 1 fimbriae were first described to mediate endocytosis via the caveola pathway (Baorto et al., 1997; Shin et al., 2000; Duncan et al., 2004), but internalization mediated by the clathrin endocytic pathway has recently been reported (Eto et al., 2008). By using inhibitors affecting each of these pathways, we found that only inhibitors affecting actin polymerization and clathrin-coated pits formation reduced BEN2908 entry, providing good indication that its internalization in avian hepatocytes and in human pneumocytes depends on the clathrin endocytic pathway.

In conclusion, our work indicates that the APEC strain BEN2908 is able to usurp cellular endocytic pathways to invade human and avian epithelial cells where it is able to survive over several days.

Acknowledgements

This work was supported by the Era-Net PathoGenoMics programm (ANR-06-PATHO-002-01 grant).

References

- Antao, E.M., Ewers, C., Gurlebeck, D., Preisinger, R., Homeier, T., Li, G., Wieler, L.H., 2009, Signature-tagged mutagenesis in a chicken infection model leads to the identification of a novel avian pathogenic *Escherichia coli* fimbrial adhesin. PLoS One 4, e7796.
- Baorto, D.M., Gao, Z., Malaviya, R., Dustin, M.L., van der Merwe, A., Lublin, D.M., Abraham, S.N., 1997, Survival of FimH-expressing enterobacteria in macrophages relies on glycolipid traffic. Nature 389, 636-639.
- Dho, M., Lafont, J.P., 1982, *Escherichia coli* colonization of the trachea in poultry: comparison of virulent and avirulent strains in gnotoxenic chickens. Avian Dis 26, 787-797.
- Dozois, C.M., Chanteloup, N., Dho-Moulin, M., Bree, A., Desautels, C., Fairbrother, J.M., 1994, Bacterial colonization and in vivo expression of F1 (type 1) fimbrial antigens in chickens experimentally infected with pathogenic *Escherichia coli*. Avian Dis 38, 231-239.
- Duncan, M.J., Li, G., Shin, J.S., Carson, J.L., Abraham, S.N., 2004, Bacterial penetration of bladder epithelium through lipid rafts. J Biol Chem 279, 18944-18951.
- Eto, D.S., Gordon, H.B., Dhakal, B.K., Jones, T.A., Mulvey, M.A., 2008, Clathrin, AP-2, and the NPXY-binding subset of alternate endocytic adaptors facilitate FimH-mediated bacterial invasion of host cells. Cell Microbiol 10, 2553-2567.
- Giard, D.J., Aaronson, S.A., Todaro, G.J., Arnstein, P., Kersey, J.H., Dosik, H., Parks, W.P., 1973, *In vitro* cultivation of human tumors: establishment of cell lines derived from a series of solid tumors. J Natl Cancer Inst 51, 1417-1423.

- 1 Gophna, U., Barlev, M., Seijffers, R., Oelschlager, T.A., Hacker, J., Ron, E.Z., 2001,
2 Curli fibers mediate internalization of *Escherichia coli* by eukaryotic cells.
3 Infect Immun 69, 2659-2665.
- 4 Kawaguchi, T., Nomura, K., Hirayama, Y., Kitagawa, T., 1987, Establishment and
5 characterization of a chicken hepatocellular carcinoma cell line, LMH. Cancer
6 Res 47, 4460-4464.
- 7 Longo-Sorbello, G.S.A., Saydam, G., Banerjee, D., Berniti, J.R., 2006. Cytotoxicity and
8 cell growth assays. In: Celis, J.E. (Ed.), Cell Biology: a Laboratory Handbook,
9 Academic Press, San Diego, pp. 315-324.
- 10 Marc, D., Arne, P., Bree, A., Dho-Moulin, M., 1998, Colonization ability and
11 pathogenic properties of a *fim*- mutant of an avian strain of *Escherichia coli*. Res
12 Microbiol 149, 473-485.
- 13 Martinez, J.J., Mulvey, M.A., Schilling, J.D., Pinkner, J.S., Hultgren, S.J., 2000, Type 1
14 pilus-mediated bacterial invasion of bladder epithelial cells. Embo J 19, 2803-
15 2812.
- 16 Moulin-Schouleur, M., Reperant, M., Laurent, S., Bree, A., Mignon-Grasteau, S.,
17 Germon, P., Rasschaert, D., Schouler, C., 2007, Extraintestinal pathogenic
18 *Escherichia coli* strains of avian and human origin: link between phylogenetic
19 relationships and common virulence patterns. J Clin Microbiol 45, 3366-3376.
- 20 Pizarro-Cerda, J., Cossart, P., 2006, Bacterial adhesion and entry into host cells. Cell
21 124, 715-727.
- 22 Provence, D.L., Curtiss, R., 3rd, 1994, Isolation and characterization of a gene involved
23 in hemagglutination by an avian pathogenic *Escherichia coli* strain. Infect
24 Immun 62, 1369-1380.

- Rodriguez-Siek, K.E., Giddings, C.W., Doetkott, C., Johnson, T.J., Fakhr, M.K., Nolan, L.K., 2005, Comparison of *Escherichia coli* isolates implicated in human urinary tract infection and avian colibacillosis. Microbiology 151, 2097-2110.
- Russo, T.A., Johnson, J.R., 2003, Medical and economic impact of extraintestinal infections due to *Escherichia coli*: focus on an increasingly important endemic problem. Microbes Infect 5, 449-456.
- Shin, J.S., Gao, Z., Abraham, S.N., 2000, Involvement of cellular caveolae in bacterial entry into mast cells. Science 289, 785-788.

Figure legends

Figure 1. Interaction of *E. coli* strains BEN2908 and DM34 with LMH and A549 cells. Adhesive and invasion abilities of strains BEN2908 (■) and DM34 (□) are expressed, on a logarithmic scale, as the percentage of adherent (A) or intracellular (B) bacteria in relation to the number of bacteria in the inoculum. Data are means $\pm$ SEM of at least three independent experiments performed in triplicate. *, $P < 0.05$; **, $P < 0.005$ by Student's *t*-test.

Figure 2. Transmission electron micrographs of epithelial cells infected with *E. coli* strain BEN2908. Ultrathin sections of LMH (A to C and G) and A549 (D to F) monolayers fixed at 2 h (A to F) or 72 h (G) post-infection were observed. Panel E shows a high magnification of cell membrane-associated bacteria of panel D. Arrowheads indicate lamellipodia (B), synchronized dividing intracellular bacteria (C), endocytic pits (E) and microvilli (F).

Figure 3. Effect of different inhibitors on internalization of *E. coli* strains BEN2908 and DM34 by epithelial cells. Invasion efficiency of LMH (A) and A549 (B) cells by strains BEN2908 (■) or DM34 (□) in the presence of inhibitors was calculated and expressed in relation to that of the untreated controls. The inhibitors are cytochalasin D (Cyt D) at 0.8 μ M, chlorpromazine (Chlor) at 14 μ M, nystatine (Nys) at 25 μ M and filipin III (Fil) at 6 μ M. Data are means $\pm$ SEM of at least three independent experiments performed in triplicate. * $P < 0.05$ ** $P < 0.0005$ versus respective control by Student's *t*-test. Internalization of DM34 was not tested in the presence of nystatine and filipin in LMH cells or of any inhibitors in A549 cells.

Figure 4. Intracellular survival of *E. coli* BEN2908 in LMH and A549 cells. Intracellular and extracellular bacteria were enumerated at different times post-infection. Data of intracellular bacteria are means of at least two independent experiments performed in duplicate. SEM not comprised within symbols are indicated by bars. Extracellular bacteria accounted for less than 2% of the total number of bacteria throughout the assay. Viability and growth of infected and non-infected cells were the same during the entire incubation period.

Figure 1

Figure 2
[Click here to download high resolution image](#)

Figure 3

Figure 4