In vitro inhibitory effect of tetrahydrocurcuminoids on *Fusarium proliferatum* growth and Fumonisin B1 biosynthesis

Journal:	<i>Food Additives and Contaminants</i>
Manuscript ID:	TFAC-2010-231.R1
Manuscript Type:	Original Research Paper
Date Submitted by the Author:	30-Sep-2010
Complete List of Authors:	COMA, véronique; Université Bordeaux1, USBB Portes, Elise; Université Bordeaux1, USBB Gardrat, Christian; Université Bordeaux1, USBB RICHARD-FORGET, Florence; INRA, MycSA Castellan, alain; Université Bordeaux1, USBB
Methods/Techniques:	Microbiology
Additives/Contaminants:	Antioxidants, Fumonisinis, Mycotoxins - fungi
Food Types:	Cereals, Cereals and grain

SCHOLARONE™
Manuscripts

In vitro inhibitory effect of tetrahydrocurcuminoids on *Fusarium proliferatum* growth and Fumonisin B₁ biosynthesis

Véronique COMA^{a*}, Elise PORTES^a, Christian GARDRAT^a, Florence RICHARD-FORGET^b, and Alain CASTELLAN^{a *}

^a Université Bordeaux 1, US2B, UMR 5103 CNRS-INRA-UBx, F-33405 Talence, France

^b INRA Bordeaux-Aquitaine, Unité MYCSA, UPR 1264, B.P. 81, F-33883 Villenave d'Ornon, France

* Corresponding authors: v.coma@us2b.u-bordeaux1.fr, phone: +33 5 4000 2913, Fax: +33 5 4000 6439

Abstract

Many plant-pathogens often produce toxic metabolites when growing on food and feed. Some antioxidative components seem to prevent fungal growth and mycotoxin formation. Recently, we synthesised a new class of powerful antioxidative compounds, *i.e.* tetrahydrocurcuminoids, and structure/antioxidant activity relationships have been established. The South West of France is known to produce large amounts of corn, which can be infected by *Fusarium* species, and particularly *F. proliferatum*. In this context, the efficiency of tetrahydrocurcuminoids, which can be obtained from natural curcuminoids, was investigated to control *in vitro* the growth of *F. proliferatum* and the production of its associated mycotoxin fumonisin B₁. The relation between structure and antifungal activity was studied. Tetrahydrocurcumin **THC1**, with two guaiacyl phenolic subunits, showed the highest inhibitory activity (measured as the radial growth on agar medium) against the *F. proliferatum* development (67% inhibition at a concentration of 13.6 $\mu\text{mol.mL}^{-1}$). The efficiencies of **THC2** (36% at a concentration of 11.5 $\mu\text{mol.mL}^{-1}$), which contains syringyl phenolic units and **THC3** (30% at a concentration of 13.6 $\mu\text{mol.mL}^{-1}$), which does not have any substituent on the aromatic rings were relatively close. These results indicate that the simultaneous presence of guaiacyl phenols and the enolic function of the β -diketone moiety play an important role in the inhibition mechanisms. The importance of this combination was confirmed by using n-propylguaiacol and acetylacetone as molecular models. Under the same conditions, ferulic acid and eugenol, other natural phenolic antioxidants, were less efficient in inhibiting fungal growth. **THC1** also reduced fumonisin B₁ production in liquid medium by approximately 35%, 50%, and 75% at concentrations of 0.8, 1.3, and 1.9 $\mu\text{mol.mL}^{-1}$, respectively. These very low inhibitory concentrations show that tetrahydrocurcuminoids could be one of the most promising biobased molecules for the control of mycotoxinogen fungal strains.

Keywords: *Fusarium proliferatum*; Fumonisin B₁; Tetrahydrocurcuminoids; Tetrahydrocurcumin; Phenolic compounds; Natural antioxidants; Antioxidants

Introduction

Fusarium species are among the most studied plant-pathogenic fungi with several species causing diseases on maize, wheat, barley, and other food and feed grain (Placinta et al 1999; Leslie et al 2006; Desjardins 2006). Climatic conditions at harvest and post-harvest treatment of grain are key determinants of fungal infection and spread. Poor post-harvest management can lead to a rapid deterioration in nutritional quality of seeds. Microbial activity can cause undesirable effects in grain, can contribute to heating and losses in dry matter through the utilization of carbohydrates, as energy source, degradation of lipids and proteins, production of volatile metabolites, and reduction of the quality of animal feed and seed (Magan and Aldred 2007). Decreased yield, as well as diminished quality and value of grains, results in significant worldwide economic losses and some methods of preservation have been proposed (Magan and Aldred 2007; Jouany 2007).

In addition, some fungi such as some *Fusarium* species, can produce mycotoxins on the crops, in the field and during the storage. These compounds are toxic, e.g. they can be lethal or carcinogens, mutagens, immunosuppressants, teratogens or oestrogens (Theumer et al 2002). *Fusarium proliferatum* is one of the most prevalent and toxigenic *Fusarium* species in freshly harvested corn in the South West of France. Fumonisin B₁ (FB₁) is generally the most abundant member of the family of mycotoxins generated by *F. proliferatum* (Leslie et al 1992).

Different strategies have been considered to reduce the contamination of human and animal food chains by fumonisins. Efficient drying of maize after harvesting is one of the most effective control strategies (Jouany 2007). However, maize is usually harvested wet (Singh et al 2001) and a delayed drying step results in a significant increase in fumonisin

contamination. At present, post-harvest strategies evaluated to date are mainly based on the use of chemical antifungal agents or modified atmospheres (Jouany 2007). Antioxidants also are known to inhibit fungal growth and associated mycotoxin production for several toxigenic *Fusarium* species (Siranidou et al 2002, Boutigny et al 2008). Among synthetic antioxidants, esters of *para*-hydroxybenzoic acid (paraben) and butylated hydroxy-anisole (BHA) reduced fungal growth and mycotoxin production by these strains (Farnochi et al 2005) whereas butylated hydroxy-toluene (BHT) was less effective on *Aspergillus* ones (Passone et al 2005, 2008, 2009). Consumer perception that the use of industrially synthesized food preservatives may be associated with potential toxicological problems has led to interest in the use of natural antioxidants such as cyclic terpenes (Dambolena et al 2008), vanillic and caffeic acids (Samapundo et al 2007), resveratrol (Aldred et al 2008) and various natural extracts (Sisti et al 2008).

Recently, we have studied a new class of natural phenolic antioxidants, tetrahydrocurcuminoids (THCs) (Castellan et al 2007; Portes et al 2007; Portes et al 2009). They are derived from curcuminoids extracted from the roots of *Curcuma longa* L., commonly called turmeric. THCs are colorless and more potent than yellow curcuminoids as antioxidants (Portes et al 2007); they could, therefore, be used in colorless foods and cosmetics, where synthetic antioxidants are commonly used (Majeed et al 1990, 2008; see also www.tetrahydrocurcuminoids.com). Curcuminoids and tetrahydrocurcuminoids involve similar mechanisms for their antioxidative action (Portes et al 2007), but tetrahydrocurcuminoids are much more photostable than curcuminoids (Castellan et al 2007). The potential antimicrobial activity of tetrahydrocurcuminoids against several Gram-positive bacteria (Venkateswarlu et al 2000) has also attracted interest because these compounds may offer protection against *Listeria monocytogenes*, a human pathogenic bacterium. According to

this promising perspective, we (Portes et al 2009) prepared films with both antibacterial and antioxidative properties from renewable sources that incorporate two efficient tetrahydrocurcuminoids derivatives, **THC1** and **THC2** (Figure 1), into a protonated chitosan film matrix.

The study described in this paper was carried out to determine the time dependence effects of three tetrahydrocurcuminoids at different doses and in two different conditions (agar solid media or liquid medium) on the growth of a particular strain of *F. proliferatum* and on the biosynthesis of the FB₁ mycotoxin. Antifungal properties will be tentatively correlated to THCs structures.

Material and Methods

Chemicals and Reagents

Compounds involved in the study are shown in Figure 1: THCs were synthesized as previously reported (Portes et al 2007); acetylacetone (**AA**), eugenol (**E**) and ferulic acid (**FA**) were commercial products obtained from Sigma-Aldrich France (Saint Quentin Fallavier); 4-n-propylguaicol (**PG**) was prepared by hydrogenation of eugenol in methanol with Pd/C (10%) under hydrogen pressure (80 bars) for 24 hours at 40° C according to Ruffin et al (2002). The solvents were obtained from Sigma-Aldrich France (Saint Quentin Fallavier) and were used as received. Fumonisin B₁ was purchased from Sigma-Aldrich (St Louis MO, USA).

Insert Figure 1

Microorganisms and Culture Conditions

The *F. proliferatum* strain, INRA 212, was from the INRA-MycSa laboratory collection. In liquid cultures, the INRA 212 strain produced predominantly FB₁. Stock cultures were

maintained on Potato dextrose agar (PDA) (Sigma-Aldrich) slants at 4°C. When inoculum was required, the *F. proliferatum* strains were grown on PDA slants at 25°C, in the dark, for 8 days and spore suspension was prepared by adding sterile distilled water to slants followed by gentle shaking. Spores concentrations were evaluated using a Thoma cell.

Antifungal Activity of Compounds

The antifungal activity of the phenolic compounds was determined as follows:

(i) A THC solution (1 mL) at a known concentration was placed on the PDA surface and the medium dried in a laminar flow hood at 25 °C for 30 min. A growth control was performed with water in place of the THCs. The bioactivity of tetrahydrofuran (THF), the solvent used to dissolve the biocides, was determined from experiments with THF in place of THCs. Five microliters of spore suspension at 10^4 spores mL⁻¹ was deposited in the middle of each Petri dish (90 mm diameter) and incubated at 25°C and 75% relative humidity (RH) for fourteen days, that corresponds to 80 % of the colonized area in THF control Petri dishes.

(ii) Fungal growth was determined by measuring the colony radial growth (RG), which was given as the average of two perpendicular diameters length measurements (in cm).

(iii) Radial Growth (RG) was used to calculate inhibition percentages as:

$$\text{Inhibition \%} = 100 (\text{RG}_{\text{THF control}} - \text{RG}_{\text{sample}}) / \text{RG}_{\text{THF control}}$$

Mycotoxin and FB₁ Analyses in Liquid Medium

Liquid-culture experiments were performed in GYEP (Glucose, Yeast Extract, Peptone) medium (Miller et al 1983). Sterile Petri dishes (55 mm diameter) containing 7 mL of GYEP medium were inoculated with the spore suspension to achieve a final concentration of 10^4 spores/mL. One milliliter of **THC1** solution, using THF/water (1v/1v) as the solvent, was added to the medium at an appropriate concentration. Cultures were incubated in the dark at

25°C and 75% RH for 25 days. Mycelium was pelleted by centrifugation (10 000 rpm for 10 min at 4°C), stored at -80°C for two days and then lyophilized and weighted. The culture medium was decanted and stored at -20°C until analyzed for FB₁.

Fumonisin were extracted from culture filtrates by using an adaptation of a described protocol (Shephard et al 1990). Briefly, 2 mL of culture filtrate was adjusted to pH 6.5 with NaOH (10 mol.L⁻¹) and fumonisins were purified by using Bond Elut Strong Anion Exchange (SAX) cartridges (Varian, Palo Alto, CA, USA). Fumonisin were eluted using 4 mL of methanol/acetic acid (99/1, v/v) and evaporated to dryness under a nitrogen stream. Dried samples were dissolved in 200 µL of methanol before high performance liquid chromatography analysis with fluorescence detection (HPLC-FLD). Ten µL of sample were derivatized with 90 µL of *o*-phthalaldehyde reagent. Twenty µL of this solution were injected into the HPLC system within one minute of derivatization. HPLC-FLD analysis was performed as previously described (Shephard et al 1990). The HPLC system consisted of an Agilent LC 1100 series apparatus equipped with a quaternary pump (G1311A), an autosampler (G1329A), a fluorimeter (G1314A) and the ChemStation chromatography manager software (Agilent, Massy, France) with an Equisorb ODS2 5 µm column (length 150 mm; i.d. 4.6 mm). Quantification of FB₁ was based on calibration with solutions prepared from commercially available standards. Results were expressed as µg of FB₁ per mg of mycelium.

Analysis of Results

Antifungal activity of THCs was evaluated using three repetitions conducted in parallel, in triplicate. The fungal radial growth was measured after 2, 3, 6, 7, 8, 9, 10 days of incubation. For mycelium and fumonisin B₁ production, data were obtained from five experiments. For each experiment, five Petri dishes were prepared according to the previously described procedure with *F. proliferatum*, using respectively water, water/THF (1v/1v), and solutions of

THC1 in water/THF at different concentrations (0.8, 1.3, 1.9 and 2.7 $\mu\text{mol.mL}^{-1}$). Another series of Petri dishes was prepared some days later followed by a third one. The experiments were repeated with three replicates per treatment being destructively sampled at days five, ten, fifteen, twenty and twenty five for analysis.

A statistical one- or two-way (treatment, incubation time) ANOVA test was made on all the data.

Results

Antifungal properties of THCs.

The effects of **THC1** at three different concentrations ranging from 2.7 to 13.4 $\mu\text{mol.mL}^{-1}$ against the *F. proliferatum* growth are reported in Figure 2. A lag-phase occurred whatever the concentration. This latent period did not exceed 2 days in all cases. After the lag-phase, the growth rates were affected by **THC1**.

Insert Figure 2

The comparison with control cultures using water, indicated that the solvent (THF) used to dissolve tetrahydrocurcuminoid plays a role in decreasing the fungal growth during the first 7 days ($p < 0.05$). Analysis of all the results displayed in Figure 2 show that the growth of *F. proliferatum* was statistically different when **THC1** was present ($p < 0.01$) and, as the THC concentrations increase, the fungal growth decreases significantly ($p < 0.05$). The inhibition factors were respectively close to 2, 4 and 6 for 2.7, 8.0 and 13.4 $\mu\text{mol.mL}^{-1}$ **THC1** concentrations, respectively. Except for the lowest **THC1** concentration (2.7 $\mu\text{mol.mL}^{-1}$), the inhibitory effect of the biocide was still detectable 10 days after the inoculation for both higher concentrations.

The efficiency of **THC1** to inhibit the *F. proliferatum* development was compared to that of, **THC2** and **THC3**, two other tetrahydrocurcuminoid derivatives (Figure 3)

Insert Figure 3

All THC derivatives inhibited the fungal growth after 10 days of incubation, and higher the concentration, higher the inhibitory properties were. The best results were obtained by **THC1** since the inhibition percentage reached 70% in cultures supplemented with a $13.4 \mu\text{mol.mL}^{-1}$ concentration. The lowest inhibition percentage, close to 30%, was obtained by **THC3**. A comparison of the three THC profile curves showed that **THC2** showed a particular behavior: the inhibition percentage was the lowest at low concentrations (8% at $2.2 \mu\text{mol.mL}^{-1}$) and was greater than that of **THC3** at higher concentrations (40% compared to 30% at $13.4 \mu\text{mol.mL}^{-1}$).

*Action of **THC1** against fumonisin B_1 production.*

Stress factors, e.g. efficient biocides, can increase the production of secondary metabolites in fungal cultures (Edreva et al; 2008). Thus, it was essential to measure the impact of the most active THC on FB_1 production. This study was performed in GYEP liquid cultures supplemented or not with **THC1**. Preliminary results were conducted in order to clarify the antifungal efficiency using four tetrahydrocurcuminoid concentrations (0.8, 1.3, 1.9, $2.7 \mu\text{mol.mL}^{-1}$) in water/THF (1v/1v). The mycelium growth in water/THF solutions, with or without **THC1**, vs incubation time is shown in Figure 4a. A comparison with control cultures in water indicate that THF plays a significant role ($p < 0.05$) in the decrease of mycelium production during the first 20 days. Statistical analysis of results showed that mycelium weights were statistically different ($p < 0.05$) in the presence and in the absence of **THC1** (water/THF), with the noticeable exception of day 15. They depended both on **THC1** treatment and incubation time ($p < 0.05$). Nevertheless, the general tendency was an increase of the mycelium production between day 5 and day 15, followed by a slow decrease after day

15. As the amount of **THC1** increased, the mycelium weight decreased ($p < 0.05$). In addition, a complete inhibition was observed with the highest **THC1** concentration ($2.7 \mu\text{mol.mL}^{-1}$).

Insert Figure 4

To investigate the specific effect of **THC1** on FB_1 biosynthesis, **THC1** concentrations $\leq 1.9 \mu\text{mol.mL}^{-1}$ were used. Fumonisin accumulation was detected as early as day 5 in water control cultures and day 10 in water/THF control cultures. This could be due to a too low fungal growth in cultures, except in water controls. Taking into account this observation, the determination of **THC1** impact on fumonisin FB_1 production will be only discussed from day 10 (Figure 4b). In the water/THF control and in the **THC1** supplemented-cultures at 0.8, 1.3, $1.9 \mu\text{mol.mL}^{-1}$, the total fumonisin production increased almost linearly, from day 10 to day 20, with calculated R^2 of 0.985, 0.999, 0.999 and 0.678, respectively. After day 20, a small decrease of the fumonisin production was observed. This could be due to a stationary production of FB_1 associated to a possible degradation or complexing of the mycotoxin with chemicals of the medium (Meca et al 2010). Statistical analysis of data indicated that all the results obtained after **THC1** treatments were significantly different from those of the control solvent ($p < 0.01$). All the **THC1** treatments had significant effects after day 10 ($p < 0.05$). For day 10, the impact of **THC1** concentrations was not significant ($p \geq 0.06$). The general tendency was a decrease in fumonisin production when **THC1** was present; as **THC1** concentrations increased, the fumonisin amounts decreased, which was statistically evidenced after 10 days of incubation.

FB_1 production measured in 15, 20 and 25 days-old cultures supplemented with **THC1** was respectively reduced by 31, 37 and 32 % (**THC1** concentration: $0.8 \mu\text{mol.mL}^{-1}$); 41, 47 and 58 % (**THC1** concentration: $1.3 \mu\text{mol.mL}^{-1}$) and close to 75% (**THC1** concentration: $1.9 \mu\text{mol.mL}^{-1}$).

Discussion

Antifungal activities of tetrahydrocurcuminoids

A comparison of the antifungal activities of **THC1**, **THC2** and **THC3** suggested that their behavior was strongly related to their structures. The presence of the phenol and methoxy groups in the *ortho* position of the benzene ring was essential in the antifungal activity of THCs. Nevertheless, a second methoxy group in the *ortho* position of the phenol group (**THC2**) led to reduce antifungal activity. Similar behavior on photochemical reactivity and antioxidant properties were observed when **THC2** was incorporated into carboxymethylcellulose (Castellan et al, 2007) or chitosan (Portes et al, 2009) solid-state matrices. The high antifungal activity of the non-phenolic **THC3** (Table 1) was surprising and suggested that the enolic part of the molecule can have an important role in the bioactivity of THCs (Sugiyama et al, 2006). We also conducted similar studies with *n*-propylguaiacol (**PG**) to approximate the action of the guaiacyl nucleus; acetylacetone (**AA**) to approximate the action of the enol group; eugenol (**E**) and ferulic acid (**FA**) to assess the role of different substituents in the *para* position of the phenol (Figure 1). The selected concentration of **AA** was equal to that of **THC1** and **THC3** (13.4 $\mu\text{mol.mL}^{-1}$). **PG**, **E** and **FA** were tested at twice the concentration (26.8 $\mu\text{mol.mL}^{-1}$), because the THCs have two benzene rings and one enol group. Results are summarized in Table 1.

Insert Table 1

The percentage inhibition of **THC3** was more than three fold higher than **AA**, which shows that the benzene rings have an important role in the antifungal mechanism. A significant antifungal activity was ascribed to **PG** (49% of growth inhibition after ten incubation days), suggesting that the guaiacyl unit played a considerable role in the bioactive properties. The biocide efficiency of **THC1** was different from the cumulative antifungal activities of **PG** and

1
2
3 **AA**, which indicated that the constituting units, **PG** and **AA**, do not well mimic the behavior
4 of **THC1**. A similar observation was made on the electronic absorption of **THC1** (Castellan
5 et al 2007).
6
7
8
9

10
11
12 Ferulic acid (**FA**) and eugenol (**E**) were introduced in this comparative study, owing to their
13 similarities in structures (Fig. 1) with parts of the THC molecule: **FA** contains both cinnamic
14 acid and guaiacyl structures; **E** has a guaiacyl unit and a non-conjugated double bond. Recent
15 papers showed some interesting data concerning these compounds: **FA** inhibits type B
16 trichothecene biosynthesis (Boutigny et al, 2009); cinnamic acid inhibits the fungal growth of
17 *Neurospora crassa* (Said et al, 2004), *Rhodotorula minuta* (Fujii et al, 2002), *F. oxysporum*
18 (Wu et al, 2008) and **E** has known antifungal activity against *Candida albicans* (Braga et al,
19 2007). In this study, **E** and **FA** did reduce the fungal growth of *F. proliferatum* by practically
20 the same inhibition percentage (*ca* 30 %) but largely lower than the inhibition induced by
21 THC derivatives. So, it seemed that the presence of either the carbon-carbon double bond or
22 the carboxylic acid function in the lateral chain of the guaiacyl unit reduced the antifungal
23 activity.
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42

43 Some differences appear when comparing antioxidant and antifungal activities of **THC1**,
44 **THC2** and **THC3**. Antioxidant properties need the presence of a phenol group in the meta- or
45 para-position of the linking chain and a phenol or a methoxy group as neighbour; moreover,
46 the involvement of the β -diketone enolic part should be weak (Portes et al 2007). Antifungal
47 properties of the THCs seem to be related to the presence of benzene rings, phenol and
48 methoxy groups in the *ortho* position of the benzene rings and also to β -diketone enol. The
49 presence of a syringyl unit (two methoxy groups near the phenol function) in the THC
50
51
52
53
54
55
56
57
58
59
60

structure appears unfavorable. Thus, it is likely that antioxidant and antifungal activities of THC₁ do not involve the same molecular mechanisms.

Action of THC₁ against fumonisin B₁ Production.

THC₁, which performed the best antifungal activity, affected the fungal growth in liquid medium and led to a complete inhibition at a relatively low concentration (2.7 µmol.mL⁻¹). When the fungal strain was not inhibited, THC₁ still influenced FB₁ production, mainly after ten days of incubation. These results suggested that both fungal growth and fumonisin biosynthesis were likely modified by THC₁. Similar results were observed with (i) artificial antioxidants, such as BHA and propylparaben, which limited the fungal growth of *Aspergillus flavus*, *Aspergillus parasitus* and *Fusarium* and the production of associated mycotoxins (Thompson 1992, 1996) and (ii) on acid-type natural antioxidants (caffeic and vanillic acids) on *Fusarium verticillioides*, *F. proliferatum* and *Aspergillus* and FB₁ metabolite (Samapundo et al 2007; Beekrum et al 2003). It is likely that the fungal membrane was affected by THC₁, as described by Torres et al (2003) and Braga et al (2007).

Tetrahydrocurcuminoids, particularly THC₁, appear as a new class of compounds that can inhibit *F. proliferatum* growth and toxic FB₁ biosynthesis at a relative low *in vitro*-concentration. The mechanisms involved in this inhibition require to be elucidated and the potentiality of *in vivo* applications has to be investigated. THC₁ is a metabolite (Pan et al, 1999) of the natural well-known non-toxic curcumin used in food, and its preparation by hydrogenation of the latter can be performed in a large scale for many industrial applications (Majeed M, Badmaev V, 2002; Seyler et al, 2003).

References

- Aldred D, Cairns-Fuller V, Magan N. (2008) Environmental factors affect efficacy of some essential oils and resveratrol to control growth and ochratoxin A production by *Penicillium verrucosum* and *Aspergillus westerdijkae* on wheat grain. J. Stored Prod. Res. 44:341-346.
- Beekrum S, Govinden R, Padayachee T, Odhav B. (2003). Naturally occurring phenols: A detoxification strategy for fumonisin B₁. Food Add. Contamin. 20:490-493.
- Boutigny AL, Barreau C, Atanosova-Penichon V, Verdal-Bonnin, MN, Pinson-Gadais L, Richard-Forget F,. (2009) Ferulic acid, an efficient inhibitor of type B trichlorothecene biosynthesis and Tri gene expression in *Fusarium* liquid cultures. Mycological Research 113: 746-753.
- Braga PC, Dal Sasso M, Culici M, Alfieri M. (2007). Eugenol and thymol, alone or in combination, induce morphological alterations in the envelope of *Candida albicans*. Fitoterapia 78:396-400.
- Castellan A, Ruggiero R, Portes E, Da Silva LG, Grelier S, Gardrat C. (2007). Photophysics and photochemistry of tetrahydrocurcuminoids. J. Photochem. Photobiol. A 190:110-120.
- Dambolena JS, Lopez AG, Canepa MC, Theumer MG, Zygodlo JA, Rubinstein HR. (2008). Inhibitory effect of cyclic terpenes (limonene, menthol, menthone and thymol) on *Fusarium verticillioides* MRC 826 growth and fumonisin B₁ biosynthesis. Toxicon 51:37-44.
- Desjardins AE. (2006). *Fusarium* mycotoxins: Chemistry, Genetics and Biology. American Association of Cereal Chemists, St. Paul, Minn., APS Press.
- Edreva A, Velikova V, Tsonev T, Dagnon S, Aktas L, Gesheva E. (2008). Stress-protective role of secondary metabolites: diversity of functions and mechanisms. Gen. Appl. Plant Physiology 34 (special issue):67-78.
- Farnochi MC, Torres AM, Magan N, Chulze SN (2005). Effects of antioxidants and competing microflora on *Fusarium verticillioides* and *F. proliferatum* populations and fumonisin production on maize grain. J. Stored Prod. Res. 41:211-219.

- Fujii T, Shimaya C, Yano A, Terado K, Sugino H, Fukuda H. (2002). Suppression of the growth of the basidiomycete yeast *Rhodotorula minuta*, by cinnamic acid. *Biotechnol. Lett.* 24:151-154.
- Glenn AE. 2007. Mycotoxigenic *Fusarium* species in animal feed. *Anim. Feed Sci. Technology* 137:213-240.
- Jouany JP. (2007). Methods for preventing, decontaminating and minimizing the toxicity of mycotoxins in feeds. *Anim Feed Sci. Technol.* 137:342-362.
- Leslie JF, Plattner RD, Desjardins AE, Klittich CJR. (1992). Fumonisin B₁ production by strains from different mating populations of *Gibberella fujikuroi* (*Fusarium* section *Liseola*). *Phytopathology* 82:341-345.
- Leslie JF, Summerell B, Bullock S. (2006). *The Fusarium Laboratory Manual*, Ames IA, Blackwell Publishing Professional.
- Magan N, Aldred D. (2007). Post-harvest control strategies: minimizing mycotoxins in the food chain. *Int. J. Food Microbiol.* 119:131-139.
- Majeed M, Badmaev V, Shimakumar U, Rajendran R. (1990). *Curcuminoids: Antioxidant Phytonutrients*. Piscataway, NJ, Nutriscience.
- Majeed M, Badmaev V. (2002) Use of tetrahydrocurcuminoids to regulate physiological and pathological events in the skin and mucosa. American patent US 6653327.
- Majeed M, Prakash L.(2008). Currying skin health with multifunctional curcuminoids. *Household and Personal Care Today.* (1):6-8.
- Meca G, Fernandez-Franzon M, Ritieni A, Font G, Ruiz MJ, Manes J. (2010). Formation of fumonisin B₁-glucose reaction products *in vitro* cytotoxicity, and lipid peroxidation on kidney cells. *J. Agric. Food Chem.* 58:1359-1365.
- Miller JD, Young JC, Trenholm HL. (1983). *Fusarium* toxins in field corn. I. Time course of fungal growth and production of deoxynivalenol and other mycotoxins. *Can. J. Bot.* 61:3080-3087.
- Pan MH, Huang TM, Lin JK. Biotransformation of curcumin through reduction and glucuronidation in mice. (1999) *Drug Metab. Dispos.* 27:486-494.

- Passone MA, Resnik S, Etcheverry MG. (2005) *In vitro* effect of phenolic antioxidants on germination, growth and aflatoxin B₁ accumulation by peanut *Aspergillus* section *Flavi*. J. Appl. Microbiol. 99:682-691.
- Passone MA, Resnik S, Etcheverry MG. 2008. The potential of food grade antioxidants in the control of *Aspergillus* section *Flavi*, interrelated mycoflora and aflatoxin B₁ accumulation on peanut grains. Food Control 19:364-371.
- Passone MA, Ruffino M, Ponzio V, Resnik S, Etcheverry MG. 2009. Postharvest control of peanut *Aspergillus* section *Flavi* populations by a formulation of food-grade antioxidants. Intern. J. Food Microbiol. 131:211-217.
- Placinta CM, D'Mello JPF, Macdonald AMC. (1999). A review of worldwide contamination of cereal grains and animal feed with *Fusarium* mycotoxins. Animal Feed Sci. Technol. 78:21-37.
- Portes E, Gardrat C, Castellan A, Coma V. (2009). Environmentally friendly films based on chitosan and tetrahydrocurcuminoid derivatives exhibiting antibacterial and antioxidant properties. Carbohydr. Polym. 76:574-584.
- Portes E, Gardrat C, Castellan A. (2007) A comparative study of the antioxidant properties of tetrahydrocurcuminoids and curcuminoids. Tetrahedron 63:9092-9099.
- Ruffin B, Grelier S, Nourmamode A, Castellan A. (2002). Attempt to approach the role of phenolic phenylpropanol structures in the photoyellowing of softwood mechanical pulps. Can. J. Chem. 80:1223-1231.
- Said S, Neves FM, Griffiths AJF. (2004). Cinnamic acid inhibits the growth of the fungus *Neurospora crassa* but is eliminated as acetophenone. Int. Biodeter. Biodegrad. 54:1-6.
- Samapundo S, De Meulenaer B, Osei-Nimoh D, Lamboni Y, Debevere J, Devlieghere F. (2007). Can phenolic compounds be used for the protection of corn from fungal invasion and mycotoxin contamination during storage? Food Microbiol. 24:465-473.
- Schmidt E, Jirovetz L, Wiecek K, Buchbauer G, Gochev V, Girova T, Stoyanova A, Geissler M. (2007). Antifungal activity of eugenol and various eugenol-containing essential oils against 38 clinical isolates of *Candida albicans*. J. Essent. Oil Bear. Plants 10:421-429.

- Seyler N, Fiandino C, Candau, D. (2003). Cosmetical or dermatological composition containing a tetrahydrocurcumin derivative and an oil bearing an amide group. European patent EP 0290561.
- Shephard GS, Sydenham EW, Thiel PG, Gelderblom WCA. (1990). Quantitative determination of fumonisins B1 and B2 high-performance liquid chromatography. *J. Liq. Chromatogr.* 13:2077-2087.
- Singh V, Yang P, Moreau RA, Hicks KB, Eckhoff SR. (2001). Effect of harvest moisture content and ambient air drying on Maize fiber oil yield and its phytosterol composition. *Starch/Stärke* 53:635-638.
- Siranidou E, Kang Z, Buchenauer H. (2002). Studies on symptom development, phenolic compounds and morphological defence responses in wheat cultivars differing in resistance to *Fusarium* head blight. *J. Phytopathol.* 150:200-208.
- Sisti M, De Santi M, Fraternale D, Ninfali P, Scoccianti V, Brandi G. (2008). Antifungal activity of *Rubus ulmifolius* Schott standardized in vitro culture. *LWT-Food science and Technology* 41:946-950.
- Sugiyama Y, Kawakishi S, Osawa T. (1996). Involvement of the β -diketone moiety in the antioxidative mechanism of tetrahydrocurcumin. *Biochem. Pharmacol.* 52:519-525.
- Theumer MG, Lopez AG, Masih DT, Chulze SN, Rubinstein HR. (2002). Immunobiological effects of fumonisin B₁ in experimental subchronic mycotoxicoses in rats. *Clin. Diagn. Lab. Immunol.* 9:149-155.
- Thompson DP. (1992) Inhibition of mycelial growth of mycotoxigenic fungi by phenolic antioxidants. *Mycologia* 84:791-793.
- Thompson DP. (1996). Inhibition of growth of mycotoxigenic *Fusarium* species by butylated hydroxyanisole and/or carvacrol. *J. Food Prot.* 59:412-415.
- Torres AM, Ramirez ML, Arroyo M, Chulze SN, Magan N. (2003). Potential use of antioxidants for control of growth and fumonisin production by *Fusarium verticillioides* and *Fusarium proliferatum* on whole maize grain. *Int. J. Food Microbiol.* 83:319-324.
- Venkateswarlu S, Rambabu M, Subbaraju GV, Satyanarayana S. (2000). Synthesis and antibacterial activity of tetrahydrocurcuminoids. *Asian J. Chem.* 12:141-144.

1
2
3 Wu HS, Raza W, Fan JQ, Sun YG, Bao W, Shen QR. (2008). Cinnamic acid inhibits growth
4 but stimulates production of pathogenesis factors by *in vitro* cultures of *Fusarium*
5 *oxysporum* f.sp. *niveum*. J. Agric. Food Chem. 56:1316-1321.
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review Only

Figure Captions

Figure 1: Chemical structures of compounds involved in the study (THCs: tetrahydrocurcuminoids; **PG**: n-propylguaiacol; **E**: eugenol; **AA**: acetylacetone; **FA**: ferulic acid)

Figure 2: *F. proliferatum* radial growth versus time of incubation in presence of **THC1** at different concentrations: (-◇-) THF; (-●-) **THC1** 2.7 $\mu\text{mol.mL}^{-1}$; (-×-) **THC1** 8.1 $\mu\text{mol.mL}^{-1}$; (-■-) **THC1** 13.4 $\mu\text{mol.mL}^{-1}$ (THF is the control-solvent, in absence of **THC1**).

Figure 3: Inhibition percentage of *F. proliferatum* growth in presence of biocides at different concentrations after 10 days incubation at 25 °C and 75 % RH: (-◆-) **THC1**; (-×-) **THC2**; (-▲-) **THC3**.

Figure 4: a) Dry weight mycelium obtained by incubation with *F. proliferatum* after 5, 10, 15, 20 and 25 days in presence of various quantities of **THC1** in liquid medium. (-□-): THF/water 1v/1v; (-△-): **THC1** 0.8 $\mu\text{mol.mL}^{-1}$; (-×-): **THC1** 1.3 $\mu\text{mol.mL}^{-1}$; (-+-): **THC1** 1.9 $\mu\text{mol.mL}^{-1}$; (-○-): **THC1** 2.7 $\mu\text{mol.mL}^{-1}$ (For clarity smoothed line were drawn between points). b) **FB₁** metabolite produced by *F. proliferatum* after 11, 15, 20 and 25 days in presence of various quantities of **THC1** in liquid medium. (-□-): THF/water 1v/1v; (-△-): **THC1** 0.8 $\mu\text{mol.mL}^{-1}$; (-×-): **THC1** 1.3 $\mu\text{mol.mL}^{-1}$; (-+-): **THC1** 1.9 $\mu\text{mol.mL}^{-1}$.

For Peer Review Only

Figure 2

For Peer Review Only

Figure 3

For Peer Review Only

Figure 4

For Peer Review Only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review Only

Figure 5

Table 1: Inhibition percentage of *F. proliferatum* radial growth after 10 days of incubation in presence of biocides (water and THF are control experiments) at 25 °C and 75 % RH.^a

Compounds ^a	Inhibition % (± SD)
Water	0 ^b
THF	0 ^b
THC1 (13.4 µmol.mL ⁻¹)	67 ± 2
THC2 (13.4 µmol.mL ⁻¹)	40 ± 5
THC3 (13.4 µmol.mL ⁻¹)	28 ± 3
AA (13.4 µmol.mL ⁻¹)	9 ± 4
PG (26.8 µmol.mL ⁻¹)	49 ± 4
E (26.8 µmol.mL ⁻¹)	31 ± 4
FA (26.8 µmol.mL ⁻¹)	29 ± 5

^a: See scheme 1; ^b:Petri dishes entirely covered by *F. proliferatum*

Figure 1