

## Derivative-based global sensitivity measures: general links with Sobol' indices and numerical tests

Matieyendou Lamboni, Bertrand Iooss, Anne-Laure Popelin, Fabrice Gamboa

► **To cite this version:**

Matieyendou Lamboni, Bertrand Iooss, Anne-Laure Popelin, Fabrice Gamboa. Derivative-based global sensitivity measures: general links with Sobol' indices and numerical tests. *Mathematics and Computers in Simulation*, Elsevier, 2013, 87, pp.45-54. <hal-00666473v2>

**HAL Id: hal-00666473**

**<https://hal.archives-ouvertes.fr/hal-00666473v2>**

Submitted on 1 Jul 2012

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Derivative-based global sensitivity measures: general links with Sobol' indices and numerical tests

M. Lamboni<sup>ab</sup>, B. Iooss<sup>c1</sup>, A.-L. Popelin<sup>c</sup>, F. Gamboa<sup>b</sup>

<sup>a</sup> *Université Paris Descartes, 45 rue des saints Pères, F-75006, France*

<sup>b</sup> *IMT, F-31062, France*

<sup>c</sup> *EDF R&D, 6 quai Watier, F-78401, France*

---

## Abstract

The estimation of variance-based importance measures (called Sobol' indices) of the input variables of a numerical model can require a large number of model evaluations. It turns to be unacceptable for high-dimensional model involving a large number of input variables (typically more than ten). Recently, Sobol and Kucherenko have proposed the Derivative-based Global Sensitivity Measures (DGSM), defined as the integral of the squared derivatives of the model output, showing that it can help to solve the problem of dimensionality in some cases. We provide a general inequality link between DGSM and total Sobol' indices for input variables belonging to the class of Boltzmann probability measures, thus extending the previous results of Sobol and Kucherenko for uniform and normal measures. The special case of log-concave measures is also described. This link provides a DGSM-based maximal bound for the total Sobol indices. Numerical tests show the performance of the bound and its usefulness in practice.

*Keywords:* Boltzmann measure; Derivative based global sensitivity measure; Global sensitivity analysis; Log-concave measure; Poincaré inequality; Sobol' indices

---

## 1. Introduction

2     With the advent of computing technology and numerical methods, computer  
3     models are now widely used to make predictions on little-known physical phenomena,  
4     to solve optimization problems or to perform sensitivity studies. These complex  
5     models often include hundreds or thousands uncertain inputs, whose uncertainties  
6     can strongly impact the model outputs (De Rocquigny *et al.* [5], Kleijnen [11], Patelli

---

<sup>1</sup>Corresponding author: bertrand.iooss@edf.fr, Phone: +33 1877969, Fax: +33 130878213

1 *et al.* [15]). In fact, it is well known that, in many cases, only a small number of  
2 input variables really act in the model (Saltelli *et al.* [19]). This number is referred  
3 to the notion of the effective dimension of a function (Caffish *et al.* [4]), which is a  
4 useful way to deal with the curse of dimensionality in practical applications.

5 Global Sensitivity Analysis (GSA) methods (Sobol [20], Saltelli *et al.* [19]) are  
6 used to quantify the influence of model input variables (and their interaction effects)  
7 on a model response. It is also an objective way to determine the effective dimen-  
8 sion by using the model simulations (Kucherenko *et al.* [12]). A first class of GSA  
9 methods, called “screening” methods, aim at dealing with a large number of input  
10 variables (from tens to hundreds). An example of screening method is the Morris’  
11 method (Morris [14]), which allows a coarse estimation of the main effects using  
12 only a few model evaluations. While taking into account the interactions between  
13 the indices, the basic form of the Morris method did not compute precise sensitiv-  
14 ity indices associated to the interactions between inputs. The second class of GSA  
15 methods are the popular quantitative methods, mainly based on the decomposition  
16 of the model output variance, which leads to the so-called variance-based methods  
17 and Sobol’ sensitivity indices. It allows computing the main and total effects (called  
18 first order and total Sobol’ indices) of each input variable, as well as interaction ef-  
19 fects. However, for functions with non linear and interaction effects, the estimation  
20 procedures become particularly expensive in terms of number of required model eval-  
21 uations. Hence, for this kind of model, variance-based methods can only be applied  
22 to a limited number of input variables (less than tens).

23 Recently, Sobol and Kucherenko [23, 24] have proposed the so-called Derivative-  
24 based Global Sensitivity Measures (DGSM), which can be seen as a kind of general-  
25 ization of the Morris screening method. DGSM seem computationally more tractable  
26 than variance-based measures, specially for high-dimensional models. They also the-  
27 oretically proved an inequality linking DGSM to total Sobol’ indices in the case of  
28 uniform or Gaussian input variables.

1 In this paper, we investigate this close relationship between total Sobol' indices  
2 and DGSM, by extending this inequality to a large class of Boltzmann probability  
3 measures. We also obtain result for the class of log-concave measures. The paper is  
4 organized as follows: Section 2 recalls some useful definitions of Sobol' indices and  
5 DGSM. Section 3 establishes an inequality between these indices for a large class  
6 of Boltzmann (resp. log-concave) probability measures. Section 4 provides some  
7 numerical simulations on two test models, illustrating how DGSM can be used in  
8 practice. We conclude in Section 5.

## 9 2. Global sensitivity indices definition

### 10 2.1. Variance-based sensitivity indices

11 Let  $Y = f(\mathbf{X})$  be a model output with  $d$  random input variables  $\mathbf{X} = (X_1, \dots, X_d)$ .  
12 If the input variables are independent (assumption A1) and  $\mathbb{E}(f^2(\mathbf{X})) < +\infty$  (as-  
13 sumption A2), we have the following unique Hoeffding decomposition (Efron and  
14 Stein [6]) of  $f(\mathbf{X})$ :

$$f(\mathbf{X}) = f_0 + \sum_j^d f_j(X_j) + \sum_{i < j}^d f_{ij}(X_i, X_j) + \dots + f_{1\dots d}(X_1, \dots, X_d) \quad (2.1)$$

$$= \sum_{u \subset \{1, 2, \dots, d\}} f_u(X_u), \quad (2.2)$$

15 where  $f_0 = \mathbb{E}[f(\mathbf{X})]$  corresponds to the empty subset;  $f_j(X_j) = \mathbb{E}[f(\mathbf{X})|X_j] - f_0$   
16 and  $f_u(X_u) = \mathbb{E}[f(\mathbf{X})|X_u] - \sum_{v \subset u} f_v(X_v)$  for any subset  $u \subset \{1, 2, \dots, d\}$ .

17 By regrouping all the terms in equation (2.1) that contain the variable  $X_j$  ( $j =$   
18  $1, 2, \dots, d$ ) in the function called  $g(\cdot)$ :

$$g(X_j, \mathbf{X}_{\sim j}) = \sum_{u \ni j} f_u(\mathbf{X}_u), \quad (2.3)$$

1 we have the following decomposition:

$$f(\mathbf{X}) = f_0 + g(X_j, \mathbf{X}_{\sim j}) + h(\mathbf{X}_{\sim j}), \quad (2.4)$$

2 where  $\mathbf{X}_{\sim j}$  denotes the vector containing all variables except  $X_j$  and  $h(\cdot) = f(\cdot) -$   
 3  $f_0 - g(\cdot)$ . Notice that this decomposition is also unique under assumptions A1 and  
 4 A2. The function  $g(\cdot)$ , itself, suffices to compute the total sensitivity indices. Indeed,  
 5 it contains all information relating  $f(\mathbf{X})$  to  $X_j$ .

**Definition 2.1.** *Assume that A1, A2 hold, let  $\mu(\mathbf{X}) = \mu(X_1, \dots, X_d)$  be the distribution of the input variables. For any non empty subset  $u \subseteq \{1, 2, \dots, d\}$ , set first*

$$D = \int f^2(\mathbf{x}) d\mu(\mathbf{x}) - f_0^2,$$

$$D_u = \int f_u^2(\mathbf{x}_u) d\mu(\mathbf{x}_u),$$

$$D_u^{tot} = \int \sum_{v \supseteq u} f_v^2(\mathbf{x}_v) d\mu(\mathbf{x}_v). \quad (2.5)$$

7 Further, the first order Sobol sensitivity indices (Sobol [20]) of  $\mathbf{X}_u$  is

$$S_u = \frac{D_u}{D}, \quad (2.6)$$

8 The total sensitivity Sobol index of  $\mathbf{X}_u$  (Homma and Saltelli [8]) is

$$S_{T_u} = \frac{D_u^{tot}}{D}. \quad (2.7)$$

9 The following proposition gives another way to compute the total sensitivity  
 10 indices.

11 **Proposition 2.1.** *Under assumptions A1 and A2, the total sensitivity indices of*  
 12 *variable  $X_j$  ( $j = 1, 2, \dots, d$ ) is obtained by the following formulas:*

$$D_j^{tot} = \int g^2(x_j, \mathbf{x}_{\sim j}) d\mu(\mathbf{x}) \quad (2.8)$$

13 and

$$D_j^{tot} = \frac{1}{2} \int [f(\mathbf{x}) - f(x'_j, \mathbf{x}_{\sim j})]^2 d\mu(\mathbf{x}) d\mu(x'_j). \quad (2.9)$$

14 **Proof 2.1.** *The first formula is an obvious consequence of equation (2.4), and it*  
 15 *is obtained by using the orthogonality of the summands in equation (2.1). Indeed,*

1  $D_j^{tot} = \int \sum_{v \supseteq j} f_v^2(\mathbf{x}_v) d\mu(\mathbf{x}_v) = \int \left[ \sum_{v \supseteq j} f_v(\mathbf{x}_v) \right]^2 d\mu(\mathbf{x}) = \int g^2(x_j, \mathbf{x}_{\sim j}) d\mu(\mathbf{x})$ . The  
2 later formula is proved in Sobol [21].

### 3 2.2. Derivative-based sensitivity indices

4 Derivative-based global sensitivity method uses the second moment of model  
5 derivatives as importance measure. This method is motivated by the fact that a  
6 high value of the derivative of the model output with respect to some input variable  
7 means that a big variation of model output is expected for a variation of the variable.  
8 This method extends the Morris method (Morris [14]). Indeed, it allows to capture  
9 any small variation of the model output due to input variables.

10 DGSM have been first proposed in Sobol and Gresham [22]. Then, they have  
11 been largely studied in Kucherenko *et al.* [13], Sobol and Kucherenko [23, 24] and  
12 Patelli *et al.* [16]. From now on, we assume that the function  $f$  is differentiable.  
13 Two kind of DGSM are defined below:

14 **Definition 2.2.** Assume that A1 holds and that  $\frac{\partial f(\mathbf{X})}{\partial x_j}$  is square-integrable (as-  
15 sumption A3). Then, for  $j = 1, 2, \dots, d$ , we define the DGSM indices by:

$$\begin{aligned} \nu_j &= \mathbb{E} \left[ \left( \frac{\partial f(\mathbf{X})}{\partial x_j} \right)^2 \right] \\ &= \int \left( \frac{\partial f(\mathbf{x})}{\partial x_j} \right)^2 d\mu(\mathbf{x}). \end{aligned} \quad (2.10)$$

16 Let  $w(\cdot)$  is be a bounded measurable function. A weighted version of the last indices  
17 is:

$$\tau_j = \int \left( \frac{\partial f(\mathbf{x})}{\partial x_j} \right)^2 w(x_j) d\mu(\mathbf{x}). \quad (2.11)$$

18 **Remark 2.1.** Sobol and Kucherenko [24] showed that, for a specific weighting func-  
19 tion  $w(x_j) = \frac{1 - 3x_j + 3x_j^2}{6}$  and for a class of linear model with respect to each input  
20 variable (following a uniform distribution over  $[0, 1]$ ), we have  $\tau_j = D_j^{tot}$ .

21 **Remark 2.2.** By bearing in mind the decomposition in equation (2.4), we can re-  
22 place in equations (2.10) and (2.11) the function  $f(\cdot)$  by  $g(\cdot)$ . In general,  $g(\cdot)$  is a

1  $d_1$  ( $d_1 \leq d$ ) dimension function, and this can drastically reduce the number of model  
 2 evaluations for the numerical computation of  $\nu$  or  $\tau$ . Thus, we have:

$$\nu_j = \int \left( \frac{\partial g(\mathbf{x})}{\partial x_j} \right)^2 d\mu(\mathbf{x}). \quad (2.12)$$

3

$$\tau_j = \int \left( \frac{\partial g(\mathbf{x})}{\partial x_j} \right)^2 w(x_j) d\mu(\mathbf{x}), \quad (2.13)$$

### 4 **3. Variance-based sensitivity indices vs. derivative-based sensitivity in-** 5 **dices**

6 As DGSM estimations need much less model evaluations than total Sobol' in-  
 7 dices estimations (Kucherenko *et al.* [13]), it would be interesting to use the DGSM,  
 8 instead of total Sobol' indices, for factors fixing setting. A formal link is there-  
 9 fore necessary to provide a mathematical relation between total Sobol' indices and  
 10 DGSM. Sobol and Kucherenko [23] have established an inequality linking these two  
 11 indices for uniform and Gaussian random variables (maximal bound for  $S_{T_j}$ ). In  
 12 this section, we extend the inequality for Sobolev' space model with the marginal  
 13 distribution of input variables belonging to the class of Boltzmann measure on  
 14  $\mathbb{R}$  (assumption A4). A measure  $\delta$  on  $\mathbb{R}$  is said to be a Boltzmann measure if  
 15 it is absolutely continuous with respect to the Lebesgue measure and its density  
 16  $d\delta(x) = \rho(x)dx = c \exp[-v(x)]dx$ . Here  $v(\cdot)$  is a continuous function and  $c$  a nor-  
 17 malizing constant. Many classical continuous probability measures used in practice  
 18 are Boltzmann measures (see de Rocquigny *et al.* [5] and Saltelli *et al.* [19]).

19 The class of Boltzmann probability measures includes the well known class of log-  
 20 concave probability measures. In this case,  $v(\cdot)$  is a convex function (assumption  
 21 A5). In other words, a twice differentiable probability density function  $\rho(x)$  is said  
 22 to be log-concave if, and only if,

$$\frac{d^2}{dx^2} [\log \rho(x)] \leq 0. \quad (3.14)$$

23 Note that the probability measure of uniform density on a finite interval is not

1 continuous on  $\mathbb{R}$ . So it cannot be considered in the class of log-concave probability  
 2 measure, nor in the class of Boltzmann probability measures.

3 The two following propositions give the formal link between Sobol' indices and  
 4 derivative-based sensitivity indices.

5 **Theorem 3.1.** *Under assumptions A1, A2, A3 and A4, we have:*

$$D_j^{tot} \leq C(\mu_j)\nu_j \quad (3.15)$$

6 with  $C(\mu_j) = 4C_1^2$  and  $C_1 = \sup_{x \in \mathbb{R}} \frac{\min(F_j(x), 1 - F_j(x))}{\rho_j(x)}$  the Cheeger constant,  $F_j(\cdot)$   
 7 the cumulative probability function of  $X_j$  and  $\rho_j(\cdot)$  the density of  $X_j$ .

8 We recall the four assumptions:

- 9 • A1: independence between inputs  $X_1, X_2, \dots, X_d$ ,
- 10 • A2:  $f \in L^2(\mathbb{R})$ ,
- 11 • A3:  $\frac{\partial f}{\partial x_j} \in L^2(\mathbb{R})$ ,
- 12 • A4: the distribution of  $X_j$  is a Boltzmann probability measure.

13 **Proof 3.1.** *The resulting inequality (3.15) is based on a one-dimensional  $L^2$ -Poincaré*  
 14 *inequality of the type  $\|u\|_{L^2} \leq C\|\nabla u\|_{L^2}$  for  $u$  a Sobolev' space function (see for ex-*  
 15 *ample [7]). It is applied here to the function  $g(\cdot)$  (equation (2.3), with  $\int g^2(x_j, \mathbf{x}_{\sim j})d\mu(\mathbf{x}) =$*   
 16  *$D_j^{tot}$  (equation (2.8)) and  $\int \left(\frac{\partial g(\mathbf{x})}{\partial x_j}\right)^2 d\mu(\mathbf{x}) = \nu_j$  (equation (2.12)). The constant*  
 17 *is obtained in Bobkov [3], and Fougères [7] for the one-dimensional Poincaré in-*  
 18 *equality. A proof of the  $d$ -dimensional Poincaré inequality is given in Bakry et al.*  
 19 *[2].*

20 **Theorem 3.2.** *Under assumptions A1, A2, A3 and A5, we have:*

$$D_j^{tot} \leq [\exp(v(m))]^2 \nu_j, \quad (3.16)$$

21 with  $C_1 = \frac{\exp(v(m))}{2}$  the Cheeger constant and  $m$  the median of the measure  $\mu_j$   
 22 (such that  $\mu(X_j \leq m) = \mu(X_j > m)$ ).

23 We recall the assumption A5: the distribution of  $X_j$  is a log-concave probability  
 24 measure.



1 **Proof 3.2.** See proof 3.1.

2 Table 1 shows Cheeger constant for some log-concave probability distributions  
3 that are used in practice for uncertainty and sensitivity analyses. We also give  
4 their medians and the functions  $v(\cdot)$ . We obtain the same results for the normal  
5 distribution  $\mathcal{N}(\mu, \sigma^2)$  similar to Sobol and Kucherenko [23] but we prove them in  
6 another way (in this case,  $v(m) = \log(\sigma)$ ). For uniform distribution  $\mathcal{U}[a, b]$ , Sobol  
7 and Kucherenko [23] obtained via direct integral manipulations the inequality  $D_j^{tot} \leq$   
8  $\frac{(b-a)^2}{\pi^2} \nu_j$ . This relation is the classical Poincaré or Wirtinger inequality (Ane *et al.*  
9 [1]).

Distribution	$v(x)$	$m$	$C_1$
Normal $\mathcal{N}(\mu, \sigma^2)$	$\frac{(x-\mu)^2}{2\sigma^2} + \log(\sigma)$	$\mu$	$\frac{\sigma}{2}$
Exponential $\mathcal{E}(\lambda)$ , $\lambda > 0$	$\lambda x - \log(\lambda)$	$\frac{\log 2}{\lambda}$	$\frac{1}{\lambda}$
Beta $\mathcal{B}(\alpha, \beta)$ , $\alpha, \beta \geq 1$	$\log [x^{1-\alpha}(1-x)^{1-\beta}]$	No expression	—
Gamma $\Gamma(\alpha, \beta)$ , scale $\alpha \geq 1$ , shape $\beta > 0$	$\log(x^{1-\alpha}\Gamma(\alpha)) + \frac{x}{\beta} + \alpha \log \beta$	No expression	—
Gumbel $\mathcal{G}(\mu, \beta)$ , scale $\beta > 0$	$\frac{x-\mu}{\beta} + \log \beta + \exp\left(-\frac{x-\mu}{\beta}\right)$	$\mu - \beta \log(\log 2)$	$\frac{\beta}{\log 2}$
Weibull $\mathcal{W}(k, \lambda)$ , shape $k \geq 1$ , scale $\lambda > 0$	$\log\left(\frac{\lambda}{k}\right) + (1-k) \log\left(\frac{x}{\lambda}\right) + \left(\frac{x}{\lambda}\right)^k$	$\lambda(\log 2)^{1/k}$	$\frac{\lambda(\log 2)^{(1-k)/k}}{k}$

Table 1: Standard log-concave probability distributions:  $v(\cdot)$  function, median  $m$  and Cheeger constant  $C_1$  (see Theorem 3.2).

10 For general log-concave measures, no analytical expressions are available for the  
11 Cheeger constant. In this latter case or in case of non log-concave but Boltzmann  
12 measure, we can estimate the Cheeger constant by numerically evaluating the ex-

1 pression  $\sup_{x \in \mathbb{R}} \frac{\min(F_j(x), 1 - F_j(x))}{\rho_j(x)}$ .

## 2 4. Numerical tests

### 3 4.1. Derivative sensitivity indices estimates

4 A classical estimator for the DGSM is the empirical one and is given below:

$$\widehat{v}_j = \frac{1}{n} \sum_{i=1}^n \left( \frac{\partial f(\mathbf{X}^{(i)})}{\partial x_j} \right)^2. \quad (4.17)$$

5 Experimental convergence properties of this estimator are given in Sobol and Kucherenko  
6 [23].

7 From definition (2.4), we know that  $\frac{\partial f(\mathbf{X}^{(i)})}{\partial x_j} = \frac{\partial g(\mathbf{X}^{(i)})}{\partial x_j}$ . Estimator of  $D_j^{tot}$   
8 (see equation (2.8)) and estimator (4.17) are based on the same function  $g(\cdot)$  and  
9 it seems that estimations of these two indices will require approximately the same  
10 number of model evaluations in order to converge towards their respective values.

11 Computation of DGSM and Sobol' indices can be performed with Monte Carlo-  
12 like algorithm, such as Latin Hypercube Sampling, quasi-Monte Carlo and Monte  
13 Carlo Markov Chain sampling. Kucherenko *et al.* [13] have shown that quasi-Monte  
14 Carlo outperforms Monte Carlo when model has a low effective dimension. Com-  
15 putation of DGSM needs model gradient estimation. For complex models, model  
16 gradient computation can easily be obtained by finite difference method. Patelli and  
17 Pradlwarter [15] proposed a Monte Carlo estimation of gradient in high dimension.  
18 They used an unbiased estimator for gradients and have shown that the number of  
19 Monte Carlo evaluations  $n \leq d$  is sufficient for gradient computations. In the worst  
20 case, their procedure requires the same number of model evaluations than the finite  
21 difference method. The method is very efficient when the model has a low effective  
22 dimension.

23 In the following Sections, we compare the estimates of the Sobol indices ( $S_j$  and  
24  $S_{T_j}$ ) and the upper bound of  $S_{T_j}$  (see inequality (3.15)). let denote  $\Upsilon_j$ , the total

1 sensitivity upper bound:

$$\Upsilon_j = C \frac{\nu_j}{D}, \quad (4.18)$$

2 where  $D$  is the variance of the model output  $f(\mathbf{X})$  and  $C = 4C_1^2$ . The goal of our  
 3 numerical tests is just to compare the differences in terms of ranking and not to  
 4 study the speed of convergence of the estimates.

#### 5 4.2. Test on the Morris function

6 As a first test, we consider the Morris function (Morris [14]) that includes 20  
 7 independent and uniform input variables. The Morris function is defined by the  
 8 following equation:

$$y = \beta_0 + \sum_{i=1}^{20} \beta_i w_i + \sum_{i<j}^{20} \beta_{i,j} w_i w_j + \sum_{i<j<l}^{20} \beta_{i,j,l} w_i w_j w_l + \sum_{i<j<l<s}^{20} \beta_{i,j,l,s} w_i w_j w_l w_s, \quad (4.19)$$

9 where  $w_i = 2 \left( x_i - \frac{1}{2} \right)$  except for  $i = 3, 5, 7$  where  $w_i = 2 \left( 1.1 \frac{x_i}{x_i + 1} - \frac{1}{2} \right)$ . The  
 10 coefficient values are:

11  $\beta_i = 20$  for  $i = 1, 2, \dots, 10$ ,

12  $\beta_{i,j} = -15$  for  $i, j = 1, 2, \dots, 6, i < j$

13  $\beta_{i,j,l} = -10$  for  $i, j, l = 1, 2, \dots, 5, i < j < l$

14 and  $\beta_{1,2,3,4} = 5$ .

15 The remaining first and second order coefficients were generated independently from  
 16 the normal distribution  $\mathcal{N}(0, 1)$  and the remaining third and fourth coefficient were  
 17 set to 0.

18 We replace the uniform distributions associated with several input variables by  
 19 different log-concave measures of the Table 1 in order to show how the bounds can  
 20 be used in practical sensitivity analysis. Table 2 shows the probability distributions  
 21 associated to each input of the Morris function.

22 We have performed some simulations that allow computing the DGSM indices  
 23 and the Sobol' indices for the 20 independent factors. Sobol' indices  $S_j$  and  $S_{T_j}$  are

Input	Probability distribution	Input	Probability distribution
X1	$\mathcal{U}[0, 1]$	X11	$\mathcal{U}[0, 1]$
X2	$\mathcal{N}(0.5, 0.1)$	X12	$\mathcal{N}(0.5, 0.1)$
X3	$\mathcal{E}(4)$	X13	$\mathcal{E}(4)$
X4	$\mathcal{G}(0.2, 0.2)$	X14	$\mathcal{G}(0.2, 0.2)$
X5	$\mathcal{W}(2, 0.5)$	X15	$\mathcal{W}(2, 0.5)$
X6	$\mathcal{U}[0, 1]$	X16	$\mathcal{U}[0, 1]$
X7	$\mathcal{U}[0, 1]$	X17	$\mathcal{U}[0, 1]$
X8	$\mathcal{U}[0, 1]$	X18	$\mathcal{U}[0, 1]$
X9	$\mathcal{U}[0, 1]$	X19	$\mathcal{U}[0, 1]$
X10	$\mathcal{U}[0, 1]$	X20	$\mathcal{U}[0, 1]$

Table 2: Probability distributions of the input variables of the Morris function

1 obtained with the principles described in Saltelli [17], i.e. using two initial Monte  
2 Carlo samples of size  $10^4$ . For more efficient convergence properties (specially for  
3 the case of small indices), the improved formulas proposed by Sobol *et al.* [25] for  
4  $S_i$  and by Saltelli *et al.* [18] for  $S_{T_i}$  are used. The approximation errors of these  
5 Monte Carlo estimates are calculated by repeating 20 times the indices estimation  
6 and the mean is taken as the estimate. With  $d = 20$  input variables, it leads to  
7  $20 \times 10^4 \times (d + 2) = 4.4 \times 10^6$  model evaluations. In fact, the size of the Monte  
8 Carlo samples have been fitted to achieve acceptable absolute errors (smaller than  
9 1%). However, the objective here is not to compare the algorithmic performances  
10 of DGSM and Sobol' indices in terms of computational cost, but just to look at the  
11 inputs ranking.

12 The total Sobol' indices are used in this paper as a reference. It shows that  
13 only the first 10 inputs have some influence. Model derivatives are evaluated for  
14 each input on a Monte Carlo sample of size  $1 \times 10^4$  by the finite-difference method  
15 (perturbation of 0.01%). Then, DGSM  $\nu_j$  require  $2.1 \times 10^5$  model evaluations.  $\Upsilon_j$   
16 is then computed using equation (4.18) where the variance of the Morris function is  
17 estimated to  $D = 991.521$ . The results are gathered in Table 3.

18 In Table 3, we can first observe that the total sensitivity upper bounds  $\Upsilon_j$  are  
19 always greater than the total sensitivity indices as expected. For each input, we

Input	$S_j$	$sd$	$S_{T_j}$	$sd$	$\nu_j$	$C$	$\Upsilon_j$
X1	0.043	0.009	0.173	0.008	2043.820	0.101	0.209
X2	0.007	0.003	0.029	0.002	2856.580	0.01	0.029
X3	0.066	0.009	0.165	0.006	31653.270	0.250	7.981
X4	0.002	0.006	0.134	0.007	2025.950	0.333	0.680
X5	0.035	0.005	0.055	0.003	4203.060	0.360	1.526
X6	0.039	0.007	0.114	0.006	1337.100	0.101	0.137
X7	0.068	0.003	0.069	0.003	6605.960	0.101	0.675
X8	0.156	0.007	0.157	0.007	1826.390	0.101	0.187
X9	0.189	0.008	0.192	0.009	2249.770	0.101	0.230
X10	0.145	0.005	0.146	0.005	1730.400	0.101	0.177
X11	0.000	0.001	0.002	0.001	22.630	0.101	0.002
X12	0.000	0.000	0.000	0.000	23.940	0.01	0.000
X13	0.000	0.001	0.001	0.000	17.670	0.250	0.004
X14	0.001	0.001	0.003	0.001	42.850	0.333	0.014
X15	0.000	0.001	0.001	0.001	19.870	0.360	0.007
X16	0.000	0.001	0.002	0.001	18.860	0.101	0.002
X17	0.000	0.001	0.002	0.001	21.400	0.101	0.002
X18	0.000	0.001	0.002	0.001	19.950	0.101	0.002
X19	0.000	0.001	0.004	0.001	54.380	0.101	0.006
X20	0.000	0.001	0.004	0.001	42.250	0.101	0.004

Table 3: Sensitivity indices (Sobol' and DGSM) for the Morris function. For the Sobol' indices  $S_j$  and  $S_{T_j}$ , 20 replicates has been used to get the standard deviation ( $sd$ ).

- 1 distinguish several situations that can occur:
- 2 1. First order and total Sobol' indices are negligible (inputs  $X11$  to  $X20$ ). In this  
3 case, we observe that the bound  $\Upsilon_j$  is always negligible. For all the inputs,  
4 this test shows the high efficiency of the bound: a negligible bound warrants  
5 that the input has no influence.
  - 6 2. First order and total Sobol' indices significantly differ from zero and have  
7 approximately the same value (inputs  $X7$  to  $X10$ ). This means that the input  
8 has some influence but no interactions with other inputs. In this case, the  
9 bound  $\Upsilon_j$  is relevant (close to  $S_{T_j}$ ), except for  $X7$ . The interpretation of the  
10 bound gives a useful information about the total influence of the input.
  - 11 3. First order Sobol' index is negligible while total Sobol' index significantly dif-  
12 fers from zero (inputs  $X1$  to  $X6$ ). In this case, the bound  $\Upsilon_j$  largely oversti-

1        mates the total Sobol' index  $S_{T_j}$  for  $X_3$ ,  $X_4$  and  $X_5$ . However, for  $X_4$ , we  
2        have  $\Upsilon_4 < 1$  and this coarse information is still usefull. For the three other  
3        inputs, the bound is relevant.

4        For two inputs ( $X_3$  and  $X_5$ ), results can be judged as strongly unsatisfactory  
5        as the bound is useless (larger than 1 which is the maximal value for a sensitivity  
6        index). We suspect that these results come from:

- 7        • the model non linearity with respect to these inputs (see equation (4.19)),
- 8        • the input distributions (exponential and Weibull).

9        The second explanation seems to be the more convincing as these types of dis-  
10        tribution can provide larger values during Monte Carlo simulations. In this case,  
11        departures from the central part of the input domain leads to uncontrolled derivative  
12        values of the Morris function. Indeed, it can be seen that  $\nu_j$  is particularly large for  
13         $X_3$  and  $X_5$ , because of high derivative values in the estimation samples. Moreover,  
14        we have no observed the same results for  $X_1$ ,  $X_2$  and  $X_4$ .

15        As a conclusion of this first test, we argue that the bound  $\Upsilon_j$  is well-suited for  
16        a screening purpose. Moreover, coupling  $\Upsilon_j$  interpretation with first order Sobol'  
17        indices  $S_j$  (estimated at low cost using a smoothing technique or a metamodel, see  
18        [19, 9]) can bring useful information about the presence or absence of interaction.  
19        For inputs following uniform, normal and exponential distributions, the bound is  
20        extremely efficient. In these particular cases, the bound is the best one and cannot  
21        be improved.

#### 22    4.3. A case study: a flood model

23        To illustrate how the Cheeger constant can be used for factors prioritization,  
24        when we use the DGSM, we consider a simple application model that simulates the  
25        height of a river compared to the height of a dyke. When the height of a river  
26        is over the height of the dyke, flooding occurs. This academic model is used as a

1 pedagogical example in Iooss [9]. The model is based on a crude simplification of  
 2 the 1D hydro-dynamical equations of Saint Venant under the assumptions of uniform  
 3 and constant flowrate and large rectangular sections. It consists of an equation that  
 4 involves the characteristics of the river stretch:

$$S = Z_v + H - H_d - C_b \quad \text{with} \quad H = \left( \frac{Q}{BK_s \sqrt{\frac{Z_m - Z_v}{L}}} \right)^{0.6}, \quad (4.20)$$

5 with  $S$  the maximal annual overflow (in meters) and  $H$  the maximal annual height  
 6 of the river (in meters).

7 The model has 8 input variables, each one follows a specific probability distribu-  
 8 tion (see Table 4). Among the input variables of the model,  $H_d$  is a design parameter.  
 9 The randomness of the other variables is due to their spatio-temporal variability, our  
 10 ignorance of their true value or some inaccuracies of their estimation. We suppose  
 that the input variables are independent.

Input	Description	Unit	Probability distribution
$Q$	Maximal annual flowrate	m <sup>3</sup> /s	Truncated Gumbel $\mathcal{G}(1013, 558)$ on $[500, 3000]$
$K_s$	Strickler coefficient	-	Truncated normal $\mathcal{N}(30, 8)$ on $[15, +\infty[$
$Z_v$	River downstream level	m	Triangular $\mathcal{T}(49, 50, 51)$
$Z_m$	River upstream level	m	Triangular $\mathcal{T}(54, 55, 56)$
$H_d$	Dyke height	m	Uniform $\mathcal{U}[7, 9]$
$C_b$	Bank level	m	Triangular $\mathcal{T}(55, 55.5, 56)$
$L$	Length of the river stretch	m	Triangular $\mathcal{T}(4990, 5000, 5010)$
$B$	River width	m	Triangular $\mathcal{T}(295, 300, 305)$

Table 4: Input variables of the flood model and their probability distributions

11

12 We also consider another model output: the associated cost (in million euros) of  
 13 the dyke presence,

$$C_p = \mathbb{1}_{S>0} + \left[ 0.2 + 0.8 \left( 1 - \exp^{-\frac{1000}{S^4}} \right) \right] \mathbb{1}_{S \leq 0} + \frac{1}{20} (H_d \mathbb{1}_{H_d > 8} + 8 \mathbb{1}_{H_d \leq 8}), \quad (4.21)$$

14 with  $\mathbb{1}_A(x)$  the indicator function which is equal to 1 for  $x \in A$  and 0 otherwise. In  
 15 this equation, the first term represents the cost due to a flooding ( $S > 0$ ) which is

1 1 million euros, the second term corresponds to the cost of the dyke maintenance  
 2 ( $S \leq 0$ ) and the third term is the investment cost related to the construction of the  
 3 dyke. The latter cost is constant for a height of dyke less than 8 m and is growing  
 4 proportionally with respect to the dyke height otherwise.

5 Sobol' indices are estimated with the same algorithms than for the Morris func-  
 6 tion, using two initial Monte Carlo samples of size  $10^5$  and 20 replicates of the  
 7 estimates. It leads to  $2 \times 10^7$  model evaluations in order to compute first order  
 8 indices  $S_j$  and total indices  $S_{T_j}$  (by taking the mean of the 20 replicates). For es-  
 9 timating the DGSM ( $\nu_j$ , weighted DGSM  $\tau_j$  and the total sensitivity upper bound  
 10  $\Upsilon_j$ ), a Sobol sequence is used with  $1 \times 10^4$  model evaluations.

11 Results of global sensitivity analysis and derivative-based global sensitivity anal-  
 12 ysis for respectively the overflow  $S$  and the cost  $C_p$  outputs are listed in Tables 5  
 13 and 6. Global sensitivity indices show small interaction among input variables for  
 14 the overflow and the cost outputs. Four input variables ( $Q$ ,  $H_d$ ,  $K_s$ ,  $Z_v$ ) drive the  
 15 overflow and the cost outputs. This variable classification will serve as reference for  
 16 comparison issue.

Input	$S_j$	$S_{T_j}$	$\nu_j$	$\tau_j$	$\Upsilon_j$
$Q$	0.343	0.353	1.296e-06	1.072	2.807
$K_s$	0.130	0.139	3.286e-03	1.033	0.198
$Z_v$	0.185	0.186	1.123e+00	1377.41	0.561
$Z_m$	0.003	0.003	2.279e-02	33.742	0.011
$H_d$	0.276	0.276	8.389e-01	23.77	0.340
$C_b$	0.036	0.036	8.389e-01	1268.90	0.105
$L$	0.000	0.000	2.147e-08	0.268	0.000
$B$	0.000	0.000	2.386e-05	1.070	0.000

Table 5: Sensitivity indices for the overflow output of the flood model.

17 Based on derivative sensitivity indices ( $\nu_j$ ) or weighted derivative sensitivity in-  
 18 dices ( $\tau_j$ ) we have obtained another subset of the most influential variables that  
 19 are  $Z_v$ ,  $C_b$ ,  $H_d$ ,  $Z_m$ . These results mean that, for example, the maximum annual  
 20 flowrate ( $Q$ ) does not have any impact on the overflow and the cost output. If we



Input	$S_j$	$S_{T_j}$	$\nu_j$	$\tau_j$	$\Upsilon_j$
$Q$	0.346	0.460	1.3906e-06	2.013	3.011e+00
$K_s$	0.172	0.269	8.5307e-03	1.926	5.129e-01
$Z_v$	0.187	0.229	1.3891e+00	1715.89	6.932e-01
$Z_m$	0.006	0.012	4.6038e-02	68.17	2.29e-02
$H_d$	0.118	0.179	1.5366e+00	44.04	6.227e-01
$C_b$	0.026	0.039	9.4628e-01	1428.69	1.180e-01
$L$	0.000	0.000	4.0276e-08	0.503	2.009e-06
$B$	0.001	0.001	4.4788e-05	2.007	5.587e-04

Table 6: Sensitivity indices for the cost output of the flood model.

1 compare these results to the global sensitivity indices, we can infer that they are  
2 obviously wrong. This is easily explained by the fact that the input variables have  
3 different unities and that the indices  $\nu_j$  and  $\tau_j$  have not been renormalized by the  
4 constant depending on the probability distribution of  $X_j$ .

5 By looking at the total sensitivity upper bound  $\Upsilon_j$ , the most influential variables  
6 are the following:  $Q$ ,  $Z_v$ ,  $H_d$ ,  $K_s$  for the overflow output and for the cost output. It  
7 gives the same subset of the most influential variables with some slight differences  
8 for the prioritization of the most influential variables. In conclusion, we state that  
9  $\Upsilon_j$  can provide correct information on input variance-based sensitivities.

## 10 5. Conclusion

11 Global sensitivity analysis, that allows exploring numerically complex model and  
12 factors fixing setting, requires a large number of model evaluations. Derivative-based  
13 global sensitivity method needs a much smaller number of model evaluations (gain  
14 factor of 10 to 100). The reduction of the number of model evaluations becomes more  
15 significant when the model output is controlled by a small number of input variables  
16 and when the model does not include much interaction among input variables. This  
17 is often the case in practice.

18 In this paper, we have produced an inequality linking the total Sobol' index and  
19 a derivative-based sensitivity measure for a large class of probability distributions

1 (Boltzmann measures). The new sensitivity index  $\Upsilon_j$ , which is defined as a con-  
2 stant times the crude derivative-based sensitivity, is a maximal bound of the total  
3 Sobol' index. It improves factors fixing setting by using derivative-based sensitivities  
4 instead of variance-based sensitivities.

5 Two numerical tests have confirmed that the bound  $\Upsilon_j$  is well-suited for a screen-  
6 ing purpose. When total Sobol' indices cannot be estimated because of a cpu time  
7 expensive model,  $\Upsilon_j$  can provide correct information on input sensitivities. Previous  
8 studies have shown that estimating DGSM with a small derivatives' sample (with  
9 size from tens to hundreds) allows to detect non influent inputs. In subsequent  
10 works, we propose to use jointly DGSM and first order Sobol' indices. With these  
11 information, an efficient methodology of global sensitivity analysis can be applied  
12 and brings useful information about the presence or absence of interaction (see Iooss  
13 *et al.* [10]).

## 14 **6. Acknowledgments**

15 Part of this work has been backed by French National Research Agency (ANR)  
16 through COSINUS program (project COSTA BRAVA noANR-09-COSI-015). We  
17 thank Jean-Claude Fort for helpful discussions and two anonymous reviewers for  
18 their valuable comments.

## 19 **References**

- 20 [1] C. Ané, S. Blachère, D. Chafaï, P. Fougères, I. Gentil, F. Malrieu, C. Roberto,  
21 G. Scheffer, Sur les inégalités de Sobolev logarithmiques, volume 10 of *Panora-*  
22 *mas et Synthèses*, Société Mathématique de France, 2000.
- 23 [2] D. Bakry, F. Barthe, P. Cattiaux, A. Guillin, A simple proof of the Poincaré  
24 inequality for a large class of probability measures including the log-concave  
25 case, *Electron. Commun. Probab.* 13 (2008) 60–6.

- 1 [3] S. Bobkov, Isoperimetric and analytic inequalities for log-concave probability  
2 measures, *The Annals of Probability* 27 (1999) 1903–21.
- 3 [4] R. Caflisch, W. Morokoff, A. Owen, Valuation of mortgage backed securities  
4 using brownian bridges to reduce effective dimension, *Journal of Computational*  
5 *Finance* 1 (1997) 27–46.
- 6 [5] E. de Rocquigny, N. Devictor, S. Tarantola (Eds.), *Uncertainty in industrial*  
7 *practice*, Wiley, 2008.
- 8 [6] B. Efron, C. Stein, The jackknife estimate of variance, *The Annals of Statistics*  
9 9 (1981) 586–96.
- 10 [7] P. Fougères, Spectral gap for log-concave probability measures on the real line,  
11 in: *Séminaire de Probabilités XXXVIII*, volume 1857 of *Lecture Notes in Math.*,  
12 Springer, Berlin, 2005, pp. 95–123.
- 13 [8] T. Homma, A. Saltelli, Importance measures in global sensitivity analysis of  
14 non linear models, *Reliability Engineering and System Safety* 52 (1996) 1–17.
- 15 [9] B. Iooss, Revue sur l’analyse de sensibilité globale de modèles numériques,  
16 *Journal de la Société Française de Statistique* 152 (2011) 1–23.
- 17 [10] B. Iooss, A.L. Popelin, G. Blatman, C. Ciric, F. Gamboa, S. Lacaze, M. Lam-  
18 boni, Some new insights in derivative-based global sensitivity measures, in:  
19 *PSAM 11 & ESREL 2012 Conference*, Helsinki, Finland.
- 20 [11] J. Kleijnen, *Design and analysis of simulation experiments*, Springer, 2008.
- 21 [12] S. Kucherenko, B. Feil, N. Shah, W. Mauntz, The identification of model ef-  
22 fective dimensions using global sensitivity analysis, *Reliability Engineering and*  
23 *System Safety* 96 (2011) 440–9.

- 1 [13] S. Kucherenko, M. Rodriguez-Fernandez, C. Pantelides, N. Shah, Monte Carlo  
2 evaluation of derivative-based global sensitivity measures, *Reliability Engineer-*  
3 *ing and System Safety* 94 (2009) 1135–48.
- 4 [14] M. Morris, Factorial sampling plans for preliminary computational experiments,  
5 *Technometrics* 33 (1991) 161–74.
- 6 [15] E. Patelli, H. Pradlwarter, Monte Carlo gradient estimation in high dimensions,  
7 *International Journal for Numerical Methods in Engineering* 81 (2010) 172–88.
- 8 [16] E. Patelli, H.J. Pradlwarter, G.I. Schuëller, Global sensitivity of structural vari-  
9 ability by random sampling, *Computer Physics Communications* 181 (2010)  
10 2072–81.
- 11 [17] A. Saltelli, Making best use of model evaluations to compute sensitivity indices,  
12 *Computer Physics Communications* 145 (2002) 280–97.
- 13 [18] A. Saltelli, P. Annoni, I. Azzini, F. Campolongo, M. Ratto, S. Tarantola, Vari-  
14 ance based sensitivity analysis of model output. design and estimator for the  
15 total sensitivity index, *Computer Physics Communications* 181 (2010) 259–70.
- 16 [19] A. Saltelli, K. Chan, E. Scott (Eds.), *Sensitivity analysis*, Wiley Series in Prob-  
17 ability and Statistics, Wiley, 2000.
- 18 [20] I. Sobol, Sensitivity analysis for non-linear mathematical models, *Mathematical*  
19 *Modelling and Computational Experiments* 1 (1993) 407–14.
- 20 [21] I. Sobol, Global sensitivity indices for non linear mathematical models and their  
21 Monte Carlo estimates, *Mathematics and Computers in Simulation* 55 (2001)  
22 271–80.
- 23 [22] I. Sobol, A. Gresham, On an alternative global sensitivity estimators, in: *Pro-*  
24 *ceedings of SAMO 1995, Belgirate*, pp. 40–2.

- 1 [23] I. Sobol, S. Kucherenko, Derivative based global sensitivity measures and the  
2 link with global sensitivity indices, *Mathematics and Computers in Simulation*  
3 79 (2009) 3009–17.
- 4 [24] I. Sobol, S. Kucherenko, A new derivative based importance criterion for groups  
5 of variables and its link with the global sensitivity indices, *Computer Physics*  
6 *Communications* 181 (2010) 1212 –7.
- 7 [25] I. Sobol, S. Tarantola, D. Gatelli, S. Kucherenko, W. Mauntz, Estimating the  
8 approximation error when fixing unessential factors in gloabl sensitivity analy-  
9 sis, *Reliability Engineering and System Safety* 92 (2007) 957–0.