Group size, grooming and fission in primates: A modeling approach based on group structure

Cédric Sueur, Jean-Louis Deneubourg, Odile Petit, Iain D. Couzin

► To cite this version:

Cédric Sueur, Jean-Louis Deneubourg, Odile Petit, Iain D. Couzin. Group size, grooming and fission in primates: A modeling approach based on group structure. *Journal of Theoretical Biology*, 2011, 273 (1), pp.156. 10.1016/j.jtbi.2010.12.035 . hal-00669205

HAL Id: hal-00669205

<https://hal.science/hal-00669205>

Submitted on 12 Feb 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Author's Accepted Manuscript

Group size, grooming and fission in primates: A modeling approach based on group structure

Cédric Sueur, Jean-Louis Deneubourg, Odile Petit, Iain D. Couzin

PII: S0022-5193(10)00690-9
DOI: doi:10.1016/j.jtbi.2010.12.035
Reference: YJTBI6304

www.elsevier.com/locate/jtbi

To appear in: *Journal of Theoretical Biology*

Received date: 29 June 2010
Revised date: 15 December 2010
Accepted date: 21 December 2010

Cite this article as: Cédric Sueur, Jean-Louis Deneubourg, Odile Petit and Iain D. Couzin, Group size, grooming and fission in primates: A modeling approach based on group structure, *Journal of Theoretical Biology*, doi:[10.1016/j.jtbi.2010.12.035](https://doi.org/10.1016/j.jtbi.2010.12.035)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Group size, grooming and fission in primates: a modeling approach based on group structure.

Cédric Sueur^{1,2,3}, Jean-Louis Deneubourg¹, Odile Petit² and Iain D. Couzin³

¹: Unit of social ecology, free university of Brussels, Brussels, Belgium

²: Ethologie des Primates, Department of Ecology, Physiology and Ethology, IPHC CNRS-UDS, Strasbourg, France

³: Department of Ecology and Evolutionary Biology, Princeton University, Princeton, U.S.A.

Corresponding author: Cédric Sueur cedric.sueur@c-strasbourg.fr, csueur@princeton.edu

Ethologie des Primates, DEPE, IPHC CNRS-UDS, 23, rue Becquerel 67087 Strasbourg Cedex, France.

Abstract

In social animals, fission is a common mode of group proliferation and dispersion and may be affected by genetic or other social factors. Sociality implies preserving relationships between group members. An increase in group size and/or in competition for food within the group can result in decrease certain social interactions between members, and the group may split irreversibly as a consequence. One individual may try to maintain bonds with a maximum of group members in order to keep group cohesion, i.e. proximity and stable relationships.

However, this strategy needs time and time is often limited. In addition, previous studies have shown that whatever the group size, an individual interacts only with certain grooming partners. There, we develop a computational model to assess how dynamics of group cohesion are related to group size and to the structure of grooming relationships. Groups' sizes after

26 simulated fission are compared to observed sizes of 40 groups of primates. Results showed
27 that the relationship between grooming time and group size is dependent on how each
28 individual attributes grooming time to its social partners, i.e. grooming a few number of
29 preferred partners or grooming equally or not all partners. The number of partners seemed to
30 be more important for the group cohesion than the grooming time itself. This structural
31 constraint has important consequences on group sociality, as it gives the possibility of
32 competition for grooming partners, attraction for high-ranking individuals as found in
33 primates' groups. It could however also have implications when considering the cognitive
34 capacities of primates.

35

36 **Keywords** social network, agent-based model, cohesion, time allocation, population structure,
37 ODD protocol

1. Introduction

Animals have to balance costs and benefits to be in close proximity to conspecifics (Krause and Ruxton 2002). On one hand, living in groups may offer the advantage of a lower predation risk and better efficiency when seeking resources. On the other hand, as group size increases, individuals may experience more within-group competition for food and have higher health risks due to the possible spread of contagious diseases (Krause and Ruxton 2002). Living in group implies interacting frequently with other group members in order to maintain group cohesion (Lehmann et al. 2007). Group cohesion may be defined using three criteria: stability, coordination and proximity. When group size or within-group competition for food increases, disadvantages may outnumber the advantages of group living, (Chapman et al. 1995; Janson and Goldsmith 1995; Ron et al. 1994). As a consequence, group cohesion decreases and the group may split either temporarily (Kerth et al. 2006; Poppa-Lisseanu et al. 2008; Wittemyer et al. 2005) or irreversibly (Henzi et al. 1997a, b; Lehman et al. 2007). In social or pre-social animals, irreversible fission is a common mode of group proliferation and dispersion. From amoeba to primates, this process may be affected by genetic or social factors (Chepko-Sade and Sade 1979; Gompper et al. 1998; Lehman et al. 2007; Seppa et al. 2008; Mehdiabadi et al. 2009; Rangel et al. 2009). In primates, irreversible group fissions rarely occur (about every five/ten years) and separation of different sub-groups often takes several months to years (Chepko-Sade and Sade 1979; Okamoto and Matsumura 2001; Van Horn et al. 2007).

In primates, grooming is considered to be the most common behavior for the maintenance of close social bonds (Schino 2001). Previous studies have shown that when an individual regularly grooms a particular partner, it seems to be more tolerant with this partner and more likely to support it during a conflict (without suggesting causality). Likewise, the partner in question typically reciprocates with the same tolerance and support (Henzi and

Barrett 1999; Dunbar et al. 2009). However, grooming needs time, and time is a limited resource (Dunbar 1992b; Lehmann et al. 2007; Majolo et al. 2008; Pollard and Blumstein 2008; Dunbar et al. 2009). In addition to maintaining social relationships, individuals need to rest, forage and move (Pollard and Blumstein 2008). As a consequence, grooming seldom exceeds 15% of day-time activity for most social species (Dunbar 1991; Lehmann et al. 2007). Some authors have investigated how an individual manages to maintain its social relationships when grooming time is limited but group size or within-group competition for food has increased (Chapman et al. 1995; Dunbar 1992b; Janson and Goldsmith 1995; Lehmann et al. 2007; Majolo et al. 2008; Pollard and Blumstein 2008; Ron et al. 1994). Lehmann et al. (2007) have suggested that when group size and the number of available partners increase, each individual will have to spend more time grooming until a certain group size for which it is impossible to maintain relationships with all group members (Dunbar 1992a; Lehmann et al. 2007; Schino et al. 2009). In this case, group cohesion – social proximity and stability - decreases and group members permanently split. This hypothesis implies that an individual tries to develop and maintain bonds with every group member, or at least, the most of conspecifics. However, other studies have shown that whatever the group size, an individual mainly interacts only with certain grooming partners, and does so much more than with other potential partners (Kudo & Dunbar 2001; Lehmann & Dunbar 2009). Individuals can therefore be considered to have a relatively fixed number of grooming partners. This is the case, for instance, in blue monkey (*Cercopithecus mitis*, Cords 2001), savanna baboons (*Papio ursinus*, Silk et al. 1999), in chimpanzees (*Pan troglodytes*, Watts 2000a,b) and in several macaque species (*Macaca sp.*, Berman et al. 2008; Lin et al. 2008; Nakamichi and Shizawa 2003). In this case, an individual allocates its grooming time to its preferred partners. Consequently, grooming time should not be dependent on group size as it was found in Majolo et al. (2008).

In this study, we wanted to investigate the dynamic of group cohesion – how group cohesion evolves, from stable groups to groups having a greater probability to fission - according to group size and group structure of grooming relationships. By inducing a variation in the group size and the distribution of grooming time in a stochastic agent-based model, we make predictions about the conditions in which a group will irreversibly split. Most of studies on the link between social structure, grooming and group size (Kudo & Dunbar 2001, Lehmann & Dunbar 2009, Lehman et al 2009) followed standard practices in social network analyses and used a criterion for distinguishing casual from meaningful relationships. Modelling permits, without using this arbitrary criterion for preferred relationships, the simulation of interactions between group members (Seyfarth, 1977; Bryson et al. 2007; Meunier et al. 2006; Sellers et al. 2007; Puga-Gonzalez et al., 2009) and also resulting sub-grouping patterns (i.e. how individuals are sub-grouped; Ramos-Fernandez et al. 2006). We attributed a specific grooming time given by each individual to each other conspecifics. We tested the following three different hypotheses for a range of group sizes (from 2 to 200 individuals). (1) An individual divides its grooming time equally among all other group members. (2) An individual does not divide its grooming time equally among all other group members (i.e. it maintains a social bond with all other group members but these bonds are different depending on the partner concerned). (3) An individual divides its grooming time among specific individuals (this number is fixed to (a) 5 and (b) 10 partners per individual, see methods for details). This social structure – grooming a specific number of partners - is suggested by several studies (Berman et al. 2008; Cords 2001; Lin et al. 2008; Nakamichi and Shizawa 2003; Silk et al. 1999, Watts 2000a,b). Once the social structure was established in the model, individuals made a decision between two states (representing the two potential sub-groups). If less than four individuals split from the main group at the simulation end, we considered the cohesion maintained (see previous works on fission: Lefebvre et al. 2003; Ron et al. 1994;

Van Horn et al. 2007). Then, we observed whether, and if so, how the group divided according to its social structure. According to general rules of cohesion or of mimetism (the probability to do a behavior depends on the number of individuals performing this behavior), we expected that if all individuals are linked together, especially by equivalent grooming relationships (hypothesis 1), the system would lead to amplification process – the more individuals join a group, and the more other ones will join it - and no splitting would be observed (Amé et al. 2006; Dussutour et al. 2005; Nicolis et al. 2003, Meunier et al. 2006). Then, group cohesion would be influenced by grooming time and group size when grooming relationships are not equal and especially when grooming is only given to a small number of partners. The simulated data were compared to observed data in order to assess which model most closely corresponds to the observed distribution of group sizes in primates (Lehmann et al. 2007). This comparison allowed us to understand which is the best rule affecting group structure and then population structure. We also tested how increased within-group competition – leading to grooming time decrease by a foraging time increase - influences group cohesion, and whether changes in group cohesion are similar according to group size. We eventually used a path analysis to investigate the relationships (direct and/or indirect) between group size, group structure and group cohesion.

2. Material and Methods

2.1 Data

2.1.1. Empirical data: We used published data about group size and grooming time in order to compare them to our simulation data. Data is taken from 40 published studies on Old World primate species/populations (see Lehman et al., 2007 for details) and are summarized in table 1.

2.1.2. *Theoretical data:* We created theoretical networks using UCINET 6.0 (Borgatti et al. 2002). Groups contained 2 - 200 individuals (2, 5, 10, 20, 40, 60 ... 200). We set networks as random (Erdos-Renyi random graph, with a linear distribution of links). The social structure is different for each generated random network. We did not set networks as scale-free since recent primate studies showed that social networks were not scale-free (i.e. with a power distribution of links) but random (see Flack et al. 2006; MacCowan et al. 2008; Sueur and Petit 2008; Kasper and Voelkl, 2009; Ramos-Fernandez et al. 2009; for studies on primate social networks; see Wasserman and Faust 1994 for social network theory). We can observe on fig.1 that some individuals are only groomed by one partner whilst other ones are groomed by 7 or 8 partners, even if one individual can only groom 5 partners. Then, some social characteristics such as dominance of individuals may be taken into account by considering the network having an Erdos-Renyi random structure. For instance, individuals groomed by a lot of partners on fig.1 might be high-ranking individuals or matriarchs. Indeed, it was shown that these individuals are more groomed than other ones (Nakamichi and Shizawa 2003; Schino 2001; Silk et al. 1999).

2.2. Definitions of parameters

We defined a bond (or a link) in a network as the time one individual groomed another one. Then the relationship is directed and does not need to be reciprocal. The grooming time per individual $T(G)_i$ was defined as the time an individual spends grooming, whatever the number of groomed partners n . According to observed data (see table 1) in this study, we considered grooming to represent no more than 15% of total day-time activity.

For hypothesis 1 (an individual equally grooms all group members) and hypothesis 2 (an individual non-equally/randomly grooms all group members), the mean strength of social relationships (i.e. dyad's social bonds) was equal to the grooming time divided by the number

of group members. For hypothesis 1, all grooming time is equal to $T(G)_i/n$. For hypothesis 2, minimum and maximum of grooming time are respectively 0.4% and 5.4% in a group of 5 individuals and 0.01% and 0.7% in a group of 200 individuals.

As far as hypothesis 3 is concerned (i.e. an individual grooms a fixed number of partners), we carried out simulations with two preferred numbers of groomed partners, $n_1 = 5$ and $n_2 = 10$. These correspond to the average and to the maximum numbers, respectively, of groomed partners found in experimental studies regardless of the group size or the species (Berman et al. 2008; Cords 2001; Lin et al. 2008; Nakamichi and Shizawa 2003; Silk et al. 1999; Watts 2000a,b; Kudo & Dunbar 2001; Lehmann & Dunbar 2009). These studies revealed that the time taken by an individual to groom a non-preferred group member could be considered unimportant compared to the time taken to groom its preferred partners. Previous studies have reported that even if individuals groomed a specific number of individuals, high-ranking individuals or matriarchs can be groomed more than others (Nakamichi and Shizawa 2003; Schino 2001; Silk et al. 1999). We took into account these results to build the theoretical group structure. Even if the number of partners an individual grooms is fixed (to 5 and 10), an individual can be groomed by more (or less) than 5 or 10 grooming partners (see fig.1 for an example).

In our model, we decided to simulate an increased within-group competition by reducing grooming time in steps of 20% (i.e. -20, -40, -60, -80, -100%). We attributed the new social bonds equally for hypothesis 1 and randomly for hypothesis 2. For hypothesis 3, we deleted one partner at each step (i.e. 4, 3, 2, 1 and 0 partner). Indeed, previous studies reported that decreased grooming time seems to have differing affects on an individual's social bonds: in the case of high within-group competition, social bonds were mainly observed amongst kin, and an attraction was observed towards the highest-ranking individuals (Berman et al. 2008;

Majolo et al. 2008; Schino 2001; Watts 2000a,b). We created the networks varying using UCINET 6.0 (Borgatti et al. 2002; Krause et al. 2007; Wey et al. 2007). Group fission was considered to have occurred when the group irreversibly split into two sub-groups (i.e. daughter groups, Ron et al. 1994) containing more than three individuals each. This criterion of three individuals was based on group fission studies (Dunbar 1988; Lefebvre et al. 2003; Ron et al. 1994; Van Horn et al. 2007). It is suggested that if the number of individuals leaving the group is inferior to three individuals (as dispersing males, females with juveniles), it is more considered as dispersion than fission.

2.3 The model

We described the model according to the ODD protocol (i.e. Overview, Design concepts and Details; Grimm et al. 2006).

Purpose

The purpose of the model is to assess how group structure in terms of group size and distribution of grooming between individuals (hypotheses 1 to 3) leads to group fission or allow group cohesion (see 2.2 for definitions). In our model, each individual has to choose between two sub-groups according to the social relationships it has with each individual in each sub-group. Then, the global variable we observed – group fission or group cohesion – is based on the sum of individual decisions.

State variables and scales

The model is based on rules of mimetism/cohesion (Markov chain process) described in several studies on collective phenomena (Amé et al. 2006; Gautrais et al. 2007; Dussutour et al. 2005). In this model, the probability of an individual joining a collective movement in one direction (the future sub-group) depends on the number but also the strength of relationships it has with the individuals already in this direction. The number of individuals, individual

identities and the network of affiliative relationships of each theoretical social network are included in the model. Then, an individual is only characterized by its affiliative relationships, based on grooming time he gave and received from its conspecifics. This model was already used in Sueur et al. (2010). In this study, authors explained how Tonkean and rhesus macaques joined a sub-group during short-term fissions. For both species, the affiliative relationships (i.e. the social network, based on contact between individuals) explain the sub-grouping patterns. Even if individuals are all connected together (one individual was at least observed once with each group member), strength of social relationships leads to fission. In this study, it means that the group fission or cohesion will not only depend on the group size but also on how individuals are connected (hypotheses 1 to 3).

Process overview and scheduling

Each group is characterized by its size (number of individuals per group) and its structure (how individuals are connected). Individuals are characterized by their social relationships depending on the three tested hypotheses and by a state S . At the start of a simulation all group members are in state $s0$ (i.e. group 0, initial group). Then, all individuals will have to choose between state $s1$ (i.e. sub-group 1) and $S2$ (i.e. sub-group 2) according to their own social relationships. This process based on social network will lead to the group cohesion or the group fission. This is the only measure we took into consideration at the end of a simulation. Simulations stop when all group members have changed from state $s0$ to states $s1$ or $s2$. Groomed partners for each group member were attributed randomly (see *Data* for details). The model was then implemented in Netlogo 3.1.4 (Wilensky 1999). We set the number of simulations to 10,000 for each hypothesis and for each set of tested parameters.

Design concepts

Emergence: the only phenomenon emerging from individual decisions in the model is the group fission or cohesion.

Fitness: we did not measure fitness of individuals in this study.

Interaction: individuals are linked to another one by the grooming time they give or they receive (see 2.2 for details).

Sensing: to change state S (1 or 2), individuals take into account an intrinsic probability λ_{iS} and the relationships they have with individuals already in the state S .

Stochasticity: the model is stochastic. At each time step, a number is randomly attributed to each individual and this number will determine if individual will change of state and for which state, according to the probabilities to be in each state.

Collectives: Collectives are represented as social groups of primates. Collectives occur as phenomena emerging from individual behaviour, specifically from the way to choose one sub-group (i.e. state) or another one according to the relationships an individual has in each sug(group (i.e. each state). The collective phenomenon emerging from this choice is the group fission (or the group cohesion).

Observation: We set the number of simulations to 10,000 for each hypothesis and for each set of tested parameters (i.e. group size). Then, for each hypothesis and each group size, we obtained a value of group cohesion (how much time the group staid cohesive up on the 10,000 simulations)

Initialization

At the start of a simulation all group members are in state $s0$ (i.e. group 0, initial group). We then induce a change of state in two randomly chosen individuals: state $s1$ (i.e. sub-group 1) for one individual and state $s2$ (i.e. sub-group 2) for the other. These two individuals are therefore the basis of the formation of the two sub-groups.

Input

At each time step, a number between 0 and 1 is randomly attributed to all other individuals i in state $s0$; when this number is lower than the theoretical probability ψ_{is_1} ($P_1 = [0, \psi_{is_1}]$) the individual changes from state $s0$ to state $s1$; when this number is comprised between ψ_{is_1} and $\psi_{is_1} + \psi_{is_2}$ ($P_2 =]\psi_{is_1}, \psi_{is_1} + \psi_{is_2}]$), then the individual changes from state $s0$ to state $s2$; however, no change of state occurs if this number is superior to $\psi_{is_1} + \psi_{is_2}$ ($P_3 =]\psi_{is_1} + \psi_{is_2}, 1]$), with $P_1 + P_2 + P_3 = 1$.

If $\Sigma r(k, i) = 0$ for an individual i , it changed of state according to its intrinsic probability λ .

The probability ψ_{is} for an individual i in state 0 to turn into state s (1 or 2) was:

$$\psi_{is} = \lambda_i + \left(\sum_{k=1}^{N-1} r(k, i)_s \right)^p$$

where λ_i was the intrinsic probability to change state. We considered that all group members had the same intrinsic probability. $\lambda_i = 0.0001$

p determined the degree of non-linearity in the response shown by individual i . The higher the value of p was, the higher was the resulting discrimination between the both directions (i.e.

the higher the individual probability ψ to go into state s) was, suggesting a deterministic

response in this study (Amé et al. 2006; Dussutour et al. 2005; Nicolis et al. 2003). $p = 5$

λ_i and p were chosen according to previous studies using similar models in primates (Meunier

et al. 2006; Petit et al. 2009; Sueur et al. 2009, 2010; Jacobs et al. in press)

$r(k, i)_s$ was the social bond of i towards k when individual k was already in state s . If individual k was not yet in state s , then $r(k, i)_s = 0$.

$\Sigma r(k, i)_s$ represented the sum of social bonds for individual i in state s .

Submodels

Hypothesis 1: (an individual grooms all group members equally):

$$r(k, i)_s = \omega_{k,i} * T(G)_i$$

where $\omega_{k,i}$ is the weight (strength of social bonds) between individual i and individual k and

$T(G)_i$ is the grooming time per individual.

For this hypothesis,

$$\forall k \in G_i, \omega_{k,i} = 1/(N-1) \quad (\text{all } r(k,i)_s \text{ were equal}).$$

N is the number of individuals in the group.

$$T(G)_i = 0.15$$

Hypothesis 2 (an individual does not groom all group members equally):

$$r(k,i)_s = \omega_{k,i} \cdot T(G)_i$$

with all $\omega_{k,i}$ were attributed randomly according to a normal distribution in order that $\omega_{k,i} > 0$

and $\sum \omega_{k,i} = 1$ for each individual of each group.

$$T(G)_i = 0.15$$

For this second hypothesis, an individual groomed all other group members but almost all the

$r(k,i)_s$ were seen to be different (see *Data* for details).

Hypothesis 3 (an individual grooms a fixed number n of partners):

$$\forall k \in G_i, \omega_{k,i} = 1/n$$

$$\forall k \notin G_i, \omega_{k,i} = 0$$

where n is the number of groomed partners (see *Definitions of parameters* for explanation)

and G_i the group of G partners groomed by the individual i .

2.4 Group structure analyses

Group structure was analyzed using the social networking approach (Krause et al. 2007; Wasserman and Faust 1994; Wey et al. 2008; Whitehead 2009). We used two indexes to determine a group structure.

Group density: number of observed bonds divided by the number of possible bonds in the group.

Group mean path length: an average number of all paths (shortest number of connections between two individuals) between all pairs of individuals in the group. For the same density (i.e. same number of social bonds in a group), group mean path length can be different. This depends on how individuals are connected (see Wasserman and Faust 1994 for details on social network theory; see Flack et al. 2006; MacCowan et al. 2008; Sueur and Petit 2008; Kasper and Voelkl, 2009; Ramos-Fernandez et al. 2009; for studies on primate social networks).

We calculated these indices using UCINET 6.0 (Borgatti et al., 2002).

2.5 Statistical analyses

The relationship between group size and group structure i.e. group mean path length and density, was determined for the three hypotheses. Curve estimation tests were used to assess whether the density and the group mean path length of a given group depends on group size, and to establish the relationship between these variables (linear, logarithmic, and exponential) (Newman et al. 2006). The same curve estimation tests were used to assess the dynamics of group cohesion for each hypothesis according to group size. In order to understand the relations between group cohesion and group size, we finally verified for each hypothesis how group cohesion is affected by group density and group mean path length.

Distributions of group sizes after simulations were compared to observed distributions (data from 40 studied groups, see *Data* for details) using the Kolmogorov-Smirnov test with

Monte-Carlo significance estimation (the number of simulations for this test was set at 10,000). This allows to assess if simulated dynamics of group cohesion/fission fits with global patterns observed in the wild. The theoretical values were obtained by dividing the number of cohesive groups (i.e. that have not split) in each group size by the total number of cohesive groups.

A Mann Whitney test was used to assess how increased within-group competition – a decrease in the grooming time - influences group cohesion within our model. The initial condition – 15% of grooming time – was compared to each other condition (12, 9, 6, 3 and 0%). Analyses were performed using SPSS 10 (SPSS Inc., Chicago, USA).

Path analysis was used to assess direct and indirect effects between group size, group structure and group cohesion. Path analyses and diagrams were carried out using AMOS5 software (AMOS Development Corporation, Spring House, PA, U.S.A.) with maximum likelihood estimations (for non-parametric analyses). This analysis was the most suitable for obtaining the best model of possible relationships and causal effects between variables (identified using AIC –Akaike Information Criterion – values, a measure a goodness of fit of a model).

α was set at 0.05. Means were $\pm$ SE (standard error). Tests are two-tailed.

3. Results

3.1. Dynamics of group cohesion according to group size

3.1.1. Relationship between group structure and group size.

Group mean path length. For both hypotheses 1 (an individual grooms all other group members equally) and 2 (an individual does not groom all other group members equally), the group mean path length was constant (1.00) whatever the group size (fig.2a). For hypothesis 3 (an individual grooms a fixed number of partners), the best fit equation between the group

mean path length and the group size was logarithmic, whatever the number of groomed partners (5 or 10) ($N = 13$ tests group sizes, $F_{1,11} \geq 195.945$, $R^2 \geq 0.947$, $P < 0.000001$; fig.2a), showing that path length and, thus connectivity between individuals highly vary for small group size (until about 40 individuals per group) whilst it is more constant for large group sizes.

Group density. For hypotheses 1 and 2, the density was constant (1.00) whatever the group size (fig.2b). For hypothesis 3, the best fit equation between the group mean path length and the group size was logarithmic, whatever the number of groomed partners (5 or 10), ($N = 13$, $F_{1,11} \geq 96.611$, $R^2 \geq 0.898$, $P < 0.000001$; fig.2b).

3.1.2. Relationship between group cohesion and group size.

For hypothesis 1, group cohesion (defined as the ratio of the number of cases where “one of the sub-groups containing three individuals or less” divided by the total number of simulations) was a constant (1.00; fig.2c). For hypothesis 2, the best fit equation between group cohesion and group size was logarithmic ($N = 13$, $F_{1,11} = 24.255$, $R^2 = 0.68$, $P = 0.0004$; fig.2c). The decrease in group cohesion was only seen to be $0.085 \pm 0.004\%$ in groups of 10 to 200 members. For hypothesis 3, whatever the number of groomed partners (5 or 10), the best fit equation between the group cohesion and the group size was exponential ($N = 13$, $F_{1,11} \geq 327.615$, $R^2 \geq 0.967$, $P < 0.000001$; fig.2c).

These results suggest that when an individual has social bonds with all other group members, whatever the quality of these social bonds, group size has little influence on group cohesion and consequently on group fission probability.

3.1.3. Relationship between group cohesion and group structure.

Group mean path length. For hypotheses 1 and 2, group cohesion was constant (1.00) whatever the group mean path length. For hypothesis 3, the best fit equation between mean path length and group cohesion is negatively linear, whatever the number of groomed partners (5 or 10) ($N = 13$, $F_{1,11} \geq 219.38$, $R^2 \geq 0.952$, $P < 0.000001$).

Group density. For hypotheses 1 and 2, the group cohesion was constant (1.00) whatever the density. For hypothesis 3, the best fit equation between density and group cohesion is logarithmic, whatever the number of groomed partners (5 or 10) ($N = 13$, $F_{1,11} \geq 471.13$, $R^2 \geq 0.977$, $P < 0.000001$).

The equation best describing the relationship between density (and the group mean path length) and group cohesion for 5 partners was similar to that calculated for 10 partners (table 2). These results suggest that group size seems to not directly influence group cohesion, but it seems to indirectly do it through group structure, and only in the case of hypothesis 3. However, this hypothesis needs to be statistically tested a test allowing to describe direct and indirect effects of variables (see. 3.2 in Results).

3.1.4. Comparison of observed group distribution and simulated group distributions.

For the observed data, and for each hypothesis, we calculated the relative distribution of group sizes. The best fit equation for the observed distribution was logarithmic ($N = 6$, $F_{1,4} = 21.29$, $R^2 = 0.852$, $P = 0.008$; fig.3). For hypotheses 1 and 2, there was no relationship between relative frequency and group size ($N = 11$, $F_{1,9} \leq 1.74$, $R^2 \leq 0.16$, $P \geq 0.215$; fig.3). Their distributions significantly differed from the observed distribution ($N_{\text{observed}} = 6$, $N_{\text{simulated}} = 11$, $z = 1.314$, $P \leq 0.021$). For hypothesis 3, whatever the number of groomed partners (5 or 10), the best fit was logarithmic ($N = 11$, $F_{1,9} \geq 720.55$, $R^2 \geq 0.98$, $P < 0.000001$; fig.3). For 5 and 10 partners, the simulated distribution did not significantly differ from the observed distribution ($N_{\text{observed}} = 6$, $N_{\text{simulated}} = 11$, $z = 0.806$, $P = 0.356$ for 5 partners; $N_{\text{observed}} = 6$,

$N_{\text{simulated}} = 11$, $z = 1.134$, $P = 0.086$ for 10 partners; fig.3). However, P - values suggested that there was less difference between the observed distribution and the distribution for 5 partners than for 10 partners. The “5 partners” condition seemed the best model to explain how an individual attributed grooming time to group-mates.

3.2. How does an increase of within-group competition affect group cohesion?

When within-group food competition increased, group members had to spend more time foraging (and therefore to move from one patch to another). As a consequence, grooming time decreased (Berman et al. 2008; Lehmann et al. 2007; Sterck et al. 1997). For hypotheses 1 and 2, a decrease in grooming time did not affect group cohesion, whatever the group size (Mann-Whitney test: $U \geq 68.5$, $P \geq 0.418$; see table 3 for means), except when grooming time is null (Mann-Whitney test: $U < 0.001$, $P < 0.00001$, table 3). For hypothesis 3, a decrease in grooming time (represented by a decrease in the number of partners) from 40% (3 partners) to 100% (0 partner), influenced group cohesion (Mann-Whitney test: $U \leq 46$, $P \leq 0.048$). This effect followed an exponential law ($N = 13$, $F_{1,11} \geq 31$, $R^2 \geq 0.74$, $P \leq 0.0001$). We can therefore conclude that according to group size, group cohesion was affected non linearly by decreased grooming time and a decreased number of partners.

3.3. Causal relations between group size, grooming time, group structure and group cohesion

The most likely causal relations among the different variables previously tested were assessed using path analyses. For hypotheses 1 and 2, the most parsimonious causal model showed that grooming time and group size did not affect group cohesion either directly or indirectly (AIC = 18; $df = 2$; $P = 1$; fig.4a). These results confirmed what we expected below. In the case of hypothesis 3, grooming time and group size affected group cohesion but did so indirectly

through the group structure and especially through the group mean path length ($AIC = 38$; $df = 2$; $P = 0.021$; see fig.4b for details about different influences).

4. Discussion

Time is a finite resource and grooming time therefore has to be limited if an individual also wishes to forage, rest or move (Dunbar 1992b; Lehmann et al. 2007; Majolo et al. 2008; Pollard and Blumstein 2008). Nevertheless, grooming is maybe the most important behavior used by primates for maintaining social relationships (Henzi and Barrett 1999; Schino 2001). As a consequence, the time an individual attributes to grooming will likely influence the quality and/or the quantity of its social bonds (Dunbar 1991, 1992b; Lehmann et al. 2007). If the grooming time of an individual decreases, the quality and/or the quantity of its social bonds should also decrease. Moreover, the structure of social networks – the number of partners per individual and the strength of these connections - may influence group fission probability and therefore group size if the group splits (Koyama 2003; Van Horn et al. 2007). As a consequence, grooming time should logically influence group cohesion and also thus group size. Our study suggests that the relationship between grooming time, group size and group cohesion depends on the way an individual distributes its grooming time to its specific social partners. If an individual grooms all its conspecifics, in an equal way or not, group size does not affect group structure and then group cohesion. If grooming time is however distributed to a specific number of partners, group structure and then group cohesion are affected.

It has previously been shown that the time primates invest in grooming increases with group size (Dunbar 1991). Nevertheless, Lehmann et al. (2007) suggested that when group size increases, each individual should spend more time grooming until a certain group size (about 40 individuals) for which individuals can not devote more time to maintain

relationships with all group members (Dunbar 1992a; Lehmann et al. 2007; Schino et al. 2009). As a consequence, group cohesion decreases and the group splits (Henzi et al. 1997a, b). This hypothesis implies that individuals should be expected to try to maintain social bonds with (i.e. to groom) all, or at least most of, group members. Our results, by contrast, suggest that if this assumption is applied (hypotheses 1 and 2 in our study), group cohesion is not influenced by grooming time, even if time for this activity decreases when group size increases. In the same way, within-group competition is not predicted to affect group cohesion when an individual grooms all its partners, but does have an effect when an individual has a specific number of partners which it grooms. This result may be explained by the fact that, as all individuals were linked to each other in one way or another (whatever the quality/strength of these social bonds), group cohesion was high and therefore the probability of the group splitting was low. This result is not dependent on our model but directly influenced by social network and sociality of individuals. For instance, Amé et al. (2006) showed that when several shelters are proposed, cockroaches (*Blatta germanica*) always aggregate together in only one shelter if this one is able to host all individuals. Even if fission is possible, the group cohesion of individuals does not lead them to split. Sueur et al. (2010) also showed that in macaques, the highest the group cohesion is (more contacts between individuals, less clustered), the lowest the probability to split is. An alternative to grooming all other group members is to groom a specific number of partners (hypothesis 3). Several authors have shown that individuals do not groom all available partners but rather a fixed number of individuals, whatever the group size (Silk et al., 1999; Cords, 2001; Nakamichi and Shizawa, 2002; Berman et al., 2008; Lin et al., 2008; Watts 2000a,v). Under this hypothesis and as found in Majolo et al. (2008), the grooming time an individual devotes to another is not directly dependent on group size. However, the study of Majolo et al. (2008) was the only one to propose this indirect link before our study and this result needs to be checked. Our model

suggests that group size, grooming time and group cohesion are linked when an individual grooms only a specific number of partners. In this scenario, group cohesion decreases when group size increases and/or grooming time decreases. Kudo & Dunbar (2001) also showed that structure of small and large groups differ. Indeed, large groups seem to be more sub-structured. This sub-grouping might be because animals deliberately invest their grooming in core coalition partners (Kudo & Dunbar 2001; Lehmann & Dunbar 2009). Indeed, a theoretical study on social network graphs showed a similar decrease of connectivity according to nodes' number (Wu, 2005). Moreover, we showed that this relationship between group cohesion and group size is logarithmic: when group size exceeded 40 individuals, group cohesion was almost null. This link was however indirect: as (1) group size and grooming time directly influenced group structure, and (2) group structure directly influenced group cohesion (in our study the inverse probability that a group splits in two sub-groups), therefore (3) group size and grooming time influenced group cohesion. To confirm this result and to understand how the group structure evolved according to both group size and the number of groomed partners, it would be interesting to study groups at the same size but with a different number of partners per individual.

Moreover, Lehmann et al. (2007) showed for instance that female dispersion and sex ratio influenced grooming: species with female philopatry spend more time grooming than species with female dispersal. Even if social characteristics such as philopatric sex or the sex ratio, that represent the variability of a social structure, are already included in the different random networks we tested with our model, we did not identify them. However, it would be interesting to assess how these characteristics influence the social network and then the group cohesion.

The distribution of the number of groups staying cohesive according to group size was similar, in our model with a fixed number of five partners, to that based on observed data.

Previous studies have confirmed that, on average, an individual preferentially grooms five partners (Berman et al. 2008; Cords 2001; Lin et al. 2008; Nakamichi and Shizawa 2003; Silk et al. 1999; Watts 2000a,b), giving further support for our hypotheses. Even if an individual grooms five preferred partners, it can still be groomed by more or less individuals than five congeners itself, since grooming is not necessarily reciprocal (Kapsalis and Berman 1994a, b; Nakamichi and Shizawa 2002; Schino 2001; Silk et al. 1999). This pattern has important consequences on group sociality: it allows competition for grooming partners, attraction to high-ranking individuals (Schino 2001) and may allow the emergence of phenomena such as the biological market (i.e., exchanges of commodities according to supply and demand; Fruteau et al. 2009). On the other hand, if an individual grooms all its partners, this kind of competition for partners could not emerge. Kinship may also constrain the relationships of individuals: they will groom their relatives. In these conditions, the two new sub-groups will be more composed on kin related individuals (Chepko-Sade and Sade, 1979; Van Horn et al., 2007). This influence did not affect however our results since groups having an influence of kinship can also be represented using random networks as the ones we used in this study (Flack et al. 2006; MacCowan et al. 2008; Sueur and Petit 2008; Kasper and Voelkl, 2009; Ramos-Fernandez et al. 2009). Grooming a specific and low number of partners could also be of interest when managing time (individuals do not need to change their grooming time when group size increases) but could also have implications when considering cognitive capacities. Several authors (Dunbar 1992a, 1996; Lehmann et al. 2007; Stevens et al. 2005) have suggested that there is a relationship between the cognitive capacities (measured as the neocortex ratio) and the number of relationships an individual can keep track of. Remembering the grooming relationships for five partners would be the easiest solution for an individual, and would be a more parsimonious process than having to remember its ties with all group members. Moreover, grooming five partners rather than all group members may

favor sub-grouping patterns and reduce within-group conflict by regrouping individuals with similar social/physiological affinities (Aureli and Schaffner 2007; Couzin and Laidre, 2009; Ramos-Fernandez et al. 2006).

We believe that a relationship may exist between group size and grooming time but that this relationship would probably be indirect: an increase in group size could result in more food competition between group members (Majolo et al. 2008). As a consequence, an individual would have to increase its foraging time and therefore decrease its grooming time. The consequence is not due to a higher number of partners, but to lower food availability. We did not directly test the relation between food competition and group size in our model, but several studies have already supported this hypothesis (Berman et al. 2008; Dittus 1988; Dunbar 1992). Moreover, group size and grooming time, elements that can be influenced by ecological factors such as food competition or predation (Lehmann et al. 2007; Majolo et al. 2008; Pollard and Blumstein 2008), did not directly influence the probability that a group would split. Social relationships directly influenced group fission probability. And these social relationships are resultants of the combined influence of group size and food competition, but also of other factors as internal or structural constraints (Thierry et al. 2004).

This study was based on grooming interactions but we suggested that similar results may be obtained for any positive interaction such as proximities, contacts or frequency of lips-making for instance. Even if our model did not test all factors having a potential influence on the structure of social relationships, it does show that the key characteristic of group cohesion and stability is group structure (Wey et al. 2007). It is interesting to note that the probability to find large group sizes decreases exponentially with group size in the context of stable groups after irreversible fission but also in fission-fusion populations (Couzin & Laidre 2009). We do not think that a group splits irreversibly in one event as it does in our model. This irreversible fission might be long, from several months to several years (Chepko-Sade

and Sade 1979; Okamoto and Matsumura 2001; Van Horn et al. 2007). We still lack of data about dynamic of fission. It should be interesting to conduct more studies about dynamical processes allowing a group to split. General principles seem to underlie rules of group cohesion at different time scales. All factors, whether social or ecological, seem to influence group cohesion through its structure. In the end, group fission probability may not depend on the sum of individual decision-making based on physiological states and/or on their complex interactions (Schino 2001; Tomasello and Call 1997), but may simply depend on the properties of the social structure, as observed in several self-organized systems (Camazine et al. 2001; Couzin and Krause 2003).

Acknowledgements

We are grateful to J. Munro for language editing and N. Poulain, biostatistician at the DEPE, IPHC, for his help on analyses. This work was supported by Wallonia Brussels International, the Belgian National Funds for Scientific Research, the Franco-American Commission, the Alsace Region and the Fyssen Foundation.

References

- Amé JM, Halloy J, Rivault C, Detrain C, Deneubourg JL (2006) Collegial decision making based on social amplification leads to optimal group formation. *Proc Nat Acad Sci U.S.A.* 103: 5835-5840.
- Aureli F, Schaffner CM (2007) Aggression and conflict management at fusion in spider monkeys. *Biol Lett.* 3: 147-149.
- Baldellou M, Adan A (1997) Time, gender, and seasonality in vervet activity: a chronobiological approach. *Primates.* 38: 31-43.

- 576 Baldellou M, Adan A (1998) Diurnal and seasonal variations in vervet monkeys' activity.
577 Psychol Rep. 83: 675-685.
- 578 Berman CM, Ogawa H, Ionica C, Yin H, Li J (2008) Variation in kin bias over time in a
579 group of Tibetan macaques at Huangshan, China: contest competition, time constraints or risk
580 response? Behaviour. 145: 863-896.
- 581 Bocian CM (1997) Niche separation of black-and-white colobus monkeys (*Colobus*
582 *angolensis* and *C. guereza*) in the Ituri Forest. Ph.D. thesis, City University of New York.
- 583 Boesch C, Boesch-Achermann H (2000) The Chimpanzees of the Taï Forest: Behavioural
584 Ecology and Evolution. Oxford: Oxford University Press.
- 585 Borgatti SP, Everett MG, Freeman LC (2002) UCINET for Windows: Software for Social
586 Network Analysis. Harvard: Analytic technologies.
- 587 Bryson JJ, Yasushi A, Lehmann H (2007) Agent-based modelling as a scientific
588 methodology: a case study analyzing primate social behaviour. Phil Trans R Soc B. 362:
589 1685-1699. doi:10.1098/rstb.2007.2061
- 590 Butynski TM (1990) Comparative ecology of blue monkeys (*Cercopithecus mitis*) in high-
591 and low-density subpopulations. Ecol Monogr. 60: 1-26.
- 592 Buzzard PJ (2004) Interspecific Competition Among *Cercopithecus campbelli*, *C. petaurista*,
593 and *C. diana* at Taï Forest, Cote d'Ivoire. New York: Columbia University Press.
- 594 Chapman CA, Chapman LJ (2000) Constraints on group size in red colobus and red-tailed
595 guenons: examining the generality of the ecological constraints model. Int J Primatol. 21:
596 565-585.
- 597 Chapman CA, Wrangham RW, Chapman LJ (1995) Ecological constraints on group-size: an
598 analysis of spider monkey and chimpanzee subgroups. Behav Ecol Sociobiol. 36 : 59-70.

- 599 Chepko-Sade BD, Sade DS (1979) Patterns of group splitting within matrilineal kinship
600 group: Study of social group structure in *Macaca mulatta* (Cercopithecidae, Primates). Behav
601 Ecol Sociobiol. 5: 67–86.
- 602 Clutton-Brock TH (1974) Activity patterns of red colobus (*Colobus badius tephrosceles*).
603 Folia Primatol. 21: 161-187.
- 604 Clutton-Brock TH (1975) Feedingbehaviour of redcolobusandblack and white colobus in East
605 Africa. Folia Primatol. 23: 165-207.
- 606 Cords M (1986) Interspecific and intraspecific variation in diet of 2 forest guenons,
607 *Cercopithecus ascanius* and *C. mitis*. J Anim Ecol. 55: 811-827.
- 608 Cords M (1995) Predator vigilance costs of allogrooming in wild blue monkeys. Behaviour.
609 132: 559-569.
- 610 Cords M (2001) Friendship among adult female blue monkeys (*Cercopithecus mitis*).
611 Behaviour. 139: 291-314.
- 612 Cords M (2002) Friendship among adult female blue monkeys (*Cercopithecusmitis*).
613 Behaviour. 139: 291-314.
- 614 Couzin ID, Krause J (2003) Self-organization and collective behaviours in vertebrates. Adv
615 Stud Anim Behav. 32: 1–75.
- 616 Couzin ID, Laidre ME (2009) Fission-fusion populations. Curr Biol. 19: R633-R635.
- 617 Dasilva GL (1989) The ecology of thewestern black and white colobus (*Colobus polykomos*
618 *polykomos*) on a riverine island in southeastern Sierra Leone. Ph.D. thesis, University of
619 Oxford.
- 620 Decker BS (1994) Effects ofhabitatdisturbance onthebehavioral ecology and demographics of
621 the Tana river red colobus (*Colobus badius rufomitratu*s). Int J Primatol. 15: 703-737.
- 622 Davies G (1984) An ecological study of the red leaf monkey (*Presbytis rubicund*) in
623 dipterocarp forests of North Borneo. Ph.D. thesis, University of Cambridge.

- 624 Dittus WP (1988) Group fission among wild toque macaques as a consequence of female
625 resource competition and environmental stress. *Anim Behav.* 36: 1626–1645.
- 626 Dunbar RIM (1974) Observations on the ecology and social organization of the green monkey
627 (*Cercopithecus sabaesus*.) in Senegal Primates. 15: 34-350.
- 628 Dunbar, RIM (1991) Functional significance of social grooming in primates. *Folia Primatol.*
629 57: 121-131.
- 630 Dunbar, RIM (1992a) Neocortex size as a constraint on group size in primates. *J Hum Evol.*
631 20: 469-493.
- 632 Dunbar RIM (1992b) Time: a hidden constraint on the behavioural ecology of baboons.
633 *Behav Ecol Sociobiol.* 31: 35-49.
- 634 Dunbar RIM (1996) Determinants of group size in primates: a general model. *Proc Brit Acad.*
635 88: 33-57.
- 636 Dunbar RIM, Dunbar EP (1974) Ecology and population dynamics of *Colobus guereza* in
637 Ethiopia. *Folia Primatol.* 21: 188-208.
- 638 Dussutour A, Deneubourg JL, Fourcassie V (2005) Amplification of preferences in a social
639 context: the case of wall-following in ants. *Proc R Soc B.* 272: 705-71.
- 640 Eley RM, Strum SC, Muchemi G, Reid GDF (1989) Nutrition, body condition, activity
641 patterns, and parasitism of free-ranging troops of Olive Baboons (*Papio Anubis*) in Kenya.
642 *Am J Primatol.* 18: 209-219.
- 643 Ellefson J (1974) A natural history of white-handed gibbons in the Malayan peninsula. In:
644 DM Rumbaugh. Eds., *Gibbon and Siamang*. Basel: Karger.
- 645 Fashing PJ (2001) Activity and ranging patterns of guerezas in the Kakamega Forest:
646 intergroup variation and implications for intragroup feeding competition. *Int J Primatol.* 22:
647 549-577.

- 648 Fawcett KA (2000) Female relationships and food availability in a forest community of
649 chimpanzees. Ph.D. thesis, University of Edinburgh.
- 650 Flack JC, Girvan M, de Waal FBM, Krakauer DC (2006) Policing stabilizes construction of
651 social niches in Primates. *Nature*. 439: 426–429.
- 652 Fossey D, Harcourt AH (1977) Feeding ecology of free-ranging mountain gorilla. *Gorilla*
653 *gorilla beringei*. In: TH Clutton-Brock. Eds., *Primate Ecology: Studies of Feeding and*
654 *Ranging Behaviour in Lemurs, Monkeys and Apes*. pp. 415-44. New York: Academic Press.
- 655 Fruteau C, Voelkl B, van Damme E, Noë R (2009) Supply and demand determine the market
656 value of food providers in wild vervet monkeys. *Proc Nat Acad Sci U.S.A.* 106: 12007-12012.
- 657 Gautrais J, Michelena P, Sibbald A, Bon R, Deneubourg J (2007) Allelomimetic
658 synchronization in Merino sheep . *Animal Behaviour* 74 (5):1443-1454
- 659 Gittins P, Raemakers J (1980) Siamang, lar und agile gibbons. In: D. J. Chivers . Eds.,
660 *Malayan Forest Primates: Ten Years' Study in Tropical Rain Forest*. pp. 63-105. New York:
661 Plenum.
- 662 Gompper ME, Gittleman JL, Wayne RK (1998) Dispersal, philopatry and genetic relatedness
663 in a social carnivore: comparing males and females. *Mol Ecol*. 7: 15163.
- 664 Grimm V, Berger U, Bastiansen F, Eliassen S, Ginot V, Giske J, Goss-Custard J, Grand T,
665 Heinz SK, Huse G, Huth A, Jepsen JU, Jørgensen C, Mooij WM, Müller B, Pe'er G, Piou C,
666 Railsback SF, Robbins AM, Robbins MM, Rossmanith E, Rüger N, Strand E, Souissi S,
667 Stillman RA, Vabø R, Visser U, DeAngelis DL (2006) A standard protocol for describing
668 individual-based and agent-based models . *Ecol Model* 198 (1-2):115-126
- 669
- 670 Henzi SP, Lycett JE, Piper SE (1997a) Fission and troop size in a mountain baboon
671 population. *Anim Behav*. 53: 525-535.

- Henzi SP, Lycett JE, Weingrill T (1997b) Cohort size and the allocation of social effort by female mountain baboons. *Anim Behav.* 54: 1235-1243.
- Henzi SP, Barrett L (1999) The value of grooming to female primates. *Primates.* 40: 47–59.
- Homewood K M (1976) Ecology and behaviour of the tana mangabey, *Cercocebus galeritus galeritus*. Ph.D. thesis, University of London.
- Howarth CJ, Wilson JM, Adamson AP, Wilson ME, Boase MJ (1986) Population ecology of the ringtailed lemur, *Lemur catta*, and the white sifaka, *Propithecus verreauxi*, at Berenty, Madagascar. *Folia Primatol.* 47: 39-48.
- Iwamoto T, Dunbar RIM (1983) Thermoregulation, habitat quality and the behavioural ecology of gelada baboons. *J Anim Ecol.* 52: 357-366.
- Jacobs A, Sueur C, Deneubourg JL & Petit O (In press). Self-organization in primates: new elements in brown lemurs. *Int J Primatol*
- Janson CH, Goldsmith ML (1995). Predicting group size in primates: foraging costs and predation risks. *Behav Ecol Sociobiol.* 36: 326-336.
- Kaplin BA, Moermond TC (2000) Foraging ecology of the mountain monkey. *Cercopithecus l'hoesti*.: implications for its evolutionary history and use of disturbed forest. *Am J Primatol.* 50: 227-246.
- Kasper C, Voelkl B (2009) A social network analysis of primate groups. *Primates.* 50: 343-356.
- Kerth G, Elbert C, Schmidtke C (2006) Group decision-making in fission-fusion societies: evidence from two-field experiments in Bechstein's bats. *Proc Roy Soc B.* 273: 2785-2790.
- Koyama NF (2003) Matrilineal cohesion and Social networks. *Int J Primatol.* 24: 797-811.
- Krause J, Ruxton GD (2002) *Living in Groups*. Oxford: Oxford University Press.
- Krause J, Croft DP, James R (2007) Social network theory in the behavioural sciences: potential applications. *Behav Ecol Sociobiol.* 62: 15-27.

- 697 Kudo H, Dunbar RIM (2001) Neocortex size and social network size in primates . Animal
698 Behaviour 62 (4):711-722
- 699 Lawes MJ (1991) Diet of Samango monkeys. *Cercopithecus mitis erythrarchus*. in the Cape
700 Vidal Dune forest, South Africa. J Zool. 224: 149-173.
- 701 Lee PC (1981) Ecological and social influences on the development of vervet monkeys. Ph.D.
702 thesis, University of Cambridge.
- 703 Lefebvre D, Ménard N, Pierre JS (2003) Modelling the influence of demographic parameters
704 on group structure in social species with dispersal asymmetry and group fission. Behav Ecol
705 Sociobiol. 53: 402-410. \$
- 706 Lehman n J, Korstjens AH, Dunbar RI (2007) Grooming, group size and social cohesion in
707 primates. Anim Behav. 74: 1617-1629.
- 708 Lehmann J, Dunbar RIM (2009) Network cohesion, group size and neocortex size in female-
709 bonded Old World primates . Proc Roy Soc B 276 (1677):4417 -4422
- 710 Li ZY, Rogers E (2004) Habitat quality and activity budgets of white-headed langurs in Fusui,
711 China. Int J Primatol. 25: 41-54.
- 712 Lin TJ, AgoramoorthyG, Huang CC, Hsu MJ (2008) Effects of Troop Size on Social
713 Relations among Male Formosan Macaques, *Macaca cyclopis*. Zoologic stud. 47: 237-246.
- 714 Mackinnon J (1974) Behavior and ecology of wild orangutans (*Pongo pygmaeus*). Anim
715 Behav. 22: 3-74.
- 716 Majolo B, De Bortoli Vizioli A, Shino G (2008) Costs and benefits of group living in
717 primates : group size effects on behaviour and demography. Anim Behav. 76: 1235-1247.
718 doi:10.1016/j.anbehav.2008.06.008
- 719 Maruhashi T (1981) Activity patterns of a troop of Japanese monkeys (*Macaca fuscata yakui*)
720 on Yakushima Island, Japan. Primates. 22: 1-14.

- 721 Matsumoto-Oda A, Oda R (1998) Changes in the activity budget of cycling female
722 chimpanzees. *Am J Primatol.* 46: 157-166.
- 723 McKey D, Waterman PG (1982) Ranging behaviour of a group of black colobus (*Colobus*
724 *satanas*) in the Douala-Edea Reserve, Cameroon. *Folia Primatol.* 39: 264-304.
- 725 McCowan B, Anderson K, Heagarty A, Cameron A (2008) Utility of social network analysis
726 for primate behavioral management and well-being. *Appl Anim Behav Sci.* 109: 396-405
- 727 Mehdiabadi NJ, Kronforst MR, Queller DC, Strassman JE (2009) Phylogeny, reproductive
728 isolation and kin recognition in the social amoeba *dictyostelium purpureum*. *Evol.* 63: 542-
729 548.
- 730 Meunier H, Leca JB, Deneubourg JL, Petit O (2006) Group movement decisions in capuchin
731 monkeys: the utility of an experimental study and a mathematical model to explore the
732 relationship between individual and collective behaviours. *Behaviour.* 143: 1511-1527. doi:
733 10.1163/156853906779366982
- 734 Nagel U (1973) Comparison of anubis baboons, hamadryas baboons and their hybrids at a
735 species border in Ethiopia. *Folia Primatol.* 19: 104-165.
- 736 Nakamachi M, Shizawa Y (2003) Distribution of grooming among adult females in a large,
737 free-ranging group of Japanese macaques. *Int J Primatol.* 24: 607-625.
- 738 Newman M, Barabasi AL, Watts DJ (2006) The structure and dynamics of networks.
739 Princeton: Princeton University Press.
- 740 Nicolis SC, Detrain C, Demolin D, Deneubourg JL (2003) Optimality of collective choices: a
741 stochastic approach. *Bull Math Biol.* 65: 795-808.
- 742 Nishida T (1990) The Chimpanzees of the Mahale Mountains. Tokyo: University of Tokyo
743 Press.

- 744 Oates JF (1977a) The guereza and its food. In: TH Clutton-Brock. Eds., Primate Ecology:
 745 Studies of Feeding and Ranging Behaviour in Lemurs, Monkeys and Apes. pp. 275-321.
 746 London: Academic Press.
- 747 Oates JF (1977b) The social life of a black-and-white colobus monkey, *Colobus guereza*. Zeit
 748 Tierpsychol. 45: 1-60.
- 749 Pollard KA, Blumstein DT (2008) Time allocation and the evolution of group size. Anim
 750 Behav. 76: 1683-1699
- 751 Pollock J (1977) The ecology and sociology of feeding in Indri indri. In: TH Cluttonbrock.
 752 Eds., Primate Ecology. pp. 37-68. London:Academic Press.
- 753 Popa-Lisseanu AG, Bontadina F, Mora O, Ibanez C (2008) Highly structured fission-fusion
 754 societies in an aerial-hawking carnivorous bat. Anim behav. 75: 471-482.
 755 doi:10.1016/j.anbehav.2007.05.011
- 756 Puga-Gonzalez I, Hildenbradt H, Hemelrijk CK (2009) Emergent patterns of social affiliation
 757 in primates, a model. Plos Comput Biol. 5, e10000630.
- 758 Okamoto K, Matsumura S (2001) Group fission in Moor macaques. Int J Primatol. 22: 481-
 759 493.
- 760 Ramos-Fernandez G, Boyer D, Gomez VP (2006) A complex social structure with fission-
 761 fusion properties can emerge from a simple foraging model. Behav Ecol Sociobiol. 60: 536-
 762 549.
- 763 Ramos-Fernandez G, Boyer D, Aureli F, Vick LG (2009) Association networks in spider
 764 monkeys. *Ateles geoffroyi*. Behav Ecol Sociobiol. 63: 999-1013.
- 765 Rangel J, Mattila HR, Seeley TD (2009) No intracolony nepotism during colony fissioning
 766 in honey bees. Proc Roy Soc B. 276: 3895-3900.
- 767 Ron T, Henzi SP, Motro U (1994) A new model of fission in primate troops. Anim Behav. 47:
 768 223-226.

- 769 Schino G (2001) Grooming, competition and social rank among female primates: a meta-
770 analysis. *Anim Behav.* 62, 265-271.
- 771 Schino G, Giuseppe F, Visalberghi E (2009) The Time Frame of Partner Choice in the
772 Grooming Reciprocation of *Cebus apella*. *Ethology.* 115: 70-76.
- 773 Sellers WI, Hill RA, Logan BS (2007) An agent-based model of group decision making in
774 baboons. *Phil Trans R Soc B.* 362: 1699-1710.
- 775 Seppa P, Fernandez-Escudero I, yllenstrand N, Pamilo P (2008) Colony fission affects kinship
776 in a social insect. *Behav Ecol Sociobiol.* 62: 589-597.
- 777 Seth PK, Seth S (1986). Ecology and behaviour of rhesus monkeys in India. In: JG Else, PC
778 Lee. Eds., *Primate Ecology and Conservation.* pp. 89-103. Cambridge: Cambridge University
779 Press.
- 780 Seyfarth RM (1977) A model of social grooming among adult female monkeys. *J Theo Biol.*
781 65: 671-698.
- 782 Silk JB, Sefarth RM, Cheney DL (1999) The structure of social relationships among female
783 savanna baboons in Moremi reserve, Botswana. *Behaviour.* 136: 679-703.
- 784 Son VD (2004) Time budgets of *Macaca fascicularis* in a Mangrove Forest, Vietnam. *Lab*
785 *Primate Newslett.* 43: 1-4.
- 786 Stanford CB (1998) Chimpanzee and Red Colobus: the Ecology of Predator and Prey.
787 Cambridge: Harvard University Press.
- 788 Starin ED (1991) Socioecology of the red colobus monkey in the Gambia with particular
789 reference to femaleemale differences and transfer patterns. Ph.D. thesis, City University of
790 New York, 406 pp.
- 791 Sterck EH., Watts DP, van Schaik CP (1997) The evolution of female social relationships in
792 nonhuman primates. *BehavEcol Sociobiol.* 41: 291-309.

- 793 Stevens JR, Cushman FA, Hauser MD (2005) Evolving the psychological mechanisms for
794 cooperation. *Ann Rev Ecol Evol Syst* .36: 499-518.
- 795 Struhsaker TT (1979) Socioecology of five sympatric monkey species in the Kibale forest,
796 Uganda. *Adv Stud Behav*. 9: 159-228.
- 797 Struhsaker TT (1980) Comparison of the behaviour and ecology of red colobus and redtail
798 monkeys in the Kibale Forest, Uganda. *Af J Ecol*. 18: 33-51.
- 799 Struhsaker TT, Leland L (1979) Socioecology of five sympatric monkey species in the Kibale
800 Forest, Uganda. *Adv Stud Behav*. 9: 159-228.
- 801 Sueur C, Petit O (2008) Organization of group members at departure of joint movements is
802 driven by social structure in macaques. *Int J Primatol*. 20: 1085-1098.
- 803 Sueur C, Petit O, Deneubourg J (2009) Selective mimetism at departure in collective
804 movements of *Macaca tonkeana*: an experimental and theoretical approach. *Anim*
805 *Behav* 78 (5):1087-1095
- 806 Sueur C, Petit O, Deneubourg J (2010) Short-term group fission processes in macaques: a
807 social networking approach . *J Exp Biol* 213 (8):1338-1346
- 808 Sugiyama Y (1976) Characteristics of the ecology of the Himalayan langurs. *J Hum Evol*. 5:
809 249-277.
- 810 Sussmann RW (1977) Feeding behaviour of *Lemur catta* and *Lemur fulvus*. In: TH
811 Cluttonbrock. Eds. ,*Primate Ecology*. pp. 1-39. London: Academic Press.
- 812 Teas J, Richie T, Taylor H, Southwick CH (1980) Population patterns and behavioural
813 ecology of rhesus monkeys. *Macaca mulatta*. in Nepal In: DG Lindburg. Eds. *The Macaques*.
814 pp. 247-262. New York: Van Nostrand Reinhold.
- 815 Tomasello M, Call J (1997) *Primate Cognition*. New York: Oxford University Press.
- 816 Tutin CEG, McGrew WC, Baldwin PJ (1983) Social organization of savanna-dwelling
817 chimpanzees, *Pan troglodytes verus*, at Mt. Assirik, Senegal *Primates* 24: 154-173.

- 818 Van Horn RC, Buchan JC, Altmann J, Alberts SC (2007) Divided destinies: group choice by
819 female savannah baboons during social group fission. *Behav Ecol Sociobiol.* 61: 1823-1837.
- 820 Van Noordwijk M A (1985) The Socioecology of Sumatran Longtailed Macaques. *Macaca*
821 *fascicularis*. II. The Behaviour of Individuals. Utrecht: University of Utrecht.
- 822 Wasserman S, Faust K (1994) Social network analysis. Cambridge: Cambridge University
823 Press.
- 824 Watts DP (2000a) Grooming between male chimpanzees at Ngogo, Kibale National Park. I.
825 Partner number and diversity and grooming reciprocity. *Int J Primatol* 21:189-210.
- 826 Watts DP (2000b) Grooming between male chimpanzees at Ngogo, Kibale National Park. II.
827 Influence of male rank on possible competition for partners *Int J Primatol* 21:211–238.
- 828 Wey T, Blumstein DT, Shen W, Jordan F (2007) Social network analysis of animal
829 behaviour: a promising tool for the study of sociality. *Anim Behav.* 75: 333-344.
- 830 White FJ (1992) Activity budgets, feeding behavior, and habitat use of pygmy chimpanzees at
831 Lomako, Zaire. *Am J Primatol.* 26: 215-223.
- 832 White FJ, Chapman CA (1994) Contrasting chimpanzees and bonobos: nearest neighbor
833 distances and choices. *Folia Primatol.* 63 181-191.
- 834 Whitehead H (2009) SOCPROG programs: analysing animal social structures. *Behav Ecol*
835 *Sociobiol.* 63: 765-778.
- 836 Whiten A (1980) The Kloss gibbon in Siberut rain forest. Ph.D. thesis, University of
837 Cambridge.
- 838 Whitesides GH (1989) Interspecific associations of Diana monkeys, *Cercopithecus diana*, in
839 Sierra Leone, West Africa: biological significance or chance? *Anim Behav.* 37: 760-776.
- 840 Wilensky U (1999) NetLogo. <http://ccl.northwestern.edu/netlogo>. Center for Connected
841 Learning and Computer-Based Modeling. Northwestern University, Evanston, IL.

Wittemyer G, Douglas-Hamilton I, Getz WM (2005) The socioecology of elephants: analysis of the processes creating multitiered social structures. *Anim Behav.* 69: 1357-1371.

Wu WC (2005) Algebraic connectivity of directed graphs. *Linear and multilinear algebra.* 53: 203-223.

Yamakoshi G (1998) Dietary responses to fruit scarcity of wild chimpanzees in Bossou, Guinea: possible implications for ecological importance of tool use. *Am J Physical Anthropol.* 106: 283-295.

Yamakoshi G (2004) Food seasonality and socioecology in Pan: are West African chimpanzees another bonobo? *Af Stud Monogr.* 25: 45-60.

Table legends

Table 1: Data (Genus, species, group size and grooming time) and references of data used in this study.

Genus	Species	Group size	Grooming time (%)	Reference(s)
<i>Avahi</i>	<i>laniger</i>	2	2	C. Harcourt (from Dunbar 1991)
<i>Cercocebus</i>	<i>galeritus</i>	27	5.5	Homewood 1976
<i>Cercopithecus</i>	<i>ascanius</i>	26.75	3.45	Struhsaker 1980; Cords 1986
<i>Cercopithecus</i>	<i>campbelli</i>	9	2.8	Buzzard 2004
<i>Cercopithecus</i>	<i>diana</i>	28.75	2.48	Whitesides 1989; Buzzard 2004
<i>Cercopithecus</i>	<i>mitis</i>	22.65	7.18	Struhsaker and Leland 1979
				Butynski 1990; Lawes 1991
				Cords 1995, 2002
				Kaplin and Moermond 2000

<i>Chlorocebus</i>	<i>aethiops</i>	19.7	9.17	Dunbar 1974; Lee 1981 Baldellou and Adan 1997, 1998
<i>Colobus</i>	<i>angolensis</i>	18	5.25	Bocian 1997
<i>Colobus</i>	<i>guereza</i>	9.04	5.52	Dunbar and Dunbar 1974 Oates 1977a, b; Bocian 1997 Fashing 2001
<i>Colobus</i>	<i>polykomos</i>	12.5	3.49	Dasilva 1989
<i>Colobus</i>	<i>satanas</i>	12	5.51	McKey and Waterman 1982
<i>Gorilla</i>	<i>gorilla</i>	11	0.09	D. Doran (from Lehman et al. 2007)
<i>Gorilla</i>	<i>G. beringei</i>	6	1	Fossey and Harcourt 1977
<i>Hylobates</i>	<i>agilis</i>	4.4	0	Gittins and Raemakers 1980
<i>Hylobates</i>	<i>klossii</i>	3.8	0	Whiten 1980
<i>Hylobates</i>	<i>lar</i>	3.4	2.1	Ellefson 1974 Gittins and Raemakers 1980
<i>Indri</i>	<i>indri</i>	4.3	1	Pollock 1977
<i>Lemur</i>	<i>catta</i>	12.2	7.18	Sussmann 1977
<i>Lemur</i>	<i>fulvus</i>	15.33	7.98	Sussmann 1977
<i>Lophocebus</i>	<i>albigena</i>	15	5.8	Struhsaker 1979
<i>Macaca</i>	<i>fascicularis</i>	82.45	7.98	Van Noordwijk 1985; Son 2004
<i>Macaca</i>	<i>fuscata</i>	36.5	10.7	Maruhashi 1981 Seth and Seth 1986
<i>Macaca</i>	<i>mulatta</i>	32	15	Teas et al. 1980
<i>Pan</i>	<i>paniscus</i>	27.8	5.7	White 1992
<i>Pan</i>	<i>t. Schweinfurthi</i>	59.2	11.67	Whrangham 1977; Nishida 1990

				White and Chapman 1994
				Matsumoto-Oda and Oda 1998
				Fawcett 2000
<i>Pan</i>	<i>t. verus</i>	40.33	8.27	Tutin et al. 1983
				Yamakoshi 1998, 2004
				Boesch and Boesch-Achermann 2000
<i>Papio</i>	<i>anubis</i>	58.8	8.3	Nagel 1973; Eley et al. 1989
<i>Papio</i>	<i>ursinus</i>	28.07	12.64	Henzi et al. 1997b
<i>Papio</i>	<i>hamadryas</i>	51	13.5	Nagel 1973
				R. Noe and H. Korstjens (from Lehman et
<i>Piliocolobus</i>	<i>badius</i>	42.5	4.5	al. 2007)
<i>Piliocolobus</i>	<i>rufomitrat</i>	16.16	0.83	Decker 1994
<i>Piliocolobus</i>	<i>temminckii</i>	26.2	5.4	Starin 1991
<i>Piliocolobus</i>	<i>tephrosceles</i>	51.67	4.99	Clutton-brock 1974, 1975; Stanford 1998
				Struhsaker and Leland 1979
				Chapman and Chapman 2000
<i>Pongo</i>	<i>pygmaeus</i>	1	0	Mackinnon 1974
<i>Presbytis</i>	<i>entellus</i>	33	4.4	Sugiyama 1976
<i>Presbytis</i>	<i>rubicunda</i>	7	0	Davies 1984
				R. Noe and H. Korstjens (from Lehman et
<i>Procolobus</i>	<i>verus</i>	3	3.58	al. 2007)
<i>Propithecus</i>	<i>verreauxi</i>	5.1	4.7	Howarth et al. 1986
<i>Trachypithecus</i>	<i>leucocephalus</i>	10	11.71	Li and Rogers 2004
<i>Theropithecus</i>	<i>gelada</i>	144.7	17.4	Iwamoto and Dunbar 1983

Table 2: equations of relationship between mean path length and group cohesion and between density and group cohesion, for 5 and 10 partners. Tests showed that the equations are similar for 5 and 10 partners. We used a comparison test for two linear regressions (transformation from logarithmic to linear for the relations between density and number of partners). We first used a Snedecor test to compare variances of each distribution ($df_1 = df_2 = 11$, $f \leq 0.75$). Then, we tested if variable ($b1_{5\text{partners}} - b1_{10\text{partners}}$, $df = 22$, $T \leq 0.27$) and variable ($\text{constant}_{5\text{partners}} - \text{constant}_{10\text{partners}}$, $df = 22$, $T \leq 0.21$) followed a Student law ($df = 22$).

Group cohesion							
N = 13	5 partners			10 partners			similar functions
	function	constant	b1	function	constant	b1	
Group mean path length	linear	1.51	- 0.45	linear	1.48	- 0.42	Yes P < 0.05
Density	logarithmic	1.03	0.23	logarithmic	1.05	0.26	Yes P < 0.05

Table 3: Influence of grooming time on mean group cohesion for each hypothesis

grooming time (% per day time)	initial time decrease (%)	mean group cohesion		
		hypothesis 1	hypothesis 2	hypothesis 3
15 (initial time)	-0	1.00	0.93	0.44
12	-20	1.00	0.93	0.33
8	-40	1.00	0.92	0.27
6	-60	1.00	0.93	0.22
3	-80	1.00	0.92	0.17
0	-100	0.00	0.00	0.00

Figure captions

Figure 1: Illustration of a social network with 40 individuals (squares). Number labels indicated the number of group members that groomed a given individual (*indegree*, Faust and Wasserman 1994). We showed from this graph that even if the number of partners groomed by the same individual was fixed at 5, an individual can be groomed by more (or less) than 5 grooming partners in the model. We built this network via Netdraw in UCINET 6.0 (Borgatti et al. 2002). Distance between individuals represents the strength of associations, and was calculated using multidimensional scaling (Whitehead 2009, Sueur and Petit 2008).

Figure 2: Influence of group size on group structure (a. on group mean path length and b. on density) and on group cohesion (c.) for each hypothesis. For hypotheses 1 and 2, where an individual groomed all other group members, there is no relationship between group size and group structure, nor between group size and group cohesion, contrary to hypothesis 3 (where an individual groomed a fixed number of partners).

Figure 3: Relative distribution of group size for observed data and under each hypothesis. The curve line is the best-fit model explaining the distribution of observed data and corresponds to an exponential law.

Figure 4: Path diagram indicating causal relations between group size, grooming time, group structure, i.e. mean strength of social relationships (a.), number of partners (b.), density and group mean path length, and group cohesion. Rectangles indicated observed variables. Arrows indicated presumed causal relations: solid arrows represented the best model based on

890 AIC value, using AMOS5; dotted arrows represented the relations we included in the analysis
891 but not selected in the best model.

Accepted manuscript

a. Hypotheses 1 and 2

a. Hypothesis 3