In reply

Michele Basso, Anna Modoni, Mauro Lo Monaco, Carlo Barone

► To cite this version:

Michele Basso, Anna Modoni, Mauro Lo Monaco, Carlo Barone. In reply. *Cancer Chemotherapy and Pharmacology*, 2011, 67 (5), pp.1191-1192. 10.1007/s00280-011-1578-z . hal-00670755

HAL Id: hal-00670755

<https://hal.science/hal-00670755>

Submitted on 16 Feb 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

In reply

Dear Editor,

We are sincerely grateful to Dr Park and colleagues for their comment on our article concerning the possible relationship between *acute* oxaliplatin-induced neurotoxicity and SK3 polymorphisms. Any remark on this topic may contribute to improve the knowledge on this complex and serious event.

In our study we found a significant association between the occurrence of *acute* oxaliplatin neurotoxicity and the presence of a shorter CAG repeat polymorphism of SK3 gene [1]. We did not conclude on the molecular mechanism relating the two events, but only speculated on a possible interference of oxaliplatin with K⁺ channels tetramerization. This fascinating speculation seems supported by a report suggesting the crucial role of the polyglutamine tract of SK3 channel in modulating both tetramer formation and channel function [2]. Taken together, the last observation and our results contribute in generating the hypothesis that we have suggested, which needs validation in a proper experimental setting, as we have clearly specified in our paper.

We agree with Dr Park and colleagues on the existence of several *in vitro* and *in vivo* studies suggesting an impairment of voltage-gated Na⁺ channels (VGNaCs) in the pathogenesis of oxaliplatin-induced neurotoxicity [3-6]. As we have also discussed in our paper, these studies showed that oxaliplatin prolongs the duration of nerve relative refractory period, assuming that this effect is a clear-cut measure of the direct inactivation kinetic of voltage-gated transient Na⁺ channels [3]. To the nerve refractoriness, however, might contribute some other factors, such as acid-base balance and Ca²⁺ ion concentration. Interestingly, *in vitro* studies, have also demonstrated that the effect of oxaliplatin on inward sodium current was mimicked by strong calcium ions chelators, suggesting an indirect effect of oxaliplatin on VGNaCs [4]. The same authors also observed that, when neuronal cells were exposed *in vitro* to oxaliplatin for prolonged interval of times, an important reduction both of depolarizing and post-hyperpolarizing phases occurred,

resulting in an increase in the duration of action potential duration [4]. This last phenomenon seems to be related to a dysfunction of SK3 channels, which play a critical role in modulating the after-hyperpolarization [7].

Accordingly, recent immunohistochemical studies have localized SK3 to presynaptic nerve terminals, where they control the after-hyperpolarization [8]. As observed in neuromyotonia, an autoimmune disease caused by autoantibodies directed against VGKC, also oxaliplatin-*acute* neurotoxicity is clinically characterized by motor nerve hyperexcitability, with a prolonged axonal after-activity following a single stimulus [9]. This means that a single nerve stimulation generates a CMAP closely followed by a high number of extra-discharges producing repeated oscillations arising in the distal portion of the peripheral nerve [1, 10-11].

The EMG counterpart of this event consists in fibrillation potentials and repetitive bursts of spontaneous activity. Interestingly, SK3 channels are also expressed in adult denervated skeletal muscles where they contribute to hyperexcitability [12]. As mentioned by Dr Park, denervation-induced muscle hyperexcitability requires SK3 expression together with modifications in the function of other membrane ion channels [13]. It should be underscored, however, that nerve and muscle hyperexcitability are distinct neurophysiologic events, in which SK3 channels might play a different role. Thus, it remains plausible that oxaliplatin might have a direct effect on SK3 channels of the pre-synaptic nerve terminal, while causing, as direct or indirect effect, also a VGNaC dysfunction.

Moreover, Dr Park and colleagues, have recently showed that, following oxaliplatin administration, sensory axons developed a significant decrease of refractoriness, suggesting a functional channelopathy of axonal Na⁺ channels, while motor axonal excitability remained unchanged [5]. The absence of motor nerve excitability is in contrast with several studies on *acute* oxaliplatin neurotoxicity, while the abnormalities in sensory nerves are frequent in *chronic* oxaliplatin-induced neurotoxicity. Thus, it cannot be excluded that changes in sensory nerve refractoriness

demonstrated by Park and colleagues might be considered an early marker of *chronic* oxaliplatin neurotoxicity rather than the expression of *acute* neurotoxicity.

Taken together, these observations suggest that *acute* oxaliplatin-related neurotoxicity has a pathophysiological mechanism that is quite different from that of *chronic* neurotoxicity. For this reason we totally agree with Dr Park that the evaluation of SK3 polymorphisms also in patients affected by chronic oxaliplatin-induced neurotoxicity might contribute to clarify this complex matter.

Michele Basso¹

Anna Modoni²

Mauro Lo Monaco²

Carlo Barone¹

¹ Medical Oncology, Catholic University of Sacred Heart, Rome, Italy

² Department of Neuroscience, Catholic University of Sacred Heart, Rome, Italy

References

1. Basso M, Modoni A, Spada D, Cassano A, Schinzari G, Lo Monaco M, Quaranta D, Tonali PA, Barone C (2010). Polimorphism of CAG motif of SK3 gene is associated with acute oxaliplatin neurotoxicity. *Cancer Chemother Pharmacol* doi: 1007/s00280-010-1466-y.
2. Frei E, Spindler I, Grissmer S, Jager H (2006). Interaction of N-terminal and C-terminal parts of small conductance Ca^{2+} activated k^{+} channels, hSK3. *Cell Physiol Biochem* 18: 165-176.
3. Adelsberger H, Quasthoff S, Grosskreutz J, Lepier A, Eckel F, Lersch C (2000). The chemotherapeutic oxaliplatin alters voltage-gated $\text{Na}(+)$ channel kinetics on rat sensory neurons. *Eur J Pharmacol* 406: 25-32.

4. Grolleau F, Gamelin L, Boisdron-Celle M, Lapied B, Phelate M, Gamelin E (2001). A possible explanation for a neurotoxic effect of the anticancer agent oxaliplatin on neuronal voltage-gated sodium channels. *J Neurophysiol* 85: 2293-2297.
5. Park SB, Lin CSY, Krishnan AV, Goldstein D, Friedlander ML, Kiernan MC (2009). Oxaliplatin-induced neurotoxicity: changes in axonal excitability precede development of neuropathy. *Brain* 132: 2712-2723.
6. Krishnan AV, Goldstein D, Friedlander M, Kiernan MC (2005). Oxaliplatin-induced neurotoxicity and development of neuropathy. *Muscle Nerve* 32: 51-60.
7. McIntyre CC, Richardson AG and Grill WM (2002). Modeling the excitability of mammalian nerve fibers: influence of afterpotentials on recovery cycle. *J Neurophysiol* 87: 995-1006
8. Obermair GJ, Kaufmann WA, Knaus HG, Flucher BE (2003). The small conductance Ca^{2+} -activated K^{+} channel SK3 is localized in nerve terminals of excitatory synapses of cultured mouse hippocampal neurons. *Eur J Neurosci* 17: 721-731.
9. Hart IK, Waters C, Vincent A, Newland C, Beeson D, Pongs O, Morris C, Newsom-Davis J (1997). Autoantibodies detected to expressed K^{+} channels are implicated in neuromyotonia. *Ann Neurol* 41: 238-246.
10. Wilson RH, Lehky T, Rebecca RT, Quinn MG, Floeter MK, Grem JL (2002). Acute oxaliplatin-induced peripheral nerve hyperexcitability. *J Clin Oncol* 20: 1767-1774.
11. Lehky TJ, Leonard GD, Wilson RH, Grem JL and Floeter MK (2004). Oxaliplatin-induced neurotoxicity: acute hyperexcitability and chronic neuropathy. *Muscle Nerve* 29: 387-392.
12. Favero M, Jiang D-J, Chiamulera C, Cangiano A, Fumagalli GF (2008). Expression of small conductance calcium-activated potassium channels (SK3) in skeletal muscle: regulation by muscle activity. *J Physiol* 19: 4763-4774.
13. Jacobson D, Herson PS, Neelands TR, Maylie J, Adelman JP (2002). SK channels are necessary but not sufficient for denervation-induced hyperexcitability. *Muscle Nerve* 26: 817-822.