Influence of whole wheat inclusion and a blend of essential oils on the performance, nutrient utilisation, digestive tract development and ileal microbiota profile of broiler chickens

Journal:	<i>British Poultry Science</i>
Manuscript ID:	CBPS-2010-077.R1
Manuscript Type:	Original Manuscript
Date Submitted by the Author:	05-Aug-2010
Complete List of Authors:	Amerah, Ahmed; Danisco Animal Nutrition Péron, Alexandre; Danisco Animal Nutrition Zaefarian, Fifi; Massey University, Institute of Food, Nutrition and Human Health Ravindran, Velmurugu; Massey University, Institute of Food, Nutrition and Human Health
Keywords:	Broilers, Nutrition, Plants

SCHOLARONE™
Manuscripts

Influence of whole wheat inclusion and a blend of essential oils on the performance, nutrient utilisation, digestive tract development and ileal microbiota profile of broiler chickens

A.M. AMERAH, A. PÉRON, F. ZAEFARIAN¹ AND V. RAVINDRAN¹

Danisco Animal Nutrition, Marlborough, SN8 1XN, UK and ¹Institute of Food, Nutrition and Human Health, Massey University, New Zealand.

Running title: Whole wheat and essential oils

Correspondence to: Ahmed Amerah, Danisco Animal Nutrition, Marlborough, SN8 1XN, UK
E-mail: Ahmed.amerah@danisco.com

Accepted for publication 12th August 2010

Abstract 1. The aim of the present experiment was to examine the influence of whole wheat inclusion and a blend of essential oils (EO; cinnamaldehyde and thymol) supplementation on the performance, nutrient utilisation, digestive tract development and ileal microbiota profile of broiler chickens.

2. The experimental design was a 2 x 2 factorial arrangement of treatments evaluating two wheat forms (ground wheat [GW] and whole wheat [WW]; 100 and 200 g/kg WW replacing GW during starter [1 to 21 d] and finisher [22 to 35 d] diets respectively) and two levels of EO inclusion (0 or 100 g/tonne diet). All dietary treatments were supplemented with 2000 xylanase units/kg feed. Broiler starter and finisher diets based on wheat and soybean meal were formulated and each diet fed *ad libitum* to 6 pens of 8 male broilers.

3. During the trial period (1-35 d), wheat form had no significant effect on weight gain or feed intake. However, WW inclusion tended to improve feed per gain. Essential oil supplementation significantly improved weight gain in both diets, but the improvements were greater in the GW diet as indicated by a significant wheat form x EO interaction.

4. Main effects of wheat form and EO on the relative weight, length and digesta content of various segments of the digestive tract were not significant. Significant interactions, however, were found for relative gizzard and caecal weights. Essential oil supplementation significantly increased the relative gizzard weight and lowered relative caecal weight in birds fed on the GW based diet, but had no effect in those fed on the WW based diet.

5. Whole wheat inclusion and EO supplementation significantly improved apparent ileal nitrogen digestibility. Apparent ileal digestible energy was not significantly influenced by the dietary treatments.

6. Ileal microbiota profiling, using denaturing gradient gel electrophoresis, showed that the ileal microbiota composition was influenced by feed form. The mean numbers of bacterial

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

species in the ileal contents of birds fed on the GW diet supplemented with EO tended to be higher than those of the ileal contents of birds fed on unsupplemented GW based diet.

7. The present data suggested that dietary addition of EO improves broiler weight gain and ileal nitrogen digestibility both in GW and WW based diets, but that the magnitude of the response to EO for weight gain was greater in GW based diet. Whole wheat feeding was found to be beneficial in terms of feed efficiency.

INTRODUCTION

The need to reduce feed costs has renewed the interest in the poultry industry in whole grain feeding, a common practice in the early days of commercial poultry production. There appears to be a general consensus that the inclusion of whole grain in broiler diets, using mixed feeding technique, will have no adverse effect on broiler performance (Preston *et al.*, 2000; Plavnik *et al.*, 2002; Amerah and Ravindran, 2008). It has been suggested that the beneficial effects of whole grain feeding on nutrient utilisation may be related to the positive effect of whole grain on gizzard development (Ferket, 2000; Svihus and Hetland, 2001). A more developed gizzard is associated with increased grinding activity, resulting in increased gut motility and a greater digestion of nutrients (Amerah *et al.*, 2007a). Higher gizzard functionality may also play a positive role in the control of bacterial populations in the gut and on gut health (Bjerrum *et al.*, 2005).

Essential oils (EO) are steam-volatile or organic-solvent extracts of plants, usually used for the pleasant odour of their essence, their flavour, their antiseptic or preservative properties, as well as their antimicrobial effect (Burt, 2004; Lee *et al.*, 2004; Wallace, 2004). The antimicrobial properties of EO have encouraged their use as a natural replacement for antibiotic growth promoters in animal feeds. In addition to the positive effects of EO against the colonisation and proliferation of pathogenic bacteria (Lee *et al.*, 2004), EO have been

shown to improve nutrient digestibility and broiler performance (Hernandez *et al.*, 2004; Lee *et al.*, 2004; Cao *et al.*, 2009).

The ban on the use of antibiotic growth promoters in feeds has encouraged nutritionists to explore alternative feed management strategies to improve the intestinal health and digestive efficiency of broiler chickens (Amerah *et al.*, 2009a). Promotion of gizzard development by whole wheat (WW) feeding is one of the strategies that have been found to prevent potentially pathogenic bacteria from entering the intestinal tract (Engberg *et al.*, 2004). Recent studies have highlighted the potential benefit of combining EO and carbohydrase enzymes (Cao *et al.*, 2009; Péron *et al.*, 2009). However, combining the effects of whole wheat inclusion on gut health and the anti-microbial properties of EO, with possible complementary improvement in broiler performance, has not yet been investigated. Therefore, the present study investigated the effects of WW inclusion and a specific blend of EO supplementation on the performance, nutrient utilisation, digestive tract parameters and ileal microbiota profile of broilers.

MATERIALS AND METHODS

Birds and housing

Experimental procedures were conducted in accordance with the Massey University Animal Ethics Committee guidelines. Day-old male broilers (Ross 308) were obtained from a commercial hatchery, individually weighed and assigned to 24 cages (8 birds per cage) in electrically heated battery brooders so that the average bird weight was similar for each cage. The birds were transferred to grower cages on d 12. The battery brooders and grower cages were housed in an environmentally controlled room with 20 h of fluorescent illumination daily. The temperature was maintained at 31°C on d 1 and gradually reduced to 22°C by 21 d of age.

Diets and conduct of the trial

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

The basal diets were based on wheat and soybean meal (Table 1). Two sets of basal diets were used: one for broiler starters (1–21 d) and the other for broiler finishers (22–35 d). The basal diets based either on ground wheat (GW) or WW (100 and 200 g/kg WW replacing GW during starter (1 to 21 d) and finisher (22 to 35 d) diets respectively). Diets were formulated to meet or exceed the Ross 308 strain recommendations for major nutrients for broilers. The basal diets were supplemented with a commercial xylanase (Danisco Xylanase, Danisco Animal Nutrition, Marlborough, UK), which provided 2000 xylanase units/kg feed. The enzyme preparation is based on endo 1,4 beta-xylanase because its main activity is produced by *Trichoderma reesei*. WW was ground in a hammer mill (Bisley’s Farm Machinery, Auckland, New Zealand) to pass through a screen size of 3 mm. Half of each formulated diet was supplemented with an encapsulated blend of EO (100 g/tonne; Enviva™ EO 101 G, Danisco Animal Nutrition, Marlborough, UK) and the other half was not supplemented with EO. The two active ingredients in the EO product were Cinnamaldehyde and Thymol. Titanium oxide (0.3%) was included in all diets as a dietary marker.

Table 1 near here

Each of the 4 dietary treatments was fed *ad libitum* to 6 replicate cages. Water was freely available throughout the trial. The diets were cold-pelleted (65–70°C). Body weight and feed intake were recorded at weekly intervals, and mortality was recorded daily, for each cage. Any bird that died was weighed and feed per gain values were calculated by dividing total feed intake by weight gain of live plus dead birds.

Digestive tract measurements

On d 35, two birds from each replicate cage were randomly selected, weighed and killed by cervical dislocation. The body cavity was immediately opened and the full and empty weights of each digestive tract segment from proventriculus to caeca were recorded. The length of each intestinal segment was determined with a flexible tape on a glass surface to prevent inadvertent stretching. The length (± 0.1 mm) of the duodenum (from the pyloric junction to

the distal most point of insertion of the duodenal mesentery), the length of the jejunum (from the distal most point of insertion of the duodenal mesentery to the junction with Meckel's diverticulum), the length of the ileum (from the junction with Meckel's diverticulum to ileo-caecal junction) and caeca (the sum of the lengths from ostium to tip of each caeca) were also determined. Following division and freeing of each of these components from any adherent mesentery, their full and empty weights (± 0.1 g) were determined along with those of the proventriculus and gizzard. Digesta were removed from the gut component by gentle squeezing for empty weight determination.

Ileal digesta collection

On d 35, 4 more birds per replicate were killed by intravenous injection with sodium pentobarbitone. Immediately after euthanasia, the ileum was excised and divided into two parts, the anterior and posterior ileum. The ileum was defined as the portion of the small intestine extending from Meckel's diverticulum to a point 40 mm proximal to the ileo-caecal junction. Contents of the posterior ileum were collected by gently flushing with distilled water into plastic containers. Digesta were pooled within a cage, freeze-dried, ground to pass through a 0.5 mm screen size and stored at -4°C in airtight containers until laboratory analysis.

Isolation of DNA and PCR-denaturing gradient gel electrophoresis (DGGE) analysis

On d 35, two birds from each replicate cage were killed by intravenous injection of sodium pentobarbitone. Approximately 1ml digesta was obtained from the middle part of the ileum and stored at -80°C until DNA extraction. Isolation of DNA and PCR-DGGE analysis were performed as described by Amerah *et al.* (2009a).

Diversity computed as the Simpsons diversity index, using the area under the densiometric curve of each band (projected on a Gaussian basis), represented population density of a particular species (band). Comparison of samples was based on the Dice

similarity coefficient and the un-weighted pair group method using arithmetic means for clustering (Hume *et al.*, 2003). The relative similarities of the groups were determined by comparing the patterns of a relatedness tree (dendogram). The degree of similarity was indicated by the percentage coefficient bar located below each dendogram. Values between 0 and 1 represented an arithmetic determination of the degree to which banding patterns were similar.

Chemical analysis

The dry matter (DM), nitrogen and gross energy (GE) were determined in samples of diets and excreta. Dry matter content was determined using standard procedures (AOAC, 2005, method 930.15). Nitrogen was determined by combustion (method 968.06; AOAC, 2005) using a CNS-200 carbon, nitrogen, and sulphur auto-analyser (LECO Corporation, St. Joseph, MI). Gross energy was determined using an adiabatic bomb calorimetry (Gallenkamp Autobomb, London, UK) standardised with benzoic acid. Diets and ileal digesta samples were assayed for titanium on a UV spectrophotometer following the method described by Short *et al.* (1996).

Calculations

Apparent ileal digestibility coefficients of nitrogen and energy were calculated using the following formula:

$$\text{Apparent ileal digestibility} = \frac{(\text{nutrient/Ti})_d - (\text{nutrient/Ti})_i}{(\text{nutrient/Ti})_d}$$

where (nutrient/Ti)_d = ratio of nutrient to titanium in the diet, and
(nutrient/Ti)_i = ratio of nutrient to titanium in the ileal digesta.

180 Statistical analysis

181 For performance parameters, cage means served as the experimental unit for statistical
182 analysis. For digestive tract measurements, individual birds were considered as the
183 experimental unit. All data were subjected to two-way ANOVA using the general linear
184 models procedure of SAS (SAS, 1997). For the microbial profile analysis, both band number
185 and Shannon Weiner index of diversity were normally distributed; thus the data were
186 analysed by three-way ANOVA for the effects of wheat form, EO inclusion and gel. The
187 latter term was introduced as, in DGGE, results vary somewhat between gels due to slight
188 variation in the denaturing gradient from gel to gel. The 12 samples from each dietary
189 treatment were divided randomly in a balanced manner among the three gels, having 4
190 random samples from each dietary treatment in a gel. The differences were considered
191 significant at $P < 0.05$ and significant differences between means were separated by the least
192 significant difference test.

193 RESULTS

194 Bird performance

195 The influence of treatments on the performance of broilers is summarised in Table 2. During
196 the starter phase (1-21 d), a significant interaction between wheat form and EO
197 supplementation was observed for weight gain. Essential oils improved the weight gain in
198 birds fed on GW based diet but had no effect in birds fed on the WW based diet. WW feeding
199 had no significant effect on feed intake, but significantly improved feed per gain. Essential oil
200 supplementation significantly increased feed intake, but had no significant effect on feed per
201 gain. There was no significant interaction between wheat form and EO supplementation for
202 feed intake and feed per gain.

203 During the finisher period (22-35 d), wheat form had significant no effect on weight
204 gain, feed intake or feed per gain. Essential oil supplementation significantly improved weight

gain both in GW and WW diets. No significant wheat form x EO interactions were observed for any of the performance parameters measured.

Over the whole trial period (1-35 d), EO supplementation significantly improved weight gain in both diets, but the magnitude of improvement was greater in birds fed on the GW diet, resulting in a significant wheat form x EO interaction. Whole wheat inclusion tended ($P = 0.06$) to improve the feed per gain.

Nutrient utilisation

The influence of treatments on ileal nitrogen and energy digestibility is summarised in Table 3. Whole wheat inclusion and EO supplementation significantly improved ileal nitrogen digestibility. A tendency ($P = 0.06$) for wheat form x EO interaction was observed for ileal energy digestibility coefficient and apparent ileal digestible energy. Essential oils supplementation tended to improve ileal energy digestibility coefficient and IDE in birds fed on the GW diet, but had no effect in those fed on the WW diet.

Tables 2 and 3 near here

Digestive tract morphology

The effects of wheat form and EO supplementation on the relative weight, length and digesta content of various digestive compartments are shown in Table 4. Treatments had no effect on most of the parameters measured. However, a significant interaction between wheat form and EO supplementation was found for relative gizzard and caecal weights. Essential oil supplementation increased the relative gizzard weight and reduced the caecal weight in birds fed on the GW diet, but had no effect in birds fed on the WW diet.

Number of bands and Simpsons diversity index of diversity

The effects of wheat form and EO supplementation on the number of bands and Simpsons diversity index of diversity are shown in Table 5. There were no significant differences in the numbers of bacterial species from the ileal microflora of birds fed on the GW diet compared with those fed on the WW diet. However, the mean numbers of bacterial species in the ileal

230 contents of birds fed on the GW diet supplemented with EO diet tended ($P = 0.07$) to be
231 higher than those fed on the unsupplemented GW diet.

Tables 4 and 5 near here

232 **Dendographic analysis**

233 Dendographic analysis showed a tendency for similarity of microbial communities on a basis
234 of feed form but not on a basis of EO supplementation. In all gels (Figures 1 to 3), the results
235 from birds fed on ground and whole wheat diets tended to be separated, and to cluster on
236 opposite sides of the dendogram.

Figures 1, 2 and 3 near here

237 **DISCUSSION**

238 In the present study, over the whole 35-d trial period, WW inclusion tended to improve feed
239 per gain, but had no effect on the weight gain or feed intake. Published data on the effects of
240 WW inclusion in broiler diets have been contradictory. Several studies have reported
241 improvements in feed per gain when WW has replaced part of the ground grain (Plavnik *et al.*,
242 2002; Wu and Ravindran, 2004; Wu *et al.*, 2004; Ravindran *et al.*, 2006), while others
243 found no improvement in feed efficiency (Preston *et al.*, 2000; Banfield and Forbes, 2001;
244 Svihus *et al.*, 2004; Amerah and Ravindran, 2008). In one study, feed efficiency was found to
245 be poorer when WW substituted part of the ground wheat (Bennett *et al.*, 2002). Several
246 factors may be contributing to these inconsistent results, including differences in methodology
247 and experimental protocols (Wu and Ravindran, 2004) or in the dietary nutrient density in the
248 basal diets (Plavnik *et al.*, 2002). Differences in wheat types/cultivars may provide a third
249 explanation (Amerah *et al.*, 2009b), as the wide variability in the nutrient composition
250 (Inglett, 1974), metabolisable energy (Hughes and Choct, 1999) and nutrient digestibility
251 (Ravindran *et al.*, 2005; Carré *et al.*, 2007) of wheat is well known.

252 In the present study, the blend of cinnamaldehyde and thymol supplementation
253 improved weight gain both in GW and WW diets, with a greater response observed in birds
254 fed on the ground wheat diet. Several studies have reported beneficial effects of EO on weight

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

gain (Bento *et al.*, 2008; Péron *et al.*, 2009; Yang *et al.*, 2009), while others reported no effect (Botsoglou *et al.*, 2002; Zhang *et al.*, 2005; Jang *et al.*, 2007). These differences have been attributed to the type of EO used and inclusion level (Cross *et al.*, 2007). Cross *et al.* (2007), comparing the effect of inclusion of 10 or 100 g/tonne of different herbs (thyme, oregano, marjoram, rosemary or yarrow), reported that dietary thyme oil or yarrow herb inclusion had the most positive effects on chick performance, while oregano herb and yarrow oil were the poorest. Hernandez *et al.* (2004) reported that an EO blend, containing oregano, cinnamon, and pepper and Labiatae extract from sage, thyme, and rosemary extracts, fed to broilers gave performance levels similar to those of the antibiotic growth promoter, Avilamycin. Improved weight gain in birds given EO in the current study may be related to the improvements in the ileal N digestibility observed with EO supplementation. The positive effect of EO on protein digestibility as well as DM digestibility has been observed in previous studies (Hernandez *et al.*, 2004; Cao *et al.*, 2009). The higher response of EO supplementation in the GW diet may be due to the observed tendency towards lower energy digestibility in these birds. These positive effects of EO on nutrient digestibility may be associated, in part, with their digestion-stimulating properties and antimicrobial effects (Kamel, 2001).

It was expected that WW inclusion would modulate the development of the digestive tract as this has been observed in most previous studies (Preston *et al.* 2000; Plavnik *et al.* 2002; Ravindran *et al.*, 2006; Amerah and Ravindran, 2008). Surprisingly, wheat form had no effect on any of the measured parameters of the digestive tract. Most studies reported increased gizzard size with WW inclusion, however, no effect of WW inclusion on relative gizzard weight or its relative digesta content was observed in the current study. The reason for this finding is not clear, but thinner gizzard muscles were observed in birds fed on the whole wheat. Thus, WW inclusion may increase gizzard volume but cause thinner gizzard muscles. Amerah *et al.* (2007b), comparing mash vs. pelleted diets in broilers, reported that the

gizzards of birds fed on mash or pelleted diets increased uniformly in size, there being no significant change either in the ratio of content to empty mass or in the ratio of gizzard length to breadth with diet. Another possible explanation is that the wheat used in this study was soft cultivar and did not stimulate the growth of the gizzard. Interestingly, EO supplementation increased gizzard weight in birds fed on the GW diet, but had no effect in birds fed on the WW diet. However, the interaction effect of EO and wheat form on caecal weight was less clear. When included in ground wheat based diet EO resulted a reduction in caecal weight compared with those fed on supplemented diet in whole wheat based diet. The reasons for these findings are unclear. In contrast, Hernandez *et al.* (2004) reported no effect of EO inclusion on digestive tract development.

One of the objectives of the present experiment was to investigate whether the inclusion of WW and EO supplementation influences the ileal microbial composition of broilers. The DGGE band patterns (Figures 1, 2 and 3) showed that ileal microbiota composition was influenced by feed form. Similarly, Amerah *et al.* (2009a) reported that ileal microbiota profiling using DGGE showed separation between birds receiving the GW-based diet and birds fed on the WW-based diet. Previous studies, using media culture techniques, have shown that WW feeding increases the intestinal counts of some *Lactobacillus* species (Engberg *et al.*, 2004) and reduces the number of *Clostridium perfringens* in the intestinal tract (Engberg *et al.*, 2004; Bjerrum *et al.*, 2005) compared with those fed on the GW-based diets.

In this study, the mean numbers of bacterial species in ileal contents of birds fed on the GW diet supplemented with EO tended to be higher than the ileal contents of birds fed on the unsupplemented GW diet. Hence, the addition of EO to GW diet appears to have induced the growth of additional species but with no accompanying overall changes in diversity. A previous study, using the same blend of EO, reported that the EO supplementation tended to

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

lower the proportion of *E. coli* and significantly increase the proportion of bifidobacteria in the caecal digesta (Tihihonen *et al.*, 2010). On the other hand, in the present study, EO supplementation of the WW diet showed no change in the mean numbers of bacterial species or DGGE band patterns, suggesting that the ileal microbiota composition was not influenced by EO supplementation. Whole wheat inclusion induced changes in DGGE band patterns, which may have masked the effect of EO on the microbial profile in these birds. Further studies with greater numbers of replicates are required to confirm this finding.

In conclusion, the findings of the present study suggest that this blend of cinnamaldehyde and thymol improves weight gain of broilers regardless of the wheat form, but the response may be higher in the ground wheat based diet. WW inclusion showed performance benefits during the starter phase, but did not result in a significant improvement for the overall period. Essential oil supplementation and WW feeding had a positive effect of on ileal nitrogen digestibility and an hypothesis that a combination of EO and WW will have additive effects on broiler gut microflora was also tested. Essential oil supplementation in birds fed on the WW diet showed no change in DGGE band patterns, suggesting that ileal microbiota composition was not influenced by EO supplementation in these birds.

REFERENCES

AMERAH, A.M. & RAVINDRAN, V. (2008) Influence of method of whole wheat feeding on the performance, digestive tract development and carcass traits of broiler chickens. *Animal Feed Science and Technology*, 147: 326-339.

AMERAH, A.M., RAVINDRAN, V., LENTLE, R.G. & THOMAS, D.G. (2007a) Feed particle size: Implications on the digestion and performance in poultry. *World's Poultry Science Journal*, 63: 439-451.

- 328 AMERAH, A.M., RAVINDRAN, V., LENTLE, R.G. & THOMAS, D.G. (2007b) Influence
329 of feed particle size and feed form on the performance, energy utilisation, digestive tract
330 development and digesta parameters of broiler starters. *Poultry Science*, 86: 2615-2623.
- 331 AMERAH, A.M., RAVINDRAN, V. & LENTLE, R.G. (2009a) Influence of insoluble fibre
332 and whole wheat inclusion on the performance, digestive tract development and ileal
333 microbiota profile of broiler chickens. *British Poultry Science*, 50: 366-375.
- 334 AMERAH, A.M., RAVINDRAN, V. & LENTLE, R.G. (2009b) Influence of wheat hardness
335 and xylanase supplementation on the performance, energy utilisation, digestive tract
336 development and digesta parameters of broiler starters. *Animal Production Science*, 49: 71-
337 78.
- 338 AOAC. (2005) Official Methods of Analysis. 18th edition. Association of Official Analytical
339 Chemists, Washington DC, USA.
- 340 BANFIELD, M.J. & FORBES, J.M. (2001) Effects of whole wheat dilution v. substitution on
341 coccidiosis in broiler chickens. *British Journal of Nutrition*, 86: 89-95.
- 342 BENNETT, C.D., CLASSEN, H.L. & RIDDELL, C. (2002) Feeding broiler chickens wheat
343 and barley diets containing whole, ground and pelleted grain. *Poultry Science*, 81: 995-1003.
- 344 BENTO, M.H.L., VAN DER KLIS, J.D. & SCHULZE, H. (2008) Efficacy of essential oils
345 on broiler growth performance in a semi-commercial scale facility. EAAP, 59th annual
346 meeting, Vilnius, pp144.
- 347 BJERRUM, L., PEDERSEN, K. & ENGBERG, R.M. (2005) The influence of whole wheat
348 feeding on salmonella infection and gut flora composition in broilers. *Avian Disease*, 49: 9-
349 15.
- 350 BOTSOGLOU, N.A., FLOROU-PANER, P., CHRISTAKI, E., FLETOURIS, D.J. & SPAIS,
351 A.B. (2002) Effect of dietary oregano essential oil on performance of chickens and on iron-
352 induced lipid oxidation of breast, thigh and abdominal fat tissues. *British Poultry Science*, 43:
353 223-230.

- 354 BURT, S. (2004) Essential oils: their antibacterial properties and potential applications in
355 foods - a review. *International Journal of Food Microbiology*, 94: 223-253.
- 356 CAO, P.H., LI, F.D., LI, Y.F., RU, Y.J., PÉRON, A. & SCHULZE, H. (2009) Essential oils
357 and feed enzymes improve nutrient utilization in broiler chickens fed a corn/soy-based diet
358 reduced in calcium and phosphorus. ASAS-CAAV Pacific Rim Conference, Beijing, China,
359 pp.14.
- 360 CARRÉ B., MIGNON-GRASTEAU S., PÉRON A., JUIN H. & BASTIANELLI D. (2007)
361 Wheat value: improvements by feed technology, plant breeding and animal genetics. *World's*
362 *Poultry Science Journal*, 63: 585-596
- 363 CROSS, D.E., MCDEVITT, R.M., HILLMAN, K. & ACAMOVIC, T. (2007) The effect of
364 herbs and their associated essential oils on performance, dietary digestibility and gut
365 microflora in chickens from 7 to 28 days of age. *British Poultry Science*, 48: 496-506.
- 366 ENGBERG, R.M., M.S. HEDEMANN, S. STEENFELDT & JENSEN, B.B. (2004) Influence
367 of whole wheat and xylanase on broiler performance and microbial composition and activity
368 in the digestive tract. *Poultry Science*, 83: 925-938.
- 369 FERKET, P. (2000) Feeding whole grains to poultry improves gut health. *Feedstuffs USA*
370 (September), pp. 12-14.
- 371 HERNANDEZ, F., MADRID, J., GARCIA, V., ORENGO, J. & MEGIAS, M.D. (2004)
372 Influence of two plant extracts on broiler performance, digestibility and digestive organ size.
373 *Poultry Science*, 83: 169-174.
- 374 HUGHES, R.J. & CHOCT, M. (1999) Chemical and physical characteristics of grains related
375 to variability in energy and amino acid availability in poultry, *Australian Journal of*
376 *Agricultural Research*, 50: 689-702.
- 377 HUME, M.E., KUBENA, L.F. EDRINGTON, T.S. DONSKEY, C.J. MOORE, R.W.
378 RICKE S.C. & NISBET, D.J. (2003) Poultry digestive microflora biodiversity as indicated by

- denaturing gradient gel electrophoresis. *Poultry Science*, 82: 1100-1107.
- INGLETT, G.E. (1974) *Wheat: Production and Utilisation*, The AVI Publishing Company, Inc., Westport, CT, USA.
- JANG, I.S., KO, Y.H., KANG, S.Y. & LEE, C.Y. (2007) Effect of a commercial essential oil on growth performance, digestive enzyme activity and intestinal microflora population in broiler chickens. *Animal Feed Science and Technology*, 134: 304-315.
- KAMEL, C. (2001) Tracing methods of action and roles of plant extracts in non-ruminants. In: *Recent Advances in Animal Nutrition* (eds.). Garnsworthy, P. C. and J. Wiseman, Nottingham University Press, Nottingham, UK.
- LEE, K. W., H. EVERTS, H. J. KAPPERT, H. WOUTERSE, M. FREHNER & A. C. BEYNEN (2004) Cinnaminaldehyde, but not thymol, counteracts the carboxymethyl cellulose-induced growth depression in female broiler chickens. *International Journal of Poultry Science*, 3: 608- 612.
- PÉRON, A., BENTO, M.H.L., SCHULZE, H., AMERAH, A.M. & JUIN, H. (2009) Effect of exogenous enzymes and essential oils on the performance of broiler chickens fed wheat-based diets. 17th European Symposium on Poultry Nutrition, Edinburgh, Scotland, pp 273.
- PLAVNIK, I., MACOVSKY, B. & SKLAN, D. (2002) Effect of feeding whole wheat on performance of broiler chickens. *Animal Feed Science and Technology*, 96: 229-236.
- PRESTON, C.M., MCCracken, K.J. & McALLISTER, A. (2000) Effect of diet form and enzyme supplementation on growth, efficiency and energy utilisation of wheat-based diets for broilers. *British Poultry Science*, 41: 324-331.
- RAVINDRAN, V. HEW, L.I. RAVINDRAN, G. & BRYDEN, W.L. (2005) Apparent ileal digestibility of amino acids in feed ingredients for broiler chickens, *Animal Science*, 81: 85–97
- RAVINDRAN, V., WU, Y.B., THOMAS, D.G. & MOREL, P.C.H. (2006) Influence of

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

whole wheat feeding on the development of gastrointestinal tract and performance of broiler chickens. *Australian Journal of Agricultural Research*, 57: 21-26.

SAS (1997) SAS/STAT® User's Guide: Statistics. Version 6.12. SAS Institute Inc., Cary.

NC.SHORT, F.J.P., GORTON, J., WISEMAN, J. & BOORMAN, K.N. (1996) Determination of titanium oxide added as an inert marker in chicken digestibility studies. *Animal Feed Science and Technology*, 59: 215–221.

SVIHUS, B. & HETLAND, H. (2001) Ileal starch digestibility in growing broiler chickens fed on a wheat-based diet is improved by mash feeding, dilution with cellulose or whole wheat inclusion. *British Poultry Science*, 42: 633-637.

SVIHUS, B., JUVIK, E., HETLAND, H., & KROGDAHL, A. (2004) Causes for improvement in nutritive value of broiler chicken diets with whole wheat instead of ground wheat. *British Poultry Science*, 45: 55-60.

TIIHONEN, K., KETTUNEN, H., BENTO, M.H.L., SAARINEN, M. LAHTINEN, S., OUWEHAND, A.C., SCHULZE, H. & RAUTONEN N. The effect of feeding essential oils on broiler performance and gut microbiota. *British Poultry Science*, **51**: 381-392.

WALLACE, R.J. (2004) Antimicrobial properties of plant secondary metabolites. *Proceedings of the Nutrition Society*, 63:621 – 629.

WU, Y.B. & RAVINDRAN V. (2004) Influence of whole-wheat inclusion and xylanase supplementation on the performance, digestive tract measurements and carcass characteristics of broiler chickens. *Animal Feed Science and Technology*, 116: 129-139.

WU, Y.B., RAVINDRAN, V., THOMAS, D.G., BIRTLES, M.J. & HENDRIKS, W.H. (2004) Influence of method of whole wheat inclusion and xylanase supplementation on the performance, apparent metabolisable energy, digestive tract measurements and gut morphology of broilers. *British Poultry Science*, 45: 385-394.

YANG, Y., IJI, P.A. & CHOCT, M. (2009) Dietary modulation of gut microflora in broiler

- 1
2
3 429 chickens: a review of the role of six kinds of alternatives to in-feed antibiotics. *World's*
4
5 430 *Poultry Science Journal*, 65: 97-114.
6
7
8 431 ZHANG, K.Y., YAN, F., KEEN, C.A. and WALDROUP, P.W. (2005) Evaluation of
9
10 432 microencapsulated essential oils and organic acids in diets for broiler chickens. *International*
11
12 433 *Journal of Poultry Science*, 4: 612-619.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1. *Composition and calculated analysis (g/kg) of the basal diets*

Ingredient as fed	Starter diet	Finisher diet
Wheat	619.4	660.5
Soybean meal 48	304.2	254.7
Soybean oil	34.98	46.03
Dicalcium phosphate	17.49	14.92
Limestone	1.166	0.964
Lysine HCl	2.41	1.56
DL-Methionine	2.54	2.04
L-Threonine	0.27	0.56
Salt	2.97	2.97
Trace mineral-vitamin premix ¹	3.00	3.00
Titanium oxide	-	3.00
Xylanase ²	+	+
<i>Calculated analysis</i>		-
Metabolisable energy, MJ/kg	12.55	12.97
Crude protein	220.0	200.0
Lysine	13.0	11.0
Methionine + cystine	9.50	8.50
Calcium	10.0	8.50
Available phosphorus	4.50	4.00

¹Supplied per kilogram of diet: antioxidant, 100 mg; biotin, 0.2 mg; calcium pantothenate, 12.8 mg; cholecalciferol, 60 µg; cyanocobalamin, 0.017 mg; folic acid, 5.2 mg; menadione, 4 mg; niacin, 35 mg; pyridoxine, 10 mg; trans-retinol, 3.33 mg; riboflavin, 12 mg; thiamine, 3.0 mg; dl-α-tocopheryl acetate, 60 mg; choline chloride, 638 mg; Co, 0.3 mg; Cu, 3 mg; Fe, 25 mg; I, 1 mg; Mn, 125 mg; Mo, 0.5 mg; Se, 200 µg; Zn, 60 mg.

² Danisco Xylanase, Danisco Animal Nutrition, Marlborough, UK. The enzyme was included at a rate of 1kg/tonne. According to the manufacturer, this inclusion rate supplied 2000 xylanase units/kg feed.

Table 2. Influence of wheat form and essential oil supplementation on the weight gain (g) feed intake (g) and feed per gain (g/g) of male broilers fed on wheat/soy-based diets¹

Wheat form	Essential oil	1-21 d			22-35 d			1-35 d		
		Weight gain	Feed intake	Feed per gain	Weight gain	Feed intake	Feed per gain	Weight gain	Feed intake	Feed per gain
Ground wheat	-	939 ^b	1312	1.399	1406	2392	1.701	2344 ^c	3704	1.580
	+	1002 ^a	1355	1.366	1448	2385	1.647	2451 ^a	3740	1.540
Whole wheat	-	986 ^a	1310	1.328	1407	2356	1.675	2393 ^b	3666	1.531
	+	986 ^a	1325	1.344	1447	2400	1.659	2433 ^a	3725	1.531
Pooled SEM		8.5	11.8	0.0137	10.5	33.9	0.0235	12.7	40.8	0.0143
Main effects										
<i>Wheat form</i>										
Ground wheat		970	1333	1.382 ^a	1427	2388	1.674	2397	3722	1.560
Whole wheat		986	1317	1.336 ^b	1427	2378	1.667	2413	3695	1.531
<i>Essential oil</i>										
-		962	1311 ^a	1.364	1407 ^a	2374	1.688	2369	3685	1.556
+		994	1340 ^b	1.355	1447 ^b	2392	1.653	2442	3732	1.536
<i>P values</i>										
Wheat form		0.08	NS	**	NS	NS	NS	NS	NS	0.06
Essential oil		***	*	NS	***	NS	NS	***	NS	NS
Wheat form x essential oil		***	NS	0.09	NS	NS	NS	*	NS	NS

¹ Each value represents the mean of 6 replicates (8 birds per replicate).

^{a,b,c} Means in a column not sharing a common superscript are significantly different at $P < 0.05$.

NS = not significant; * = $P < 0.05$; ** = $P < 0.01$; *** = $P < 0.0001$.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 3. *Influence of wheat form and essential oil supplementation on ileal digestibility coefficients of nitrogen (IDCN) and gross energy (IDCGE), and apparent ileal digestible energy (AIDE, MJ/kg DM) of male broilers fed on wheat-soy diets (1-35 d)¹*

Wheat form	Essential oils	IDCN	IDCGE	AIDE
Ground wheat	-	0.840	0.696	13.23
	+	0.861	0.738	14.02
Whole wheat	-	0.861	0.748	14.19
	+	0.865	0.736	13.96
Pooled SEM		0.006	0.013	0.253
Main effects				
Wheat form				
Ground wheat		0.850 ^a	0.717	13.62
Whole wheat		0.863 ^b	0.741	14.08
Essential oil				
-		0.850 ^a	0.722	13.71
+		0.863 ^b	0.737	13.99
<i>P</i> values				
Wheat form		*	0.08	0.09
Essential oil		*	NS	NS
Wheat form x essential oil		NS	0.06	0.06

¹ Each value represents the mean of 6 replicates (8 birds per replicate).
^{a,b} Means in a column not sharing a common superscript are significantly different at *P* < 0.05.
NS = not significant; * = *P* < 0.05.

Table 4. *Influence of wheat form and essential oil supplementation on gross morphology of the digestive tract of male broilers fed on wheat-soy diets (1-35 d)¹*

Wheat form	Ground wheat		Whole wheat		SEM	P values		
Essential oil	-	+	-	+		Wheat form (WF)	Essential oil (EO)	WF x EO
Relative empty weights (g/kg BW)								
Pancreas	1.67	1.63	1.68	1.75	0.059	NS	NS	NS
Gizzard	6.25 ^b	7.07 ^a	7.13 ^a	6.50 ^{ab}	0.223	NS	NS	**
Small intestine ²	16.89	15.90	16.60	16.80	0.420	NS	NS	NS
Caeca	0.985 ^{ab}	0.906 ^b	0.947 ^{ab}	1.031 ^a	0.038	NS	NS	*
Relative digesta content (g/kg BW)								
Gizzard	1.57	2.10	1.94	2.61	0.389	NS	NS	NS
Small intestine ²	33.56	34.51	31.69	31.60	1.70	NS	NS	NS
Caeca	1.635	1.594	1.621	1.624	0.204	NS	NS	NS
Relative length (cm/kg BW)								
Small intestine ²	88.00	80.36	83.12	84.01	2.10	NS	NS	NS
Caeca	8.66	8.44	8.78	8.70	0.215	NS	NS	NS

^{a,b}Means in a row not sharing a common superscript are significantly different at $P < 0.05$.

¹Each value represents the mean $\pm$ SEM of 12 birds.

²Small intestine = duodenum + jejunum + ileum.

Table 5. *Influence of wheat form and essential oil supplementation on overall species diversity*

Feed Form	Essential oil	Band number	Shannon Weiner
Ground wheat	-	13±0.6	1.03±0.17
Ground wheat	+	16±0.7 ¹	0.96±0.11
Whole wheat	-	14±0.5	0.73±0.18
Whole wheat	+	14±0.8	0.94±0.14

¹Tended ($P = 0.07$) to be different from unsupplemented ground wheat diet

Figure 1. Denaturant gradient gel electrophoresis profile (Gel 1) from ileal digesta of broilers as influenced by wheat form (whole wheat [WW] and ground wheat [GW]) and essential oils (EO). Relative similarity of band patterns of two denaturing gradient gel electrophoresis gels is indicated by their grouping on the dendrogram and the percentage coefficient (bar).

Figure 2. Denaturant gradient gel electrophoresis profile (Gel 2) from ileal digesta of broilers as influenced by wheat form (whole wheat [WW] and ground wheat [GW]) and essential oils (EO). Relative similarity of band patterns of two denaturing gradient gel electrophoresis gels is indicated by their grouping on the dendrogram and the percentage coefficient (bar).

Figure 3. Denaturant gradient gel electrophoresis profile (Gel 3) from ileal digesta of broilers as influenced by wheat form (whole wheat [WW] and ground wheat [GW]) and essential oils (EO). Relative similarity of band patterns of two denaturing gradient gel electrophoresis gels is indicated by their grouping on the dendrogram and the percentage coefficient (bar).