Changes in the nutrient ratios and phytoplankton community after declines in nutrient concentrations in an eutrophic semi-enclosed bay in Hong Kong

Alle A.Y. Lie, C. Kim Wong, Jackie Y.C. Lam, J.H. Liu, Y.K. Yung

► To cite this version:

Alle A.Y. Lie, C. Kim Wong, Jackie Y.C. Lam, J.H. Liu, Y.K. Yung. Changes in the nutrient ratios and phytoplankton community after declines in nutrient concentrations in an eutrophic semi-enclosed bay in Hong Kong. *Marine Environmental Research*, 2011, 71 (3), pp.178. 10.1016/j.marenvres.2011.01.001 . hal-00673201

HAL Id: hal-00673201

<https://hal.science/hal-00673201>

Submitted on 23 Feb 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Title: Changes in the nutrient ratios and phytoplankton community after declines in nutrient concentrations in an eutrophic semi-enclosed bay in Hong Kong

Authors: Alle A.Y. Lie, C. Kim Wong, Jackie Y.C. Lam, J.H. Liu, Y.K. Yung

PII: S0141-1136(11)00003-1

DOI: [10.1016/j.marenvres.2011.01.001](https://doi.org/10.1016/j.marenvres.2011.01.001)

Reference: MERE 3497

To appear in: *Marine Environmental Research*

Received Date: 17 June 2010

Revised Date: 5 January 2011

Accepted Date: 12 January 2011

Please cite this article as: Lie, A.A., Wong, K., Lam, J.Y., Liu, J.H., Yung, Y.K. Changes in the nutrient ratios and phytoplankton community after declines in nutrient concentrations in an eutrophic semi-enclosed bay in Hong Kong, *Marine Environmental Research* (2011), doi: 10.1016/j.marenvres.2011.01.001

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 Changes in the nutrient ratios and phytoplankton community after declines in nutrient
2 concentrations in an eutrophic semi-enclosed bay in Hong Kong

3

4 Alle A. Y. Lie^{1^*}, C. Kim Wong¹, Jackie Y. C. Lam², J.H. Liu² & Y. K. Yung²

5

6 ¹Department of Biology, The Chinese University of Hong Kong, Shatin, Hong Kong

7 ²Water Policy and Science Group, Hong Kong Government Environmental Protection

8 Department, Revenue Tower, 5 Gloucester Road, Wan Chai, Hong Kong

9 [^]Current address: Department of Biological Sciences, University of Southern

10 California, 3616 Trousdale Parkway, Los Angeles, 90089-0371, CA, USA

11

12 ^{*}Corresponding author:

13 email: alie@usc.edu

14 Tel.: +852 26096773; Fax: +852 26035391

15 Abstract

16

17 Tolo Harbour is a landlocked bay with poor tidal flushings in the northeastern part of
18 Hong Kong. During the 1980s, excessive nutrient loading led to dramatic increase in
19 nutrient concentrations, accompanied by lower N:P ratios, higher algal biomass and
20 shifts in the phytoplankton community. We studied the effects of nutrient loading
21 reduction measures on nutrient concentrations, nutrient ratios and phytoplankton
22 dynamics in Tolo Harbour by comparing data collected before the full implementation
23 of nutrient loading reduction measures (1986–1997) to those after the implementation
24 (1998–2008). Such measures led to declines in nutrient concentrations, changes in
25 N:P and N:Si ratios, lower chlorophyll-*a* concentrations and fewer algal blooms.
26 Diatoms were the most abundant phytoplankton group in Tolo Harbour both before
27 and after declines in nutrient concentrations. The density of dinoflagellates did not
28 change, but substantial increase in other algal group abundance was recorded.

29

30 Keywords: Nutrients, Nutrient ratios, Phytoplankton, Eutrophication, Coastal waters,
31 Tolo Harbour.

1. Introduction

Eutrophication is a common consequence of anthropogenic activities on water bodies, and is widely regarded as the most wide-spread water quality problem globally (Carpenter et al., 1999; Smith et al., 1999; Smith, 2003). The most common effects of eutrophication are increases in phytoplankton productivity and biomass, which in turn produce other adverse effects such as decreases in water transparency, increases in the frequency of bottom hypoxia or anoxia and loss of various aquatic organisms (Nixon, 1995; Smith et al., 1999; Smith, 2003). Numerous recent studies have linked eutrophication to algal blooms which cause ecological and economical damages (e.g. Anderson et al., 2008; Cloern, 2001; Smith et al., 1999). Nutrient loadings may cause changes in the nutrient ratios (Justić et al., 1995; Turner, 2002), and such changes may affect the species composition of the phytoplankton community and higher trophic levels (Officer and Ryther, 1980; Philippart et al., 2000; Smith, 1983; Turner et al., 1998; Turner, 2002). Although eutrophication in freshwater systems has been studied extensively and intensively during the past few decades, with many studies reporting on the effects of increased or decreased nutrient concentrations in lakes (e.g. Battarbee et al., 2005; Carvalho and Kirika, 2003; Jeppesen et al., 2005; Jeppesen et al., 2007; Mehner et al., 2001), similar kinds of

51 studies, with long-term monitoring data, are relatively scarce in marine coastal
52 ecosystems (Examples summarized by Smith et al., 1999).

53 Hong Kong is one of the most densely populated cities in the world. Intensive
54 urbanization and industrialization during the past few decades has dramatically
55 increased nutrient input into Hong Kong's coastal seas (Lee et al., 2006). The concern
56 on the coastal water quality led to the initiation of a marine water quality monitoring
57 program by the Hong Kong Environmental Protection Department in the 1980s (Lee
58 et al., 2006). One of the intensely monitored zones is Tolo Harbour, a landlocked bay
59 in the northeastern corner of Hong Kong.

60 Tolo Harbour consists of a shallow inner bay, with a mean depth of 6–7 m, and a
61 narrow tidal channel which opens into the neighboring Mirs Bay. Urban development
62 and large-scale reclamation around Tolo Harbour began in the 1970s. Sewage
63 discharge from the expanding population caused a sharp increase in nutrient input
64 (Lam and Ho, 1989; Wear et al., 1984). Hodgkiss and Yim (1995) estimated a daily
65 nitrate and phosphate loading of 32,460 kg and 3,190 kg respectively into Tolo
66 Harbour. In addition to nutrient inputs, the bottleneck topography of the Tolo Harbour
67 leads to poor tidal exchange rates and persistence of high nutrient concentrations (Lee
68 et al., 2006). Increased nutrient concentrations produced algal blooms, with
69 dinoflagellates such as *Noctiluca scintillans* being the most common causative species

during the 1980s and 1990s (Lam and Ho, 1989; Liu and Wong, 2006). Although *Noctiluca* are generally considered as heterotrophic (Kjørboe and Titelman, 1998), strains of *Noctiluca miliaris* with photosynthetic symbionts have been found in Southeast Asia (Sweeney, 1971; Hansen et al., 2004), which may benefit from increased nutrient concentrations (Sriwoon et al., 2008). In addition, the increased nutrient concentrations would also favor the growth of diatoms and other prey for *N. scintillans* (Liu and Wong, 2006).

Wear et al. (1984) reported mean concentrations of $\text{NO}_x\text{-N}$ and $\text{PO}_4\text{-P}$ in the inner Tolo Harbour to be $0.05\ \mu\text{M}$ and $0.01\ \mu\text{M}$ respectively during 1974–1976. By early 1990s, annual mean concentrations have increased to $19.99\ \mu\text{M}$ for inorganic nitrogen and $1.94\ \mu\text{M}$ for $\text{PO}_4\text{-P}$ (Hong Kong Environmental Protection Department, 1991). Moreover, $\text{PO}_4\text{-P}$ values as high as $10.65\ \mu\text{M}$ have been reported in Tolo Harbour in 1978–1979 (Hodgkiss and Chan, 1983). Decreasing trends in N:P ratio were recorded during the 1980s and early 1990s (Hodgkiss and Chan, 1997; Yung et al., 1997), and was considered to be caused by excessive P input (Hodgkiss and Chan, 1983; Hodgkiss and Yim, 1995). Some researchers noted that the observed changes in nutrient parameters had caused a shift in the species composition of the phytoplankton community, with diatoms gradually replaced by dinoflagellates, (Chan and Hodgkiss, 1987; Hodgkiss and Ho, 1997; Lam and Ho, 1989), although a review of the data

89 collected by the Hong Kong Environmental Protection Department did not reveal such
90 a shift in the phytoplankton during 1982–1992 (Yung et al., 1997).

91 The Hong Kong Government initiated the Tolo Harbour Action Plan in 1986 to
92 halt the declining water quality of Tolo Harbour. Two of the most important
93 components of the action plan were the modification of existing sewage treatment
94 works processes (implemented in 1992) and the Tolo Harbour Effluent Export
95 Scheme (Chau, 2007). Commissioned in the mid-1990s and became fully operational
96 in 1998, the Tolo Harbour Effluent Export Scheme allows treated sewage from two
97 large sewage treatment works to be exported via pipelines for discharge outside of
98 Tolo Harbour. Nitrogen input into Tolo Harbour was reduced to $\sim 1000 \text{ kg day}^{-1}$ by
99 2000 (Hong Kong Environmental Protection Department, 2006).

100 In this study, data collected from four sampling stations in Tolo Harbour during
101 the period from 1986–2008 were analyzed to provide information on the long-term
102 changes in the hydrography and phytoplankton dynamics. Conditions before
103 (1986–1997) and after (1998–2008) the full implementation of the Tolo Harbour
104 Effluent Export Scheme were compared to: (1) determine whether declines in nutrient
105 concentrations had led to changes in the nutrient ratios, and (2) identify shifts in the
106 phytoplankton community that may be related to changes in the nutrient
107 concentrations and ratios. Overall changes of the hydrography and phytoplankton

community in Tolo Harbour after declines in nutrient concentrations were also compared to those reported in other ecosystems after recovery from eutrophication.

2. Materials and Methods

All data were provided by the Hong Kong Environmental Department. Four sampling stations in Tolo Harbour, TM3, TM4, TM6 and TM8 (Fig. 1), were sampled frequently, mostly once each month, but irregularly from 1982–1992. Monthly means were calculated for each station. Physico-chemical parameters, including temperature, salinity, pH and dissolved oxygen concentration (DO), were recorded at 1 m below the water surface, mid-depth in the water column and 1 m above the bottom by a SEACAT 19*plus* CTD water profiler (Sea-BIRD Electronics). Water samples were also collected at the three depths using Niskin-type water bottles for the determination of BOD₅, chlorophyll-*a* concentration, suspended solids concentration, and the concentrations of NO₃-N, NH₄-N, total inorganic nitrogen (TIN), total nitrogen (TN), PO₄-P, total phosphorus (TP) and SiO₂. The data from the three depths were averaged into depth-averaged data. Values from the bottom depth were also analyzed for DO and BOD₅ to provide information on hypoxia at the water bottom. Measurement of BOD₅, nutrient concentrations, chlorophyll-*a* concentration and suspended solids

concentration were carried out by the Hong Kong Government Laboratory using procedures described by APHA (American Public Health Association et al. 1998). Data on algal bloom occurrences were obtained from the Hong Kong Agriculture, Fisheries and Conservation Department, and were independent from our data. Such data on algal bloom occurrences was based on sighting reports of seawater discoloration, and may therefore underestimate algal bloom occurrences. It also does not contain taxonomic information regarding the algal bloom.

(Insert Fig. 1 here)

Water samples from 1 m below the water surface were preserved in Lugol's iodine solution (1: 100 ml). In the laboratory, phytoplankton were examined in an Utermöhl counting chamber under an inverted microscope, and the densities of diatoms, dinoflagellates and other algal groups (Algae that were neither diatoms nor dinoflagellate) were counted. Phytoplankton data for TM6 was only available after 1993, while data collection for other stations have began in 1986 (TM8) and 1988 (TM3 and TM4).

Two-way ANOVA was used to test for temporal differences before and after nutrient loading reduction, as well as the spatial differences among sampling stations

in various water quality and phytoplankton parameters. Tukey's test was used for multiple comparison between sites. Kendall's correlation was used to test the correlation between phytoplankton and water quality variables.

3. Results

In general, physico-chemical, biological and nutrient parameters showed similar changes among the four sampling stations after nutrient loading reduction (Table 1). Temperature, salinity, bottom DO and TIN/PO₄ increased at all stations after nutrient loading reduction, while pH, BOD₅, chlorophyll-*a* concentrations, all nutrient concentrations and TIN/SiO₂ decreased. DO and SiO₂ concentration did not differ significantly between the two time periods and among the stations respectively ($p > 0.05$). All other parameters showed significant differences both temporally and spatially, with most p values < 0.001 . Comparison among sampling stations revealed a general trend of improvement in water quality, marked by changes such as decreasing nutrient concentrations and increasing DO and secchi depth, from TM3 to TM8.

(Insert Table 1 here)

N (including $\text{NO}_3\text{-N}$, $\text{NH}_4\text{-N}$, total inorganic N and total N) and P (including $\text{PO}_4\text{-P}$ and total P) concentrations were high, especially at TM3 and TM4, prior to nutrient loading reduction (Fig. 2). Concentrations decreased noticeably within a year after the full implementation of the sewage export scheme, and have remained at relatively low levels in recent years. N and P concentrations differed markedly among stations before nutrient loading reduction, but the differences became less apparent afterwards. SiO_2 concentrations did not appear to differ among stations before nutrient loading reduction. NO_3/NH_4 fluctuated widely during the entire study period, but the value was generally lowest at TM3 and highest at TM8 (Fig. 3). In contrast, TIN/PO_4 and TIN/SiO_2 showed much clearer increasing and decreasing trends respectively (Fig. 3). TIN/PO_4 decreased gradually from between 1986 and 1996, but have returned to approximately the 16:1 Redfield N:P ratio after declines in nutrient concentrations. Declines in nutrient concentrations reduced the among-station differences in TIN/SiO_2 and the values at all stations were well below the 1:1 Redfield ratio for N:Si.

(Insert Fig. 2 and Fig. 3 here)

Chlorophyll-*a* concentrations and densities of different phytoplankton groups exhibited varying trends temporally (Figs. 4 and 5 respectively). The lack of apparent

184 similar trends between chlorophyll-*a* concentrations and phytoplankton densities may
185 be due to the fact that chlorophyll-*a* data were depth-averaged, while phytoplankton
186 data were from the surface only. In general, the concentrations of chlorophyll-*a* and
187 densities of all phytoplankton groups were highest at TM3 and lowest at TM8.
188 Dinoflagellate density at TM3 increased to two major peaks of $> 10,000$ cells ml^{-1} in
189 1996 and 1998, and decreased to $< 1,000$ cells ml^{-1} in recent years. Other algal groups
190 occurred at densities of < 5000 cells ml^{-1} before 1996, but increased rapidly and
191 considerably after 1996. Density at TM3 remained $> 6,000$ cells ml^{-1} after declines in
192 nutrient concentrations. As diatoms were the most abundant group, the density of total
193 phytoplankton exhibited temporal and spatial patterns more similar to those of
194 diatoms than those of dinoflagellates and other algal groups. Declines in nutrient
195 concentrations did not affect the density of dinoflagellates significantly, but densities
196 of diatoms, other algal groups and total phytoplankton all showed significant increase
197 after declines in nutrient concentrations. Despite a significant increase in total
198 phytoplankton density, the number of algal blooms recorded in Tolo Harbour dropped
199 significantly from a mean of 15.8 before nutrient loading reduction to 7.6 (Paired
200 t-test, $p = 0.02$) afterwards. No specific phytoplankton group or genus identified in the
201 study could be concluded to be restricted to a particular station. Diatom genera
202 common in all four stations and with average densities $> 1,000$ cells ml^{-1} in TM3 and

> 600 ml⁻¹ in TM8 included *Cerataulina*, *Chaetoceros*, *Pseudo-nitzschia*, *Leptocylindrus* and *Thalassiosira*. *Prorocentrum* and *Scrippsiella* were the most abundant dinoflagellate genera in all stations. Average density was > 1,000 cells ml⁻¹ in TM3 and > 250 cells ml⁻¹ in TM8 for *Prorocentrum*, and > 700 cells ml⁻¹ in TM3 and > 100 cells ml⁻¹ in TM8 for *Scrippsiella*. The dinoflagellates *Gymnodinium*, *Gyrodinium* and *Protoperidinium* were also frequently found at all stations, but their average densities were much lower. Small flagellates represented the most abundant and frequently recorded other algal group in the samples.

(Insert Fig. 4 and Fig. 5 here)

Hong Kong is characterized by a warm wet season from April to September, and a relatively dry and cool season for the rest of the year. However, when plots of seasonal means of nutrient concentrations and phytoplankton densities were made, no apparent and general trend could be identified (Data not shown).

Kendall's correlation did not reveal any strong or obvious trends between phytoplankton variables and various parameters (Table 2). Results for TM3 and TM4 seemed to oppose those for TM6 and TM8 at times. For instance, correlation between phytoplankton variables and DO and BOD₅ were usually positive at TM3 and TM4,

but negative at TM6 and TM8, and vice versa with the results for SiO₂. It is also not uncommon to find contradicting results between phytoplankton cell densities and chlorophyll-*a* concentrations due to the different trends exhibited by these two parameters. Correlation between phytoplankton variables and nutrient variables were mostly negative in all stations except TM6. Most phytoplankton variables at TM3 and TM4 correlated significantly and negatively with nutrient concentrations. In TM6, chlorophyll-*a* concentrations correlated negatively with NO₃-N and NH₄-N concentrations, while dinoflagellate densities correlated positively with total N, PO₄-P and total P concentrations. NO₃/NH₄ correlated negatively with the densities of all groups of phytoplankton at TM3, and positively with diatom densities at TM6 and TM8. TIN/PO₄ correlated positively with other algal group densities in all stations. TIN/SiO₂ correlated negatively with phytoplankton densities in all stations, but exhibited especially strong negative correlations with the densities of dinoflagellates, other algal groups and total phytoplankton at TM3 and TM4. Kendall's correlations between the number of algal blooms and all parameters were also determined. Surprisingly, the number of algal blooms did not correlate significantly with phytoplankton densities, nutrient concentrations and nutrient ratios. It correlated weakly with pH ($r = 0.313$) and bottom DO ($r = 0.305$) ($p < 0.05$), and strongly with BOD₅ ($r = 0.513$) and chlorophyll-*a* concentration ($r = 0.417$) ($p < 0.01$).

(Insert Table 2 here)

4. Discussion

4.1. Temporal and spatial variations in marine water quality and phytoplankton densities

While spatially there is an apparent pattern of better water quality in TM6 and TM8, all parameters show high temporal variations and do not exhibit an apparent trend except changes due to nutrient loading reduction measures. Our data set is unable to explain the cause of specific events such as the relatively lower nutrient concentrations during 1989 (Fig. 2), and the rise in phytoplankton densities in 1995 (Fig. 5), which may be due to isolated algal bloom events that inflated the annual mean value. Both of these parameters may also be affected by factors such as water current, weather conditions and anthropogenic activities that are not available in our data set.

Changes in various physico-chemical and biological parameters, especially nutrient concentrations, after the implementation of the Tolo Harbour Action Plan

suggested that nutrient loading reduction measures had led to water quality improvement in Tolo Harbour. While Hu et al. (2001) warned that recovery after nutrient loading reduction in Tolo Harbour may be slowed due to nutrients released into the water column from contaminated sediments, N and P concentrations appeared to decrease quite rapidly (Fig. 2) after the full implementation of the Tolo Harbour Effluent Export scheme. Surprisingly, although chlorophyll-*a* concentrations had decreased after declines in nutrient concentrations, the densities of both diatoms and other algal groups have increased. This was similar to the observation reported earlier by Yung et al. (1997) that there was an increase in the density of phytoplankton, especially that of other algal groups, when chlorophyll-*a* concentrations had not increased. As other algal groups in both Yung et al. (1997) and this study included mainly unidentified small flagellates and cryptomonads, an increase in the abundance of these small organisms may not be accompanied by an increase of chlorophyll-*a* concentrations. This is somewhat supported by the fact that correlation between density of other algal group and chlorophyll-*a* concentrations was recorded only at TM8, while diatoms and dinoflagellates correlated strongly and positively with chlorophyll-*a* concentrations at all stations.

Water quality measured in terms of nutrients and chlorophyll-*a* concentration has always been better in the channel region than in the inner part of the harbour (Hu et al.,

2001; Lam and Ho, 1989; Yung et al., 1997), and a seaward decrease in nutrient concentrations from the inner harbour to the channel was evident from our results. A similar spatial pattern, with water quality improving from TM3 to TM8, is recorded in other water quality indicators such as secchi depth, DO and BOD₅. Due to its proximity to the open sea and the much lower pollution levels, TM8 was less affected by eutrophication and subsequent declines in nutrient concentrations compared to stations in the inner part of the harbour. An interesting observation was that the mean values of NO₃/NH₄ decreased at TM3, TM4 and TM6, but increased at TM8 after declines in nutrient concentrations. This may be caused by a relatively higher decline of NH₄-N concentrations than that of NO₃-N in TM8. The maximum release rate of NH₄-N from contaminated sediments into the water column in Tolo Harbour was found to be more than sixfolds higher than that of NO₃-N (Hu et al. 2001), and although Hu et al. (2001) did not specify specific nutrient release rates in various parts of Tolo Harbour, it may be possible that comparatively less NH₄-N was released back into the water column from sediments in the less polluted outer parts of the Tolo Harbour.

Spatial differences in nutrient loading and tidal flushing may have led to the observed differences in how phytoplankton in different stations had responded to declines in nutrient concentrations. Correlation between phytoplankton densities and

nutrient concentrations was poor and usually insignificant at TM6 and TM8 in the outer part of the harbour. In contrast, phytoplankton densities at TM3 and TM4 tended to correlate strongly and negatively with nutrient concentrations, suggesting that nutrients were removed rapidly by phytoplankton in the inner part of Tolo Harbour.

4.2. Changes in nutrient ratios and shifts in the phytoplankton community

Since it is believed that some phytoplankton prefer to use NH_4 over NO_3 (Dortch 1990), a decrease in NO_3/NH_4 may favor phytoplankton growth. However, the lack of consistent correlation between NO_3/NH_4 and phytoplankton density suggests that the phytoplankton community in Tolo Harbour did not have specific preference for either NO_3 or NH_4 .

Decreases in nutrient concentrations following nutrient loading reduction measures had altered the values of TIN/PO_4 and TIN/SiO_2 significantly. Hodgkiss and Ho (1997) related the increased abundance of dinoflagellates in Tolo Harbour to a decrease in N:P ratios. It must be noted that no evidence of a decrease in dinoflagellate density has been detected in our data, despite a significant increase in TIN/PO_4 after declines in nutrient concentrations. In general, the density of dinoflagellates correlate neither with TIN/PO_4 nor with most physico-chemical

parameters other than Secchi depth and BOD₅, both of which may be dependent variables affected by the dinoflagellate density. Hence, the density of dinoflagellate in Tolo Harbour may not be easily predicted from nutrient concentrations or N:P ratios alone. While our analysis has not established a strong link between nutrient concentrations and dinoflagellate densities, several investigators have reported an increasing dominance of dinoflagellates during the 1980s and early 1990s when nutrient concentrations in Tolo Harbour were at their highest (Chan and Hodgkiss, 1987; Hodgkiss and Chan, 1983; Hodgkiss and Ho, 1997; Lam and Ho, 1989). The density of dinoflagellates was at a peak in 1996 when TIN/PO₄ was very low (Figs. 3 & 4). The density of dinoflagellates reached another peak in 1998, due primarily to an extremely dense population of *Prorocentrum* (122,467 cells ml⁻¹) at TM3 in April. But dinoflagellate densities have remained relatively low since 2005, while TIN/PO₄ increased. Hodgkiss and Ho (1997) also noted a decrease in N:P ratios and an increase in algal blooms from 1981–1990. Our data showed no significant correlation between these two variables, although an increase in TIN/PO₄ and a decrease in the number of algal blooms were found after declines in nutrient concentrations. Nevertheless, while our data did not contain information about *Noctiluca* blooms, Liu and Wong (2006) showed that *Noctiluca scintillans* were still common in Tolo Harbour after declines in nutrient concentrations, and *Noctiluca* blooms still persisted.

Compared to the decrease in N concentrations, the decrease in SiO₂ concentrations was less dramatic. Although a significant decrease in depth-averaged SiO₂ concentrations was recorded after declines in nutrient input ($p = 0.045$), surface SiO₂ concentrations did not changed significantly ($p = 0.849$). Overall, TIN/SiO₂ tended to decrease at all sampling stations. Diatoms have a physiological N:Si ratio of 1:1 (Turner, 2002), and an environmental ratio of > 1 has been reported to limit diatom growth and shift a diatom-dominated community to one dominated by non-diatoms, particularly flagellates (Cloern, 2001; Egge and Aksnes, 1992; Paul et al., 2008; Turner et al. 1998). While diatom densities did correlate weakly with TIN/SiO₂ in TM3 and TM4, diatoms were the most abundant algal group in these two stations and Tolo Harbour (Lam and Ho, 1989; Yung et al., 1997) before declines in nutrient concentrations, when TIN/SiO₂ at TM3 and TM4 were frequently > 1 . This suggests that although N:Si ratios may have some effect on diatom densities, it is not enough to shift diatom dominance in Tolo Harbour. In addition, the Si concentrations in Tolo Harbour may not be limiting to diatom growth. Egge and Aksnes (1992) reported that a Si concentration of $> 2 \mu\text{M}$ was enough to induce consistent dominance by diatoms in Norway. While this low threshold concentration may not apply in the warm subtropical waters of Hong Kong, the lowest SiO₂ concentration in this study was $\sim 10 \mu\text{M}$ at TM8, fivefolds higher than the threshold concentration

reported in Norway, and may be sufficient to support diatom growth regardless of
TIN/SiO₂. It should be noted that the maintenance of dominance by diatoms may not
necessarily be beneficial both ecologically and economically. Although centric
diatoms have been suggested to be the more desirable phytoplankton group when
compared to non-diatom groups, such as dinoflagellates, due to their higher nutritional
value and low toxicity (Hecky and Kilham, 1988; Officer and Ryther, 1980), pennate
diatoms, including those belonging to the genus *Pseudo-nitzschia*, are also common in
Tolo Harbour (Lam and Ho, 1989). Some species of *Pseudo-nitzschia* have been
known to release the toxin domoic acid (Lefebvre et al., 2002; Sekula-Wood et al.,
2009), and a connection between *Pseudo-nitzschia* blooms and eutrophication has
been proposed (Parsons and Dortch, 2002; Schnetzer et al., 2007). Fortunately,
Pseudo-nitzschia spp. may prefer high nitrate waters (Paul et al., 2008; Carter et al.,
2005), so declines in nutrient concentrations in Tolo Harbour should discourage
Pseudo-nitzschia blooms.

Although our data does not contain information on the size structure of the
phytoplankton community, changes in nutrient concentration or ratios may cause a
shift from larger phytoplankton species or cells to smaller ones. Stolte and Riegman
(1995) had theorized that eutrophication favors large phytoplankton species because
their larger storage capacity allows them to better compete under high nutrient

374 concentrations. Indeed, a shift towards an eutrophic environment has been shown to
375 favor large diatoms in the western Wadden Sea (Stolte et al., 1994; Philippart et al.,
376 2000), and Qiu et al. (2010) reported strong negative correlations between $\leq 3 \mu\text{m}$
377 chlorophyll-*a* and nutrient concentrations. The half-saturation constant for P uptake
378 has been reported to be as high as 4.85–6.67 μM at 20°C for certain large diatom
379 species (Nishikawa et al. 2009 and 2010), much higher than the means of TP in Tolo
380 Harbour after nutrient loading reduction (Table 2). Declines in nutrient concentrations
381 in Tolo Harbour may therefore favor smaller cells and species that are more
382 competitive in a less eutrophic condition. Qiu et al. (2010) reported significant
383 positive correlation between $< 20 \mu\text{m}$ chlorophyll *a* concentrations and N/P ratios.
384 Our data also show a strong and positive correlation between TIN/PO_4 and other algal
385 group densities, caused mainly by an increase in the density of small flagellates and
386 *Cryptomonas*. Recent studies in Tolo Harbour have revealed that cells $< 5 \mu\text{m}$
387 accounted for $> 25\%$ of the chlorophyll-*a* biomass (Lie and Wong, 2010; Wong et al.,
388 2010). In addition, phytoplankton size structures in many aquatic ecosystems have
389 also been known to be affected by climate change, particularly by increases in water
390 column stratification due to increased temperature (Finkel, 2007; Smol et al., 2005;
391 Winder et al., 2010). If indeed such a shift to smaller phytoplankton is occurring in
392 Tolo Harbour, it would help to explain the increase in phytoplankton density as

chlorophyll-*a* concentrations decrease.

4.3. Comparison with the recovery from eutrophication of other ecosystems

The effects of declines in nutrient concentrations had been intensively studied in freshwater lakes (e.g. Jeppesen et al., 2002; Jeppesen et al., 2005; Søndergaard et al., 2002), and most have reported improvements in water quality. Water quality improvement is mainly achieved through the decrease in phytoplankton biomass. This decrease can occur as a direct response to declines in nutrient concentrations, but it can be further facilitated through increased grazing pressure from zooplankton grazers (Edmondson, 1994) and the recolonization of submerged macrophytes, which has been shown to substantially affect the success rates of water quality improvement in lakes (Mehner et al., 2002). While there may be differences between freshwater and marine systems in their responses to declines in nutrient concentrations (Cloern, 2001; Smith et al., 1999), examples of dramatic improvements in once eutrophicated marine ecosystems such as Hillsborough Bay, Florida (Johansson and Lewis, 1992) and the coastal waters of Denmark (Rask et al., 1999) had both involved the recolonization of macrophytes such as seagrasses (Hillsborough Bay) and eelgrass (Denmark). However, an early study in Tolo Harbour has shown that most of the harbour was

covered by extremely fine sediments that support little plant life (Trott, 1972).

Therefore, even with the decline of nutrient concentrations and increase in water

transparency, the spread of macrophytes in Tolo Harbour is unlikely.

In regards to the zooplankton grazers in Tolo Harbour, a study comparing the copepod community before and after declines in nutrient concentrations in Tolo Harbour revealed a decrease in copepod density, from 6,392 ind. m⁻³ in 1987–1991, to 4,810 ind. m⁻³ in 2003–2004 (Wong et al., 1993; Zhang and Wong, submitted). The mesozooplankton community in Tolo Harbour was found to be heavily preyed upon by larval and juvenile stages of the black seabream, *Acanthopagrus schlegeli*, and the Japanese seaperch, *Lateolabrax japonicus*, which use Tolo Harbour as a nursery habitat (Nip et al. 2003). While presently there is no detailed description on the fish population in Tolo Harbour, the abundance of small planktivorous fishes will most likely increase when increased fishing pressure from overfishing of larger piscivorous fishes reduces the predation pressure on the small fishes. Such changes will in turn increase the predation pressure on mesozooplankton grazers, reducing their abundance and delaying any further decrease of phytoplankton abundance.

Although every water mass may have different responses to declines in nutrient concentrations (Carvalho et al., 1995; Smith et al., 1999), it is obvious that trophic interactions play a significant role in determining the outcome (Carpenter et al., 1995,

Manca and Ruggiu, 1998; Proulx et al., 1996; Smith et al., 1999). Increase in macrophytes and zooplankton grazers can further facilitate the decrease in phytoplankton biomass (Edmondson, 1994; Mehner et al., 2002), but the presence of planktivores and piscivores can affect the abundance of zooplankton grazers. Therefore, an appropriate balance between piscivores, planktivores and zooplankton grazers should be maintained to ensure the long-term stability of the recovered ecosystem (Mehner et al. 2002). Unfortunately, there is a lack of long-term monitoring on other communities, such as zooplankton and nekton, in Tolo Harbour. As there is insufficient information on the responses of various communities to changes in nutrient loading, the phytoplankton community alone is not a reliable indicator of the trophic status of the whole ecosystem (Philippart et al. 2007). Future efforts in water quality monitoring and eutrophic coastal restoration should therefore focus not only on nutrient concentrations and phytoplankton biomass and densities, but also other important components of the aquatic ecosystem such as zooplankton and nekton communities.

Acknowledgements

The authors would like to express their thanks to the staff of the Hong Kong Environmental Protection Department and the Government Chemists for collecting

and analyzing the samples. We are also grateful to the Director of Environmental Protection for his permission to use the data. The opinions in this paper are those of the authors and do not necessarily reflect the views or policies of the Hong Kong Government. Suggestions and comments given by two anonymous reviewers are greatly appreciated.

References

- American Public Health Association, American Water Works Association, Water Environment Federation, 1998. Standard methods for the examination of water and wastewater, 20th Edition. American Public Health Association, Washington DC.
- Anderson, D.M., Burkholder, J.M., Cochlan, W.P., Glibert, P.M., Gobler, C.J., Heil, C.A., Kudela, R.M., Parsons, M.L., Rensel, J.E.J., Townsend, D.W., Trainer, V.L., Vargo, G.A., 2008. Harmful algal blooms and eutrophication: Examining linkages from selected coastal regions of the United States. *Harmful Algae* 8, 39-53.
- Battarbee, R.W., Anderson, N.J., Jeppesen, E., Leavitt, P.R., 2005. Combining palaeolimnological and limnological approaches in assessing lake ecosystem

- 469 response to nutrient reduction. *Freshwater Biology* 50, 1772-1780.
- 470 Carpenter, S.R., Christensen, D.L., Cole, J.J., Cottingham, K.L., He, X., Hodgson,
471 J.R., Kitchell, J.F., Knight, S.E., Pace, M.L., Post, D.M., Schindler, D.E.,
472 Voichick, N., 1995. Biological control of eutrophication. *Environmental Science*
473 *and Technology* 29, 784-786.
- 474 Carpenter, S.R., Ludwig, D., Brock, W.A., 1999. Management of eutrophication for
475 lakes subject to potentially irreversible change. *Ecological Applications* 9,
476 751-771.
- 477 Carter, C.M., Ross, A.H., Schiel, D.R., Howard-Williams, C., Hayden, B., 2005. *In*
478 *situ* microcosm experiments on the influence of nitrate and light on
479 phytoplankton community composition. *Journal of Experimental Marine*
480 *Biology and Ecology* 326, 1-13.
- 481 Carvalho, L., Beklioglu, M., Moss, B., 1995. Changes in a deep lake following
482 sewage diversion - a challenge to the orthodoxy of external phosphorus control
483 as a restoration strategy? *Freshwater Biology* 34, 399-410.
- 484 Carvalho, L., Kirika, A., 2003. Changes in shallow lake functioning: response to
485 climate change and nutrient reduction. *Hydrobiologia* 506-509, 789-796.
- 486 Chau, K.W., 2007. Integrated water quality management in Tolo Harbour, Hong Kong:
487 a case study. *Journal of Cleaner Production* 15, 1568-1572.

- 488 Chau, Y.K., Abesser, R.A., 1958. A preliminary study of the hydrology of Hong Kong
489 Territorial Waters. Hong Kong University Fisheries Journal 2, 43-58.
- 490 Cloern, J.E., 2001. Our evolving conceptual model of the coastal eutrophication
491 problem. Marine Ecology Progress Series 210, 223-253.
- 492 de Benardi, R., Calderoni, A., Mosello, R., 1996. Environmental problems in Italian
493 lakes, and lakes Maggiore and Orta as successful examples of correct
494 management leading to restoration. Verhandlungen der Internationale
495 Vereinigung für Theoretische und Angewandte Limnologie 26, 123-138.
- 496 Dortch, Q., 1990. The interaction between ammonium and nitrate uptake in
497 phytoplankton. Marine Ecology Progress Series 61, 183-201.
- 498 Edmondson, W.T., 1994. Sixty years of Lake Washington: a curriculum vitae. Lake
499 Reservoir Management 10, 75-84.
- 500 Egge, J.K., Aksnes, D.L., 1992. Silicate as regulating nutrient in phytoplankton
501 competition. Marine Ecology Progress Series 83, 281-289.
- 502 Finkel, Z.V., 2007. Does size matter? The evolution of modern marine food webs. In:
503 Falkowski, P.G., Knoll, A.H. (Eds.), The evolution of aquatic photoautotrophs.
504 Academic Press, New York. Pp. 333-350.
- 505 Hansen, P.J., Miranda, L., Azanza, R., 2004. Green *Noctiluca scintillans*: a

- 506 dinoflagellate with its own greenhouse. Marine Ecology Progress Series 275,
507 79-87.
- 508 Hecky, R.E., Kilham, P., 1988. Nutrient limitation of phytoplankton in freshwater and
509 marine environments: A review of recent evidence on the effects of enrichment.
510 Limnology & Oceanography 33, 796-822.
- 511 Hodgkiss, I.J., Chan, B.S.S., 1983. Pollution studies in Tolo Harbour, Hong Kong.
512 Marine Environmental Research 10, 1-44.
- 513 Hodgkiss, I.J., Ho, K.C., 1997. Are changes in N:P ratios in coastal waters the key to
514 increased red tide blooms? Hydrobiologia 352, 141-147.
- 515 Hodgkiss, I.J., Yim, W.W., 1995. A case study of Tolo Harbour, Hong Kong. In:
516 McComb, A.J. (Ed.) Eutrophic shallow estuaries and lagoons. CRC Press,
517 Florida. Pp. 41-57.
- 518 Hong Kong Environmental Protection Department, 2009. Marine Water Quality in
519 Hong Kong in 2008. The Government of the Hong Kong Special Administrative
520 Region, Hong Kong.
- 521 Hong Kong Environmental Protection Department, 1991. Marine Water Quality in
522 Hong Kong in 1990. The Government of the Hong Kong Special Administrative
523 Region, Hong Kong.

- 524 Hu, W.F., Lo, W., Chua, H., Sin, S.N., Yu, P.H.F., 2001. Nutrient release and sediment
525 oxygen demand in a eutrophic land-locked embayment in Hong Kong.
526 *Environment International* 26, 369-375.
- 527 Jeppesen, E., Jensen, J.P., Søndergaard, M., 2002. Response of phytoplankton,
528 zooplankton and fish to re-oligotrophication: an 11-year study of 23 Danish
529 lakes. *Aquatic Ecosystem Health and Management* 5, 31-43.
- 530 Jeppesen, E., Meerhoff, M., Jacobsen, B.A., Hansen, R.S., Søndergaard, M., Jensen,
531 J.P., Lauridsen, T.L., Mazzeo, N., Branco, C.W.C., 2007. Restoration of shallow
532 lakes by nutrient control and biomanipulation – the successful strategy varies
533 with lake size and climate. *Hydrobiologia* 581, 269-285.
- 534 Jeppesen, E., Søndergaard, M., Jensen, J.P., Havens, K.E., Anneville, O., Carvalho, L.,
535 Coveney, M.F., Deneke, R., Dokulil, M.T., Foy, B., Gerdeaux, D., Hampton,
536 S.E., Hilt, S., Kangur, K., Köhler, J., Lammens, E.H.H.R., Lauridsen, T.L.,
537 Manca, M., Miracle, M.R., Moss, B., Nöges, P., Persson, G., Phillips, G.,
538 Portielje, R., Romo, S., Schelske, C.L., Straile, D., Tatrai, I., Willén, E. &
539 Winder, M., 2005. Lake responses to reduced nutrient loading - an analysis of
540 contemporary long-term data from 35 case studies. *Freshwater Biology* 50,
541 1747-1771.
- 542 Johansson, J.O.R., Lewis, R.R., 1992. Recent improvements in water quality and

- 543 biological indicators in Hillsborough Bay, a highly impacted subdivision of
544 Tampa Bay, Florida, USA. In: Volenweider, R.A., Marchetti, R., Viviani, R.
545 (Eds.), Marine coastal eutrophication. The response of marine transitional
546 systems to human impact: Problems and perspectives for restoration. Science of
547 the total environment supplement 1992. Elsevier Science, Amsterdam. Pp.
548 1199-1216.
- 549 Justić, D., Rabalais, N.N., Turner, R.E., 1995. Stoichiometric nutrient balance and
550 origin of coastal eutrophication. Marine Pollution Bulletin 30, 41-46.
- 551 Kiørboe, T., Titelman, J., 1998. Feeding, prey selection and prey encounter
552 mechanisms in the heterotrophic dinoflagellate *Noctiluca scintillans*. Journal of
553 Plankton Research 20, 1615-1636.
- 554 Lam, C.W.Y., Ho, K.C., 1989. Phytoplankton characteristics of Tolo Harbour. Asian
555 Marine Biology 6, 5-18.
- 556 Lee, J.H.W., Harrison, P.J., Kuang, C., Yin, K., 2006. Eutrophication dynamics in
557 Hong kong Coastal Waters: Physical and Biological Interactions. In Wolanski, E.
558 (Ed.), Environment in Asia Pacific Harbours. Springer, the Netherlands. Pp.
559 187-206.
- 560 Lefebvre, K.A., Silver, M.W., Coale, S.L., Tjeerdema, R.S., 2002. Domoic acid in
561 planktivorous fish in relation to toxic *Pseudo-nitzschia* cell densities. Marine

- 562 Biology 140, 625-631.
- 563 Lie, A.A.Y., Wong, C.K. 2010. Selectivity and grazing impact of
564 microzooplankton on phytoplankton in two subtropical semi-enclosed
565 bays with different chlorophyll concentrations. *Journal of Experimental*
566 *Marine Biology and Ecology*, in press.
- 567 Liu, X., Wong, C.K., 2006. Seasonal and spatial dynamics of *Noctiluca scintillans* in a
568 semi-enclosed bay in the northeastern part of Hong Kong. *Botanica Marina* 49,
569 145-150.
- 570 Manca, M., Ruggiu, D., 1998. Consequences of pelagic food-web changes during a
571 long-term lake oligotrophication process. *Limnology & Oceanography* 43,
572 1368-1373.
- 573 Mehner, T., Benndorf, J., Kasprzak, P., Koschel, R., 2002. Biomanipulation of lake
574 ecosystems: successful applications and expanding complexity in the underlying
575 science. *Freshwater Biology* 47, 2453-5465.
- 576 Mehner, T., Kasprzak, P., Wysujack, K., Laude, U., Koschel, R., 2001. Restoration of
577 a stratified lake (Feldberger Haussee, Germany) by a combination of nutrient
578 load reduction and long-term biomanipulation. *International Review of*
579 *Hydrobiology* 2, 253-265.
- 580 Nip, T.H.M., Ho, W., Wong, C.K., 2003. Feeding ecology of larval and juvenile black

- 581 seabream (*Acanthopagrus schlegeli*) and Japanese seaperch (*Lateolabrax*
 582 *japonicus*) in Tolo Harbour, Hong Kong.
- 583 Nishikawa, T., Tarutani, K., Yamamoto, T., 2009. Nitrate and phosphate uptake
 584 kinetics of the harmful diatom *Eucampia zodiacus* Ehrenberg, a causative
 585 organism in the bleaching of aquacultured *Porphyra* thalli. Harmful Algae 8,
 586 513-517.
- 587 Nishikawa, T., Tarutani, K., Yamamoto, T., 2010. Nitrate and phosphate uptake
 588 kinetics of harmful diatom *Coscinodiscus wailesii*, a causative organism in the
 589 bleaching of aquacultured *Porphyra* thalli. Harmful Algae 9, 563-567.
- 590 Nixon, S. W., 1995. Coastal marine eutrophication: A definition, social causes, and
 591 future concerns. Ophelia 41, 199-219.
- 592 Officer, C.B., Ryther, J.H., 1980. The possible importance of silicon in marine
 593 eutrophication. Marine Ecology Progress Series 3, 83-91.
- 594 Parsons, M.L., Dortch, Q., 2002. Sedimentological evidence of an increase in
 595 *Pseudo-nitzschia* (Bacillariophyceae) abundance in response to coastal
 596 eutrophication. Limnology & Oceanography 47, 551-558.
- 597 Paul, J.T., Ramaiah, N., Sardesai, S., 2008. Nutrient regimes and their effect on
 598 distribution of phytoplankton in the Bay of Bengal. Marine Environmental
 599 Research 66, 337-344.

- 600 Philippart, C.J.M., Beukema, J.J., Cadée, G.C., Dekker, R., Goedhart, P.W., van Iperen,
601 J.M., Leopold, M.F., Herman, P.M.J., 2007. Impacts of nutrient reduction on
602 coastal communities. *Ecosystems* 10, 95-118.
- 603 Philippart, C.J.M., Cadée, G.C., van Raaphorst, W., Riegman, R., 2000. Long-term
604 phytoplankton-nutrient budgets, and denitrification potential. *Limnology &*
605 *Oceanography* 45, 131-144.
- 606 Proulx, M., Pick, F.R., Mazumder, A., Hamilton, P.B., Lean, D.R.S., 1996. Effects of
607 nutrients and planktivorous fish on the phytoplankton of shallow and deep
608 aquatic ecosystems. *Canadian Journal of Fisheries and Aquatic Sciences* 77,
609 1556-1572.
- 610 Qiu, D., Huang, L., Zhang, J., Lin, S., 2010. Phytoplankton dynamics in and near the
611 highly eutrophic Pearl River Estuary, South China Sea. *Continental Shelf*
612 *Research* 30, 177-186.
- 613 Rask, N., Pedersen, S.T., Jensen, M.H., 1999. Response to lowered nutrient discharges
614 in the coastal waters around the island of Funen, Denmark. *Hydrobiologia* 363,
615 69-81.
- 616 Reckhow, K.H., Chapra, S.C., 1999. Persistent organic pollutants (POPs): State of the
617 science. *Environmental Pollution* 100, 209-221.
- 618 Sekula-Wood, E., Schnetzer, A., Benitez-Nelson, C.R., Anderson, C., Berelson, W.M.,

- 619 Brzezinski, M.A., Burns, J.M., Caron, D.A., Cetinic, I., Ferry, J.L., Fitzpatrick,
620 E., Jones, B.H., Miller, P.E., Morton, S.L., R.A., Siegel, D.A., Thunell, R., 2009.
621 Rapid downward transport of the neurotoxin domoic acid in coastal waters.
622 Nature Geoscience 2, 272-275.
- 623 Schnetzer, A., Miller, P.E., Schaffner, R.A., Stauffer, B.A., Jones, B.H., Weisberg,
624 S.B., DiGiacomo, P.M., Berelson, W.M., Caron, D., 2007. Blooms of
625 *Pseudo-nitzschia* and domoic acid in the San Pedro Channel and Los Angeles
626 harbor areas of the Southern California Bight, 2003-2004. Harmful Algae 6,
627 372-387.
- 628 Smith, V.H., 1983. Low nitrogen to phosphorus ratios favor dominance by blue-green
629 algae in lake phytoplankton. Science 221, 669-671.
- 630 Smith, V.H., 1998. Cultural eutrophication of inland, estuarine, and coastal waters. In:
631 Pace, M.L., Groffman, P.M. (Eds.) Successes, limitations and frontiers in
632 ecosystem science. Springer, New York. Pp. 7-49.
- 633 Smith, V.H., 2003. Eutrophication of freshwater and coastal marine ecosystems. A
634 global problem. Environmental Science and Pollution Research 10, 126-139.
- 635 Smith, V.H., Tilman, G.D., Nekola, J.C. 1999. Eutrophication: impacts of excess
636 nutrient inputs on freshwater, marine, and terrestrial ecosystems. Environmental
637 Pollution 100, 179-196.

- 638 Smol, J.P., Wolfe, A.P., Birks, H.J.B., Douglas, M.S.V., Jones, V.J., Korhola, A.,
 639 Pientzl, R., Rühland, K., Sorvari, S., Antoniades, D., Brooks, S.J., Fallu, M.A.,
 640 Hughes, M., Keatley, B.E., Laing, T.E., Michelutti, N., Nazarova, L., Nyman,
 641 M., Paterson, A.M., Perren, B., Quinlan, R., Rautio, M., Saulnier-Talbot, E.,
 642 Siltone, S., Solovieva, N., Weckström, J., 2005. Climate-drive regime shifts in
 643 the biological communities of arctic lakes. *Proceedings of the National*
 644 *Academy of Sciences* 102, 4397-4402.
- 645 Søndergaard, M., Jensen, J.P., Jeppesen, E., Møller, P.H., 2002. Seasonal dynamics in
 646 the concentrations and retention of phosphorus in shallow Danish lakes after
 647 reduced loading. *Aquatic Ecosystem Health and Management* 5, 19-23.
- 648 Sriwong, R., Pholpunthin, P., Lirdwitayaprasit, T., Kishino, Furuya, K., 2008.
 649 Population dynamics of green *Noctiluca scintillans* (Dinophyceae) associated
 650 with the monsoon cycle in the upper gulf of Thailand. *Journal of Phycology* 44,
 651 605-615.
- 652 Stolte, W., McCollin, T., Noordeloos, A.A.M., Riegman, R., 1994. Effect of nitrogen
 653 source on the size distribution within marine phytoplankton populations. *Journal*
 654 *of Experimental Marine Biology and Ecology* 184, 83-97.
- 655 Stolte, W., Riegman, R., 1995. Effect of phytoplankton cell size on transient-state
 656 nitrate and ammonium uptake kinetics. *Microbiology* 141, 1221-1229.

- 657 Sweeney, B.M., 1971. Laboratory studies of a green *Noctiluca* from New Guinea.
658 Journal of Phycology 7, 53-58.
- 659 Trott, L.B., 1972. Marine ecology in Tolo Harbour, Hogn Kong. Chung Chi Journal 11,
660 26-32.
- 661 Turner, R.E., 2002. Element ratios and aquatic food webs. Estuaries 25, 694-703.
- 662 Turner, R.E, Qureshi, N., Rabalais, N.N., Dortch, Q., Justić, D., Shaw, R.F., Cope, J.,
663 1998. Fluctuating silicate:nitrate ratios and coastal plankton food webs.
664 Proceedings of the National Academy of Sciences 95, 13048-13051.
- 665 Wear, R.G., Thompson, G.B., Stirling, H.P., 1984. Hydrography, nutrients and
666 plankton in Tolo Harbour, Hong Kong. Asian Marine Biology 1, 59-75.
- 667 Winder, M., Reuter, J.E., Schladow, S.G., 2009. Lake warming favors small-sized
668 planktonic diatom species. Proceedings of the Royal Society B 276, 427-435.
- 669 Wong, C.K., Chan , A.L.C., Chen, Q.C., 1993. Planktonic copepods of Tolo Harbour,
670 Hong Kong. Crustaceana 64, 76-84.
- 671 Wong, C.K., Tang, C.C.H., Lie, A.A.Y., Lam, J.Y.C., Yung, Y.K. 2010. Size structure
672 and taxonomic composition of phytoplankton in Hong Kong waters. In: Tow,
673 T.T., Yusup, Y., Fizri, F.F.A. (Eds.) Proceedings International Conference on
674 Environmental Research and Technology (ICERT 2010). Universiti Sains
675 Malaysia, Malaysia. Pp. 560-564.

676 Wu, R.S.S., 1988. Marine pollution in Hong Kong: a review. *Asian Marine Biology*
677 5:1-23.

678 Yung, Y.K., Wong, C.K., Broom, M.J., Ogden, J.A., Chan, S.C.M., Leung, Y., 1997.
679 Long-term changes in hydrography, nutrients and phytoplankton in Tolo
680 Harbour, Hong Kong. *Hydrobiologia* 352, 107-115.

681 Zhang, G., Wong, C.K. Changes in the planktonic copepod community in a landlocked
682 bay in the subtropical coastal waters of Hong Kong during recovery from
683 eutrophication. *Hydrobiologia* submitted.

684

685 **Figure Legends**

686 **Fig. 1.** A map of Hong Kong and Tolo Harbour showing the location of the four
687 sampling stations.

688

689 **Fig. 2.** Temporal variations in the annual means of total inorganic nitrogen, total
690 nitrogen, total phosphorus and SiO_2 concentration at the four sampling stations from
691 1986–2008. The shaded area represents the period before nutrient loading reduction .

692

693 **Fig. 3.** Temporal variations in the annual means of NO_3/NH_4 , TIN/PO_4 and TIN/SiO_2
694 at the four sampling stations from 1986–2008. The shaded area represents the period

before nutrient loading reduction. The dashed lines indicate the Redfield N:P and N:Si ratios of 16:1 and 1:1 respectively.

Fig. 4. Temporal variations in the annual means of chlorophyll *a* at the four sampling stations from 1986–2008. The shaded area represents the period before nutrient loading reduction.

Fig. 5. Temporal variations in the annual means of diatoms, dinoflagellates, other algal groups and total phytoplankton densities at the four sampling stations from 1986–2008. The shaded area represents the period before nutrient loading reduction. Monitoring began in 1988 for TM3 and TM4, 1994 for TM6 and 1986 for TM8.

Research highlights

Our study investigates the long-term changes in physico-chemical parameters, including nutrient concentration, nutrient ratios, and phytoplankton before and after nutrient reduction in an eutrophic bay in Hong Kong. Such study with long-term data relating environmental factors and phytoplankton is rather uncommon in marine ecosystems. We believe that such study of how anthropogenic causes led to changes in the phytoplankton community fits the aims and scope of your journal.

Table 1. Mean values ($\pm$ standard deviation) of various physico-chemical, biological and nutrient ratios. “All stations” refers to the mean of all four sampling stations. “86–97” indicates the period before nutrient loading reduction (i.e. 1986–1997), and “98–08” indicates the period after nutrient loading reduction (i.e. 1998–2008). “Overall” refers to the entire study period of 1986–2008. Values are depth-averaged values except for “Bottom DO” and “Bottom BOD₅”.

Parameter		Station				
		TM3	TM4	TM6	TM8	All stations
Temperature (°C)	86–97	23.06 $\pm$ 4.33	22.88 $\pm$ 4.32	22.55 $\pm$ 4.14	22.20 $\pm$ 4.02	22.67 $\pm$ 4.19
	98–08	23.93 $\pm$ 4.13	23.68 $\pm$ 4.13	23.36 $\pm$ 4.01	22.92 $\pm$ 3.87	23.47 $\pm$ 4.02
	Overall	23.47 $\pm$ 4.25	23.26 $\pm$ 4.24	22.93 $\pm$ 4.09	22.55 $\pm$ 3.96	23.06 $\pm$ 4.12
pH	86–97	8.31 $\pm$ 0.20	8.29 $\pm$ 0.23	8.26 $\pm$ 0.19	8.24 $\pm$ 0.17	8.27 $\pm$ 0.18
	98–08	8.21 $\pm$ 0.20	8.16 $\pm$ 0.21	8.16 $\pm$ 0.20	8.16 $\pm$ 0.15	8.17 $\pm$ 0.17
	Overall	8.26 $\pm$ 0.20	8.23 $\pm$ 0.23	8.21 $\pm$ 0.20	8.20 $\pm$ 0.17	8.22 $\pm$ 0.19
Salinity	86–97	30.75 $\pm$ 1.55	30.77 $\pm$ 1.66	31.44 $\pm$ 1.51	31.96 $\pm$ 1.06	31.22 $\pm$ 1.39
	98–08	31.18 $\pm$ 1.56	31.29 $\pm$ 1.48	31.78 $\pm$ 1.21	32.31 $\pm$ 1.07	31.64 $\pm$ 1.28
	Overall	30.96 $\pm$ 1.57	31.02 $\pm$ 1.60	31.61 $\pm$ 1.38	32.13 $\pm$ 1.08	31.42 $\pm$ 1.35
DO (mg l ⁻¹)	86–97	6.86 $\pm$ 1.93	6.85 $\pm$ 1.86	6.45 $\pm$ 1.61	6.36 $\pm$ 1.65	6.63 $\pm$ 1.60
	98–08	6.98 $\pm$ 1.37	6.70 $\pm$ 1.33	6.26 $\pm$ 1.33	6.06 $\pm$ 1.34	6.50 $\pm$ 1.22
	Overall	6.92 $\pm$ 1.68	6.78 $\pm$ 1.62	6.36 $\pm$ 1.48	6.22 $\pm$ 1.51	6.57 $\pm$ 1.43
Bottom DO (mg l ⁻¹)	86–97	4.97 $\pm$ 2.44	4.78 $\pm$ 2.42	4.83 $\pm$ 2.34	5.10 $\pm$ 2.56	4.92 $\pm$ 2.18
	98–08	6.30 $\pm$ 1.86	5.86 $\pm$ 1.88	5.30 $\pm$ 1.92	5.19 $\pm$ 2.02	5.66 $\pm$ 1.73
	Overall	5.61 $\pm$ 2.28	5.29 $\pm$ 2.24	5.05 $\pm$ 2.16	5.15 $\pm$ 2.31	5.28 $\pm$ 2.01
BOD ₅ (mg l ⁻¹)	86–97	2.55 $\pm$ 0.99	2.44 $\pm$ 0.89	1.81 $\pm$ 0.72	1.18 $\pm$ 0.60	2.00 $\pm$ 0.65
	98–08	1.98 $\pm$ 0.75	1.73 $\pm$ 0.56	1.38 $\pm$ 0.47	0.89 $\pm$ 0.56	1.49 $\pm$ 0.45
	Overall	2.28 $\pm$ 0.93	2.10 $\pm$ 0.83	1.60 $\pm$ 0.65	1.04 $\pm$ 0.60	1.76 $\pm$ 0.62
Bottom BOD ₅ (mg l ⁻¹)	86–97	1.33 $\pm$ 0.65	1.38 $\pm$ 0.79	0.95 $\pm$ 0.56	0.71 $\pm$ 0.40	1.09 $\pm$ 0.50
	98–08	1.35 $\pm$ 0.66	1.24 $\pm$ 0.60	0.89 $\pm$ 0.56	0.56 $\pm$ 0.42	1.01 $\pm$ 0.41
	Overall	1.34 $\pm$ 0.66	1.32 $\pm$ 0.71	0.92 $\pm$ 0.56	0.63 $\pm$ 0.42	1.05 $\pm$ 0.46

Secchi Depth (m)	86–97	1.83 ± 0.66	1.82 ± 0.67	2.23 ± 0.79	3.49 ± 1.39	2.35 ± 0.63
	98–08	1.96 ± 0.69	2.10 ± 0.67	2.59 ± 0.88	3.41 ± 1.27	2.51 ± 0.69
	Overall	1.89 ± 0.67	1.89 ± 0.67	2.40 ± 0.85	3.45 ± 1.33	2.43 ± 0.66
Suspended solids (mg l ⁻¹)	86–97	3.41 ± 2.62	3.63 ± 3.11	3.03 ± 2.37	3.20 ± 3.20	3.32 ± 2.38
	98–08	3.20 ± 3.97	2.42 ± 1.36	2.07 ± 1.01	2.26 ± 1.39	2.49 ± 1.30
	Overall	3.31 ± 3.33	3.05 ± 2.51	2.57 ± 1.90	2.75 ± 2.54	2.92 ± 1.98
Chl <i>a</i> (µg l ⁻¹)	86–97	9.63 ± 6.87	9.82 ± 7.43	6.25 ± 4.80	3.38 ± 6.21	7.27 ± 4.80
	98–08	9.33 ± 6.28	7.58 ± 4.37	5.20 ± 2.90	3.13 ± 2.17	6.31 ± 3.15
	Overall	9.49 ± 6.58	8.75 ± 6.25	5.75 ± 4.03	3.26 ± 4.72	6.81 ± 4.12
NO ₃ -N (µM)	86–97	3.21 ± 3.17	3.40 ± 3.74	2.29 ± 2.15	1.52 ± 1.43	2.60 ± 2.30
	98–08	1.37 ± 2.36	1.42 ± 2.28	1.27 ± 1.95	1.45 ± 1.75	1.38 ± 1.87
	Overall	2.33 ± 2.96	2.45 ± 3.28	1.80 ± 2.11	1.48 ± 1.59	2.02 ± 2.19
NH ₄ -N (µM)	86–97	10.13 ± 6.37	8.27 ± 5.49	5.61 ± 3.95	3.56 ± 3.24	6.89 ± 4.25
	98–08	4.94 ± 4.39	4.40 ± 4.67	3.26 ± 2.53	2.57 ± 1.76	3.79 ± 3.09
	Overall	7.65 ± 6.08	6.42 ± 5.46	4.48 ± 3.54	3.08 ± 2.68	5.41 ± 4.04
Total inorganic N (µM)	86–97	14.31 ± 8.79	12.65 ± 8.00	8.82 ± 5.27	5.76 ± 3.69	10.38 ± 5.86
	98–08	6.77 ± 6.27	6.37 ± 6.16	5.10 ± 3.64	4.64 ± 2.49	5.72 ± 4.28
	Overall	10.71 ± 8.55	9.64 ± 7.82	7.04 ± 4.92	5.22 ± 3.22	8.15 ± 5.66
Total N (µM)	86–97	51.53 ± 19.03	48.58 ± 18.83	41.39 ± 19.00	35.10 ± 26.40	44.15 ± 18.05
	98–08	28.08 ± 20.20	25.98 ± 19.14	22.25 ± 16.91	19.04 ± 15.90	23.84 ± 17.64
	Overall	40.32 ± 22.81	37.77 ± 22.06	32.23 ± 20.39	27.42 ± 23.40	34.44 ± 20.52
PO ₄ -P (µM)	86–97	1.32 ± 0.78	1.32 ± 0.72	0.93 ± 0.49	0.59 ± 3.69	1.04 ± 0.54
	98–08	0.30 ± 0.23	0.31 ± 0.24	0.30 ± 0.16	0.33 ± 0.16	0.31 ± 0.17
	Overall	0.84 ± 0.77	0.84 ± 0.74	0.63 ± 0.49	0.46 ± 0.30	0.69 ± 0.55
Total P (µM)	86–97	3.59 ± 2.13	3.43 ± 1.58	2.75 ± 1.37	2.23 ± 1.42	3.00 ± 1.43
	98–08	1.14 ± 0.61	1.03 ± 0.53	0.94 ± 0.38	0.87 ± 0.31	0.99 ± 0.38
	Overall	2.42 ± 2.01	2.28 ± 1.69	1.89 ± 1.37	1.58 ± 1.25	2.04 ± 1.47

SiO ₂ (μM)	86–97	13.56 ± 8.41	13.96 ± 7.86	13.90 ± 6.40	12.97 ± 5.03	13.60 ± 6.47
	98–08	12.64 ± 7.87	13.24 ± 7.38	12.68 ± 6.26	12.44 ± 5.84	12.75 ± 6.27
	Overall	13.12 ± 8.16	13.61 ± 7.63	13.32 ± 6.35	12.72 ± 5.43	13.19 ± 6.38
NO ₃ /NH ₄	86–97	0.35 ± 0.33	0.75 ± 3.50	0.72 ± 1.44	0.73 ± 0.93	0.64 ± 1.08
	98–08	0.30 ± 0.56	0.41 ± 0.72	0.69 ± 2.10	1.16 ± 2.73	0.64 ± 1.34
	Overall	0.32 ± 0.45	0.58 ± 2.58	0.71 ± 1.78	0.94 ± 2.01	0.64 ± 1.21
TIN/PO ₄	86–97	12.12 ± 6.06	10.39 ± 4.67	10.44 ± 5.13	11.69 ± 7.30	11.16 ± 5.01
	98–08	24.99 ± 18.58	23.16 ± 16.93	18.69 ± 12.95	15.44 ± 10.09	20.57 ± 12.91
	Overall	18.27 ± 15.00	16.49 ± 13.94	14.39 ± 10.52	13.49 ± 8.93	15.66 ± 10.70
TIN/SiO ₂	86–97	1.37 ± 1.26	1.13 ± 1.00	0.74 ± 0.64	0.47 ± 0.26	0.93 ± 0.71
	98–08	0.73 ± 0.78	0.62 ± 0.65	0.49 ± 0.53	0.41 ± 0.27	0.56 ± 0.52
	Overall	1.07 ± 1.11	0.89 ± 0.89	0.62 ± 0.60	0.44 ± 0.27	0.75 ± 0.65

Table 2. Kendall's correlation between phytoplankton variables and various physico-chemical, biological and nutrient ratios parameters at four sampling stations in Tolo Harbour. "Dino" represents dinoflagellates, "Others" represents algal groups other than diatoms or dinoflagellates, "Total" represents total phytoplankton cell counts. "+/-" represents statistical significance at the 0.05 level; "++/--" represents statistical significance at the 0.01 level. Tested values were depth-averaged values except for "Bottom DO" and "Bottom BOD₅".

Parameter	Station														
	TM3					TM4					TM6				
	Diatom	Dino	Others	Total	Chl- <i>a</i>	Diatom	Dino	Others	Total	Chl- <i>a</i>	Diatom	Dino	Others	Total	Chl- <i>a</i>
Temperature	++	-		++		++			++		++			++	
pH			--		++			--	-	++					++
Salinity					-					--		-	-	-	--
DO		++			++				+		--			-	
Bottom DO		++	++	+		-	+	++			--			--	-
BOD ₅		++	--		++		++	--		++		++	+	++	
Bottom BOD ₅		+			++	-	++		-	+				+	
Secchi Depth	--	--		--	--	--		+	--	--	--	-		--	--
Suspended solids		++		+	++		+			++				++	++
NO ₃ -N	--	--	--	--	-	--		--	--	-				+	
NH ₄ -N	--		--	--		--	-	--	--		-			--	--

Total inorganic N	--	-	--	--	--	--	-	--	--	--		--		--
Total N	--		--	--	+	--		--	--	++	++	++	--	
PO ₄ -P	--	-	--	--	-	--	-	--	--		+		-	-
Total P	--		--	--		--		--	--	++	++	-	--	--
SiO ₂	--				-	--					+		+	+
NO ₃ /NH ₄	-	--	--	--					-		+	++		+
TIN/PO ₄			++	++				++	-		++		+	+
TIN/SiO ₂	-	--	--	--		-	--	--	--			-		--

Fig. 1.

Fig. 2.

Fig. 3.

Fig. 4.

Figure 5.