Accepted Manuscript

Title: Application of primary haemocyte culture of *Penaeus monodon* in the assessment of cytotoxicity and genotoxicity of heavy metals and pesticides

Authors: Seena Jose, P. Jayesh, A. Mohandas, Rosamma Philip, I. S. Bright Singh

PII: S0141-1136(11)00002-X

DOI: [10.1016/j.marenvres.2010.12.008](https://doi.org/10.1016/j.marenvres.2010.12.008)

Reference: MERE 3496

To appear in: *Marine Environmental Research*

Received Date: 29 October 2010

Revised Date: 23 December 2010

Accepted Date: 26 December 2010

Please cite this article as: Jose, S., Jayesh, P., Mohandas, A., Philip, R., Singh, B. Application of primary haemocyte culture of *Penaeus monodon* in the assessment of cytotoxicity and genotoxicity of heavy metals and pesticides, *Marine Environmental Research* (2011), doi: 10.1016/j.marenvres.2010.12.008

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Application of primary haemocyte culture of *Penaeus monodon* in the assessment of cytotoxicity and genotoxicity of heavy metals and pesticides

Seena Jose^a, P. Jayesh^a, A. Mohandas^a, Rosamma Philip^b and I. S. Bright Singh^{a*}

^aNational Centre for Aquatic Animal Health, Cochin University of Science and Technology, Fine Arts Avenue, Kochi 682016, India

^bDepartment of Marine Biology, Microbiology and Biochemistry, School of Marine Sciences, Cochin University of Science and Technology, Fine Arts Avenue, Kochi 682016, India

Corresponding author. Address: National Centre for Aquatic Animal Health, Cochin University of Science and Technology, Fine Arts Avenue, Kochi 682016, India. Tel.: +91 484 2381120; fax: +91 484 2381120.

E-mail address: bsingh@md3.vsnl.net.in (I. S. Bright Singh).

ABSTRACT

Lack of shrimp cell lines has hindered the study of pollutants which adversely affects shrimp health and its export value. In this context a primary haemocyte culture developed from *Penaeus monodon* was employed for assessing the cytotoxicity and genotoxicity of two heavy metal compounds, cadmium chloride and mercuric chloride and two organophosphate insecticides, malathion and monocrotophos. Using MTT assay 12 h IC_{50} values calculated were $31.09 \pm 16.27 \mu M$ and $5.52 \pm 1.16 \mu M$ for cadmium chloride and mercuric chloride and $59.94 \pm 52.30 mg l^{-1}$ and $186.76 \pm 77.00 mg l^{-1}$ for malathion and monocrotophos respectively. Employing Comet assay, DNA damage inflicted by these pollutants on haemocytes were evaluated and the pollutants induced DNA damage in $> 60\%$ of the cells. The study suggested that haemocyte culture could be used as a tool for quantifying cytotoxicity and genotoxicity of aquaculture drugs, management chemicals and pollutants.

Keywords: Aquaculture, *Penaeus monodon*, primary haemocyte culture, heavy metals, organophosphate pesticides, cytotoxicity, genotoxicity, MTT assay, Comet assay

1. Introduction

Cell lines are widely utilized for assessing the toxicity of major aquatic pollutants such as heavy metals and pesticides. Effluents from industries and runoff from agricultural fields may contain such pollutants, which on reaching aquatic environments lead to impairment of the ecosystem. Needless to say, the bio-accumulation and persistence of these pollutants in the aquatic environment constitute a serious threat to biological life and also to human beings indirectly by virtue of their involvement in the food chain (Binelli and Provini, 2004).

Among heavy metals, cadmium and mercury are the most highly toxic elements owing to their persistence in the environment. They are redox inactive which challenge antioxidant defense by binding to thiols in cells such as reduced glutathione (GSH) (Stohs and Bagchi, 1995; Ercal et al., 2001). They cause oxidative and nitrosative stress (Stohs and Bagchi, 1995; Ercal et al., 2001; Pompella et al., 2003) and damage macromolecules in cells. Apoptosis or necrotic cell deaths are the usual consequences (Pulido and Parrish, 2003). Toxicity of pesticides on non - target organisms and ecosystems is of world wide concern (Pimentel et al., 1998). Among pesticides, organophosphate pesticides are stable enough to remain in their toxic form for relatively shorter periods of time (Poovala et al., 1998). They act primarily by inhibiting the enzyme acetylcholine esterase at cholinergic junctions of the nervous system (Gallo and Lawryk, 1991) leading to functional changes at the tissue/organism level (Pope et al., 2005).

Heavy metals and pesticides are known genotoxicants, as they have the capacity to interact with and damage the structure of the DNA molecule, often with a concomitant adverse effect on biological integrity (Shugart, 1995; 1998). Many genotoxicants may cause DNA strand breakage by the formation of free radicals or abasic sites which can result in the breakage of phosphodiester linkages within the DNA molecule (Shugart, 2000).

Attempts to develop cell cultures from marine invertebrates date back about a century (Gomot, 1971; Rannou, 1971). Yet for unidentified reasons all efforts to develop continuous cell lines from these organisms including crustaceans have been ineffective so far (Mothersill and Austin, 2000; Rinkevich, 2005). *Penaeus monodon* is one among the important cultured crustacean in the world. Lack of shrimp cell lines has been hampering progress of research in testing the effects of drugs and management chemicals prior to their administration in aquaculture systems. This has also hindered the study on the effects of pollutants such as heavy metals and pesticides in the aquatic environment, which adversely affects shrimp health and its export value. Even though development of primary cell cultures (Chen et al., 1986; Fan and Wang, 2002; Maeda et al., 2003) and their application in the study of viral pathogens (Wang et al., 2000; Maeda et al., 2004; Jiang et al., 2005) have been reported, their utility in the assessment of cytotoxicity and genotoxicity of heavy metals and pesticides has not been attempted. In the study undertaken here primary haemocyte culture of *P. monodon* was utilized for examining cytotoxicity and genotoxicity of two heavy metal compounds, cadmium chloride ($\text{CdCl}_2 \cdot 2\frac{1}{2} \text{H}_2\text{O}$) and mercuric chloride (HgCl_2) and two organophosphate insecticides, malathion ($\text{C}_{10}\text{H}_{19}\text{O}_6\text{PS}_2$) and monocrotophos ($\text{C}_7\text{H}_{14}\text{NO}_5\text{P}$) which reach aquaculture systems

through run off. Malathion and monocrotophos are also used as insecticides in paddy fields and aquaculture ponds. Comet assay was employed to evaluate the DNA damage caused by these pollutants. The work was conducted to assess the usefulness of primary haemocyte culture for the study of cytotoxicity and genotoxicity of aquatic pollutants.

2. Materials and methods

2.1. Experimental animals

WSSV and Monodon Baculo virus (MBV) negative *P. monodon* larvae obtained from a local hatchery were stocked and reared in a recirculating shrimp rearing system integrated with nitrifying bioreactors (Kumar et al., 2009) in sea water having a salinity of 15 g l⁻¹. Water quality was maintained by the addition of a probiotic DetrodigestTM (NCAAH, India) to manage detritus and EnterotrophicTM (NCAAH, India) to control *Vibrio*. Water quality parameters measured were pH (7.5 to 8.5), temperature (25 ± 1 °C), NH₄⁺ - N (< 0.1 mg l⁻¹) and alkalinity (60-75 mg l⁻¹). The larvae were fed three times daily with commercially available pelleted feed having 40% protein (Higashimaru, India) made of fish meal, wheat gluten, squid meal, clam meal, wheat flour, yeast, soy lecithin, krill meal, marigold flower petal extract, refined fish oil, cholesterol, minerals and vitamins. They were confirmed WSSV negative by nested PCR (WSSV detection kit, Genei, India) when they grew to 8-12 g after a culture period of three months, and used for all experiments.

2.2. Development of primary haemocyte culture

The animals were sacrificed by immersing in crushed ice and disinfected by maintaining in 800 mg l⁻¹ sodium hypochlorite solution prepared in ice cold -sea water (salinity 15 g l⁻¹) for 10 minutes. Subsequently, they were washed 5 times in sterile ice cold sea water, dipped in 70% alcohol and rinsed in ice-cold sea water. Haemolymph was withdrawn aseptically using capillary tubes containing 100 µl anticoagulant (Tris HCl 0.01 M (pH 7), Sucrose 0.25 M, Tri Sodium Citrate 0.1 M) from rostral sinus and diluted to obtain 5 x 10⁵ cells ml⁻¹ using modified 2X L-15 (Leibovitz) medium supplemented with 2% glucose, MEM vitamins (1X), tryptose phosphate broth (2.95 g l⁻¹), 20% FBS, N-phenylthiourea (0.2 mM), 0.06 µg ml⁻¹ chloramphenicol, 100 µg ml⁻¹ streptomycin and 100 IU ml⁻¹ penicillin, and aliquots of 200 µl were dispensed into 96 well microplates (Greiner Bio-One) and incubated at 25 °C (Jose et al., 2010) for 12 hours. Haemolymph from 5 animals were pooled and diluted to seed a microwell plate.

2.3. Cytotoxicity of cadmium chloride (CdCl₂·2 ½ H₂O) and mercuric chloride (HgCl₂)

Cadmium chloride and mercuric chloride were used as representatives of aquatic pollutants. Stock solutions of cadmium chloride (Qualigens) and mercuric chloride (Merck) were prepared in cell culture medium. A range finding test was carried out before the definitive test and different concentrations of the compounds were added to the wells to get the final concentrations ranging from 0.49 to 500 µM for cadmium chloride and 0.49 to 62.5 for mercuric chloride respectively. Replicates were maintained for each concentration. Cells with the growth medium were kept as control. After 12 hours incubation, wells were observed under phase contrast microscope (Leica, Germany) and

MTT assay was performed. Percentage inhibition of cells at each concentration of the compounds was calculated based on the formula,

Percentage inhibition of haemocytes = $[100 - (\text{Average absorbance (MTT assay) of haemocytes at a particular concentration of the compound} / \text{Average absorbance of control haemocytes without the compound}) \times 100]$.

2.4. Cytotoxicity of malathion [S-(1, 2-dicarboethoxyethyl) O, O-dimethyl phosphorodithioate]; ($C_{10}H_{19}O_6PS_2$)) and monocrotophos (E)-(dimethyl-1-methyl-3-(methylamino)-3-oxo-1-propenyl phosphate); ($C_7H_{14}NO_5P$)

Commercially available preparations of malathion (50% w/w) and monocrotophos (53% w/w) were used for toxicity studies. Insecticides were dissolved in DMSO and diluted with medium to get stock concentrations of 500 mg l⁻¹ and 1000 mg l⁻¹ of malathion and monocrotophos respectively. Concentrations were fixed based on the range finding test and different concentrations of the compounds were added to the wells to get final concentrations ranging from 0.76 to 185.2 mg l⁻¹ for malathion and 0.98 to 1000 mg l⁻¹ for monocrotophos maintaining replicates for each concentration. Cells with the growth medium were kept as control. After 12 hours incubation wells were observed under phase contrast microscope (Leica, Germany) and MTT assay was performed. Percentage inhibition of cells at each concentration of the compounds was calculated.

2.5. MTT assay

The assay is a colorimetric method based on the determination of cell viability utilizing the reaction of a tetrazolium salt (3-(4, 5-dimethylthiazol-2-yl)-2, 5-diphenyl tetrazolium bromide, MTT) with the mitochondria of metabolically active cells. The reduction of the tetrazolium salt by nicotinamide adenine dinucleotide dehydrogenase (NADH) and nicotinamide adenine dinucleotide phosphate dehydrogenase (NADPH) within the cells produces insoluble purple formazan crystals, which are later solubilized yielding a purple-colored solution (Mosmann, 1983).

After replacing the medium, 50 μ l of MTT (Sigma) solution (5 mg ml⁻¹ in PBS; 720 mOsm) were added to each well and incubated for 5 hours in dark. Blank consisted of medium alone with MTT added. After incubation, the medium was removed and MTT-formazan crystals were dissolved in 200 μ l dimethylsulfoxide. Absorbance was recorded immediately at 570 nm in a micro plate reader (TECAN Infinite Tm, Austria).

2.6. Comet assay

Haemocytes were exposed to cadmium chloride (140 μ M), mercuric chloride (17 μ M), malathion (60 mg l⁻¹) and monocrotophos (186 mg l⁻¹) for 4 hours. These concentrations were chosen because more than 80% viability of cells was recorded after 4 hour exposure by trypan blue viability assay as dead cells might give rise to false positive results in Comet assay. After the exposure to the compounds, cells were detached using a cell scraper (Greiner Bio-One), pelletised, washed and suspended in PBS. Negative control consisted of cells without any treatment. Added 150 μ l normal agarose solutions (Sigma, USA) (0.75% in PBS) to a frosted slide and covered with a cover slip (22 x 50mm) to get a uniform layer, and allowed to solidify. To prepare the second layer, 10 μ l

cell suspension was mixed with 70 μ l low melting agarose (Sigma, USA) (0.75% in PBS) and, the suspension was layered over the one prepared initially after removing the cover slip. The preparation was kept undisturbed after replacing the cover slip until agarose got solidified. A third layer of low melting agarose of the above strength was applied over the second layer after removing the cover slip and replaced the same subsequently. The slides were treated with pre chilled lysis solution (2.5 M NaCl, 100 mM EDTA, 10 mM Tris, 1.5 M NaOH, pH 10 with 1% Triton X 100 and 10% DMSO) for 1 hour at 4 °C after removing the cover slips. On completion of the process, the slides were incubated in electrophoresis buffer (300 mM NaOH and 1mM EDTA, pH>13) for 20 min at 4 °C for unwinding of DNA, and were electrophoresed for 40 min at 25 V and 300 mA in the same buffer. Subsequently, the slides were neutralized in 0.4 M Tris (pH 7.5), stained with ethidium bromide (2 μ g ml⁻¹) and observed under fluorescent microscope (Olympus, Germany) at 400X magnification, and the cells were photographed. For each sample, 25 cells from each slide were analysed for DNA damage using Comet ScoreTM (TriTek Corp., Summerduck, USA), of the Comet scoring software. Experiments were repeated four times. Percentage of nuclei with tails, tail length, percentage DNA in the tail and tail moment (arbitrary units) were used to estimate DNA damage. Tail moment is the tail length multiplied by fraction of DNA in the tail.

2.7. Statistical analysis

The results in the figures are average values of 4 replicates $\pm$ standard deviation. The results of the cytotoxicity assay were analyzed by probit analysis using the SPSS

software (SPSS Inc., USA). The effects of treatments were statistically analyzed by analysis of variance (ANOVA). Differences were considered significant at $P < 0.05$.

3. Results

3.1. Cytotoxicity of cadmium chloride and mercuric chloride

Control haemocytes appeared spherical or elliptical initially, and within a few hours of culturing two distinct morphological types such as round to elliptical and small and large spindle shaped cells, with granules, could be seen in the modified 2X L-15 medium (Fig. 1). At higher concentrations of cadmium chloride ($>125 \mu\text{M}$) and mercuric chloride ($>15.6 \mu\text{M}$) round cells were few and spindle shaped cells were totally absent. More round cells and few spindle shaped cells were present at lower concentrations ($<15.6 \mu\text{M}$ for cadmium chloride and $<1.95 \mu\text{M}$ for mercuric chloride) (Fig 1). At still lower concentrations number of spindle cells was higher. Toxicity of cadmium chloride and mercuric chloride in primary haemocyte culture in terms of percentage inhibition of cells determined through MTT assay is presented in Figs 2 and 3. IC_{50} values were $31.09 \pm 16.27 \mu\text{M}$ and $5.52 \pm 1.16 \mu\text{M}$ ($7.10 \pm 3.72 \text{ mg l}^{-1}$ and $1.50 \pm 0.31 \text{ mg l}^{-1}$) for cadmium chloride and mercuric chloride respectively.

3.2. Cytotoxicity of malathion and monocrotophos

At higher concentrations of malathion and monocrotophos ($>61.7 \text{ mg l}^{-1}$ and $>166 \text{ mg l}^{-1}$ respectively) only round cells were present and the number of spindle shaped cells

Fig 1

Fig 2,
3, 4,

increased as the concentrations decreased (Fig 1). Toxicity of malathion and monocrotophos in primary haemocyte culture in terms of percentate inhibition of cells determined hrough MTT assay is presented in Figs 4 and 5. IC₅₀ value of malathion was found to be $59.94 \pm 52.30 \text{ mg l}^{-1}$ and of that of monocrotophos $186.76 \pm 77.00 \text{ mg l}^{-1}$. Malathion was found to be non toxic at $\leq 6.86 \text{ mg l}^{-1}$ and monocrotophos at $\leq 15.6 \text{ mg l}^{-1}$.

3.3. Genotoxicity of cadmium chloride, mercuric chloride, malathion and monocrotophos

All pollutants tested induced DNA strand breakage and thereby Comets in the haemocytes (Fig 6). In the negative control most of the nuclei were circular without any visible tail while short tails were observed in $11.46 \pm 7.89\%$ of cells (Table 1). Percentage of cells with tails were significantly higher ($P < 0.05$) than the control for cadmium chloride, mercuric chloride, malathion and monocrotophos ($> 60\%$). Tail length of the Comets in mercuric chloride treatment was the lowest (23.09 ± 6.83 pixels) while for cadmium chloride, malathion and monocrotophos length of the tail ranged from 28-33 pixels. Percentage tail DNA for all the pollutants were significantly higher ($P < 0.05$) but no significant differences ($P > 0.05$) were obtained between that of cadmium chloride and mercuric chloride, and malathion and monocrotophos. Tail moment was found to be lesser for mercuric chloride (11.82 ± 5.50) ($P < 0.05$) when compared to the other three and the values were highest for cadmium chloride (17.17 ± 6.22). In the control tail moment was significantly lower (0.04 ± 02) ($P < 0.05$).

Fig
6

Table
1

4. Discussion

Deleterious effects of environmental pollutants may result from direct toxic action on the tissues or from subtle alterations in the homeostatic mechanisms such as the immune system (Auffret and Oubella, 1997). The immune defense of shrimp is comprised of cellular and humoral - mediated mechanisms, in which haemocytes play a key role and this makes haemocyte culture an ideal system for evaluating the effects of environmental pollutants. Primary cell cultures of penaeid shrimp have been developed from different tissues such as lymphoid (Chen and Wang, 1999; Lang et al., 2002), heart (Tong and Miao, 1996; Chen and Wang, 1999), ovary (Shimizu et al., 2001; Maeda et al., 2003) and embryo (Toullec et al., 1996; Fan and Wang, 2002) while haemocyte culture was attempted only by a few researchers in *P. vannamei*, *P. aztecus* (Ellender et al., 1992), *P. japonicus* (Itami et al., 1999) and *P. chinensis* (Jiang et al., 2005). In the primary haemocyte culture developed, round to elliptical cells could be regarded as hyaline cells and small and large spindle shaped cells with granules could be regarded as granular or semi-granular cells based on the morphology and presence of granules. The metabolic activity of the culture was almost the same during the second, third and fourth days and the activity started declining only subsequently. The culture remained viable for upto 8 days. Even though dividing cells were not observed, DNA synthesis was detected in $22 \pm 7\%$ of cells by means of 5-Bromo-2'-deoxyuridine (BrdU) assay at 24 hours (Jose et al., 2010). Studies on the application of haemocyte culture or any other primary cultures of shrimp in cytotoxicity and especially the genotoxicity of drugs, management chemicals and pollutants in aquaculture systems have been rarely cited in literature. The *P. monodon* haemocyte culture has been applied for assessing cytotoxicity of N-methyl-1-hydroxyphenazine, a proposed aquaculture drug (Preetha et al., 2009) and benzalkonium

chloride, an aquaculture management chemical (Jose et al., 2010) employing MTT assay. Sung et al. (2003) examined the effects and toxicity of phthalate esters on the isolated haemocytes of *Macrobrachium rosenbergii* utilizing haemocytic adhesion, pseudopodia formation, superoxide anion production and phenoloxidase activity.

Toxicity of cadmium chloride and/or mercuric chloride has been evaluated in vivo on different aquatic organisms such as fishes (Pereira et al., 2009), sea urchin (Filosto et al., 2008), shrimp (Lee et al., 2000; Espericueta et al., 2001) and crab (Botton, 2000). In our study toxicity was tested in vitro on primary haemocyte culture of *P. monodon* and 12 hour IC_{50} values for cadmium chloride and mercuric chloride were $31.09 \pm 16.27 \mu M$ and $5.52 \pm 1.16 \mu M$, respectively. Mercuric chloride was found to be more toxic than cadmium chloride and concentration dependent increase was observed in the toxicity of both heavy metals in haemocytes. Espericueta et al. (2001) reported the LC_{50} values as 2.49 and $1.23 mg l^{-1}$ for cadmium and mercury, respectively in post larvae of *Litopenaeus vannamei* after 96 hour exposure while in *Limulus polyphemus* embryos IC_{50} was reported to be $>1000 mg l^{-1}$ and $12.80 mg l^{-1}$ for Cd and Hg after 24 hour exposure (Botton, 2000). In the present study, MTT assay which measured the metabolic activity as well as viability of cells was utilized to estimate the cytotoxicity rather than trypan blue assay which measured only the viability. IC_{50} value of malathion was found to be $59.94 \pm 52.30 mg l^{-1}$ and that of monocrotophos was $186.76 \pm 77.00 mg l^{-1}$. Cytotoxicity increased with concentration for both the pesticides, and malathion was more toxic when compared to monocrotophos. Similarly, concentration dependant increase in the toxicity of malathion was detected in grass carp cell line (Chen et al., 2006) and human liver carcinoma cell line (Moore et al., 2010). IC_{50} of malathion ranged from $37.94 \pm 1.93 mg$

l⁻¹ for 12 h to 3.04 ± 0.27 mg l⁻¹ for 72 hours in grass carp cells (Chen et al., 2006). In the nauplii (48 h old) of *Artemia salina* 24 h LC₅₀ of monocrotophos was estimated to be 262.68 ± 17.30 mg l⁻¹ (Rao et al., 2007). These values are comparable to those in the present study. Semi-granular or granular cells were more sensitive to heavy metals and pesticides than the hyaline cells and the number of these cells increased as the pollutant concentration decreased. According to Holmblad and Soderhall (1999) semi-granular cells appear to be the most sensitive and react first during an immune response, by degranulating. This degranulation stimulates the granular cells also to degranulate as well. The advantage of using haemocyte culture over explant cultures for cytotoxicity studies is the easiness in quantification of cells to be seeded to maintain uniform cell number in micro plates, a prerequisite for quantal assays, and the rapidity in obtaining monolayer of cells.

Different methods have been developed for detecting DNA strand damage such as micronucleus and sister-chromatid exchange assays which are based on the enumeration of downstream aberrations after DNA damage (Lee and Steinert, 2003). A more recent method is the Comet assay which detects the DNA strand breaks and alkali labile sites by measuring the migration of DNA from immobilized nuclear DNA (Singh et al., 1988). A large number of studies report that Comet assay is more sensitive when compared to sister chromatid exchanges or micronucleus test (Mitchelmore and Chipman, 1998; Lee and Steinert, 2003; Frenzilli et al., 2009). The main advantages of the Comet assay include the collection of data at the level of the individual cell, the need for a small number of cells per sample and the sensitivity for detecting DNA damage (Lee and Steinert, 2003; Jha, 2004). Tail length is the basic parameter used for quantifying DNA

damage in the Comet assay. However, the tail length levels off after migrating to a certain distance. After the maximum length is attained, the Comet tail will not be extending in length, however grow in intensity as compared to that with the head portion (Bowden et al., 2003). Therefore, in addition to tail length, percentage tail DNA and tail moment were also calculated using Comet scoring software.

Comet assay has become one of the most popular tools for detecting DNA strand breaks in aquatic animals (Frenzilli et al., 2009) and has been employed to assess the DNA damage in aquatic vertebrates such as *Cyprinus carpio* (Arabi et al., 2004), *Carassius auratus* (Masuda et al., 2004), *Danio rerio* (Kosmehl et al., 2008) and *Liza aurata* (Pereira et al., 2009), and invertebrates such as *Limnoperna fortunei* (Villela et al., 2007), *Dreissena polymorpha* (Juhel et al., 2007), *Polycelis feline* (Horvat et al., 2005), *Palaemonetes pugio* (Lee et al., 2000; Lee et al., 2008), *Mytilus edulis* (Rank and Jensen, 2003; Cheung et al., 2006) and *Crassostrea virginica* (Bissett et al., 2009). For in vivo studies in aquatic invertebrates such as oysters, mussels, clams and shrimp, cells from haemolymph, embryos, gills and digestive glands were used for Comet assay (Lee and Steinert, 2003). In vitro studies of genotoxicants were reported in haemocytes, gill cells and digestive gland cells from various mollusks namely *Mytilus edulis*, *Nassarius tegula* and *Musculista senhousia* (Sastre et al., 1997; Mitchelmore et al., 1998; Wilson et al., 1998). Studies of genotoxicants in shrimp cell cultures were not reported while Lee et al. (2000) studied the effects of genotoxicants such as chromium (III) chloride, sodium chromate, mercuric chloride, and 2-methyl-1,2-naphthoquinone (MNQ) on grass shrimp, *Palaemonetes pugio*, embryos employing Comet assay. DNA damage in a variety of aquatic animals has been associated with reduced growth, abnormal development and

reduced survival of embryos, larvae and adults (Steinert, 1999; Lee et al., 2000). In grass shrimp embryos Comet assay was used to provide a sensitive measure of DNA strand breakage exposed to genotoxicants, highway run off sediments, sediments with coal ash and phototoxicants (Lee et al., 2000; 2008; Lee and Kim, 2002; Kim and Lee, 2004).

In the concentrations tested cadmium chloride and mercuric chloride induced Comets in 72% and 61% of the cells respectively after an exposure over a period of 4 hours. Genotoxicity of cadmium was reported in rat hemispheres and cerebellum (Klimova and Misurova, 2002) in rat Leydig cells (Yang et al., 2003), in *Palaemonetes pugio* embryos (Hook and Lee, 2004), in *Mytilus edulis* (Pruski and Dixon, 2002; Emmanouil et al., 2007) and in *Litopenaeus vannamei* (Chang et al., 2009). The mechanism of metal induced carcinogenesis is still unknown, but one possible pathway may involve the interaction of metals with DNA, either directly or indirectly (Hartmann and Speit, 1994). Cadmium induces an oxidative DNA damage leading to DNA strand breaks, DNA-protein cross-linking and inhibitions of DNA repair (Dally and Hartwig, 1997; Valverde et al., 2001). Genotoxicity of mercury was reported in *Cyprinus carpio* gill cells (Arabi et al., 2004), blood cells of *Liza aurata* (Pereira et al., 2009), human salivary gland tissue cells and lymphocytes (Schmid et al., 2007) and *Palaemonetes pugio* embryo (Lee et al., 2000). HgCl_2 have a deleterious effect on the membrane integrity and glutathione (GSH) content of gill cell suspensions of carp, *Cyprinus carpio* L. and mercury caused DNA strand breaks at 3000 μM (Arabi et al., 2004). HgCl_2 has been reported to cause DNA damage in *P. pugio* embryos at concentrations as low as 0.003 μM after a 12 hour exposure period (Lee et al., 2000). DNA damage was reported by Pereira et al. (2009) in golden grey mullet, *Liza aurata* due to environmental exposure

of mercury. Clastogenic effects mostly associated with the spindle mechanism disturbance and the generation of reactive oxygen species, accompanied by glutathione depletion contribute to the genotoxicity of mercury (de Flora et al., 1994). The present study demonstrates the potential of cadmium chloride and mercuric chloride to induce DNA damage in primary haemocyte culture.

Giri et al. (2002) evaluated the genotoxicity of malathion using chromosome aberration, sister chromatid exchange and sperm abnormality assays in mice. Genotoxicity was detected in human liver carcinoma cells at 24 mM exposure of malathion. In *Channa punctatus*, sublethal concentrations of malathion (Kumar et al., 2010) and monocrotophos (Ali and Kumar, 2008) induced dose dependant DNA damage in gill, kidney and lymphocytes were reported during a study period of 1-29 days and 1-21 days respectively. Jamil et al. (2005) reported genotoxic effect of monocrotophos in human lymphocytes in vitro employing Comet assay. A concentration dependant increase in the DNA damage was detected in the erythrocytes of *Tilapia mossambica* (Banu et al., 2001) and the gill cells of the bivalve, *Meretrix ovum* (Revankar and Shyama, 2009) after the exposure to monocrotophos. Both malathion and monocrotophos are organophosphorous compounds capable of phosphorylation and alkylation. Although phosphorylation is responsible for the inhibition of acetylcholineesterase and produces corresponding neurotoxicity in target and non-target species, other biological activities such as mutation appear to be due to the alkylation properties (Eto and Ohkawa, 1970). The heterocyclic bases of the nucleic acids may serve as nucleophilic agents for such a reaction, and alkylation may take place on nitrogen atoms possessing high electron density. Alkylation is more likely with the methyl ester group than ethyl and higher alkyl

ester groups, and phosphate esters such as monocrotophos are more reactive than the phosphorothionate esters such as malathion. As organophosphorous insecticides are chemical alkylating agents (Wild, 1975), alkylation of DNA bases either directly or indirectly via protein alkylation is probably involved in the DNA damage (Mohan, 1973). According to Wild (1975) the phosphorous moiety in the organophosphates appears to be a good substrate for nucleophilic attack and it causes phosphorylation of DNA, which is an instance of DNA damage. Malathion used in the present study is a commercial grade product and may contain impurities formed during manufacturing and storage. Among the impurities are malaoxon formed from malathion by oxidation and isomalathion formed by isomerization (Berkman et al., 1993). Malaoxon and isomalathion have been reported to be mutagenic (Flessel et al., 1993; Pluth et al., 1996, Blasiak et al., 1999). These two compounds also could contribute to the genotoxicity of malathion. In the present study malathion and monocrotophos, which are the two commonly used insecticides in India, have proved to be genotoxic to *P. monodon* primary haemocyte culture.

In conclusion, the study has demonstrated the use of primary shrimp haemocyte culture for testing the cytotoxicity of aquatic pollutants such as heavy metals and pesticides employing MTT assay. The primary cell culture system could also be employed to assess the DNA damage induced by genotoxic pollutants in the aquatic environment. The present study confirms the genotoxic potential of two heavy metal compounds, cadmium chloride and mercuric chloride and two organophosphate insecticides, malathion and monocrotophos, and this forms the first report of the demonstration of genotoxicity of pollutants using shrimp cell culture as a model system.

402

403 **Acknowledgements**

404 This research was supported by Department of Biotechnology, Government of India
405 (BT/PR3509/AAQ/03/173/2000; BT/PR8050/AAQ/03/289/2006). The first author thanks
406 DBT for Fellowship.

407

408 **References**

- 409 Ali, D., Kumar, S., 2008. Long term genotoxic effect of monocrotophos in different
410 tissues of fresh water fish *Channa punctatus* (Bloch) using alkaline single cell gel
411 electrophoresis. Science of the Total Environment 405, 345-350.
- 412 Arabi, M., 2004. Analyses of impact of metal ion contamination on Carp (*Cyprinus*
413 *carpio* L.) gill cell suspensions. Biological Trace Element Research 100, 229-245.
- 414 Auffret, M., Oubella, R., 1997. Haemocyte aggregation in the oyster *Crassostrea gigas*:
415 in vitro measurement and experimental modulation by xenobiotics. Comparative
416 Biochemistry and Physiology Part A: Physiology 118, 705-710.
- 417 Banu, B.S., Danadevi, K., Rahman, M.F., Ahuja, Y.R., Kaiser, J., 2001. Genotoxic effect of
418 monocrotophos to sentinel species using comet assay. Food and Chemical Toxicology
419 39, 361-366.
- 420 Berkman, C.E., Thompsun, C.M., Perrin, S.R., 1993. Synthesis, absolute configuration, and
421 analysis of malathion, malaxon, and isomalathion enantiomers. Chemical Research in
422 Toxicology 718-723.

- 423 Binelli, A., Provini, A., 2004. Risk for human health of some POPs due to fish from Lake
424 Iseo. *Ecotoxicology and Environmental Safety* 58, 139-145.
- 425 Bissett, Jr. W., Smith, L., Thompson, J.A., 2009. Geostatistical analysis of DNA damage
426 in oysters, *Crassostrea virginica*, in Lavaca Bay, Texas. *Ecotoxicology* 18, 69-74.
- 427 Blasiak, J., Jalszynski, P., Trzeciak, A., Szyfter, K., 1999. In vitro studies on the
428 genotoxicity of the organophosphorus insecticide malathion and its two
429 analogues. *Mutation Research* 445, 275-283.
- 430 Botton, M.L., 2000. Toxicity of cadmium and mercury to horseshoe Crab (*Limulus*
431 *polyphemus*) embryos and larvae. *Bulletin of Environmental Contamination and*
432 *Toxicology* 64, 137-143.
- 433 Bowden, R.D., Buckwalter, M.R., McBride, J.F., Johnson D.A., Murray, B.K., O'Neill,
434 K.L., 2003. Tail profile: a more accurate system for analyzing DNA damage using
435 the Comet assay. *Mutation Research* 537, 1-9.
- 436 Chang, M., Wang, W.N., Wang, A.L., Tian, T.T., Wan, P., Zheng, Y., Liu, Y., 2009. Effects of
437 cadmium on respiratory burst, intracellular Ca²⁺ and DNA damage in the white shrimp
438 *Litopenaeus vannamei*. *Comparative Biochemistry and physiology - Part C:*
439 *Toxicology and Pharmacology* 149, 581-586.
- 440 Chen, S.N., Chi, S.C., Kou, G.H., Liao, I.C., 1986. Cell culture from tissues of grass
441 prawn, *Penaeus monodon*. *Fish Pathology* 21, 161-166.

- 442 Chen, S.N., Wang, C.S., 1999. Establishment of cell culture systems from penaeid shrimp
443 and their susceptibility to white spot disease and yellow head viruses. Methods in
444 Cell Science 21, 199-206.
- 445 Chen, X.Y., Shao, J.Z., Xiang, L.X., Liu, X.M., 2006. Involvement of apoptosis in
446 malathion-induced cytotoxicity in a grass carp (*Ctenopharyngodon idellus*) cell
447 line. Comparative Biochemistry and Physiology Part C 142, 36-45.
- 448 Cheung, V.V., Depledge, M.H., Jha, A.N., 2006. An evaluation of the relative sensitivity
449 of two marine bivalve mollusc species using the Comet assay. Marine
450 Environmental Research 62, S301-S305.
- 451 Dally, H., Hartwig, A., 1997. Induction and repair inhibition of oxidative DNA damage
452 by nickel (ii) and cadmium chloride (II) in mammalian cells. Carcinogenesis 18,
453 1021-1026.
- 454 De Flora, S., Bennicelli, C., Bagnasco, M., 1994. Genotoxicity of mercuric compounds.
455 A review. Mutation Research 317, 57-79.
- 456 Ellender, R.D., Najafabadi, A.K., Middlebrooks, B.L., 1992. Observations on the primary
457 culture of haemocytes of *Penaeus* species. Journal of Crustacean Biology 12, 178-
458 185.
- 459 Emmanouil, C., Sheehan, T.M.T., Chipman, J.K., 2007. Macromolecule oxidation and DNA
460 repair in mussel (*Mytilus edulis* L.) gill following exposure to Cd and Cr (VI).
461 Aquatic Toxicology 82, 27-35.

- 462 Ercal, N., Gurer-Orhan, H., Aykin-Burns, N., 2001. Toxic metals and oxidative stress
463 Part I: Mechanisms involved in metal induced oxidative damage. Current Topics
464 in Medicinal Chemistry 1, 529-539.
- 465 Espericueta, M.G.F., Voltolina, D., Lopez, J.I.O., 2001. Acute toxicity of cadmium, mercury,
466 and lead to whiteleg shrimp (*Litopenaeus vannamei*) postlarvae. Bulletin of
467 Environmental Contamination and Toxicology 67, 580-586.
- 468 Eto, M., Ohkawa, H., 1970. Biochemical toxicology of insecticides. Academic Press, New
469 York, p23.
- 470 Fan, T.J., Wang, X.F., 2002. In vitro culture of embryonic cells from the shrimp, *Penaeus*
471 *chinensis*. Journal of Experimental Marine Biology and Ecology 267, 175-184.
- 472 Filosto, S., Roccheri, M.C., Bonaventura, R., Matranga, V., 2008. Environmentally
473 relevant cadmium concentrations affect development and induce apoptosis of
474 *Paracentrotus lividus* larvae cultured in vitro. Cell Biology and Toxicology 24,
475 603-610.
- 476 Flessel, P., Quintana, P.J.E., Hooper, K., 1993. Genetic toxicity of malathion, a review.
477 Environmental and Molecular Mutagenesis 22, 7-17.
- 478 Frenzilli, G., Nigro, M., Lyons, B.P., 2009. The Comet assay for the evaluation of
479 genotoxic impact in aquatic environments. Mutation Research 681, 80-92.
- 480 Gallo, M.A., Lawryk, N.J., 1991. Organic phosphorous pesticides, in: Hayes, W.J., Laws,
481 E.R. (Eds.), Handbook of Pesticide Toxicology. Academic Press, NewYork, pp.
482 5-13.

- 483 Gomot, L., 1971. The organotypic culture of invertebrates other than insects, in: Vago, C.
484 (Ed.), Invertebrate Tissue Culture. Academic Press, New York, pp. 41-36.
- 485 Giri, S., Prasad, S.B., Giri, A., Sharma, G.D., 2002. Genotoxic effects of malathion: an
486 organophosphorus insecticide, using three mammalian bioassays in vivo.
487 Mutation Research – Genetic Toxicology and Environmental Mutagenesis 514,
488 223-231.
- 489 Hartman, A., Speit, G., 1994. Comparative investigations of the genotoxic effects of
490 metals in the single cell gel (SCG) assay and the sister chromatid exchange (SCE)
491 test. Environmental and Molecular Mutagenesis 23, 299-305.
- 492 Holmblad, T., Soderhall, K., 1999. Cell adhesion molecules and antioxidative enzymes in
493 a crustacean, possible role in immunity. Aquaculture 172, 111-123.
- 494 Hook, S.E., Lee, R.F., 2004. Interactive effects of UV, benzo[α] pyrene, and cadmium on
495 DNA damage and repair in embryos of the grass shrimp *Palaemonetes pugio*.
496 Marine Environmental Research 58, 735-739.
- 497 Horvat, T., Kalafatic, M., Kopjar, N., Kovacevic, G., 2005. Toxicity testing of herbicide
498 nor-flurazon on an aquatic bioindicator species—the planarian *Polycelis felina*
499 (Daly), Aquatic Toxicology 73, 342–352.
- 500 Itami, T., Maeda, M., Kondo, M., Takahashi, Y., 1999. Primary culture of lymphoid
501 organ cells and hemocytes of kuruma shrimp, *Penaeus japonicus*. Methods in Cell
502 Science 21, 237-244.

- 503 Jamil, K., Shaik, A.P., Mahboob, M., Krishna, D., 2005. Effect of organophosphorus and
 504 organochlorine pesticides (monochrotophos, chlorpyrifos, dimethoate, and
 505 endosulfan) on human lymphocytes in vitro. *Drug and Chemical Toxicology* 27,
 506 133-144.
- 507 Jha, A.N., 2004. Genotoxicological studies in aquatic organisms: An overview. *Mutation*
 508 *Research* 552, 1-17.
- 509 Jiang, Y.S., Zhan, W.B., Wang, S.B., Xing, J., 2005. Development of primary
 510 shrimp hemocyte cultures of *Penaeus chinensis* to study white spot syndrome
 511 virus (WSSV) infection. *Aquaculture* 253, 114-119.
- 512 Jose, S., Mohandas, A., Philip, R., Bright Singh, I.S., 2010. Primary haemocyte
 513 culture of *Penaeus monodon* as an in vitro model for white spot syndrome
 514 virus titration, viral and immune related gene expression and cytotoxicity
 515 assays. *Journal of Invertebrate Pathology*. In Press,
 516 doi:10.1016/j.jip.2010.08.006.
- 517 Juhel, G., O'Halloran, J., Culloty, S.C., O'riordan, R.M., Davenport, J., O'Brien, N.M.,
 518 James, K.F., Furey, A., Allis, O., 2007. In vivo exposure to microcystins induces
 519 DNA damage in the haemocytes of the zebra mussel, *Dreissena polymorpha*, as
 520 measured with the Comet assay. *Environmental and Molecular Mutagenesis* 48,
 521 22-29.
- 522 Kim, G.B., Lee, R.F., 2004. Effects of genotoxic compounds on DNA and development
 523 of early and late grass shrimp embryo stages. *Marine Environmental Research*.
 524 57, 329-338.

- 525 Klimova, S., Misurova, E., 2002. The effect of cadmium and ionizing radiation on DNA
526 and RNA content in the rat hemispheres and cerebellum. Bulletin of Veterinary
527 Institute of Pulley 46, 127-137.
- 528 Kosmehl, T., Hallare, A.V., Braunbeck, T., Hollert, H., 2008. DNA damage induced by
529 genotoxicants in zebrafish (*Danio rerio*) embryos after contact exposure to freeze-
530 dried sediment and sediment extracts from Laguna Lake (The Philippines) as
531 measured by the Comet assay. Mutation Research 650, 1–14.
- 532 Kumar, R., Nagpure, N.S., Kushwaha, B., Srivastava, S.K., Lakra, W.S., 2010.
533 Investigation of the genotoxicity of malathion to freshwater teleost fish *Channa*
534 *punctatus* (Bloch) using the micronucleus test and comet assay. Archives of
535 Environmental Contamination and Toxicology 58, 123-130.
- 536 Kumar, V.J.R., Achuthan, C., Manju, N.J., Philip, R., Bright Singh, I.S., 2009. Activated
537 packed bed bioreactor for rapid nitrification in brackish water hatchery systems.
538 Journal of Industrial Microbiology and Biotechnology 36, 355-365.
- 539 Lang, G.H., Nomura, N., Wang, B.Z., Matsumura, M., 2002. Penaeid (*Penaeus*
540 *japonicus*) lymphoid cells replicate by cell division in vitro. In Vitro Cellular and
541 Developmental Biology - Animal 38, 142-145.
- 542 Lee, R., Kim, G.B., 2002. Comet assays to assess DNA damage and repair in grass
543 shrimp embryos exposed to phototoxicants. Marine Environmental Research 54,
544 465-469.
- 545 Lee, R.F., Kim, G.B., Maruya, K.A., Steinert, S.A., Oshima, Y., 2000. DNA strand
546 breaks (Comet assay) and embryo development effects in grass shrimp

- 547 (*Palaemonetes pugio*) embryos after exposure to genotoxicants. Marine
548 Environmental Research 50, 553-557.
- 549 Lee, R.F., Niencheski, L.F.H., Brinkley, K., 2008. DNA strand breaks in grass shrimp
550 embryos exposed to highway runoff sediments and sediments with coal fly ash.
551 Marine Environmental Research 66, 164-165.
- 552 Lee, R.F., Steinert, S., 2003. Use of the single cell gel electrophoresis/ Comet Assay for
553 detecting DNA damage in aquatic (marine and freshwater) animals. Mutation
554 Research 544, 43-64.
- 555 Maeda, M., Mizuki, E., Itami, T., Ohba, M., 2003. Ovarian primary culture of the kuruma
556 shrimp *Marsupenaeus japonicus*. In Vitro Cellular and Developmental Research -
557 Animal 39, 208-212.
- 558 Maeda, M., Saitoh, H., Mizuki, E., Itami, T., Ohba, M., 2004. Replication of white spot
559 syndrome virus in ovarian primary cultures from the kuruma shrimp, *Marsupenaeus*
560 *japonicus*. Journal of Virological Methods 116, 89-94.
- 561 Masuda, S., Deguchi, Y., Masuda, Y., Watanabe, T., Nukaya, H., Terao, Y., Takamura,
562 T., Wakabayashi, K., Kinae, N., 2004. Genotoxicity of 2-[2-(acetylamino)-4-
563 [bis(2-hydroxyethyl)amino]-5-methoxyphenyl]-5-amino-7-bromo-4-chloro-2H
564 benzotriazole (PBTA-6) and 4-amino-3,30-dichloro-5,40-dinitro-biphenyl
565 (ADDB) in goldfish (*Carassius auratus*) using the micronucleus test and the
566 Comet assay. Mutation Research 560, 33-40.

- 567 Mitchelmore, C.L., Birmelin, C., Livingstone, D.R., Chipman, J.K., 1998. Detection of
568 DNA strand breaks in isolated mussel (*Mytilus edulis* L.) digestive gland cells
569 using the “Comet” assay. *Ecotoxicology and Environmental Safety* 41, 51–58.
- 570 Mitchelmore, C.L., Chipman, J.K., 1998. DNA stand breakage in aquatic organisms and
571 the potential value of the Comet assay in environmental monitoring. *Mutation*
572 *Research* 399, 135-147.
- 573 Mohan, G., 1973. Comparison of the mutagenic activity of eight organophosphorus
574 insecticides in *E. coli*. *Mutation Research* 21, 196.
- 575 Moore, P.D., Yedjou, C.G., Tchounwou, P.B., 2010. Malathion induced oxidative stress,
576 cytotoxicity, and genotoxicity in human liver carcinoma (HepG₂) cells.
577 *Environmental Toxicology* 25, 221-226.
- 578 Mothersill, C., Austin, B., 2000. *Aquatic Invertebrate Cell Culture*. Springer, Berlin.
- 579 Mosmann, T., 1983. Rapid colorimetric assay for cellular growth and survival:
580 application to proliferation and cytotoxicity assays. *Journal of Immunological*
581 *Methods* 65, 55-63.
- 582 Pereira, C.S.A., Guilherme, S.I.A.G., Barroso, C.M.M., Verschaeve, L., Pacheco, M.G.,
583 Mendo, S.A.L.V., 2009. Evaluation of DNA Damage Induced by environmental
584 exposure to mercury in *Liza aurata* using the Comet assay. *Archives of*
585 *Environmental Contamination and Toxicology* 58, 112-122.
- 586 Pimentel, D., Greine, A., Bashore, T., 1998. Economic and environmental costs of
587 pesticide use, in: Rose, J. (Ed.), *Environmental Toxicology: Current*
588 *Developments*. Gordon and Breach Science Publishers, UK, pp. 121-187.

- 589 Pluth, J.M., Nicklas, J.A., O'Neill, J.P., Albertini, R.J., 1996. Increased frequency of
590 specific genomic deletions resulting from in vitro malathion exposure. *Cancer*
591 *Research* 56, 2393-2399.
- 592 Pompella, A., Visvikis, A., Paolicchi, A., DeTata, V., Casini, A.F., 2003. The changing
593 faces of glutathione, a cellular protagonist. *Biochemical Pharmacology* 66, 1499–
594 1503.
- 595 Poovala, V.S., Kanji, V.K., Tachikawa, H., Salahudeen, A.K., 1998. Role of oxidant
596 stress and antioxidant protection in acephate induced renal tubular cytotoxicity.
597 *Toxicological Sciences* 46, 403-409.
- 598 Pope, C.N., Liu, J., Karanth, S., 2005. Pharmacology and toxicology of cholinesterase
599 inhibitors: uses and misuses of a common mechanism of action. *Environmental*
600 *Toxicology and Pharmacology* 19, 433-446.
- 601 Preetha, R., Jose, S., Prathapan, S., Vijayan, K.K., Jayaprakash, N.S., Philip, R., Bright
602 Singh, I.S., 2010. An inhibitory compound produced by *Pseudomonas* with
603 effectiveness on *Vibrio harveyi*. *Aquaculture Research* 41, 1452-1461.
- 604 Pruski, A.M., Dixon, D.R., 2002. Effects of cadmium on nuclear integrity and DNA
605 repair efficiency in the gill cells of *Mytilus edulis* L. *Aquatic Toxicology* 57, 127-
606 137.
- 607 Pulido, M.D., Parrish, A.R., 2003. Metal-induced apoptosis: mechanisms. *Mutation*
608 *Research* 533: 227–241.
- 609 Rank, J., Jensen, K., 2003. Comet assay on gill cells and haemocytes from the blue
610 mussel *Mytilus edulis*. *Ecotoxicology and Environmental Safety* 54, 323-329.

- 611 Rannou, M., 1971. Cell cultures of invertebrates other than mollusks and arthropods, in:
612 Vago C., (Ed.), Invertebrate Tissue Culture. Academic Press, New York, pp.385-
613 410.
- 614 Rao, V.J., Kavitha, P., Jakka, N.M., Sridhar, V., Usman, P.K., 2007. Toxicity of
615 organophosphates on morphology and locomotor behavior in brine shrimp,
616 *Artemia salina*. Archives of Environmental Contamination and Toxicology 53,
617 227-232.
- 618 Revankar, P.R., Shyama, S.K., 2009. Genotoxic effects of monocrotophos, an
619 organophosphorous pesticide, on an estuarine bivalve, *Meretrix ovum*. Food and
620 Chemical Toxicology 47, 1618-1623.
- 621 Rinkevich, B., 2005. Marine invertebrate cell cultures: New millennium trends. Marine
622 Biotechnology 7, 429-439.
- 623 Sastre, M., Steinert, S., Streib-Montee, R., 1977. Single cell gel/Comet assay applied to
624 the analysis of pollution-induced damage in the mud welk *Nassarius tegula* and
625 the ribbed mussel *Musculista senhousia*, in: Abstracts of the 18th Annual Meeting
626 of the Society of Environmental Toxicology and Chemistry. Society of
627 Environmental Toxicology and Chemistry, Florida, p. 77.
- 628 Schmid, K., Sassen, A., Staudenmaier, R., Kroemer, S., Reichl, F.X., Harreus, U., Hagen,
629 R., Kleinsasser, N., 2007. Mercuric dichloride induces DNA damage in human
630 salivary gland tissue cells and lymphocytes. Archives of Toxicology 81, 759-767.

- 631 Shimizu, C., Shike, H., Klimpel, K.R., Burns, J.C., 2001. Hemolymph analysis and
 632 evaluation of newly formulated media for culture of shrimp cells (*Penaeus*
 633 *stylirostris*). In Vitro Cellular and Developmental Biology Animal 37, 322-329.
- 634 Shugart, L.R., 1995. Biomarkers of DNA damage, in: de Serres, F.J., Bloom, A.D. (Eds.)
 635 Ecotoxicity and Human Health. Lewis Publishers Inc., Boca Raton, 123-141.
- 636 Shugart, L.R., 1998. Structural damage to DNA in response to toxicant exposure, in:
 637 Forbes, V.E. (Ed.), Genetics and Ecotoxicology. Taylor & Francis, Philadelphia,
 638 pp. 151-168.
- 639 Shugart, L.R., 2000. DNA damage as a biomarker of exposure. Ecotoxicology 9, 329-
 640 340.
- 641 Singh, N.P., McCoy, M.T., Tice, R.R., Schneider, E.L., 1988. A simple technique for the
 642 quantitation of low levels of DNA damage in individual cells. Experimental Cell
 643 Research. 175, 184-191.
- 644 Steinert, S.A., 1999. DNA damage as bivalve biomarker and as an environmental
 645 assessment tool. Biomarkers 4, 492- 496.
- 646 Stohs, S.J., Bagchi, D., 1995. Oxidative mechanisms in the toxicity of heavy metals. Free
 647 Radical Biology and Medicine 18, 321-336.
- 648 Sung, H.H., Kao, W.Y., Su, Y.J., 2003. Effects and toxicity of phthalate esters to
 649 hemocytes of giant freshwater prawn, *Macrobrachium rosenbergii*. Aquatic
 650 Toxicology 64, 25-37.
- 651 Tong, S.I., Miao, H.Z., 1996. Attempts to initiate cell cultures from *Penaeus chinensis*
 652 tissues. Aquaculture 147, 151-157.

- 653 Toullec, J.Y., Crozat, Y., Patrois, J., Porcheron, P., 1996. Development of primary cell
654 cultures from the penaeid shrimps *Penaeus vannamei* and *P. indicus*. Journal of
655 Crustacean Biology 16, 643-649.
- 656 Valverde, M., Trejo, C., Rojas, E., 2001. Is the capacity of lead acetate and cadmium
657 chloride to induce genotoxic damage due to direct DNA-metal interaction?
658 Mutagenesis 16, 265-270.
- 659 Villela, I.V., de Oliveira, I.M., Silveira, J.C., Dias, J.F., Henriques, J.A., da Silva, J.,
660 2007. Assessment of environmental stress by the micronucleus and Comet assays
661 on *Limnoperna fortunei* exposed to Guaíba hydrographic region samples (Brazil)
662 under laboratory conditions. Mutation Research 628, 76–86.
- 663 Wang, C.H., Yang, H.N., Tang, C.Y., Lu, C.H., Kou, G.H., Lo, C.F., 2000. Ultrastructure
664 of white spot syndrome virus development in primary lymphoid organ cell
665 cultures. Diseases of Aquatic Organisms 41, 91-104.
- 666 Wild, D., 1975. Mutagenicity studies on organophosphorus insecticide. Mutation
667 Research 32, 133-150.
- 668 Wilson, J.T., Pascoe, P.L., Parry, J.M., Dixon, D.R., 1998. Evaluation of the Comet assay
669 as a method for the detection of DNA damage in the cells of a marine
670 invertebrate, *Mytilus edulis* L. (Mollusca: Pelecypoda). Mutation Research 399,
671 87–95.

672 Yang, J.M., Arnush, M., Chen, Q.Y., Wu, X.D., Pang, B., Jiang, X.Z., 2003. Cadmium-
673 induced damage to primary cultures of rat Leydig cells. *Reproductive Toxicology*
674 17, 553-560.

675

676

Research Highlights

- Primary haemocyte culture was developed from *Penaeus monodon*.
- Cytotoxicity and genotoxicity determination of aquatic pollutants was conducted.
- IC₅₀ values were calculated using MTT assay.
- All heavy metals and pesticides tested induced DNA strand breakage in haemocytes.

Table 1

DNA damage induced by heavy metals and pesticides in the primary haemocyte culture of *P. monodon*. Mean values and standard deviation obtained from average of 25 cells per experiment (Total 4 experiments for each compound).

Parameters	Control	Cadmium chloride (140 μ M)	Mercuric chloride (17 μ M)	Malathion (60 mg l ⁻¹)	Monocrotophos (186 mg l ⁻¹)
% cells with tail	11.46 $\pm$ 7.89	72.20 $\pm$ 7.46	60.83 $\pm$ 17.72	67.75 $\pm$ 13.55	61.55 $\pm$ 12.98
Tail length (pixels)	1.3 $\pm$ 0.50	33.47 $\pm$ 9.24	23.09 $\pm$ 6.83	29.13 $\pm$ 7.75	28.17 $\pm$ 7.87
% DNA in tail	2.04 $\pm$ 1.25	44.21 $\pm$ 4.76	44.72 $\pm$ 7.25	40.81 $\pm$ 9.10	38.85 $\pm$ 13.68
Tail moment	0.04 $\pm$ 0.03	17.17 $\pm$ 6.23	11.82 $\pm$ 5.50	14.36 $\pm$ 5.56	13.86 $\pm$ 5.74

Fig.1. Primary haemocytes after 12 hour exposure to A) Cadmium chloride (15.63 μM), B) Mercuric chloride (1.95 μM), C) Malathion (61.7 mg l^{-1}), D) Monocrotophos (100 mg l^{-1}), and E) Control.

Fig.2. Toxicity of cadmium chloride in primary haemocyte culture in terms of percentage inhibition of cells determined through MTT assay (n=4).

Fig.3. Toxicity of mercuric chloride in primary haemocyte culture in terms of percentage inhibition of cells determined through MTT assay (n=4).

Fig.4. Toxicity of malathion in primary haemocyte culture in terms of percentage inhibition of cells determined through MTT assay (n=4).

Fig.5. Toxicity of monocrotophos in primary haemocyte culture in terms of percentage inhibition of cells determined through MTT assay (n=4).

Fig.6. Comets formed by haemocytes after exposure to A) Cadmium chloride, B) Mercuric chloride, C) Malathion, D) Monocrotophos, and E) Control.

Fig.1.

Fig.2.

Fig.3.

Fig.4.

Fig.5

Fig.6.