Novel H1N2 swine influenza reassortant strain in pigs derived from the pandemic H1N1/2009 virus

Ana Moreno, Livia Di Trani, Silvia Faccini, Gabriele Vaccari, Daniele Nigrelli, M. Beatrice Boniotti, Emiliana Falcone, Arianna Boni, Chiara Chiapponi, Enrica Sozzi, et al.

► To cite this version:

Ana Moreno, Livia Di Trani, Silvia Faccini, Gabriele Vaccari, Daniele Nigrelli, et al.. Novel H1N2 swine influenza reassortant strain in pigs derived from the pandemic H1N1/2009 virus. *Veterinary Microbiology*, 2011, 149 (3-4), pp.472. 10.1016/j.vetmic.2010.12.011 . hal-00683507

HAL Id: hal-00683507

<https://hal.science/hal-00683507>

Submitted on 29 Mar 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Title: Novel H1N2 swine influenza reassortant strain in pigs derived from the pandemic H1N1/2009 virus

Authors: Ana Moreno, Livia Di Trani, Silvia Faccini, Gabriele Vaccari, Daniele Nigrelli, M. Beatrice Boniotti, Emiliana Falcone, Arianna Boni, Chiara Chiapponi, Enrica Sozzi, Paolo Cordioli

PII: S0378-1135(10)00584-5
DOI: doi:10.1016/j.vetmic.2010.12.011
Reference: VETMIC 5123

To appear in: *VETMIC*

Received date: 21-9-2010
Revised date: 18-11-2010
Accepted date: 7-12-2010

Please cite this article as: Moreno, A., Di Trani, L., Faccini, S., Vaccari, G., Nigrelli, D., Boniotti, M.B., Falcone, E., Boni, A., Chiapponi, C., Sozzi, E., Cordioli, P., Novel H1N2 swine influenza reassortant strain in pigs derived from the pandemic H1N1/2009 virus, *Veterinary Microbiology* (2010), doi:10.1016/j.vetmic.2010.12.011

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Novel H1N2 swine influenza reassortant strain in pigs derived from the pandemic H1N1/2009 virus

Ana Moreno¹, Livia Di Trani², Silvia Faccini¹, Gabriele Vaccari², Daniele Nigrelli¹, M. Beatrice Boniotti¹,
Emiliana Falcone², Arianna Boni², Chiara Chiapponi¹, Enrica Sozzi¹, Paolo Cordioli¹

¹ *Istituto Zooprofilattico Sperimentale della Lombardia e dell'Emilia Romagna (IZSLER), Via Bianchi, 9,
25124-Brescia, Italy*

² *Department of Veterinary Public Health and Food Safety, Istituto Superiore di Sanità, V.le Regina Elena,
299, 0016- Rome, Italy*

* Corresponding author. Mailing address: Laboratory of Virology, Department of Animal Health, IZSLER, Via
Bianchi, 9, 25124 – Brescia (Italy). Phone: +39 030 2290361. Fax: +39 030 2290535. E-mail:
anamaria.morenomartin@bs.izs.it

Abstract

Swine influenza monitoring programs have been in place in Italy since the 1990's and from 2009 testing for the pandemic H1N1/2009 virus (H1N1pdm) virus was also performed on all the swine samples positive for type A influenza. This paper reports the isolation and genomic characterization of a novel H1N2 swine influenza reassortant strain from pigs in Italy that was derived from the H1N1 pdm virus.

In May 2010, mild respiratory symptoms were observed in around 10% of the pigs raised on a fattening farm in Italy. Lung homogenate taken from one pig showing respiratory distress was tested for Influenza type A and H1N1pdm by two real time RT-PCR assays. Virus isolation was achieved by inoculation of lung homogenate into specific pathogen free chicken embryonated eggs (SPF CEE) and applied onto Caco-2 cells and then the complete genome sequencing and phylogenetic analysis was performed from the CEE isolate.

The lung homogenate proved to be positive for both influenza type A (gene M) and H1N1pdm real time RT-PCRs. Virus isolation (A/Sw/It/116114/2010) was obtained from both SPF CEE and Caco-2 cells. Phylogenetic analysis showed that all of the genes of A/Sw/It/116114/2010, with the exception of neuraminidase (NA), belonged to the H1N1pdm cluster. The NA was closely related to two H1N2 double reassortant swine influenza viruses (SIVs), previously isolated in Sweden and Italy. NA sequences for these three strains were clustering with H3N2 SIVs. The emergence of a novel reassortant H1N2 strain derived from H1N1pdm in swine in Italy raises further concerns about whether these viruses will become established in pigs. The new reassortant not only represents a pandemic (zoonotic) threat but also has unknown livestock implications for the European swine industry.

34

Keywords: swine influenza, reassortant H1N2 strain, pandemic H1N1/2009 virus, pig, Italy

36 1- Introduction

37 Influenza type A viruses are currently circulating in the European pig population with high prevalence of
 38 H1N1, H1N2 and H3N2 subtypes. In April 2009 a new virus, identified as pandemic H1N1/2009 strain of
 39 influenza A (H1N1pdm), was reported to have caused a human influenza outbreak in Mexico (Centers for
 40 Disease Control and prevention, 2009, pp. 400-2). By June 2009 the H1N1pdm had rapidly spread among
 41 humans, leading the WHO to the declaration of the first influenza pandemic of the 21st century (WHO, phase
 42 of pandemic alert, 2009). The virus is a quadruple reassortant and contains mostly genes from the triple
 43 reassortant H1N1 viruses found in swine in North America as well as two genes, Neuraminidase (NA) and
 44 Matrix (M) from Influenza viruses found in swine in Europe and Asia (Schnitzer et al., 2009). In Italy, swine
 45 influenza monitoring programs, based on genome detection, virus isolation and sequencing of all respiratory
 46 forms, have been in place since the nineties, especially in North Italy where more than 75% of the swine
 47 industry is located. Responding to the emergence of the new pandemic virus and consequently to the recent
 48 reports on human to animal transmission (Hofshagen et al., 2009; OIE, 2009), testing for the H1N1pdm was
 49 also performed on all the swine samples which resulted positive to influenza type A. In the last two years,
 50 systematic virological surveillance of influenza viruses in pigs has led to the isolation of 67 swine influenza
 51 viruses (SIVs) belonging to the following subtypes: 24 H1N1, 25 H1N2, 18 H3N2 SIVs (Moreno et al., 2010,
 52 pp.73), 2 H1N1pdm viruses (Moreno et al., 2010, pp. 52-56) and one new reassortant strain derived from
 53 H1N1pdm. This paper reports the first isolation of a new H1N2 reassortant strain on a Italian pig farm. The
 54 molecular characterization and phylogenetic analysis of the new reassortant strain revealed that all the
 55 genes except the neuraminidase (NA) belonged to the H1N1pdm viruses and the NA gene was closely
 56 related to two H1N2 SIV strains, previously isolated in Sweden (A/Sw/Sweden/1021/2009) and Italy
 57 (A/sw/Italy/58769/2010).

58 2- Materials and methods

59 2.1 - Clinical samples

60 In May 2010, a clinical outbreak was reported on a pig fattening farm with 4000 heads located in the
 61 province of Mantua (North Italy). The clinical signs observed were coughing, dyspnoea, fever, anorexia and
 62 depression with 10% morbidity, 0,5% mortality. This farm used a continuous flow system of management
 63 with monthly introduction of animals and fortnightly departures to the slaughterhouse. SIV vaccination
 64 program had not been in place at the farm. One dead pig was submitted to the laboratory for respiratory
 65 disease diagnostic tests.

66 2.2 - Influenza type A antigen detection

67 Lung homogenate was screened for the presence of influenza A viruses by M gene real time RT-PCR (rt RT-
 68 PCR) (Spackman et al., 2002). Positive samples were further tested for the H1N1pdm by rt RT-PCR,
 69 according to the CDC procedure (Centers for Disease Control and prevention, 2009).

70 2.3 - Detecting swine respiratory pathogens

71 The lung homogenate was cultured using different media for the most common swine respiratory bacterial
 72 pathogens. The presence of Porcine reproductive and respiratory syndrome virus (PRRSV), Porcine
 73 circoviruses type 2 (PCV2) and *Mycoplasma hyopneumoniae* was determined by using either RT-PCR,
 74 multiplex PCR and PCR assays respectively, as previously described (Calsamiglia et al., 1999; Ouadani et
 75 al., 1999; Suarez et al., 1994).

76 2.4 - Virus isolation and subtype determination

77 The novel SIV isolate was obtained from a 10% lung homogenate applied onto Caco-2 cells and further
 78 inoculated through the allantoic sac route of 9-11 day-old SPF chicken embryonated eggs (CEE). Culture
 79 supernatant (CS) after observed cytopathic effect and allantoic fluid (AF) were tested with haemagglutination
 80 assay (HA) using chicken erythrocytes performed as described (Swine influenza, OIE Manual, 2010) and for
 81 influenza type A and H1N1pdm by real time RT-PCR assays. The subtype of the isolate was determined
 82 from culture supernatant, allantoic fluid as well as from lung tissues by three multiplex RT-PCR assays using
 83 primers to amplify H1, H3, N1 and N2 SIVs (Chiapponi et al., 2003) and primers described by CDC to amplify
 84 fragment 4 of HA gene of H1N1pdm (WHO sequencing primers and protocol, 2009).

85 2.5 - Genome sequencing and phylogenetic analysis

86 For genomic sequencing viral RNA was extracted from allantoic fluid of A/sw/It/116114/2010 using QIAamp
 87 ViralRNA Mini Kit (Qiagen, Hilden, Germany). SuperScript III One-Step RT-PCR System with Platinum Taq
 88 High Fidelity (Invitrogen Carlsbad, CA, USA) was used to perform RT-PCR of 46 amplicons representing the
 89 entire viral genome using the Influenza A H1N1pdm genome Primer Set (Applied Biosystems, Foster City,
 90 CA, USA). Sequencing reactions were performed using BigDye Terminator Cycle Sequencing kit v3.1 and
 91 resolved by an ABI 3130 DNA automatic sequencer. DNA sequences were combined and edited using the
 92 Lasergene sequencing analysis software package (DNASTAR, Madison, WI). Multiple sequence alignments
 93 were made using ClustalW and maximum parsimony phylogenetic trees were created using MEGA4
 94 (Tamura et al., 2007). Each tree is a consensus of 500 bootstrap replicates.

95 The identification as a novel reassortant strain was confirmed by full length sequencing of HA and NA genes
 96 performed directly on lung homogenates.

97 2.6 - Nucleotide sequence accession numbers

98 GenBank accession numbers of the complete genome of A/sw/Italy/116114/2010 were assigned CY067659
 99 to CY067666. GenBank accession numbers of A/sw/Italy/58769/2010 were assigned HM771274 to
 100 HM771279, HQ168025 and HQ168026.

101 3- Results

102 3.1 - Virus isolation and subtype determination

103 Lung homogenates from one pig carcass showing purple areas of consolidation in the apical lobes of the
 104 lungs, interlobular oedema, and fibrinous pleuritis resulted positive for influenza type A and H1N1pdm by real
 105 time RT-PCR assays. The clinical specimen was further inoculated in SPF CEE and onto Caco-2 cells for
 106 virus isolation.

107 The presence of influenza A and the H1N1pdm was detected on AF and Caco-2 CS by real time RT-PCRs.
 108 A multiplex RT-PCR specific for subtype determination was further used to subtype both AF and CS that
 109 unexpectedly resulted to be H1pdmN2. The presence of the H1pdmN2 virus in the lung tissues was also
 110 confirmed by RT-PCR and was the only H-N combination detected in the sample. The isolate was named
 111 A/Sw/It/116114/2010. Respiratory pathogens such as bacteria, *M. hyopneumoniae*, PRRSV or PCV2, which
 112 are involved in the porcine respiratory complex, were not observed. Only a *Streptococcus suis* was isolated
 113 from the lungs and therefore could be the cause of the fibrinous pleuritis.

114 3.2 - Genomic sequencing and phylogenetic analysis

115 HA and NA sequences were obtained from the clinical specimen and compared with the Influenza
 116 sequences available in Genbank, confirming the results of subtype determination obtained by the RT-PCR
 117 and real time RT-PCR assays. Full genome sequencing was also obtained from the virus isolate, confirming
 118 the results for HA and NA obtained from the original sample.

119 Phylogenetic analysis showed that all the genes of A/Sw/It/116114/2010, with the exception of
 120 neuraminidase (NA), belonged to the H1N1pdm clusters. In particular H1 showed more than 99% homology
 121 to H1pdm sequences. The phylogenetic tree of the HA gene shows that A/Sw/It/116114/2010 grouped
 122 together with H1N1pdm sequences in cluster 7, as most of the H1N1pdm human isolates in Italy (Potdar et
 123 al., 2010; Valli et al., 2010) (figure n.1). By contrast, NA was closely related to H1N2 SIV double reassortant
 124 viruses, isolated in Sweden (A/Sw/Sweden/1021/2009) (Bálint et al., 2009) and recently also in Italy
 125 (A/Sw/It/58769/2010) with 93,8% and 95,5% homology respectively. In the phylogenetic tree, the NA gene of
 126 these two H1N2 SIVs grouped together with A/Sw/It/116114/2010 and were located in the cluster of H3N2
 127 SIVs (figure n.2). This Italian strain was isolated in 2010 from piglets with respiratory lesions of SI, raised on
 128 a breeder farm located in North Italy.

Full genome sequencing of A/Sw/It/116114/2010 confirms a very high similarity (from 99-100%) through the seven genes to H1N1pdm circulating in human and animal species. Comparison with the H1N1pdm sequences available in GenBank shows in A/Sw/It/116114/2010 three unique amino-acid (aa) changes respectively in HA (N260K) (H3 numbering) and NP (A85V and G404S) proteins, not yet associated with any well characterized phenotype marker of influenza viruses. Two other aa changes of interest were also found. The first one in PB2 protein (K340N), frequently co-occurred with the potentially virulent HA-D222G mutation (WHO, 2009), although the latter is absent in A/Sw/It/116114/2010. The second aa change is the E67K substitution in NS2 protein, previously correlated with a possible modification of the protein function in viral nucleocapsid export from the nucleus (Wu et al., 2009). Comparison between NA proteins of A/Sw/It/116114/2010 and A/Sw/It/58769/2010 revealed changes in different phylogenetically important regions (PIRs) (Fanning et al., 2000): PIR-A-Y40C and C42R, PIR-F N339H, PIR-G D346N, PIR-K N401D. Combinations of HA and NA subtypes need to be functional to recognize the same type of sialic acid modifications to allow smooth interplay of the two proteins, which is important for the viral cycle (Wagner et al., 2002).

Complete genome sequences of the novel H1N2 strain were compared to those of A/California/4/2009 H1N1pdm and A/Sw/It/58769/2010 H1N2 SIV and the amino acid differences were reported in table n.1.

4 - Discussion

The identification of the new H1N2 reassortant virus demonstrates once again the important role played by pigs as mixing vessels for animal and human influenza, providing a place for reassortment and host adaptation to take place. These results show that the introduction of H1N1pdm in the swine population has provided opportunities for reassortment with the risk that the H1N1pdm virus could alter transmissibility and increase virulence. The success of interspecies transmission of influenza virus depends on the viral gene constellation. Successful transmission between species can follow genetic reassortment with a progeny virus containing a specific gene constellation with the ability to replicate in the new host. The new reassortant virus could have the capability to replicate efficiently in humans, considering its specific gene constellation. This is characterized by 7 genes belonging to H1N1pdm, well-adapted to humans and the NA gene closely related to H1N2 SIVs.

The isolation of H1N1pdm from pigs in different countries also raises further concerns about the possible establishment of H1N1pdm in the swine population. This could lead to a scenario where a co-circulation of influenza A viruses within swine that could act as mixing vessels for the reassortment between influenza viruses of both mammal and avian origin, with unknown implications for public health. All these facts could

have a great impact in the Northern part of Italy where over 70% of swine and 65% of poultry are raised and several epidemics of avian influenza, H5 and H7 subtypes, occurred in the last ten years (Alexander et al., 2007).

The immune selection is not considered important in pigs due to the continuous availability of susceptible pigs. Previous studies reported that human H1N1 and H3N2 viruses could be frequently transmitted to pigs but fail to persist (Van Reeth, 2007). This could be explained by the fact that virus strains with different antigenic characteristics could be disadvantaged compared to the well-adapted established viruses which are circulating among the susceptible population. However, some cases, such as the emergence of the H1N2 subtype in UK with a HA closely related to human H1N1 viruses (Brown et al., 1998) and subsequently spread in the European pig population, suggest that the genes of human viruses may persist after reassortment with one or more influenza viruses in pigs and following adaptation to pigs may often be associated with clinical disease. Moreover, the recent emergence and persistence of reassortant H1N2 strains in Italy since 2003, with a NA closely related to recent H3N2 human viruses (Moreno et al., 2010, pp.73), could be another example supporting these hypotheses.

In the case we are reporting, after the isolation of the reassortant strain the three following groups of animals sent to the slaughterhouse were tested for influenza type A. Nasal swabs, collected from 60 pigs, were tested by M gene real time RT-PCR and resulted negative.

The unusual circulation of H1N1pdm in swine herds worldwide raises concern about the generation and transmission to humans of virus reassortants with key molecular determinants of pathogenicity lacking in the current H1N1pdm (Neumann et al., 2009).

Recently, swine influenza surveillance in Hong Kong has given rise to the isolation of a reassortant virus containing seven genes derived from SIVs circulating in China, and the H1N1pdm NA gene (Vijaykrishna et al., 2010).

These independent isolation events underline the importance of surveillance and complete genetic characterization of influenza viruses isolated in the swine population worldwide.

Conflicts of interest statement

All authors have declared that no conflicts of interest exist.

Acknowledgments

We would like to thanks Dr. Leonardo James Vinco and Manola Adella for excellent technical assistance.

The study was partially funded by Network of Excellence for Epizootic Disease Diagnosis and Control (EPIZONE) an European Union funded research project officially started on 1st of June 2006 and supported by the European Union Sixth Research Framework Programme (<http://www.epizone-eu.net>).

References

- 1 Alexander, D.J., 2007. An overview of the epidemiology of avian influenza. *Vaccine* 25, 5637-44.
- 2 Bálint, A., Metreveli, G., Widén, F., Zohari, S., Berg, M., Isaksson, M., Renström, L.H.M., Wallgren, P., Belák, S., Segall, T., Kiss, I., 2009. The first Swedish H1N2 swine influenza virus isolate represents an uncommon reassortant. *Virol. J.* 6, 180-6.
- 3 Brown, I.H., Harris, P.A., McCauley, J.W., Alexander, D.J., 1998. Multiple genetic reassortment of avian and human influenza A viruses in European pigs, resulting in the emergence of an H1N2 virus of novel genotype. *J. Gen. Virol.* 79, 2947-55.
- 4 Calsamiglia, M., Pijoan, C., Trigo, A., 1999. Application of a nested polymerase chain reaction assay to detect *Mycoplasma hyopneumoniae* from nasal swabs. *J. Vet. Diagn. Invest.* 11, 246-251.
- 5 Centers for Disease Control and Prevention, 2009. Swine influenza A (H1N1) Infection in two children – Southern California, March-April 2009. *MMWR* 2009; 58:400-2.
- 6 Centers for Disease Control and Prevention, 2009. CDC protocol of realtime RTPCR for influenza A (H1N1), rev 2. Oct 2009; [cited 2009 October 6]; Available from <http://www.who.int/csr/resources/publications/swineflu/realtimeptpcr/en/index.html>
- 7 Chiapponi, C., Fallacara, F., Foni, E., 2003. Subtyping of H1N1, H1N2 and H3N2 swine influenza viruses by two multiplex RT-PCR. In Paolo Martelli, Sandro Cavarani, Antonio Lavazza (ed), *PRRS, PMWS, Swine Influenza. 4th International Symposium on emerging and re-emerging pig disease*; Rome, Italy; pp258-259. Available from <http://www.unipr.it/arpa/facvet/dip/dipsa/ric/prrs2003/258-259.pdf>
- 8 Fanning, T.G., Reid, A.H., Taubenberger, J.K., 2000. Influenza A virus neuraminidase: regions of the protein potentially involved in virus-host interactions. *Virology* 276(2), 417-23.
- 9 Hofshagen, M., Gjerset, C.E., Tarpai, A., Brun, E., Dannevig, B., Bruheim, T., Fostad, I.G., Iversen, B., Hungnes, O., Lium, B., 2009. Pandemic influenza A (H1N1) v: human to pig transmission in Norway?. *Euro Surveill* 2009 [serial on the internet], Nov 2009; [cited 2009 November 12]; 14(45) article 5; available from: <http://www.eurosurveillance.org/ViewArticle.aspx?ArticleId=19406>
- 10 Moreno, A., Barbieri, I., Chiapponi, C., Sozzi, E., Foni, E., Zanoni, M.G., Lelli, D., Cordioli, P., 2010. Genetic evolution of Italian H1N2 swine influenza viruses during 1998-2009: presence of new reassortant strains, abstr. p73. In: *Proceedings of the 4nd annual meeting EPIZONE (Network of Excellence for*

- 220 Epizootic Disease Diagnosis and Control) "Need for Speed", 7-10 June, Saint Malo, France; available
221 from <http://www.epizone-eu.net/publicdocs/Downloads/Abstract%20Book%20AM%20France.pdf>
- 222 11Moreno, A., Di Trani, L., Alborali, L., Vaccari, G., Barbieri, I., Falcone, E., Sozzi, E., Puzelli, S., Ferri, G.,
223 Cordioli, P., 2010. First pandemic H1N1 outbreak from a pig farm in Italy. TOVJ. 4, 52-56.
- 224 12Neumann, G., Noda, T., Kawaoka, Y., 2009. Emergence and pandemic potential of swine-origin H1N1
225 influenza virus. Nature 458(7249), 931-939.
- 226 13OIE World Animal Health Information Database: follow up report [serial on the internet]. Influenza A H1N1,
227 Canada: report n.1, June 2009; [cited 2009 June 11]; Argentina: report n.1, July 2009; [cited 2009 July 1];
228 England: report n.1, Sept 2009; [cited 2009 September 25]; Norway: report n.1, Oct 2009; [cited 2009
229 October 13]; China: report n.1, Nov 2009; [cited 2009 November 13];. Available from
230 <http://www.oie.int/wahis/>.
- 231 14Ouardani, M., Wilson, L., Jetté, R., Montpetit, C., Dea, S., 1999. Multiplex PCR for detection and typing of
232 porcine circoviruses. J Clin Microbiol. 37(12), 3917-3924.
- 233 15Potdar, V.A., Chadha, M.S., Jadhav, S.M., Mullick, J., Cherian, S.S., Mishra, A.C., 2010. Genetic
234 characterization of the influenza A pandemic (H1N1) 2009 virus isolates from India. PLoS One 15,5(3),
235 e9693.
- 236 16Schnitzler, S.U., Schnitzler, P., 2009. An update on swine-origin influenza virus A/H1N1: a review. Virus
237 Genes 39, 279-292.
- 238 17Spackman. E., Senne, D.A., Myers, T.J., Bulaga, L.L., Garber, L.P., Perdue, M.L., Lohman, K., Daum,
239 L.T., Suarez, D.L., 2002. Development of a real-time reverse transcriptase PCR assay for type A influenza
240 virus and the avian H5 and H7 hemagglutinin subtypes. J Clin Microbiol. 40(9), 3256-60.
- 241 18Suárez, P., Zardoya, R., Prieto. C., Solana, A., Tabarés, E., Bautista, J.M., Castro, J.M., 1994. Direct
242 detection of the porcine reproductive and respiratory syndrome (PRRS) virus by reverse polymerase chain
243 reaction (RT-PCR). Arch Virol. 135, 89-99.
- 244 19Swine influenza, 15.06.2010 posting date. Chapter 2.8.8, OIE Manual of diagnostic tests and vaccines for
245 terrestrial animals, 2010. Available from:
246 http://www.oie.int/eng/normes/mmanual/2008/pdf/2.08.08_SWINE_INFLUENZA.pdf
- 247 20Tamura, K., Dudley, J., Nei, M., Kumar, S., 2007. MEGA4: Molecular Evolutionary Genetics Analysis
248 (MEGA) software version 4.0. Mol Biol Evol 24(8), 1596-9.

- 249 21Valli, M.B., Meschi, S., Selleri, M., Zaccaro, P., Ippolito, G., Capobianchi, M.R., Menzo, S., 1020.
250 Evolutionary pattern of pandemic influenza (H1N1) 2009 virus in the late phases of the 2009 pandemic.
251 PLoS Curr Influenza, Mar 3.
- 252 22Van Reeth, K., 2007. Avian and swine influenza viruses: our current understanding of the zoonotic risk.
253 Vet res. 38, 243-260.
- 254 23Vijaykrishna, D., Poon L.L.M., Zhu, H.C., Ma, S.K., Li, O.T.W., Cheung, C.L., Smith, G.J.D., Peiris, J.S.M.,
255 Guan, Y., 2010. Reassortment of pandemic H1N1/2009 influenza A virus in swine. Science 18,1529.
- 256 24Wagner, R., Matrosovich, M., Klenk, H.D., 2002. Functional balance between haemagglutinin and
257 neuraminidase in influenza virus infections. Rev Med Viro. 12(3),159-66.
- 258 25WHO, 2009. Preliminary review of D222G amino acid substitution in the haemagglutinin of pandemic
259 influenza A (H1N1) 2009 viruses. Available from:
260 [http://www.who.int/csr/resources/publications/swineflu/cp165_2009_2812_review_d222g_amino_acid_sub](http://www.who.int/csr/resources/publications/swineflu/cp165_2009_2812_review_d222g_amino_acid_substitution_in_ha_h1n1_viruses.pdf)
261 [stitution_in_ha_h1n1_viruses.pdf](http://www.who.int/csr/resources/publications/swineflu/cp165_2009_2812_review_d222g_amino_acid_substitution_in_ha_h1n1_viruses.pdf)
- 262 26WHO, 15.05.2009 posting date. Sequencing primers and protocol. Available from
263 http://www.who.int/csr/resources/publications/swineflu/GenomePrimers_20090512.pdf
- 264 27WHO, 2009. Current WHO phase pf pandemic alert. Available from:
265 http://www.who.int/csr/disease/avian_influenza/phase/en/index.html
- 266 28Wu, R., Zhang, H., Yang, K., Liang, W., Xiong, Z., Liu, Z., Yang, X., Shao, H., Zheng X., Chen, M., Xu, D.,
267 2009. Multiple amino acid substitutions are involved in the adaptation of H9N2 avian influenza virus to
268 mice. Vet Microbiol. 138(1-2), 85-91.
- 269

270 Table n.1 - Amino acid substitutions in HA and internal proteins of H1N1pdm viruses A/California/04/2009 -
 271 A/Sw/It/290272/09 and H1N2 reassortant A/Sw/It/116114/10.

Gene segment	Residue number	A/California/04/2009 (H1N1pdm)	A/Sw/It/290272/09 (H1N1pdm)	A/Sw/It/116114/10 (H1N2 reassortant virus)
PB2	340	K	K	N
	405	S	T	S
PB1	386	K	R	K
	535	I	V	I
	728	V	V	I
	736	K	G	K
PA	14	I	V	V
	224	P	S	S
	256	K	Q	K
HA	100	P	S	S
	214	T	A	A
	220	S	T	T
	277	N	N	K
	391	E	K	E
	544	V	V	I
	85	A	A	V
NP	100	V	I	I
	101	D	N	D
	375	D	D	N
	404	G	G	S
	123	I	V	V
NS1	123	I	V	V
NS2	67	E	E	K

272

273

274 Figure n.1 - Phylogenetic relationship of the HA gene of the swine reassortant isolate
275 (A/sw/Italy/116114/2010 H1N2) with database retrieved pandemic H1N1/2009 viruses. The unrooted tree
276 was created by maximum parsimony method and bootstrapped with 500 replicates. Only bootstrap values
277 higher than 50% are shown.

278

279 Figure n.2 - Phylogenetic relationship of the NA gene of the swine reassortant H1N2 Italian isolate
280 (A/sw/Italy/116114/2010 H1N2) with other Italian SIV viruses and database retrieved H1N2 and H3N2 swine
281 and H3N2 human viruses. The unrooted tree was generated as described in the legend of figure n.1.

Figure n.1 - Phylogenetic relationship of the HA gene of the swine reassortant isolate (A/sw/Italy/116114/2010 H1N2) with database retrieved p H1N1 viruses. The unrooted tree was created by maximum parsimony method and bootstrapped with 500 replicates. Only bootstrap values higher than 50% are shown.

Figure n.2 - Phylogenetic relationship of the NA gene of the swine reassortant H1N2 Italian isolate (A/sw/Italy/116114/2010 H1N2) with other Italian SIV viruses and database retrieved H1N2 and H3N2 swine and H3N2 human viruses. The unrooted tree was generated as described in the legend of figure n.1.