Post mortem development of meat quality as related to changes in cytoskeletal proteins of chickens muscles

Jolanta Tomaszewska-Gras, Jacek Kijowski

► To cite this version:

Jolanta Tomaszewska-Gras, Jacek Kijowski. Post mortem development of meat quality as related to changes in cytoskeletal proteins of chickens muscles. *British Poultry Science*, 2011, 52 (02), pp.189-201. 10.1080/00071668.2011.561281 . hal-00686677

HAL Id: hal-00686677

<https://hal.science/hal-00686677>

Submitted on 11 Apr 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Post mortem development of meat quality as related to changes in cytoskeletal proteins of chickens muscles

Journal:	<i>British Poultry Science</i>
Manuscript ID:	CBPS-2009-280.R1
Manuscript Type:	Original Manuscript
Date Submitted by the Author:	02-Jun-2010
Complete List of Authors:	Tomaszewska-Gras, Jolanta; Poznan University of Life Sciences, Department of Food Quality Management Kijowski, Jacek; Poznan University of Life Sciences, Department of Food Quality Management
Keywords:	Biochemistry, Broilers, Meat, chicken meat ageing , SDS-PAGE & Western blotting, cytoskeletal proteins

SCHOLARONE™
Manuscripts

CBPS-2009-280

Edited Lewis November 2010, MacLeod January 2011

***Post mortem* development of meat quality as related to changes in cytoskeletal proteins of chicken muscles**

J. TOMASZEWSKA-GRAS, F. J.G. SCHREURS¹ AND J. KIJOWSKI

Faculty of Food Science and Nutrition , Poznan University of Life Sciences, Wojska Polskiego 31, 60-624 Poznan, Poland, ¹Institute for Animal Science and Health (ID-DLO), Lelystad, The Netherlands

Running title: Meat quality and cytoskeletal proteins

Correspondence to: Dr J. Tomaszewska-Gras, Faculty of Food Science and Nutrition, Poznan University of Life Sciences, Wojska Polskiego 31, 60-624 Poznan, Poland.
E-mail: gras@up.poznan.pl

Accepted for publication 13th October 2010

Abstract 1. A procedure was developed to separate high and medium molecular weight myofibrillar proteins from chicken muscular tissue with a high resolution by flat bed sodium-dodecyl-sulphate polyacrylamide gel electrophoresis (SDS-PAGE) and subsequent detection by either a general protein stain or Western blotting. These procedures were used to analyse the degradation process of cytoskeletal proteins in chicken breast and leg muscles during meat ageing.

2. This study demonstrates the degradation of all the examined cytoskeletal proteins: titin, nebulin and desmin as well as vinculin, a protein composing the costamere structure. All the examined proteins were found to be degraded during ageing of chicken breast and leg muscles.

3. Degradation of titin, nebulin and desmin started at 3 h *post mortem* in breast muscle. Intact titin and nebulin disappeared within 1 d. Intact desmin and vinculin were not detectable after 3 d *post mortem*. In leg muscle, the degradation process of all the examined proteins evolved much more slowly than in breast chicken muscles.

4. The changes observed in shear force, myofibrillar fragmentation and cooking loss were related to changes in cytoskeletal proteins and used to identify marker proteins or degradation products for the purpose of monitoring the development of meat ageing. The ageing process was faster in breast muscle than in leg muscle.

5. Significant correlations were found between degradation processes of titin, nebulin, and desmin and shear force, as well as myofibril fragmentation index of breast and leg muscles.

INTRODUCTION

The development of meat quality during ageing is dependent on the *post mortem* loss of longitudinal and lateral integrity of the muscular tissue, as a result of proteolytic degradation of cytoskeletal proteins resulting in tenderisation of meat (Young *et al.*, 1980; Quali, 1992). During *post mortem* ageing of meat, the major contractile proteins of muscle, *i.e.* actin and

1
2
3 51 myosin, together comprising approximately 80% of the myofibrillar mass, seem not to be
4
5 52 affected by proteolysis (Samejima and Wolfe, 1976; Yates *et al.*, 1983; Bandman and Zdanis,
6
7 53 1988). The most important structural components involved in meat ageing are considered to
8
9 54 be the cytoskeletal proteins, such as titin, nebulin and desmin (Koohmaraie *et al.*, 1984; Fritz
10
11 55 and Greaser, 1991; Robson, 1995) and vinculin (Taylor *et al.*, 1995). Titin, also called
12
13 56 connectin, is an extremely large monomeric protein found in striated muscle (Maruyama *et*
14
15 57 *al.*, 1977; Wang *et al.*, 1979) which forms a set of longitudinal filaments spanning from the
16
17 58 M-line region to the Z-line. Nebulin is considered to be a long molecule of molecular weight
18
19 59 of approx. 800 kDa which runs parallel to, and in close association with, the thin filaments.
20
21 60 Both titin and nebulin are largely responsible for the longitudinal integrity of the muscular
22
23 61 fibre. Desmin is a protein of approximately 53 kDa, present in intermediate filaments
24
25 62 connecting adjacent myofibrils at their Z-line levels (Robson, 1995) and one of the many
26
27 63 proteins composing the cell membrane skeleton, named costameres (Taylor *et al.*, 1995).
28
29 64 Desmin is considered to be the most important protein responsible for the lateral integrity of
30
31 65 the muscle fibre (Robson, 1995). Vinculin is one of the most important proteins composing
32
33 66 the costamere structure with a molecular weight of approx. 126 kDa and belongs to a chain of
34
35 67 proteins involved in forces of the myofibrils to the anchorites of the muscle (Taylor *et al.*,
36
37 68 1995).
38
39
40
41
42
43
44
45

46 69 The kinetics of *post mortem* proteolytic degradation of these key cytoskeletal proteins is
47
48 70 of crucial importance for meat quality development. In the 1980s, most of the SDS-PAGE
49
50 71 studies were focused on the extremely large proteins, titin and nebulin, and their breakdown
51
52 72 (Lusby *et al.*, 1983; Paterson and Parrish, 1987). Depending on the molecular weight of
53
54 73 myofibrillar proteins, different percentages of total acrylamide and crosslinker have been used
55
56 74 throughout literature to produce gels of various porosities and with good separating
57
58 75 capabilities. For the extremely large myofibrillar proteins, such as titin and nebulin, with
59
60

76 molecular weights of approximately 2800 and 800 kDa, respectively, highly porous
77 polyacrylamide gels have been utilised. These gels are extremely difficult to handle,
78 especially during Western blotting or staining, due to their fragile nature. The gels described
79 in literature for the separation of these high molecular weight proteins, without exception,
80 were of the 'vertical slab-gel type' (Studier, 1973; Porzio and Pearson, 1977; Paterson and
81 Parrish, 1987; Fritz *et al.*, 1989; Granzier and Wang, 1993).

82 Because of the limited sensitivity and separation of proteins by SDS-PAGE, it was
83 necessary to search for a new, more sensitive technique to detect proteins and their
84 degradation products, which are present in the muscle at very low concentrations.
85 Cytoskeletal proteins and some of their degradation products separated on these gels may be
86 detected very sensitively and specifically by Western blotting, first introduced by Burnette
87 (1981), however, chicken myofibrillar proteins are very difficult to separate (Chou *et al.*,
88 1994). Bandman and Zdanis (1988) successfully used Western blotting to study protein
89 breakdown in beef with antibodies to myosin and titin. This immunochemical method has also
90 been used in studies on titin and nebulin breakdown in bovine meat by Fritz and Greaser
91 (1991) and in studies on the breakdown of desmin in chicken muscle by Schreurs (1991).
92 Taylor *et al.* (1995) used Western blotting and electron microscopy to demonstrate that the
93 first change occurring in *post mortem* beef muscle was the disintegration of the N2-line and
94 costameres.

95 Presently, little is known about the changes in cytoskeletal proteins of poultry meat, and
96 so the purpose of this study was to investigate the degradation of cytoskeletal proteins in
97 chicken muscle during *post mortem* ageing. Sample preparation, gel casting and Western
98 blotting were optimised for chicken muscle tissue. The utilised gels were ultra-thin, highly
99 porous polyacrylamide gels on a plastic backing film, and were used in a flat-bed gel-
100 electrophoresis set-up. This procedure resulted in very high-resolution electropherograms for

1
2
3 101 high molecular weight proteins. The procedure was adapted to medium molecular weight
4
5 102 myofibrillar proteins, such as vinculin and desmin, using precast, commercially available
6
7
8 103 SDS-PAGE gels of 7.5 and 12.5 % total acrylamide, resulting in a comparably high
9
10 104 resolution.

11
12 105 In order to be able to relate changes observed in the cytoskeletal proteins to meat quality
13
14 106 characteristics, myofibrillar fragmentation as well as cooking loss and shear force were
15
16
17 107 measured in both breast and leg muscles.

18
19
20 108 MATERIALS AND METHODS

21
22 109 Water was always of Milli-Q (Millipore, The Netherlands) quality. Standard laboratory
23
24 110 chemicals were obtained from Merck and were of analytical grade, unless stated otherwise.
25
26
27 111 Precast ExcelGel SDS Homogeneous 7.5% (no. 80-1260-01) and gradient 8-18% gels,
28
29 112 ExcelGel SDS buffer strips (no. 17-1342-01) and Coomassie brilliant blue were obtained
30
31 113 from Pharmacia. Acrylamide/bisacrylamide and tetraethylenemethylene-diamine (TEMED)
32
33 114 were obtained from BDH. Reinforced nitro-cellulose membranes, pore size 0.45 µm, for
34
35 115 Western blotting were obtained from Schleicher and Schuell.

36
37
38 116 **Animals and sampling techniques**

39
40
41 117 A total of 210 6-week-old Ross commercial broilers were obtained on two occasions from a
42
43 118 local slaughterhouse, transported to the Institute in typical commercial broiler crates, and
44
45 119 subsequently slaughtered at the in-house processing plant of the Institute for Animal Science
46
47 120 and Health (ID-DLO, Lelystad). Birds were stunned in a water bath stunner set at 100 V and
48
49 121 50 Hz for approximately 10 s, and the jugular veins severed with a rotating knife, but were
50
51 122 not scalded, plucked or eviscerated. After an approximate 3-min bleeding time, birds were
52
53 123 skinned and deboned. The initial (0 h) samples for electrophoresis were taken immediately
54
55 124 after excision of the muscles, frozen in liquid nitrogen, and stored at -80°C. Subsequently,
56
57
58 125 skinned breast and leg parts were wrapped in paper towels saturated with an antimicrobial
59
60

solution (10 mM sodium azide, 100 mg/kg chloramphenicol, 10 mg/kg riframycin) and kept in a cold room at 4°C for sampling at 1, 3 and 6 h and further at 1, 2, 3, 5, and 7 d of *post mortem* storage. Samples for SDS-PAGE and Western blotting, as well as for the measurement of the myofibrillar fragmentation index (MFI), were taken repeatedly from the same 9 carcasses at the above mentioned times *post mortem*.

The cooking loss and shear force measurements were carried out at 8 sample times on 12 different carcasses per sampling time (96 animals in total). The experiment described above was repeated two times, which resulted in 210 animals being used in all the measurements: 18 carcasses (9 carcasses x 2 experiments) for SDS- PAGE, Western blotting and MFI, and 192 carcasses (12 carcasses x 8 periods x 2 experiments) for shear force, pH and cooking loss.

Sample preparation

Whole muscle samples (breast, *pectoralis major*, and thigh muscle, *biceps femoris*) for SDS-PAGE and Western Blotting were taken (approximately 2 g) at the above times *post mortem*, frozen in liquid nitrogen and stored at -80°C until analysis. On the day of analysis, samples were again cooled in liquid nitrogen and subsequently pulverised using a liquid nitrogen cooled mortar and pestle. The resulting powder was dissolved into a volume of sample buffer according to Fritz and Greaser (1991), containing 25 mM Tris-HCl pH = 6.8, 8 M urea, 2 M thiourea, 3% (w/v) SDS, 75 mM dithiothreitol (DTT) and bromophenol blue, to yield a protein concentration of 1 mg/ml. Samples were heated for 20 min at 50°C and subsequently cooled in tap water to room temperature and centrifuged at 12000 g for 5 min.

Polyacrylamide gel preparation

Commercially available precast ExcelGels (12.5 x 260 x 0.5 mm) with homogeneous 7.5% and a gradient of 8 to 18% total acrylamide were used. For the separation of high molecular weight proteins, such as titin and nebulin, special gels of 3.2% total acrylamide (bisacrylamide:acrylamide, 1:37) were prepared; 3.2% homogeneous polyacrylamide gels

were cast in gel cassettes of 12.5 x 260 x 1.0 mm with sample wells of 5 x 5 x 1.0 mm (= 25 µl), containing a GelBond PAG plastic backing film obtained from Pharmacia. The casting solution was composed of 3.2 % acrylamide, 0.08 % N, N'-methylene-bisacrylamide, 375 mM Tris.HCl pH 8.8, 0.1% SDS, 0.03 mg/ml ammonium persulphate and 0.4% TEMED. After polymerisation and disassembly of the gel cassettes, the gel was covalently bound to the plastic backing film and could be easily removed from the gel casting moulds and positioned on top of the cooling plate of the Multiphor II electrophoresis unit.

Electrophoresis

Flat bed SDS-PAGE was performed with a horizontal Pharmacia LKB Multiphor II System. Muscle samples were loaded on 7.5% and 12.5 % gels (10 µg of protein) and on 3.2% gel (20 µg of protein). On the 7.5% and 12.5% gels, several aliquots of a mixture of Pharmacia low molecular weight (LMW) and high molecular weight HMW-SDS (Pharmacia, 17-0615-01) calibrators were applied to estimate the molecular weight of different bands. Onto gels used for subsequent western blotting also a mixture of the above mentioned standards was applied as well as samples prepared from muscle tissue that were frozen immediately after slaughter; ExcelGel buffer strips (245 x 4.5 mm) were used as anodic and cathodic buffers. The strip buffer system formed, together with the gel buffer, a discontinuous buffer system. High- and low-molecular weight standards from Pharmacia were used to identify protein molecular weights. Electrophoresis was performed at the constant current of 50 mA with the power supply set at a maximum of 600 V and a maximum power of 30 W. The power supply was equipped with a volt-hour (VH) integrator and electrophoresis was extended until 500 VH per gel had been recorded.

Fixation and staining

Immediately after electrophoresis, gels were immersed in a fixing solution (40% ethanol, 10% acetic acid) for 30 min. Subsequently, gels were incubated in the staining solution of 0.05%

1
2
3 176 Coomassie brilliant blue R-250 (PhastGelBlueR, Pharmacia, no. 17-0518-01) in 25% ethanol,
4
5 177 8% acetic acid for 30 min, after which, gels were destained in the destaining solution (25%
6
7 178 ethanol, 8% acetic acid) until a clear background was obtained. Destained gels were soaked in
8
9 179 the preserving solution (10% acetic acid, 10% glycerol), and then dried covered with a clear
10
11 180 cellophane preserving film.
12

15 181 **Western blotting and detection**

16
17 182 Immediately after electrophoresis, as described above, gels were separated from the backing
18
19 183 film and transferred to a nitro-cellulose membrane, soaked in transfer buffer (25 mM Tris-192
20
21 184 mM glycine, pH = 8.3, 0.1% w/v SDS and 10% v/v methanol). Subsequently, proteins were
22
23 185 transferred from the gel onto the nitro-cellulose membrane by a semi-dry transfer method
24
25 186 using a constant voltage setting of 500 V for 120 min, and then blotted membranes were
26
27 187 incubated with blocking solution (Phosphate buffered saline (PBS) containing 0.1% Tween-
28
29 188 20 and 5% non-fat dry milk) for 30 min at 37°C to saturate protein binding places on nitro-
30
31 189 cellulose. Saturated membranes were later immersed in diluted primary antibody in
32
33 190 incubation buffer (PBS containing 0.1% Tween-20 and 0.1% non-fat dry milk) and incubated
34
35 191 overnight at 4°C in a moist environment. Antibodies and dilutions used were as follows:
36
37 192 Rabbit polyclonal antibody raised against titin at the Institute for Animal Science and Health
38
39 193 diluted 1:2 500
40
41 194 Rabbit polyclonal antibody to desmin, Sigma D-8281, diluted 1:400
42
43 195 Mouse monoclonal antibody to nebulin, Sigma N-9891, diluted 1:400
44
45 196 Mouse monoclonal antibody to vinculin, Sigma V-9131, diluted 1:200.
46
47 197 After overnight incubation, membranes were washed 3 times with incubation buffer to
48
49 198 remove excess antibody. Subsequently, membranes were incubated with a secondary antibody
50
51 199 alkaline phosphatase conjugate, either anti-rabbit (Boehringer Mannheim No. 1214-632;
52
53 200 diluted 1:2 000) or anti-mouse (Sigma A-3688; diluted 1:1 000), for 3 h at room temperature.
54
55
56
57
58
59
60

The detection of alkaline phosphatase, and thus antigen-antibody complexes, was carried out for 5 min using a 0.45 mg/ml solution of 5-bromo-4-chloro-3-indolyl phosphate/nitro blue tetrazolium (Sigma No. B-5655) chromogenic substrate. The reaction was stopped by quickly rinsing the membrane in water.

Densitometry

Gels and blots were scanned on a Hewlett Packard 6100C tabletop scanner with 400 dpi, which is comparable to a resolution of 63.6 µm. Image scans were subsequently processed as densitometer traces on a tabletop computer equipped with the Quantiscan programme from Biosoft. From the analysis of lanes containing calibrator samples, a standard molecular weight calibration curve was prepared. For the analysis of high molecular weight proteins, titin and nebulin, a molecular weight calibration curve was prepared using bands appearing in the fresh muscle sample lane. For this purpose, molecular weight of intact titin was set to 2800 kDa, intact nebulin to 800 kDa, and myosin heavy chain to 212 kDa.

Titin and nebulin contents were calculated by conversion to the known quantity of the internal standard (β-galactosidase) introduced to the meat sample according to the method published by Tomaszewska-Gras *et al.* (2002). A total of 18 gels were run (9 carcasses x 2 experiments). For gel densitometry analysis, a total of 13 (best stained, undisturbed) gels were selected for each carcass from the two experiments.

Myofibrillar fragmentation index (MFI)

Repeated MFI measurements were carried out according to Olson *et al.* (1976). In the MFI method, absorbance was measured in a suspension of isolated myofibrils. This index defines the degree of myofibril fragmentation to shorter fragments during meat ageing. This method consists of the following stages: myofibril extraction, determination of myofibrillar protein concentration and determination of the index. Samples (4 g) were weighed in two replications from each cooled breast (*P. major*) and thigh muscles (three thigh muscles mixed together: *m.*

226 *quadriceps femoris*, *biceps femoris* and *m. semitendinosus*), and 40 ml MFI buffer chilled to
227 2°C were added to each sample. The composition of MFI buffer: 75 mM KCl, 10 mM
228 K₂HPO₄, 5 mM EDTA, 1 mM NaN₃, 1 mM PMSF, 1mM 1,4-disulfanylbuthane-2,3-diol, pH =
229 7.0. Samples were homogenised at a high rotational speed for 30 s and next centrifuged (1000
230 g for 15 min at 2°C). In order to remove sarcoplasmic proteins, the supernatant was decanted
231 and the remaining myofibril precipitate 40 ml MFI buffer added and myofibrils washed. After
232 myofibril precipitate was obtained, 20 ml MFI buffer were added. The suspension was filtered
233 through fine sieves (1 mm mesh size) in order to remove connective tissue. In such prepared
234 myofibril suspension, protein concentration was determined by the biuret method. A total of
235 0.25 ml was collected from the myofibril suspension in two replications and 0.75 ml MFI
236 buffer was added, to which 4 ml biuret reagent were added and the sample mixed. Samples
237 were placed in the dark for 30 min. Simultaneous with sample analyses, a standard curve was
238 prepared from bovine albumin (BSA) at concentrations of 0, 2.5, 5.0, 7.5 and 10 mg/ml.
239 Absorbance was read on a spectrophotometer at a wavelength of 540 nm. Based on the
240 standard curve and the size of absorbance, the protein concentration was determined in each
241 sample. Samples at a protein concentration of 0.5 mg/ml were prepared in two replications for
242 MFI measurements. After samples were mixed, their absorbance was measured at a
243 wavelength of 540 nm. The fragmentation index was calculated by multiplying the result by
244 200.

245 **pH measurement**

246 Changes in hydrogen ion concentration were measured directly in breast and thigh muscles
247 using a stiletto electrode of a CG - 837 pH-meter by Schott.

248 **Cooking loss and shear force**

249 Breast halves and legs designated for shear force measurements were weighed, vacuum sealed
250 and heated to a core temperature of 85°C in a boiling water bath and, immediately after

1
2
3 251 heating, brought back to 25°C in a water bath. Immediately before shear force measurement,
4
5 252 samples were removed from the sealing bag and blotted dry with a paper towel. Subsequently,
6
7
8 253 samples were re-weighed and weight differences before and after heating recorded as cooking
9
10 254 loss.

11
12
13 255 **Shear force measurements**

14
15 256 The cooked and chilled muscle samples were excised with a special scalpel, longitudinally to
16
17 257 the muscle fibres to reach sample dimensions: 1 x 1 x 5 cm. Breast muscle (*P. major*) and
18
19 258 three thigh muscles (*m. quadriceps femoris*, *biceps femoris*, *m. semitendinosus*) were taken for
20
21 259 shear force analysis. Measurements of shear force were carried out on an Adamel Lhomargy
22
23 260 instrument (Division Instruments S.A., Paris, France) with a single-knife Warner-Bratzler
24
25 261 attachment. Shear force values were expressed in N/cm².

26
27
28
29 262 **Statistical analysis**

30
31 263 All results obtained in the two experiments were statistically analysed and no effect of flock
32
33 264 found. Repeated measures ANOVA was carried out on the MFI data using the general linear
34
35 265 models module of the SPSS/Windows® statistical package. For cooking loss and shear force
36
37 266 measurements, a one-way ANOVA was used. Correlation coefficients of the relationships of
38
39 267 MFI, cooking loss and shear force with the optical density classification of the different
40
41 268 proteins bands on the Western blots was carried out with the Spearman rank correlation
42
43 269 module after the classification were converted to scores in the following manner: - = 0, ± = 1,
44
45 270 + = 2, ++ = 3, +++ = 4. Significant differences were identified at *P* < 0.05.

46
47
48
49
50 271 **RESULTS AND DISCUSSION**

51
52
53 272 **SDS-PAGE and Western blotting**

54
55 273 Results of SDS-PAGE and Western blotting analyses are shown in Figures 1, 2, 3 and 4.
56
57 274 Figure 1 shows results of SDS-PAGE for meat proteins on 3.2% (self-made) and Figure 2 on
58
59 275 7.5% (precast) total acrylamide gels. The lower gel concentration was used for the separation

of proteins larger than 200 kDa in size. Bands of titin and nebulin from 7.5% gels were determined quantitatively (Table 1) according to the method published by Tomaszewska-Gras *et al.* (2002). Figures 3 and 4 show results of Western blots from 7.5% total acrylamide gels with electropherograms of meat proteins developed with polyclonal anti-titin and monoclonal anti-nebulin antibodies respectively, and from 12.5% total acrylamide gels with polyclonal anti-desmin and monoclonal anti-vinculin antibodies respectively. Western blotting analysis was performed to examine degradation products of proteins and their molecular weight. Table 2 shows results of semi-quantitative scoring of banding patterns that will be discussed below.

Titin degradation was analysed on the basis of SDS-PAGE with a 3.2% gel (Figure 1), a 7.5% gel (Figure 2) and Western blotting (Figure 3). The results from 3.2 SDS PAGE gel at death showed that titin in breast and leg meat was visible as a closely-spaced doublet T1 and T2; this is in agreement with other studies (Wang, 1982; Kimura *et al.*, 1984). With increasing time *post mortem*, the amount of titin T1 decreased and the amount of titin T2 increased in all analysed samples. In breast meat, this phenomenon was most noticeable 6 h *post mortem*, when the T2 band was much more intensive than in the 0-h sample. This suggests that T2 was raised from the degradation of T1. The T1 to T2 conversion was completed in all samples 24 h *post mortem*, when T1 was no longer detectable, whereas T2 remained present for 7 d. Leg muscle samples show a comparable conversion of T1 into T2. Although the same amounts of protein were applied to gels in case of breast muscle samples, leg muscle samples showed a larger number of smaller titin degradation products, appearing as faint banding patterns between titin and nebulin, than was the case for breast muscle samples.

Tables 1 & 2 and Figures 1, 2, 3 & 4 near here

The analysis of quantitative changes in titin and nebulin cytoskeletal proteins was carried out by SDS-PAGE on 7.5% gel. For gel densitometry analysis, a total of 13 (best stained, undisturbed) gels were selected for each carcass (breast and leg) from the two

1
2
3 301 experiments. The losses of intact titin and nebulin were determined quantitatively using a
4
5 302 known quantity of the internal standard, which had been introduced to each sample of meat
6
7
8 303 (Tomaszewska-Gras *et al.*, 1999, 2002). The analytical results are presented in Table 1. Titin
9
10 304 content decreased significantly in breast muscle after 1 d, whereas it was after 2 and 3 d of
11
12 305 refrigerated storage in thigh muscles. The observation of titin bands on gel revealed that the
13
14 306 disappearance of the T1 form of titin, both in breast and thigh muscles, resulted in a
15
16 307 significant decrease of the titin quantity over the studied period. In case of nebulin, significant
17
18 308 differences in its content were noted in breast muscle after 24 h and in thigh muscles after 48
19
20 309 h of storage. It was also observed on gels that a significant decrease of nebulin content was
21
22 310 associated with the total degradation of its native form of 800 kDa to the form of 600 kDa.
23
24
25
26

27 311 The Western blotting analysis with polyclonal antibodies to titin was used to detect
28
29 312 degradation products of titin. On the titin blot of breast muscle (Figure 3), 5 clearly visible
30
31 313 titin degradation products were identified, referred to as Ta, Tb, Tc, Td and Te. Components
32
33 314 Ta, Tc, Td and Te with molecular weights of approx. 1700, 1100, 900 and 700 kDa
34
35 315 respectively, already appeared early *post mortem* (at 3 h) as very faint bands. The relative
36
37 316 amount of these components increased gradually until the second day *post mortem*, when the
38
39 317 highest concentration was recorded except for Tc, of which the largest relative amounts were
40
41 318 present during the first day after slaughter. The fifth component Tb, with a molecular weight
42
43 319 of approx. 1400 kDa, appeared at 3 d *post mortem* just below titin T2. The T1 form was
44
45 320 already degraded at that time. A more complex pattern of titin degradation products appeared
46
47 321 in leg muscle, with some very distinct bands. At least 6 bands could be identified, designated
48
49 322 Ta, Tb, Tc, Td, Te and Tf, with approximate molecular weights of 1900, 1400, 1100, 900, 700
50
51 323 and 500 kDa respectively. The Ta, as well as the intact titin band, gradually decreased during
52
53 324 the measuring period. All the lower molecular weight bands Tc, Td, Te, and Tf gradually
54
55 325 increased until an intensity plateau was reached after 72 h. No disappearance of degradation
56
57
58
59
60

bands was observed within 7 d *post mortem*, with the exception of the Tc band. It was faintly present from 0 h on and increasing in optical density during aging, but almost disappeared in the 7-d sample. The Tb component appeared in the 24-h sample and remained as a faint band up to 7 d *post mortem*. The presence of 1700 and 400 kDa degradation bands in chicken breast muscle had previously been reported (Kimura *et al.*, 1984), the first probably corresponding to the Ta band. Chou *et al.* (1994) identified a 1200 kDa component in chicken leg meat, but not in breast meat. In our study, corresponding 1100 kDa bands were identified in both muscle types. Tanabe *et al.* (1992) suggested that this component originates from T1 and is composed of the Z-line side part of the T1 molecule.

SDS-PAGE and Western blotting results prepared from 7.5% gels with monoclonal antibodies to nebulin, as shown in Figure 4, demonstrated that the degradation process of nebulin in breast muscle started after 3 h as a small, diffuse smearing in the nebulin band and became more intense with the increasing storage time. The cleavage of nebulin into two bands was observed 6 h after slaughter; intact nebulin with a molecular weight of approximately 800 kDa and an immunoreactive degradation product with a molecular weight of approximately 500 kDa, designed Na. After 72 h of storage, a smaller degradation product, designed Nb, was found on blots with a molecular weight of approx. 400 kDa. After 5 and 7 d of storage, these two bands remained visible. In leg muscle, after 6 h only a small fraction of the original intact nebulin was left. After 24 h *post mortem*, most of the nebulin was degraded into a smaller cleavage product, designed Na, with a molecular weight of 500 kDa. Within 72 h *post mortem*, virtually all nebulin was degraded into smaller, no longer immunoreactive, fragments. Only small amounts of two fragments, the Na fragment and a new cleavage product designed Nb, with an apparent molecular weight of approximately 400 kDa, remained visible. On some blots the latter proteolytic fragment appeared as a closely spaced doublet.

Based on blots, it may be concluded that the breakdown of nebulin was most prominent between 3 and 48 h, when intact nebulin was degraded and disappeared. This fact is consistent with previous reports, where the degradation of nebulin was analysed only by SDS-PAGE (Paxhia and Parrish, 1988; Chou *et al.*, 1994).

Figure 4 shows also the results of Western blotting with polyclonal antibodies to desmin. The results showed that in breast muscle, desmin degradation process started within 3 h storage. Below the native band (53 kDa) of desmin, a faint band of approx. 47 kDa appeared, referred to as component Da. A new band was visible on the blot after 6 h ageing, designated Db, with a molecular weight of approx. 40 kDa. These two bands remained very strong until the second day. The most remarkable degradation of desmin in breast muscle occurred between 6 and 72 h *post mortem*. After this time, intact desmin disappeared completely and only three bands of products, *i.e.* Da (47 kDa), Db (40 kDa) and Db (34 kDa), remained until the seventh day of storage. In leg muscle, desmin degradation was much slower, with only very faint degradation visible after 24 h. At 48 h, the density of the native molecule started to become faint and clearly defined degradation products started to appear. The degradation pathway seems to be equal to that of the breast muscle, although much retarded. In other studies of desmin degradation by Western blotting analysis bovine meat was used (Hwan and Bandman, 1989; Taylor *et al.*, 1995). It has been established that the degradation of desmin in the bovine semimembranosus muscle, a fast-ageing muscle type, occurred between 24 and 72 h, while in bovine *biceps femoris*, a slow-ageing muscle type, desmin degradation was much slower.

Degradation of vinculin was completed in breast muscle within 3 d *post mortem*, when intact vinculin was invisible on Western blots (Figure 4). In studies on vinculin degradation, it was difficult to establish the time when degradation started as a result of the use of a monoclonal antibody. The antibody is too specific to be able to pick up degradation products.

In leg muscle, vinculin degradation was much slower, as is shown by the intact molecule still being present vaguely 7 d *post mortem*. A comparison of chicken muscle to the fast-ageing bovine semimembranosus muscle (Taylor *et al.*, 1995) showed that vinculin disappeared at the same time.

In conclusion, it may be stated that on titin and desmin blots, the intact protein molecule and immunoreactive degradation products were visible, progressing in time *post mortem*. Monoclonal antibodies against nebulin also recognised some breakdown, as it was shown by the decrease in intact protein. The degradation pathway of nebulin generates some degradation products that are still recognised by the monoclonal antibody. The monoclonal anti-vinculin antibody was not capable of detecting degradation products of the native protein. This means that, in order to detect degradation products, the use of polyclonal antibodies is still preferred. Desmin and vinculin bands are not made visible on gels by Coomassie staining. Desmin constitutes about 0.18% of skeletal muscle protein (Pearson and Young, 1989). Vinculin has been calculated to comprise about 0.005% of breast muscle proteins (Maruyama, 1985). Both these proteins are easily visualised on Western blots.

Table 3 near here

Myofibrillar fragmentation index (MFI) during *post mortem* storage

MFI is a measure of the average length of myofibrils and is significantly related to tenderness, *i.e.* with shorter myofibrils the index and meat tenderness values are higher (Taylor *et al.*, 1995). Most studies on MFI have related this index at a specified time *post mortem* and most of them were conducted on beef. In this study we monitored changes in MFI in breast and leg chicken muscles during 7 d of storage. Table 3 shows the results of MFI measurement with the time *post mortem*. In breast muscle (*P. major*), MFI increases during *post mortem* ageing until it reaches plateau values after 24 h. In leg muscles (three muscles: *biceps femoris*, *quadriceps femoris*, *semitendinosus* considered as a whole), MFI increases gradually during

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

the whole *post mortem* period that was investigated. Values recorded in leg muscle are approximately 0.5 the values observed in breast muscle. Differences in absolute MFI values indicate a scatter trend of breast muscle myofibrils different from the scatter trend for leg muscle myofibrils.

The highest increase of MFI value in breast muscle was recorded 24 h after slaughter and reached approximately 100% of the initial value. In thigh muscles, MFI increased by approx. 60% of the initial value after 48 h of storage. This indicates much slower and less intensive myofibrillar degradation in leg than in breast muscles, which may be the effect of smaller amounts of proteolytic enzymes in thigh muscles. These results are consistent with the results of the SDS-PAGE and Western blotting experiments (Figures 1, 2, 3 and 4) and results reported in literature. In beef, MFI increases much more quickly in predominantly white muscles rather than in red muscles (Olson *et al.*, 1976). Similar results were observed by Taylor *et al.* (1995) on beef, where MFI index for the 'fast-aging' *semimenbranosus* muscle was higher than for the 'slow aging' *biceps femoris* muscle. Ultramicroscopic changes related to *post mortem* proteolytic degradation proceed much more quickly in chicken breast muscle than in leg muscle (Hay *et al.*, 1973).

Earlier, it was generally believed that the increase in MFI after slaughter is caused by the weakening of the Z-disk, but electron micrographs (Taylor *et al.*, 1995) of myofibrils used in the MFI assay showed that Z-disks in these myofibrils were intact and that rupture of myofibrils in the MFI assay occurred in the I-band (N2-line) and not at the Z-disk. This suggests that observed *post mortem* fragmentation of myofibrils is due to degradation of titin and nebulin, which are constituents of the N2-line (Funatsu *et al.*, 1990).

Table 4 near here

Development of tenderness during *post mortem* storage of breast and leg muscles

Table 4 shows the results of shear force measurements in different muscles with time *post mortem*. Breast muscles (*P. major*) were tenderer than leg muscles throughout the

investigation period. In breast muscle, the course of shear force changes is different from that in the investigated leg muscles because of differences in fibre type and different amounts of intramuscular connective tissue. It has previously been shown that in leg muscles (*biceps femoris*), thickness of perimysium is about two times higher than in breast muscles (Liu *et al.*, 1996). Within 24 h *post mortem*, breast muscle exhibits its final shear force, however, the maximum shear force is reached at 6 h *post mortem*. The leg muscle shear force proceeds much more slowly, but its maximum is reached much earlier at 3 h *post mortem*. This indicates a faster rigor development in leg muscle than in breast muscle, but a slower *post mortem* ageing in the three leg muscles investigated. This is consistent with the results of the SDS-PAGE experiments (Figures 1, 2, 3 and 4) and the results reported in literature (Olson *et al.*, 1976; Hay *et al.*, 1973; Chou *et al.*, 1994).

In studies on MFI, it was stated that in breast muscle MFI values reached the plateau at 24 h *post mortem*, whereas the fragmentation process in leg muscles was finished at 48 h *post mortem*. Based on the maximum toughness of breast muscle, 95% of all tenderisation occurred 24 h *post mortem*. On the other hand, 81%, 89.5% and 86% of all tenderisation occurred within 48 h of slaughter in three leg muscles (*biceps femoris*, *quadriceps femoris*, *semitendinosus*).

Table 5 near here

Cooking loss during *post mortem* storage

Table 5 shows the results of cooking loss measurements with time *post mortem*. Cooking loss observed in breast muscle gradually increases after slaughter. In leg muscle, cooking loss first increased quickly, but at 6 h *post mortem*, it decreased again. The course of cooking loss changes in leg muscle suggests a relationship with *rigor mortis* development. During the first 6 h after slaughter there is a drop in pH and an increase of shear force, when *rigor mortis* occurs, the myofibrillar proteins are approaching the isoelectric point; this leads to a loss of water holding capacity. However, the gradual increase of cooking loss in breast muscle

1
2
3 450 throughout the investigation period suggests that there is a relationship between cooking loss
4
5 451 and *post mortem* proteolytic breakdown of myofibrillar proteins. It is generally accepted that
6
7
8 452 the cooking loss of meat is inversely correlated with pH, with a range of 5.0 – 6.5 being of
9
10 453 practical importance; any alteration of pH has a large influence on the water holding capacity
11
12 454 of myofibrillar proteins (Pearson and Young, 1989). However, since the pH decline, as well
13
14
15 455 as other *rigor mortis*-related factors, does not play a role in *post rigor* meat, the gradual
16
17 456 increase of cooking loss has to be explained by some other mechanism. The degradation of
18
19
20 457 the myofibrillar structure may play a role in the increase in cooking loss.

21
22 458 **Relationship between the degradation process of cytoskeletal proteins and meat**
23
24 459 **tenderness, MFI and cooking loss**

25
26
27 460 In order to identify possible candidate marker fragments, semi-quantitative scores for relative
28
29 461 intensity of different proteins and cleavage products on Western blots were converted to
30
31 462 numerical scores as described in the Materials and Methods. Subsequently, the Spearman rank
32
33 463 correlation coefficients were calculated for the relationships between these density scores and
34
35 464 MFI, cooking loss and shear force measurements. For leg meat, shear force average was
36
37 465 calculated for the three muscles tested. This was possible because shear forces were corrected
38
39
40 466 for sample size. Table 6 shows the results of these calculations.

Table 6 near here

41
42
43 467 In breast muscle, the degradation of titin as detected on Western blot did not correlate
44
45 468 well with meat quality attributes. The trend for of fragment Tc, however, showed very high
46
47 469 correlation coefficients with both cooking loss and shear force measurements. The decline of
48
49
50 470 optical density of intact desmin as well as the course of optical density of the Dc fragments
51
52 471 showed a high correlation with MFI, cooking loss and shear force, although the correlation of
53
54 472 intact desmin with MFI was lower. The Db fragment showed a high correlation with MFI, but
55
56 473 lower than the non-significant correlations with cooking loss and shear force measurements.
57
58
59 474 The nebulin decline correlated highly with cooking loss and, to a lesser extent, with shear
60

force measurements. Nebulin fragment Na was correlated only with MFI. There was only a 5% correlation fragment of Nb with either cooking loss or shear force. The decrease of vinculin showed no significant correlation with any of the traits.

In leg muscle tissue, titin fragments Ta, Tb, Td, Te and Tf showed significant correlations with MFI, cooking loss and shear force measurements. Only the decrease in the contents of intact titin and the Tc fragment bands showed low correlations. The decrease of intact desmin content and the increase in contents of fragments Db and Dc correlated significantly with the course of changes in meat quality traits. The decrease of intact nebulin content and the increase in fragment Na content correlated significantly with meat quality parameters. The Nb fragment was only significantly correlated with shear force. The decrease in the level of intact vinculin molecules was significantly correlated with MFI and highly correlated with shear force.

Conclusions

It may be stated that the degradation of specific myofibrillar cytoskeletal proteins, as studied in the present work, follows the pattern of *post mortem* tenderisation and, as such, may be utilised as an indicator of meat ageing. Titin, nebulin and desmin degradations follow a course that is consistent with myofibrillar fragmentation and shear force. However, it has to be taken into account that processes involved in *rigor mortis* development, such as pH decrease, may interfere with these relationships. More research is needed to identify specific proteolytic fragments of titin, nebulin and desmin, which can serve as specific markers for the estimation of the degree of proteolysis. Once these components are identified, specific probes may be developed to quickly and easily monitor the course of meat ageing. We used Western blot analysis to identify some possible candidate protein fragments that might prove useful for this purpose. However, since Western blotting is a technique that may only be used to identify qualitative variations between different muscle samples it is unsuitable as a method for

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

measuring quantitatively the course of *post mortem* proteolytic degradation of cytoskeletal proteins. Immunochemical techniques such as ELISA, however, are of a quantitative nature and future research will thus have to be carried out to develop similar methods, possibly in the form of 'dip-stick' methods or bio-sensors, using the above mentioned specific probes, to monitor meat ageing in-line at the processing plant.

ACKNOWLEDGEMENTS

The authors wish to express their gratitude to H. Goedhart, H.G.M. Reimert, C. van Cruijningen and J.K. Waltmann for their expert technical support. The Dutch Poultry and Egg Board financially supported part of this work.

REFERENCES

BANDMAN, E. & ZDANIS, D. (1988) An immunological method to assess protein degradation in post-mortem muscle. *Meat Science*, **22**: 1-19.

BURNETTE, W.N. (1981) "Western Blotting": Electrophoretic transfer of proteins from sodium dodecyl sulfate -polyacrylamide gels to unmodified nitrocellulose and radiographic detection with antibody and radioiodinated protein. *Analytical Biochemistry*, **112**: 195-203.

CHOU, R-G.R., TSENG, T-F., LIN, K-J. & YANG, J-H. (1994) Post-mortem changes in myofibrillar proteins of breast and leg muscles from broilers, spent hens and Taiwanese country chickens. *Journal of the Science of Food Agriculture*, **65**: 297-302.

FRITZ, J.D. & GREASER, M.L. (1991) Changes in titin and nebulin in postmortem bovine muscle revealed by gel electrophoresis, western blotting and immunofluorescence microscopy. *Journal of Food Science*, **56**: 607-610.

FRITZ, J.D., SWARTZ, D.R. & GREASER, M.L. (1989) Factors affecting polyacrylamide gel electrophoresis and electroblotting of high-molecular weight myofibrillar proteins. *Analytical Biochemistry*, **180**: 205-210.

- 524 FUNATSU, T., HIGUCHI, H. & ISHIWAT, S.I. (1990) Elastic filaments in skeletal muscle
525 revealed by selective removal of thin filaments with plasma gelsolin. *Journal of Cell Biology*,
526 **110**: 53-62.
- 527 GRANZIER, H.L.M. & WANG, K. (1993) Gel electrophoresis of giant proteins:
528 Solubilization and silver-staining of titin and nebulin from single muscle fiber segments.
529 *Electrophoresis*, **14**: 56-64.
- 530 HAY, J.D., CURRIE, R.W., WOLFE, F.H. & SANDERS, E.J. (1973) Effect of post mortem
531 aging on chicken muscle fibrils. *Journal of Food Science*, **38**: 981-986.
- 532 HWAN, S-F. & BANDMAN, E. (1989) Studies of desmin and α -actinin degradation in
533 bovine semitendinosus muscle. *Journal of Food Science*, **54**: 1426-1430.
- 534 KIMURA, S., YOSHIDOMI, H. & MARUYAMA, K. (1984) Proteolytic fragments of
535 connectin cause aggregation of myosin filaments but not of actin filaments. *Journal of*
536 *Biochemistry*, **96**: 1947-1950.
- 537 KOOHMARAE, M., KENNICK, W.H., ELGASIM, E.A. & ANGLEMIER, A.F. (1984)
538 Effect of postmortem storage on muscle protein degradation: analysis by SDS-polyacrylamide
539 gel electrophoresis. *Journal of Food Science*, **49**: 292-293.
- 540 LIU, A., NISHIMURA, T. & TAKAHASHI, K. (1996) Relationship between structural
541 properties of intramuscular connective tissue and toughness of various chicken skeletal
542 muscle. *Meat Science*, **43**: 43-49.
- 543 LUSBY, M.L., RIDPATH, J.F., PARRISH, F.C. JR. & ROBSON, R.M. (1983) Effect of
544 postmortem storage on degradation of the myofibrillar protein titin in bovine longissimus
545 muscle. *Journal of Food Science*, **48**: 1787-1795.
- 546 MARUYAMA, K. (1985). Myofibrillar cytoskeletal proteins of vertebrate striated muscle, in:
547 LAWRIE, R. (Ed) *Developments in Meat Science-3*, pp. 25-49 (London, Elsevier Applied
548 Science Publishers).

- 549 MARUYAMA, K., MATSUBARA, S., NATORI, K., NONOMURA, Y., KIMURA, S.,
550 OHASHI, K., MURAKAMI, F., HAND, S. & EGUCHI, G. (1977) Connectin, an elastic
551 protein of muscle. *Journal of Biochemistry*, **82**: 317-337.
- 552 OLSON, D.G., PARRISH JR., F.C. & STROMER, M.H. (1976) Myofibril fragmentation and
553 shear resistance of three bovine muscles during postmortem storage. *Journal of Food Science*,
554 **41**: 1036-1041.
- 555 PATERSON, B.C. & PARRISH JR., F.C. (1987) SDS-PAGE conditions for detection of titin
556 and nebulin in tender and tough bovine muscle. *Journal of Food Science*, **52**: 509-510.
- 557 PAXHIA, J.M. & PARRISH JR., F.C. (1988) Effect of post mortem storage on titin and
558 nebulin in pork and poultry light and dark muscles. *Journal of Food Science*, **53**: 1599-1601.
- 559 PEARSON, A.M. & YOUNG, R.B. (1989) Cytoskeletal and Other Proteins of the Myofibrils,
560 in: PEARSON, A.M. & YOUNG, R.B. (Ed) *Muscle and Meat Biochemistry*, pp. 131-181
561 (San Diego, Academic Press).
- 562 PORZIO, M.A. & PEARSON, A.M. (1977) Improved resolution of myofibrillar proteins with
563 sodium dodecyl sulfate-polyacrylamide gel electrophoresis. *Biochimica et Biophysica Acta*,
564 **490**: 27-34.
- 565 QUALI, A. (1992) Proteolytic and physicochemical mechanism involved in meat texture
566 development. *Biochimie*, **74**: 251-258.
- 567 ROBSON, R.M. (1995) Myofibrillar and cytoskeletal structures and proteins in mature
568 skeletal muscle cells, in: QUALI, A., DEMEYER, D.I. & SMULDERS, F.J. (Eds) *Expression*
569 *of Tissue Proteinases and Regulation of Protein Degradation as Related to Meat Quality*, pp.
570 267-281 (Utrecht, ECCEAMST).
- 571 SAMEJIMA, K. & WOLFE, F.H. (1976) Degradation of myofibrillar protein components
572 during post-mortem ageing of chicken muscle. *Journal of Food Science*, **41**: 250-254.

- 1
2
3 573 SCHREURS, F.J.G. (1991) An immunological method to assess proteolytic breakdown of
4
5
6 574 desmin, a structural myofibrillar protein, in broiler breast meat. *Proceedings of 10th*
7
8 575 *European Symposium on the Quality of Poultry Meat*, Doorwerth pp. 69-76.
9
10 576 STUDIER, F.W. (1973) Analysis of bacteriophage T7 early RNA's and protein on slab gels.
11
12 577 *Journal of Molecular Biology*, **79**: 237-248.
13
14
15 578 TANABE, R., NAKAI, H. & TAKAHASHI, K. (1992) Post mortem changes in skeletal
16
17 579 muscle connectin. *Proceedings of 38th International Congress of Meat Science and*
18
19 580 *Technology*, Vol 3, Clermont-Ferrand pp. 427-432.
20
21
22 581 TAYLOR, R.G., GEESINK, G.H., THOMPSON, V.F., KOOHMARAIE M. & GOLL, D.E.
23
24 582 (1995) Is Z-disk degradation responsible for postmortem tenderization? *Journal of Animal*
25
26 583 *Science*, **73**: 1351-1367.
27
28
29 584 TOMASZEWSKA-GRAS, J., KIJOWSKI, J. & SCHREURS, F. (1999) Titin and nebulin
30
31 585 degradation in relation to myofibrillar fragmentation index and tenderness in broiler meat.
32
33 586 *Proceedings of 14th European Symposium on the Quality of Poultry Meat*, Bologna pp. 143-
34
35 587 150.
36
37
38 588 TOMASZEWSKA-GRAS, J., SCHREURS, F. & KIJOWSKI, J (2002) Quantitative
39
40 589 determination of titin and nebulin in poultry meat by SDS-PAGE with an internal standard.
41
42 590 *Meat Science*, **62**: 61-66
43
44
45 591 WANG, K. (1982) Purification of titin and nebulin. *Methods Enzymology*, **85**: 264-274.
46
47
48 592 WANG, K, MCCLURE, J. & TU, A. (1979) Titin: major myofibrillar components of striated
49
50 593 muscle. *Proceedings of National Academy of Science, U.S.A.*, **72**: 3698-3702.
51
52
53 594 YATES, L.D., DUTSON, T.R., CALDWELL, J. & CARPENTER, Z.L. (1983) Effect of
54
55 595 temperature and pH on the post-mortem degradation of myofibrillar proteins. *Meat Science*,
56
57 596 **9**: 157-162.
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

597 YOUNG, O.A., GRAAFHUIS, A.E. & DAVEY, L. (1980) Post mortem changes in
598 cytoskeletal muscle. *Meat Science*, **5**: 41-55.

For Peer Review Only

Table 1. *Changes in amounts of titin and nebulin determined by 7.5% SDS-PAGE, quantitative method with internal standards (breast, pectoralis major; leg, biceps femoris)*

Time <i>post mortem</i> (h)	Titin content as percentage of total protein content		Nebulin content as percentage of total protein content	
	Breast n = 13	Leg n = 13	Breast n = 13	Leg n = 13
0	5.7±0.49 ^a	5.8±0.84 ^a	1.6±0.14 ^a	1.6±0.33 ^a
1	5.9±0.59 ^a	5.8±0.56 ^{ab}	1.7±0.17 ^a	1.4±0.30 ^a
3	5.8±0.51 ^a	5.5±0.63 ^{ab}	1.7±0.16 ^a	1.3±0.29 ^a
6	5.7±0.77 ^a	5.6±0.48 ^{ab}	1.6±0.24 ^a	1.4±0.40 ^a
24	4.6±0.62 ^b	5.5±0.48 ^{ab}	1.4±0.15 ^b	1.5±0.38 ^b
48	4.4±0.32 ^b	5.1±0.51 ^b	1.3±0.19 ^b	1.1±0.33 ^b
72	4.6±0.39 ^b	4.0±0.56 ^c	1.2±0.15 ^b	0.8±0.23 ^c
120	4.6±0.49 ^b	3.6±0.66 ^c	1.1±0.23 ^c	0.7±0.22 ^c
168	4.3±0.44 ^b	3.5±0.56 ^c	0.9±0.21 ^c	0.6±0.16 ^c

Group averages and standard deviations are shown.

Group averages not sharing a common superscript differ significantly within columns at $P \leq 0.05$.

n = number of samples from the same 13 carcasses.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Table 2. Relative optical densities of bands on Western blots (number of blots n=3) in chicken breast (*P. major*) and leg (*biceps femoris*) muscles

	Fragment	Mw	Time post mortem (h)								
			0	1	3	6	24	48	72	120	168
Breast (<i>P. major</i>)	Titin	2800	++	++	++	++	++	++	++	++	+
	Ta	1700	+	++	++	++	+++	+++	+++	++	++
	Tb	1400	-	-	-	-	-	-	±	++	++
	Tc	1100	+	++	++	++	+	±	-	-	-
	Td	900	+	+	++	++	++	+++	++	++	+
	Te	700	-	±	+	+	++	+++	++	+++	+
	Desmin	53	+++	+++	+++	+++	++	+	±	-	-
	Da	47	-	-	±	+	++	++	+	±	+
	Db	40	-	-	-	±	+	+	++	+	+
	Dc	30-37	-	-	-	-	±	±	+	+	+
	Nebulin	800	++	++	++	+	+	-	-	-	-
	Na	500	-	-	-	±	±	+	++	±	±
	Nb	400	-	-	-	-	-	-	±	+	+
	Vinculin	126	±	+	+	+	+	+	±	-	-
Leg (<i>biceps femoris</i>)	Titin	2800	+	++	++	++	+	+	+	+	±
	Ta	1700	++	++	++	++	++	+	+	+	+
	Tb	1400	-	-	-	-	±	+	+	+	+
	Tc	1100	±	±	±	±	±	+	+	+	±
	Td	900	-	±	±	±	+	+++	+++	+++	+++
	Te	700	±	+	+	+	+	++	++	++	++
	Tf		-	-	-	±	±	+	+	+	+
	Desmin	53	+++	+++	+++	+++	+++	++	+	±	±
	Da	47		±	±	±	±	+	+	+	±
	Db	40	-	-	-	-	-	+	+	+	±
	Dc	30-37	-	-	-	-	-	±	+	+	+
	Nebulin	800	+++	++	++	+	+	-	-	-	-
	Na	500	-	-	-	-	±	+	±	+	±
	Nb	400	-	-	-	-	-	-	±	±	±
	Vinculin	126	++	++	++	++	++	++	+	+	±

Table 3. *Changes in myofibrillar fragmentation index (MFI) with time post mortem in breast (*P. major*) and leg muscles (considered as a whole)*

Time post mortem (h)	MFI	
	Breast (n = 18)	Leg (n = 18)
0	30.0 ± 7.9 ^a	18.0 ± 3.7 ^a
1	35.3 ± 9.0 ^b	20.5 ± 3.6 ^a
3	40.6 ± 10.7 ^c	23.3 ± 3.5 ^b
6	52.6 ± 9.1 ^d	25.1 ± 3.3 ^{bc}
24	62.8 ± 6.3 ^e	26.2 ± 4.5 ^{cd}
48	66.3 ± 4.8 ^e	30.7 ± 5.7 ^e
72	66.0 ± 5.0 ^e	30.1 ± 2.3 ^e
120	65.6 ± 6.8 ^e	28.7 ± 3.8 ^{de}
168	62.9 ± 9.4 ^e	33.5 ± 4.7 ^f

Group averages and standard deviations are shown.

Group averages not sharing a common superscript differ significantly within columns at $P \leq 0.05$.

n = number of muscle samples, measurements in 18 separate muscle samples, each made in duplicate.

Table 4. Shear force in *P. major*, *biceps femoris*, *quadriceps femoris* and *semitendinosus* chicken muscles with time post mortem

Time post mortem (h)	Shear force (N/cm ²)			
	Breast (<i>pectoralis major</i>) (n = 24)	Leg (<i>biceps femoris</i>) (n = 24)	Leg (<i>quadriceps femoris</i>) (n = 24)	Leg (<i>semitendinosus</i>) (n = 24)
0	35.8 ± 5.3 ^c	78.5 ± 11.1 ^c	116.9 ± 18.0 ^a	83.0 ± 19.0 ^b
3	46.6 ± 9.0 ^b	104.5 ± 18.9 ^a	92.6 ± 9.8 ^b	91.9 ± 19.4 ^a
6	93.3 ± 23.2 ^a	89.5 ± 13.5 ^b	95.7 ± 14.5 ^b	71.8 ± 19.2 ^c
24	17.9 ± 6.8 ^d	45.4 ± 6.8 ^d	56.9 ± 12.1 ^c	41.4 ± 8.6 ^d
48	15.8 ± 3.8 ^d	44.3 ± 6.1 ^d	39.1 ± 6.7 ^d	34.7 ± 6.7 ^{de}
72	16.3 ± 5.3 ^d	37.5 ± 5.7 ^e	35.4 ± 6.0 ^{de}	28.5 ± 6.0 ^{ef}
120	15.6 ± 3.0 ^d	34.3 ± 6.7 ^{ef}	32.7 ± 6.4 ^{de}	27.2 ± 3.8 ^{ef}
168	14.3 ± 3.3 ^d	30.0 ± 6.1 ^f	29.9 ± 7.1 ^e	24.6 ± 5.5 ^f

Group averages and standard deviations are shown
Group averages not sharing a common superscript differ significantly within columns at $P \leq 0.05$.
n = number of muscle samples, measurements in 24 separate muscle samples, each made in duplicate.

Table 5. *Changes in pH and cooking loss of breast and leg muscles during post mortem storage at 4°C*

Time <i>post mortem</i> (h)	pH		Cooking loss (%)	
	Breast n = 24	Leg n = 24	Breast n = 24	Leg n = 24
0	6.3±0.10 ^a	6.5±0.16 ^a	4.8±0.96 ^a	8.7±0.88 ^c
3	6.0±0.25 ^b	6.2±0.24 ^{bc}	6.2±1.14 ^b	9.8±1.73 ^d
6	5.6±0.25 ^c	6.1±0.13 ^c	6.9±0.81 ^{bc}	11.5±1.92 ^e
24	5.4±0.13 ^d	6.1±0.25 ^c	7.2±0.96 ^c	8.2±1.31 ^{bc}
48	5.6±0.19 ^{cd}	6.1±0.08 ^c	8.8±1.60 ^d	7.4±1.17 ^{ab}
72	5.5±0.15 ^{cd}	6.2±0.18 ^{bc}	8.8±1.57 ^d	6.9±1.36 ^a
120	5.6±0.14 ^c	6.2±0.18 ^{bc}	9.5±2.02 ^d	7.9±1.52 ^{bc}
168	5.8±0.19 ^e	6.3±0.10 ^{ab}	9.3±1.76 ^d	8.0±1.47 ^{bc}

Group averages and standard deviations are shown.

Group averages not sharing a common superscript differ significantly within columns at $P \leq 0.05$.

n = number of muscle samples, measurements in 24 separate muscle samples, each made in duplicate.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 6. Correlation coefficients determined for dependencies between changes in cytoskeletal proteins and qualitative parameters of poultry meat stored for 7 d at 4°C

Breast meat (<i>P. major</i>)					Leg meat				
SDS- PAGE		MFI	Cooking loss	Shear force	SDS- PAGE		MFI	Cooking loss	Shear force
Western blotting	Titin	-0.910**	-0.861**	0.909**	Western blotting	Titin	-0.790*	0.581	0.880**
	Nebulin	-0.770*	-0.868**	0.780*		Nebulin	-0.877**	0.577	0.922**
	Titin	-0.137	-0.415	0.577		Titin	-0.671*	0.619	0.770*
	Ta	0.597	0.177	0.013		Ta	-0.866**	0.873**	0.873**
	Tb	0.508	0.830*	-0.784*		Tb	-0.913**	-0.913**	-0.913**
	Tc	-0.728*	-0.845**	0.914**		Tc	0.548	-0.845**	-0.507
	Td	0.671*	0.180	0.069		Td	0.940**	-0.856**	-0.894**
	Te	0.898**	0.708*	-0.519		Te	0.913**	-0.809*	-0.861**
	Desmin	-0.796*	-0.963**	-0.945**		Tf	0.935**	-0.823*	-0.926**
	Da	0.713*	0.317	-0.210		Desmin	-0.817**	0.728*	0.932**
	Db	0.896**	0.777*	-0.664		Da	0.671*	-0.782*	-0.574
	Dc	0.802**	0.919**	-0.882**		Db	0.783*	-0.913**	-0.756*
	Nebulin	-0.935**	-0.931**	0.823*		Dc	0.820**	-0.809*	-0.913**
	Na	0.922**	0.617	-0.447		Nebulin	-0.949**	0.805*	0.907**
	Nb	0.508	0.830*	-0.784		Na	0.802**	-0.819*	-0.756*
	Vinculin	-0.173	-0.384	0.546		Nb	0.639	-0.620	-0.845**
						Vinculin	-0.677*	0.550	0.866**

Correlations significant at * $P \leq 0.05$ level (2-tailed), ** $P \leq 0.01$ level (2-tailed).

Figure 1. SDS-PAGE of 3.2% gel of chicken muscles (breast and leg) at 0, 1, 3, 6, 24, 48, 72, 120 and 168 h post mortem and stored at the temperature 4°C.

Figure 2. SDS-PAGE of 7.5% gel of chicken muscles samples (breast and leg) at 0, 1, 3, 6, 24, 48, 72, 120 and 168 h post mortem when stored at 4° C with internal standard (GAL - β -galactosidase) S – marker proteins: myosin (212 kDa), α -macroglobulin (170kDa), β -galactosidaze (116 kDa), transferrin (76 kDa) and glutamic dehydrogenase (53 kDa).

Figure 3. Western blotting of titin at 0, 1, 3, 6, 24, 48, 72, 120 and 168 h post mortem using polyclonal antibody to titin in breast and leg chicken muscles. M – SDS- PAGE of sample at '0' time.

Figure 4. Western blotting of three cytoskeletal proteins at 0, 1, 3, 6, 24, 48, 72, 120 and 168 h post mortem using polyclonal antibody to desmin and monoclonal antibody to nebulin and vinculin in breast and leg chicken muscles. M – SDS- PAGE of sample at '0' time. S - SDS- PAGE of sample with molecular weight marker proteins.