Infant diet and subcutaneous fat mass in early childhood. The Generation R Study

Vincent Jaddoe, Büşra Durmuş, Lamise Ay, Liesbeth Duijts, Henriëtte A Moll, Anita Cs Hokken-Koelega, Hein Raat, Albert Hofman, E Steegers

► To cite this version:

Vincent Jaddoe, Büşra Durmuş, Lamise Ay, Liesbeth Duijts, Henriëtte A Moll, et al.. Infant diet and subcutaneous fat mass in early childhood. The Generation R Study. European Journal of Clinical Nutrition, 2011, 10.1038/ejcn.2011.174 . hal-00687021

HAL Id: hal-00687021

<https://hal.science/hal-00687021>

Submitted on 12 Apr 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Infant diet and subcutaneous fat mass in early childhood.

The Generation R Study

Running title: infant diet and subcutaneous fat

Büşra Durmuş, MSc ^{1,2,3}, Lamise Ay, MD, PhD ^{1,2,3}, Liesbeth Duijts, MD, PhD ^{1,2,3},
Henriëtte A. Moll, MD, PhD ², Anita C.S. Hokken-Koelega, MD, PhD ², Hein Raat, MD, PhD ⁴,
Albert Hofman, MD, PhD ³, Eric A.P. Steegers, MD, PhD ⁵,
Vincent W.V. Jaddoe, MD, PhD ^{1,2,3}

¹ The Generation R Study Group, Erasmus Medical Center, Rotterdam,
The Netherlands;

² Department of Pediatrics, Erasmus Medical Center, Rotterdam, The Netherlands;

³ Department of Epidemiology, Erasmus Medical Center, Rotterdam, The Netherlands;

⁴ Department of Public Health, Erasmus Medical Center, Rotterdam, The Netherlands;

⁵ Department of Obstetrics and Gynecology, Erasmus Medical Center, Rotterdam,
The Netherlands.

Correspondence:

Dr. Vincent. W.V. Jaddoe, MD, PhD, The Generation R Study Group (AE-006),
Erasmus Medical Center, PO Box 2040, 3000 CA Rotterdam, The Netherlands
E-mail: v.jaddoe@erasmusmc.nl; Phone: +31107043405; Fax: +31107044645

Funding: The first phase of the Generation R Study is made possible by financial support from the Erasmus Medical Center, Rotterdam, the Erasmus University Rotterdam and the Netherlands Organization for Health Research and Development. The study sponsors had no role in study design, data analysis, interpretation of data, or writing this report. Vincent Jaddoe received an additional grant from the Netherlands Organization for Health Research and Development (ZonMw 90700303, 916.10159). Büşra Durmuş was supported by an unrestricted grant from Danone Research.

1 **ABSTRACT**

2 **Background:** Breastfeeding has a protective effect on childhood obesity, but the influences
3 on body composition in early childhood are not known.

4 **Objectives:** To assess whether the duration and exclusiveness of breastfeeding, and the
5 timing of introduction of solid foods are associated with the subcutaneous fat mass in early
6 childhood.

7 **Methods:** This study was embedded in a population-based prospective cohort study among
8 779 children. Peripheral (biceps, triceps) and central (suprailiacal and subscapular)
9 subcutaneous fat mass was measured as skinfold thickness at the ages of 1.5, 6 and 24
10 months.

11 **Results:** Breastfeeding duration was not associated with subcutaneous fat mass at the age
12 of 1.5 months. Shorter breastfeeding was associated with higher peripheral and total
13 subcutaneous fat mass at the age of 6 months (p value for trend <0.05), but not at the age of
14 24 months. As compared to children who were exclusively breast fed for 4 months, those
15 who were non-exclusively breast fed had a higher central fat mass at the age of 24 months
16 (p value for trend <0.01). Timing of introduction of solid foods was not associated with
17 subcutaneous fat mass.

18 **Conclusion:** Our results suggest that a shorter duration and non-exclusive breastfeeding
19 affect early body composition during the first 2 years of life. Follow up studies at older ages
20 are needed to explore the long-term consequences.

Keywords: infant feeding, obesity, breastfeeding, solid foods, skinfold, pediatrics

21

22 INTRODUCTION

23 Several studies showed a protective effect of breastfeeding on the risk of overweight in
24 children and adults (Owen *et al.*, 2005; Owen *et al.*, 2005; Harder *et al.*, 2005, Arenz *et al.*,
25 2004). Although the effect estimates for the associations of breastfeeding with mean body
26 mass index are generally small, they seem to be consistent (Owen *et al.*, 2005; Owen *et al.*,
27 2005; Harder *et al.*, 2005; Arenz *et al.*, 2004; Bergmann *et al.*, 2003). An inverse dose-
28 dependent association has been shown, suggesting that longer duration of breastfeeding is
29 associated with a lower body mass index (Harder *et al.*, 2005; Arenz *et al.*, 2004). Some
30 studies have also suggested that early weaning may increase body mass index in childhood,
31 but results are inconsistent (Wilson *et al.*, 1998; Wright *et al.*, 2004; Forsyth *et al.*, 1993;
32 Morgan *et al.*, 2004). Body mass index provides only information about body weight,
33 whereas it does not distinguish between fat and lean mass (Wells and Fewtrell, 2006; Nevill
34 *et al.*, 2006). Since an unfavorable fat distribution may be stronger related to cardiovascular
35 and metabolic diseases, it is important to explore the associations of breastfeeding with
36 measures of fat distribution. Only a few studies have examined the relationships between
37 breastfeeding in infancy and direct measures of adiposity in childhood, but no consistent
38 associations were observed (Burdette *et al.*, 2006; Gale *et al.*, 2007). Skinfold thickness is a
39 valid measurement for subcutaneous fat mass assessment in epidemiological studies (Wells
40 and Fewtrell, 2006; Nevill *et al.*, 2006). These measurements are quick and simple to obtain
41 in most age groups, including young infants (Wells and Fewtrell, 2006). We have previously
42 shown that birth weight is associated with subcutaneous fat mass in early childhood (Ay *et*
43 *al.*, 2009). Subcutaneous fat mass also tends to track throughout early childhood (Ay *et al.*,
44 2009).

45 We examined in a population-based prospective cohort study among 779 children,
46 the associations of breastfeeding duration and exclusivity, and timing of introduction of solid
47 foods with peripheral, central and total subcutaneous fat mass in early childhood.

48

49

50 **METHODS**

51 **Design**

52 This study was embedded in the Generation R Study, a population-based prospective cohort
53 study of pregnant women and their children from fetal life onwards in Rotterdam, the
54 Netherlands (Jaddoe *et al.*, 2010). Enrollment in the study was aimed at early pregnancy
55 (gestational age <18 weeks) but was possible until birth of the child. All children were born
56 between April 2002 and January 2006, and form a prenatally enrolled birth-cohort that is
57 currently followed until young adulthood. Additional detailed assessments of postnatal growth
58 and development were conducted in a subgroup of 1106 Dutch mothers and their children
59 from late pregnancy (Jaddoe *et al.*, 2010). Between February 2003 and April 2005, all
60 pregnant mothers participating in the Generation R study, who met this criterion, were
61 approached for additional measurements (Jaddoe *et al.*, 2010). Of all approached women,
62 80% agreed to participate in the subgroup study. The study protocol was approved by the
63 Medical Ethical Committee of the Erasmus Medical Centre, Rotterdam. Written informed
64 consent was obtained from all parents.

65

66 **Population for analysis**

67 From the total of 1106 children, 1039 children participated in at least one of the postnatal
68 assessments at the ages of 1.5, 6 and 24 months (Figure 1). Children without complete
69 information on breastfeeding (n = 225) were excluded from the analyses. Of the remaining
70 814 live births with complete data on breastfeeding, skinfold measurements were measured
71 in at least one of the three visits in 799 children. Next, twins (n = 20) were excluded from the
72 analyses since twins are correlated and may differ from other children in the relation between
73 breastfeeding and subcutaneous fat mass. Missing skinfold measurements were mainly due
74 to crying behavior. No differences in child and maternal characteristics were found between
75 children with and without skinfold measurements (n = 779 versus n = 1106).

76

Duration and exclusiveness of breastfeeding and solids foods introduction

Information about breastfeeding initiation and continuation was obtained from delivery reports and postal questionnaires at the ages of 2, 6 and 12 months after birth, as previously described (Durmuş *et al.*, 2011). Mothers were asked whether they ever breast fed their child (yes; no) and at what age they quit breastfeeding. Subsequently, breastfeeding duration was categorized into three groups: 1) never; 2) less than 4 months; 3) 4 months or longer.

Duration of exclusive breastfeeding was defined by using information about at what age other types of milk and/or solids were introduced in the first 6 months of life, according to a short food frequency questionnaire. The information about duration and exclusiveness of breastfeeding was combined and categorized into the following three categories: 1) never; 2) non-exclusive breast fed until 4 months; and 3) exclusive breast fed until 4 months. Never indicates infants who were never breast fed. Non-exclusive indicates infants receiving both breastfeeding, and formula feeding or solids during the first four months. Exclusive indicates infants who have been breast fed, without any other milk, solids or fluids during the first four months. Information about timing of introduction solid foods like fruit- and vegetable snacks was obtained from the same short food frequency questionnaire. Starting solid foods was defined as the age that a fruit- or vegetable snack was given for the first time (< 4 months; 4-5 months; and > 5 months).

Subcutaneous fat mass measurements and anthropometrics

Subcutaneous fat mass was measured as skinfold thicknesses (SFT) in millimeters at the ages of 1.5, 6 and 24 months on the left side of the body at four different sites (biceps, triceps, suprailiacal and subscapular) according to standard procedures by using a skinfold caliper (Slim Guide, Creative Health Products) (Lohman *et al.*, 1991). Four well-trained medical assistants performed all measurements (WHO, 2006). The consensus between and among observers for the medical assistants was analyzed using the intraclass correlation coefficient (ICC) (Bland and Altman, 1986; Shrout and Fleiss, 1979). Intraobserver ICC was 0.88 and interobserver ICC was 0.76. Total subcutaneous fat mass was calculated from the

sum of biceps SFT + triceps SFT + suprailiacal SFT + subscapular SFT. Central subcutaneous fat mass was calculated from the sum of suprailiacal SFT + subscapular SFT. Peripheral subcutaneous fat mass was calculated from the sum of triceps SFT + biceps SFT (Ketel *et al.*, 2007; Birmingham *et al.*, 1993). Body length at the age of 1.5 and 6 months was measured in supine position to the nearest millimeter using a neonatometer and in 24-month olds height was measured in standing position by a Harpenden stadiometer (Holtain Limited, Dyfed, U.K.). Weight was measured to the nearest grams in naked infants at the age of 1.5 and 6 months by using an electronic infant scale (SECA) and in 24-month olds by a mechanical personal scale (SECA).

Covariates

Gestational age at birth, birth weight and sex were obtained from midwife and hospital registries at birth. Information about highest attained maternal educational level and parity were obtained at enrollment in the study. Educational level of the parents was defined according to the classification of Statistics Netherlands (Statistics Netherlands, 2004). Information on maternal smoking during pregnancy (yes; no) was retrieved from the prenatal questionnaire. Maternal height and weight were measured at enrollment while the mother stood without shoes and heavy clothing, and body mass index was calculated (kg/m^2).

Statistical analysis

Differences in baseline characteristics between the breastfeeding variables ever (never; ever), duration (< 4 months; >4 months) and exclusivity (non-exclusive breast fed until 4 months; exclusively breastfed until 4 months) were compared with Student's t-test for continuous variables and Chi-square tests for categorical variables. The associations of breastfeeding (never; ever), breastfeeding duration (never; < 4 months; and > 4 months), breastfeeding exclusivity (never; non-exclusive until 4 months; and exclusive until 4 months), and timing of introduction of fruit-and vegetable snacks (< 4 months; 4- 5 months; and > 5 months) with peripheral, central, and total subcutaneous fat mass at the ages of 1.5, 6 and 24 months, were assessed using multiple linear regression models. The reference groups of these regression models were ever breast fed, breastfeeding duration of >4 months, exclusive breast fed until 4 months, respectively. In these models, breastfeeding categories were used as predictor variables and peripheral, central and total fat as outcome variables. The models were adjusted for potential confounders including child's age at visit, sex, maternal education, maternal body mass index, smoking and parity, gestational age, birth weight, current height and observer. The models focused on the role of timing of introduction of solid foods, were additionally adjusted for breastfeeding duration. Tests for trends were performed by treating each categorized variable as a continuous term and by entering the variable into the fully adjusted linear regression model. To handle missing values in covariates, we performed multiple imputations by generating 5 independent datasets for all analyses using the Markov Chain Monte Carlo method. Imputations were based on the relationships between all covariates included in this study. All measures of association are presented with their 95% confidence intervals (CI). Statistical analyses were performed using the Statistical Package of Social Sciences version 17.0 for Windows (SPSS Inc., Chicago, IL, USA).

Results

Subject characteristics are given in Table 1. Of the total group of 779 children, 87.9% had ever been breastfed with a mean duration of 4.5 months (range 0.5 – 12.0). Differences between the breastfeeding categories are given in the Supplementary material (Tables S1, S2, S3). As compared to mothers who never breast fed their children, those who breast fed their children had more frequently a higher educational level, were more likely to consume alcohol during pregnancy but less likely to smoke (Table S1). Mothers who breast fed their children for more than 4 months, compared to mothers who breast fed their children shorter than 4 months, were older and had a lower body mass index (Table S2). Mothers who exclusively breast fed their children had a lower weight, were more likely to have more than one child and had children with a higher birth weight (Table S3).

The associations of ever breastfeeding, breastfeeding duration and exclusivity with peripheral, central and total fat mass at the ages of 1.5, 6 and 24 months are shown in Table 2. Breastfeeding duration was not associated with subcutaneous fat mass at the age of 1.5 months. Shorter breastfeeding was associated with higher peripheral and total subcutaneous fat mass at the age of 6 months (p value for trend <0.05), but not at the age of 24 months. As compared to children who were breast fed exclusively for 4 months, those who were never breast fed had a higher peripheral and total subcutaneous fat mass at the age of 6 months and higher central fat mass at the age of 24 months (p value for trend <0.01).

Table 3 presents the associations between age at introduction of solid foods and peripheral, central and total fat mass at the ages of 1.5, 6 and 24 months. As compared to children who received solid foods after the age of 5 months, those receiving solid foods before the age of 5 months, tended to have increased subcutaneous fat mass measures, but these associations were not significant.

DISCUSSION

Main findings

Results from this study suggest that a shorter duration and non-exclusive breast fed affect early body composition during the first 2 years of life. A shorter duration of breastfeeding was associated with higher peripheral and total subcutaneous fat mass at the age of 6 months, whereas non-exclusive breast fed was associated with higher central subcutaneous fat mass at the age 24 months. Early introduction of solid foods was not associated with subcutaneous fat mass measures.

Strengths and weaknesses

The most important strengths of this study are the population-based prospective design with a relatively large number of subjects being studied from early pregnancy onwards, and information about a large number of potential confounders available. Our analyses were based on 799 children with skinfold thickness measurements. Furthermore, information was available about duration and exclusivity of breastfeeding. Some methodological issues need to be considered. Of the total group of 1039 children, breastfeeding information was available in 78%. This non-response would lead to biased effect estimates if the associations of breastfeeding duration and exclusivity with skinfold thickness measurements would be different between those included and not included in the analyses. This seems unlikely because biased estimates in cohort studies mainly arise from loss to follow-up rather than from non-response at baseline (Nohr *et al.*, 2006). Information about breastfeeding was prospectively collected by questionnaires without direct reference to any skinfold measurement. Although assessing breastfeeding by questionnaires seems to be a valid method, misclassification may occur (Li *et al.*, 2005). We estimated breastfeeding exclusivity according to whether the child received breastfeeding without any other infant formula, milk or solids according to the short food frequency questionnaire. This definition does not cover the strict criteria used by the World Health Organization (WHO), which suggest that even the

use of water in combination with breastfeeding does not fulfill the definition of exclusivity. However, we did ask for the most commonly introduced solids and fluids. Furthermore, in The Netherlands it is not common that children receive breastfeeding in combination with the use of water to prevent dehydration. Therefore, we consider our measurement of exclusive breast fed as a good proxy for exclusive breast fed according to the World Health Organization criteria. Our definition of solid foods included only fruit or vegetable snacks. However in the first months of life it is not likely that other products were introduced. We created our specific duration, exclusivity and solid foods categories based on the collected data and growth measurements. These categories do not enable direct comparison with the widely used categories (6 months) of the WHO. Furthermore, our study group was ethnically homogenous and the mothers were highly educated. We were not able to assess the effect of breastfeeding on subcutaneous fat mass development in children with different ethnic and social backgrounds. Finally, we used skinfold thickness as a measure of subcutaneous fat mass because of the limited use of body mass index as a direct measure of adiposity in early childhood. Skinfold thickness provide a simple, easy, and quick yet highly informative assessment of regional fatness in most age groups and can be used in large-scale epidemiological studies (Wells and Fewtrell, 2006). In general, intraobserver and interobserver error are low compared to between-subject variability, but in obese children accuracy and precision are poorer (Wells and Fewtrell, 2006; Nevill *et al.*, 2006). Furthermore, skinfold thickness has a limitation in assessing lean and fat mass of the whole body (Wells and Fewtrell, 2006).

Comparison with other studies

Several studies have shown that breastfeeding is associated with a lower risk of later overweight and obesity (Owen *et al.*, 2005; Owen *et al.*, 2005; Harder *et al.*, 2005, Arenz *et al.*, 2004). Body mass index is a poor outcome due to its low predictive value and lack of information about fat distribution (Wells *et al.*, 2007). Studies focusing on the association between breastfeeding and body composition instead of body mass index did show

inconsistent associations (Burdette *et al.*, 2006; Gale *et al.*, 2007). This may be due to differences in body measurements, ages and samples sizes. A study among adult males from Brazil did not show an association between breastfeeding and adult body fat, measured by skinfolds and fat mass using a bio-impedance scale (Victoria *et al.*, 2003), while a large study in the UK reported a negligible protective effect of breastfeeding duration for more than 6 months on mean body fat measured with Dual Energy X-ray Absorptiometry (DEXA) in children aged 9-10 years (Toschke *et al.*, 2007). Any association was attenuated after adjustment for confounders. One study in Southampton reported a graded association between shorter breastfeeding duration and higher DXA- derived fat mass in children aged 4 years (Robinson *et al.*, 2009). In contrast, two studies, assessing the associations of breastfeeding and direct measures of body composition using DEXA at ages of 2 and 5 years, did not show any association (Burdette *et al.*, 2006; Tulldahl *et al.*, 1999). We used skinfold thickness as measures of subcutaneous fat mass at younger ages, but the results are in line with these previous studies. Only at the age of 6 months we found that shorter breastfeeding duration leads to higher peripheral and total fat mass. Our study was the first study that used both breastfeeding duration and exclusivity, to examine the effect on subcutaneous fat mass. We showed that non- exclusive breast fed was associated with higher peripheral and total subcutaneous fat mass at the age of 1.5 months and higher central fat mass at the age of 24 months. We cannot explain these differences. However, central fat mass might be stronger related to adverse cardiovascular and metabolic health outcomes (Fox *et al.*, 2007; Pischon *et al.*, 2008). Although the effect estimates were small, exclusive breast fed might affect subcutaneous fat mass development in early childhood.

The European Society for Pediatric Gastroenterology, Hepatology and Nutrition (ESPGHAN) Nutrition Committee recommended in 2008 that complementary foods (solid foods and liquids other than breastfeeding or formula) should not be introduced before 4 months and not later than 6 months (Agostoni *et al.*, 2008). Many studies in industrialized countries showed non-adherence to these recommendations (Noble and Emmett, 2006; Scott *et al.*, 2009). In our Dutch study population, we found that most mothers introduced

solid foods mainly after the age of 4 months. There is conflicting evidence about the relation between timing of introduction of solid foods and adiposity in childhood. A few studies did not show differences in adiposity between early and delayed introduction of solid foods (Kramer, 1981; Zive *et al.*, 1992). One recent study suggested that a diet based on fruit, vegetables, and home-prepared foods in the first year of life is associated with a higher lean and lower fat mass measured with DXA at the age of 4 years (Robinson *et al.*, 2009). The same study group reported a greater gain in both weight and skinfold thickness between 6 and 12 months among infants who received the same diet with fruit, vegetables and home-prepared foods at the age of 6 months (Baird *et al.*, 2008). We did not observe significant associations between early introduction of solid foods and subcutaneous fat mass measures.

Conclusion

Our results suggest that a shorter duration and non-exclusive breast fed affect early body composition during the first 2 years of life. Early introduction of solid foods was not associated with subcutaneous fat mass measures. Follow-up studies are needed to assess whether breastfeeding duration and exclusivity affect subcutaneous fat mass and other measures of body composition at older ages.

Acknowledgments

The Generation R Study is conducted by the Erasmus Medical Center in close collaboration with the School of Law and Faculty of Social Sciences of the Erasmus University Rotterdam, the Municipal Health Service Rotterdam area, Rotterdam, the Rotterdam Homecare Foundation, Rotterdam and the Stichting Trombosedienst and Artsenlaboratorium Rijnmond (STAR), Rotterdam. We gratefully acknowledge the contribution of participating mothers, general practitioners, hospitals, midwives and pharmacies in Rotterdam.

Conflict of interest

The authors declare no conflict of interest.

REFERENCES CITED

- Agostoni C, Decsi T, Fewtrell M, Goulet O, Kolacek S, Koletzko B et al; ESPGHAN Committee on Nutrition. (2008). Complementary feeding: a commentary by the ESPGHAN Committee on Nutrition. *J Pediatr Gastroenterol Nutr* **46**, 99-110.
- Arenz S, Rückerl R, Koletzko B, von Kries R (2004). Breast-feeding and childhood obesity--a systematic review. *Int J Obes Relat Metab Disord* **28**, 1247-56.
- Ay L, Van Houten VA, Steegers EA, Hofman A, Witteman JC, Jaddoe VW et al. (2009). Fetal and postnatal growth and body composition at 6 months of age. *J Clin Endocrinol Metab* **94**, 2023-30.
- Ay L, Hokken-Koelega AC, Mook-Kanamori DO, Hofman A, Moll HA, Mackenbach JP et al. (2008). Tracking and determinants of subcutaneous fat mass in early childhood: the Generation R Study. *Int J Obes (Lond)* **32**, 1050-9.
- Baird J, Poole J, Robinson S, Marriott L, Godfrey K, Cooper C et al; Southampton Women's Survey Study Group. (2008). Milk feeding and dietary patterns predict weight and fat gains in infancy. *Paediatr Perinat Epidemiol* **22**, 575-86.
- Bergmann KE, Bergmann RL, Von Kries R, Böhm O, Richter R, Dudenhausen JW et al. (2003). Early determinants of childhood overweight and adiposity in a birth cohort study: role of breast-feeding. *Int J Obes Relat Metab Disord* **27**, 162-72.
- Birmingham B, Dyer AR, Shekelle RB, Stamler J (1993). Subscapular and triceps skinfold thicknesses, body mass index and cardiovascular risk factors in a cohort of middle-aged employed men. *J Clin Epidemiol* **3**, 289-302.
- Bland JM, Altman DG (1986). Statistical methods for assessing agreement between two methods of clinical measurement. *Lancet* **1**, 307-10.
- Burdette HL, Whitaker RC, Hall WC, Daniels SR (2006). Breastfeeding, introduction of complementary foods, and adiposity at 5 y of age. *Am J Clin Nutr* **83**, 550-8.
- Durmuş B, van Rossem L, Duijts L, Arends LR, Raat H, Moll HA et al. (2011). Breast-feeding and growth in children until the age of 3 years: the Generation R Study. *Br J Nutr* **31**, 1-9.
- Forsyth JS, Ogston SA, Clark A, Florey CD, Howie PW (1993). Relation between early introduction of solid food to infants and their weight and illnesses during the first two years of life. *BMJ* **306**, 1572-6.

317 Fox CS, Massaro JM, Hoffmann U, Pou KM, Maurovich-Horvat P, Liu CY et al. (2007). Abdominal
 318 visceral and subcutaneous adipose tissue compartments: association with metabolic risk
 319 factors in the Framingham Heart Study. *Circulation* **116**, 39-48.
 320 Gale CR, Javaid MK, Robinson SM, Law CM, Godfrey KM, Cooper C (2007). Maternal size in
 321 pregnancy and body composition in children. *J Clin Endocrinol Metab* **92**, 3904-11.
 322 Harder T, Bergmann R, Kallischnigg G, Plagemann A (2005). Duration of breastfeeding and risk of
 323 overweight: a meta-analysis. *Am J of Epidemiol* **163**, 397-403.
 324 Jaddoe VW, van Duijn CM, van der Heijden AJ, Mackenbach JP, Moll HA, Steegers EA et al. (2010).
 325 The Generation R Study: design and cohort update 2010. *Eur J Epidemiol* **25**, 823-41.
 326 Ketel IJ, Volman MN, Seidell JC, Stehouwer CD, Twisk JW, Lambalk CB (2007). Superiority of
 327 skinfold measurements and waist over waist-to-hip ratio for determination of body fat
 328 distribution in a population-based cohort of Caucasian Dutch adults. *Eur J Endocrinol* **6**, 655-
 329 61.
 330 Kramer MS (1981). Do breast-feeding and delayed introduction of solid foods protect against
 331 subsequent obesity? *J Pediatr* **98**, 883-7.
 332 Li R, Scanlon KS, Serdula MK (2005). The validity and reliability of maternal recall of breastfeeding
 333 practice. *Nutr Rev* **63**, 103-10.
 334 Lohman TG, Roche AF, Martorell R (1991). Anthropometric Standardization Reference Manual:
 335 Abridged Edition. *Human Kinetics Books: Champaign, IL*.
 336 Morgan JB, Lucas A, Fewtrell MS (2004). Does weaning influence growth and health up to 18
 337 months? *Arch Dis Child*. **89**, 728-33.
 338 Nevill AM, Stewart AD, Olds T, Holder R (2006). Relationship between adiposity and body size reveals
 339 limitations of BMI. *Am J Phys Anthropol* **129**, 151-6.
 340 Noble S, Emmett P (2006). Differences in weaning practice, food and nutrient intake between
 341 breast- and formula-fed 4-month-old infants in England. *J Hum Nutr Diet* **19**, 303-13.
 342 Nohr EA, Frydenberg M, Henriksen TB, Olsen J (2006). Does low participation in cohort studies
 343 induce bias? *Epidemiology* **17**, 413-18.
 344 Owen CG, Martin RM, Whincup PH, Davey-Smith G, Gillman MW, Cook DG (2005). The effect of
 345 breastfeeding on mean body mass index throughout life: a quantitative review of published
 346 and unpublished observational evidence. *Am J Clin Nutr* **82**, 1298-1307.

347 Owen CG, Martin RM, Whincup PH, Smith GD, Cook DG (2005). The effect of infant feeding on the
 348 risk of obesity across the life course: a quantitative review of published evidence. *Pediatrics*
 349 **115**, 1367-1377.

350 Pischon T, Boeing H, Hoffmann K, Bergmann M, Schulze MB, Overvad K et al. (2008). General and
 351 abdominal adiposity and risk of death in Europe. *N Engl J Med* **359**, 2105-2120.

352 Robinson SM, Marriott LD, Crozier SR, Harvey NC, Gale CR, Inskip HM et al; Southampton Women's
 353 Survey Study Group. (2009). Variations in infant feeding practice are associated with body
 354 composition in childhood: a prospective cohort study. *J Clin Endocrinol Metab* **94**, 2799-805.

355 Scott JA, Binns CW, Graham KI, Oddy WH (2009). Predictors of the early introduction of solid foods in
 356 infants: results of a cohort study. *BMC Pediatr* **22**, 9:60.

357 Shrout PE, Fleiss JL (1979). Intraclass correlations: uses in assessing rater reliability. *Psychol Bull*
 358 **86**, 420-8.

359 Standaard onderwijsindeling 2003. (2004) Voorburg/ Heerlen: Statistics Netherlands.

360 Toschke AM, Martin RM, von Kries R, Wells J, Smith GD, Ness AR (2007). Infant feeding method and
 361 obesity: body mass index and dual-energy X-ray absorptiometry measurements at 9-10 y of
 362 age from the Avon Longitudinal Study of Parents and Children (ALSPAC). *Am J Clin Nutr* **85**,
 363 1578-85.

364 Tulldahl J, Pettersson K, Andersson SW, Hulthén L (1999). Mode of infant feeding and achieved
 365 growth in adolescence: early feeding patterns in relation to growth and body composition in
 366 adolescence. *Obes Res* **7**, 431-7.

367 Victora CG, Barros F, Lima RC, Horta BL, Wells J (2003). Anthropometry and body composition of 18
 368 year old men according to duration of breast feeding: birth cohort study from Brazil. *BMJ*
 369 **327**, 901.

370 Wells JC, Fewtrell MS (2006). Measuring body composition. *Arch Dis Child* **91**, 612-7.

371 Wells JC, Chomtho S, Fewtrell MS (2007). Programming of body composition by early growth and
 372 nutrition. *Proc Nutr Soc* **66**, 423-34.

373 WHO Multicentre Growth Reference Study Group (2006). Reliability of anthropometric measurements
 374 in the WHO Multicentre Growth Reference Study. *Acta Paediatr* **450**, 8-46.

375 Wilson AC, Forsyth JS, Greene SA, Irvine L, Hau C, Howie PW (1998). Relation of infant diet to
 376 childhood health: seven year follow up of cohort of children in Dundee infant feeding study.

377 *BMJ* **316**, 21-5.

378 Wright CM, Parkinson KN, Drewett RF (2004). Why are babies weaned early? Data from a

379 prospective

380 population based cohort study. *Arch Dis Child* **89**, 813-6.

381 Zive MM, McKay H, Frank-Spohrer GC, Broyles SL, Nelson JA, Nader PR (1992). Infant-feeding

382 practices and adiposity in 4-y-old Anglo- and Mexican-Americans. *Am J Clin Nutr* **55**, 1104-8.

383

Legend for Figure 1

Flow chart of participants in study

Table 1 Subject characteristics ^a

	Total (n = 779)
Maternal characteristics	
Age (years)	32.3 (24.9 – 37.9)
Height (cm)	170.8 (6.3)
Weight (kg)	72.0 (13.4)
Body Mass Index (kg/m ²)	24.7 (4.4)
Highest educational level (%)	
Low	1.8
Moderate	34.5
High	63.7
Smoking in pregnancy (%)	
Ever	22.2
Never	77.8
Alcohol consumption in pregnancy (%)	
Ever	70.1
Never	29.9
Parity (%)	
0	62.5
≥1	37.5
Birth characteristics	
Males (%)	51
Gestational age (weeks)	40.3 (37.1 – 42.1)
Weight (grams)	3515 (549)
Small for gestational age (<5%)	4.9

Low birth weight (<2500 g) (%)	3.6
Preterm birth (%)	4.1
Breastfeeding	
Ever (%)	87.5
Duration (months)	4.5 (0.0 – 12.0)
Solid foods (fruit- or vegetables)	
< 4 months introduced (%)	7.7
> 4-5 months introduced (%)	60.4
> 5 months introduced (%)	31.9

^aValues are means (SD), percentages or medians (90% range) for variables with skewed distribution. Of the total group, data were missing on maternal anthropometrics (n = 5), maternal education (n = 9), maternal smoking (n = 98), maternal alcohol use (n = 93), parity (n = 1), breastfeeding duration (n = 141), introduction solid foods (n = 153).

Table 2 Associations of ever breastfeeding, breastfeeding duration and exclusivity of breastfeeding with subcutaneous skinfolds (mm) at different ages in early childhood^{a b}

[illegible]

4 months ^c

(n = 210)

<i>P for trend</i>	<i>P = 0.07</i>	<i>P = 0.41</i>	<i>P = 0.16</i>	<i>P < 0.01</i>	<i>P = 0.13</i>	<i>P = 0.02</i>	<i>P = 0.38</i>	<i>P < 0.01</i>	<i>P = 0.06</i>
--------------------	-----------------	-----------------	-----------------	---------------------------	-----------------	------------------------	-----------------	---------------------------	-----------------

^aValues are unstandardized regression coefficients (95% confidence interval).

^b* $P < 0.05$ and ** $P < 0.01$ using multiple linear regression models. Models are adjusted for child's age at visit, sex, birth weight, maternal education, maternal body mass index, smoking and parity, gestational age, current height and observer.

^cNon-exclusive breastfeeding until 4 months includes partial until 4 months, partial thereafter; and partial until 4 months, not thereafter. Exclusive breastfeeding until 4 months includes exclusive until 6 months; exclusive until 4 months, partial thereafter; exclusive until 4 months, not thereafter.

Table 3 Age at introduction of solid foods and subcutaneous skinfolds (mm) in early childhood ^{a b c}

Solid foods (fruit- or vegetables)	Age 6 months			Age 24 months		
	Peripheral fat (mm) (95% CI)	Central fat (mm) (95% CI)	Total fat (mm) (95% CI)	Peripheral fat (mm) (95% CI)	Central fat (mm) (95% CI)	Total fat (mm) (95% CI)
N = 779						
	N = 486	N = 484	N = 484	N = 476	N = 469	N = 469
< 4 months (n = 48)	0.23 (-1.11, 1.56)	0.96 (-0.30, 2.21)	1.18 (-1.04, 3.40)	1.66 (-0.11, 3.42)	-0.09 (-1.39, 1.21)	1.29 (-1.46, 4.04)
> 4-5 months (n = 378)	-0.06 (-0.79, 0.67)	0.45 (-0.24, 1.14)	0.40 (-0.82, 1.61)	0.82 (-0.09, 1.73)	0.66 (-0.01, 1.33)	1.46 (0.05, 2.88)*
>5 months (n = 200)	Reference	Reference	Reference	Reference	Reference	Reference
<i>P for trend</i>	<i>P = 0.89</i>	<i>P = 0.10</i>	<i>P = 0.31</i>	<i>P = 0.03</i>	<i>P = 0.29</i>	<i>P = 0.08</i>

^aValues are unstandardized regression coefficients (95% confidence interval) using multiple linear regression models.

^bModels are adjusted for child's age at visit, sex, maternal education, maternal body mass index, smoking and parity, gestational age, birth weight, current height, observer and breastfeeding duration.

^c * $P < 0.05$

Figure 1 Flow chart of participants in study