Microsatellite genotyping reveals diversity within populations of , the secondary symbiont of tsetse flies

Oumarou Farikou, Flobert Njiokou, Gérard Cuny, Anne Geiger

► To cite this version:

Oumarou Farikou, Flobert Njiokou, Gérard Cuny, Anne Geiger. Microsatellite genotyping reveals diversity within populations of , the secondary symbiont of tsetse flies. *Veterinary Microbiology*, 2011, 150 (1-2), pp.207. 10.1016/j.vetmic.2011.01.021 . hal-00687331

HAL Id: hal-00687331

<https://hal.science/hal-00687331>

Submitted on 13 Apr 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Title: Microsatellite genotyping reveals diversity within populations of *Sodalis glossinidius*, the secondary symbiont of tsetse flies

Authors: Oumarou Farikou, Flobert Njiokou, Gérard Cuny, Anne Geiger

PII: S0378-1135(11)00040-X
DOI: doi:10.1016/j.vetmic.2011.01.021
Reference: VETMIC 5159

To appear in: *VETMIC*

Received date: 19-9-2010
Revised date: 8-1-2011
Accepted date: 24-1-2011

Please cite this article as: Farikou, O., Njiokou, F., Cuny, G., Geiger, A., Microsatellite genotyping reveals diversity within populations of *Sodalis glossinidius*, the secondary symbiont of tsetse flies, *Veterinary Microbiology* (2010), doi:10.1016/j.vetmic.2011.01.021

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Short Communication

Microsatellite genotyping reveals diversity within populations of *Sodalis glossinidius*, the secondary symbiont of tsetse flies

Running title: *Sodalis glossinidius* genotyping using microsatellite assay

Oumarou Farikou^{a,b}, Flobert Njiokou^b, Gérard Cuny^a and Anne Geiger^{a,*}

^aUMR 177, IRD-CIRAD, CIRAD TA A-17/G, Campus International de Baillarguet, 34398 Montpellier Cedex 5, France

^bUniversity of Yaoundé I, Faculty of Science, BP 812, Yaoundé, Cameroon

* Corresponding Author: Dr Anne Geiger

Address: UMR 177, IRD-CIRAD, CIRAD TA A-17/G, Campus International de Baillarguet, 34398 Montpellier Cedex 5, France.

Phone: +33 (0)4 67 59 39 25

Fax: +33 (0)4 67 59 38 94

E-mail: anne.geiger@ird.fr

Abstract

The aim of this study was to develop a PCR-based microsatellite genotyping method for identifying genetic diversity in *Sodalis glossinidius*, a symbiont associated with tsetse fly infection by trypanosomes causing human and animal trypanosomiasis. Allelic polymorphism at three loci, investigated on 40 fly gut extracts, evidenced eight alleles and the existence of five genotypes. This novel approach was shown to be efficient and suitable for routine large-scale genotyping of *S. glossinidius* present in the biologically complex tsetse fly extracts; it could favor progress in the fields of diagnosis, epidemiology, population genetics, and fly/symbiont/trypanosome interactions.

Keywords: *Sodalis glossinidius*, symbiont, microsatellites, genotyping, tsetse fly, Nagana, sleeping sickness

1. Introduction

Tsetse flies transmit trypanosomes responsible for human African trypanosomiasis (HAT) as well as the animal disease (Nagana). HAT affects a wide range of people in sub-Saharan Africa and is fatal if untreated, and Nagana is estimated to cost African agriculture US\$4.5 billion per year (Reinhardt, 2002). The available drugs are unsatisfactory (Barrett, 2006), while resistance is increasing (Matovu et al., 2001). Consequently, investigations for novel strategies must continue. Trypanosome transmission needs the parasite to establish in the tsetse fly midgut and to mature into an infective form before being transmitted to a novel mammalian host (Van den Abbeele et al., 1999). Tsetse flies harbor three different symbiotic bacteria among which *Sodalis glossinidius* is suspected of being involved in the fly's vector competence (Cheng and Aksoy, 1999; Dale and Welburn, 2001; Welburn and Maudlin, 1999). Recently, this bacterium was shown to be associated with infection of field populations of tsetse flies by trypanosomes (Farikou et al., 2010). Furthermore, genetic diversity was evidenced in *S. glossinidius* populations from insectary flies, suggesting that fly infection by trypanosomes could be favored by the presence of symbiont-specific genotype(s) (Geiger et al., 2007). Thus, a large-scale evaluation of the *S. glossinidius* genetic diversity in field populations of flies is needed. The AFLP technique previously used on DNA extracted from insectary fly hemolymph is not well adapted to routinely typing field samples because the hemolymph volume of such individual fly is too low, and the gut extracts, including the bacteria, are too complex. We therefore aimed to develop a Variable Number Tandem Repeat (VNTR) approach known to be highly discriminant (Le Flèche et al., 2006; Lindstedt, 2005; Whatmore et al., 2006). As the microsatellite approach has never been used to investigate the genetic diversity of *S. glossinidius*, we had to determine whether: a) it would be appropriate to distinguish genetically distinct *Sodalis* strains, b) the microsatellite approach could work on complex biological extract samples, and c) it would be suitable for routine analyses.

2. Materials and methods

Glossina palpalis gambiensis individuals were chosen from flies that were field sampled in different areas of Burkina Faso. Pupae were collected from these flies. After adult emergence, the population was maintained in a level 2 containment insectary at 23°C and 80% relative humidity, without any selection. Individuals used in the present work were randomly chosen. The flies were dissected in a drop of sterile 0.9% saline solution. Midguts from each fly were then separately transferred into microfuge tubes containing ethanol (95°) for further symbiont analyses. The instruments were carefully cleaned after the dissection of each fly to prevent contamination. The microfuge tubes were stored at –20°C until use.

DNA was extracted from midgut using the classical protocol (Navajas et al., 1998). Briefly, tissues were homogenized with a pestle in a cetyl trimethyl ammonium bromide (CTAB) buffer (CTAB 2%; 0.1 M Tris, pH 8; 0.02 M EDTA pH 8; 1.4 M NaCl) and incubated at 60°C for 30 min. The DNA was extracted from the lysis mixture with chloroform/isoamyl alcohol (24/1; V/V) and precipitated by adding isopropanol (V/V). After centrifugation (10,000 g, 15 min), the pellet was rinsed with 70% ethanol, air-dried, and resuspended in distilled sterile water. The DNA samples were stored at –20°C until PCR amplification processing.

S. glossinidius microsatellite repeat sequences were selected in a data bank (GenBank, accession number AP008232) (Toh et al., 2006) using the Vector NTI software; primer pairs were designed for the sequences flanking such regions and tested to ensure that they specifically amplify *S. glossinidius* and not host DNA. Table 1 shows the primers selected to perform the present study.

One µl of DNA was used for subsequent DNA amplification. Polymerase chain reactions (PCRs) were carried out using a DNA thermal cycler (PE Applied Biosystems, Foster City, CA, USA) in 20-µl final volumes, containing 4 pmol of each primer, 0.2 mM of each

deoxyribonucleotide, 1X incubation buffer with 1.5 mM MgCl₂ (Quantum Appligène, Ilkirch, France) and 0.5 units of Taq DNA polymerase (Quantum Appligène). Samples were first denatured at 94°C for 3 min and then processed through 40 cycles consisting in a denaturation step at 94°C for 30 s, an annealing step at 55°C for 30 s, and an extension step at 72°C for 1 min. The final elongation step was lengthened to 5 min at 72°C. PCR products were checked by 2% agarose gel electrophoresis and visualized under UV light after ethidium bromide staining (0.5 µg/ml). Allele bands were then resolved in nondenaturing acrylamide gels (10%), after loading 2-5 µl PCR products, and revealed by ethidium bromide staining.

In order to assess whether the amplicons actually corresponded to *S. glossinidius* microsatellites, bands representing the different alleles were purified using the Wizard DNA Clean Up System (Promega Corporation, Madison, WI, USA). Purified PCR fragments were cloned into pGEM-T Easy Vector (Promega Corporation). Clones of each cloned allele were purified with the Wizard Plus SV Minipreps DNA Purification System Kit (Promega, Charbonnière, France) and sequenced by GATC Biotech (Konstanz, Germany), allowing the allele size to be determined as well as the exact number of microsatellite repeats (GenBank accession numbers HQ842611-HQ842618).

3. Results

Figure 1 shows an example of an electrophoretic separation on a 10% polyacrylamide gel of different alleles; it demonstrates the development of the experimental process to be successful. Table 1 summarizes the characteristics of the three microsatellite markers selected to investigate the polymorphism within the *S. glossinidius* samples from 40 *G. palpalis gambiensis* flies, and to test the genotyping method: sequence of the repeats, part of the flanking sequences used as primers, their location on the *S. glossinidius* genome, and finally the number and size of the alleles evidenced at each locus. The repeat sequence of the ADN_g 2/5 marker is (AC). At this locus, two alleles that differ in the number of repeats, either five or seven, were identified among the 40 samples; their size, including the flanking sequences, was 176 and 180 bp, respectively. At the locus carrying the ADN_g 21/22 marker, four alleles were characterized that differ in the number of repeats, six, seven, nine, or 11, of the repeated unit (GCC), and in their total length, 163, 166, 172, and 178 bp, respectively. Finally, two alleles were identified at the ADN_g 15/16 locus; they differed in the number, five or seven, of their (AGG) unit repeat, and in their size, 106 or 112 bp. The combination of alleles amplified at the three loci on each sample (fly midgut extract) characterizes the genotype of the *Sodalis* strain the fly harbors. Significant diversity was evidenced among the *S. glossinidius* strains from the 40 tsetse flies. Table 2 presents the combination of alleles, identified by their size expressed as base pairs, for the 40 strains. Five allelic combinations, i.e., five genotypes, were found, one of which was much more frequent (62.5%) than the four others (22.5, 5, 5, and 5% respectively).

4. Discussion

One of the challenges in the study of *S. glossinidius* is the difficulty encountered when attempting to detect genetic diversity among strains. Previous studies have inferred phylogenetic relatedness among strains based on similarity at the 16S rDNA gene (Aksoy et al., 1997). However, single-gene phylogenetics are unreliable for resolving close relationships (Baldo et al., 2006). As shown in the introduction, to progress in the understanding of the role played by the symbiont *S. glossinidius* in the tsetse's capacity to transmit trypanosomes, the availability of an efficient and accurate genotyping tool is needed; the previously used AFLP method (Geiger et al., 2007) is not well suitable for routine genotyping of symbiont populations from field flies: an alternative and effective new method had to be developed.

From a general point of view, it appears that the *S. glossinidius* developed microsatellite method fulfils the performance criteria for a genotyping assay (Bricker and Ewalt, 2005), i.e., typing ability, reproducibility, stability, and discriminating power which is even better when the investigation is extended to several loci. Substantial genetic diversity was therefore observed among the 40 flies: eight alleles at the three loci investigated. These eight alleles were shown to be combined differently in the 40 individuals and resulted in the identification of five genotypes. It should be noted that the approach is actually quite appropriate to genetic diversity analyses of microorganisms present in complex biological samples such as *S. glossinidius* included in tsetse flies midgut extracts. It requires a small quantity of biological material, and the alleles can be analyzed by simple polyacrylamide gel electrophoresis; standardization and quality control are easy to achieve, and the recorded data can be easily coded. The proposed technical approach will be useful for large scientific uses, genetic population structure, epidemiology etc. It could possibly also be used for detecting mixed infections: given that *S. glossinidius* is haploid, the detection of more than one band (one

allele) at a given locus will mean that more than one *S. glossinidius* genotype is present in the sample analyzed and the existence of a mixed infection can be assessed in the fly sampled. To conclude, the microsatellite genotyping process we have adapted is an important tool for routine and effective diagnosis of *Sodalis* strains and will advance the elucidation of the role played by the symbiont in the vector competence of the tsetse fly, and finally in human and animal trypanosomiasis.

Conflict of interest statement

There is no conflict of interest with respect to funding or any other issue.

Acknowledgments

This study has been supported by the World Health Organization (WHO) and the International Atomic Energy Agency (IAEA).

O. Farikou is a PhD student supported by the Institut de Recherche pour le Développement (IRD).

References

- Aksoy, S., Chen, X., Hypsa, V., 1997. Phylogeny and potential transmission routes of midgut-associated endosymbionts of tsetse (*Diptera:Glossinidae*). *Insect Mol. Biol.* 6, 183-90.
- Baldo, L., Bordenstein, S., Wernegreen, J.J., Werren, J.H., 2006. Widespread recombination throughout *Wolbachia* genomes. *Mol. Biol. Evol.* 23, 437-449.
- Barrett, MP., 2006. The rise and fall of sleeping sickness. *Lancet.* 367, 1377-1378.
- Bricker, B.J., Ewalt, D.R., 2005. Evaluation of the HOOOF-print assay for typing *Brucella abortus* strains isolated from cattle in the United States: results with four performance criteria. *BMC Microbiol.* 5, 37.
- Cheng, Q., Aksoy, S., 1999. Tissue tropism, transmission and expression of foreign genes *in vivo* in midgut symbionts of tsetse flies. *Insect Mol. Biol.* 8, 125-132.
- Dale, C., Welburn, SC., 2001. The endosymbionts of tsetse flies: manipulating host-parasite interactions. *Int. J. Parasitol.* 31, 628-631.
- Farikou, O., Njiokou, F., Mbida Mbida, JA., Njitchouang, GR., Djeunga, HN., Asonganyi, T., Simarro, P.P., Cuny, G., Geiger, A., 2010. Tripartite interactions between tsetse flies, *Sodalis glossinidius* and trypanosomes-an epidemiological approach in two historical human African trypanosomiasis foci in Cameroon. *Infect. Genet. Evol.* 10, 115-21.

Geiger, A., Ravel, S., Mateille, T., Janelle, J., Patrel, D., Cuny, G., Frutos, R., 2007. Vector competence of *Glossina palpalis gambiensis* for *Trypanosoma brucei* s.l. and genetic diversity of the symbiont *Sodalis glossinidius*. *Mol. Biol. Evol.* 24, 102-9.

Le Flèche, P., Jacques, I., Grayon, M., Al Dahouk, S., Bouchon, P., Denoeud, F., Nockler, K., Neubauer, H., Guilloteau, L.A., Vergnaud, G., 2006. Evaluation and selection of tandem repeat loci for a *Brucella* MLVA typing assay. *BMC Microbiol.* 6, 9.

Lindstedt, B.A., 2005. Multiple-locus variable number tandem repeats analysis for genetic fingerprinting of pathogenic bacteria. *Electrophoresis* 26, 2567-2582.

Matovu, E., Seebeck, T., Enyaru, JCK., Kaminsky, R., 2001. Drug resistance in *Trypanosoma brucei* spp., the causative agents of sleeping sickness in man and nagana in cattle. *Microbes Infect.* 3, 763-770.

Navajas, M., Lagnel, J., Gutierrez, J., Boursot, P., 1998. Species-wide homogeneity of nuclear ribosomal ITS2 sequences in the spider mite *Tetranychus urticae* contrasts with extensive mitochondrial COI polymorphism. *Heredity.* 80, 742-752.

Reinhardt, E., 2002. Travailler ensemble: la mouche tsé-tsé et la pauvreté rurale. [Internet]. Available from: www.un.org/french/pubs/chronique/2002/numero2/0202p17_la_mouche_tse_tse.html. *Chronique ONU*, ONU Editor-September 02.

266 Toh, H., Weiss, BL., Perkin, SAH., Yamashita, A., Oshima, K., Hattori, M., Aksoy, S., 2006.
 267 Massive genome erosion and functional adaptations provide insights into the symbiotic
 268 lifestyle of *Sodalis glossinidius* in the tsetse host. *Genome Res.* 16, 149-156.
 269
 270 Van den Abbeele, J., Claes, Y., Bockstaele, D., Ray, D., Coosemans, M., 1999. *Trypanosoma*
 271 *brucei* spp. development in the tsetse fly: characterization of the post-mesocyclic stages in the
 272 foregut and proboscis. *Parasitology* 118, 469-478.
 273
 274 Welburn, SC., Maudlin, I., 1999. Tsetse-trypanosome interactions: rites of passage. *Parasitol.*
 275 *Today* 15, 399-403.
 276
 277 Whatmore, A.M., Shankster, S.J., Perrett, L.L., Murphy, T.J., Brew, S.D., Thirlwall, R.E.,
 278 Cutler, S.J., Macmillan, A.P., 2006. Identification and characterization of variable-number of
 279 tandem-repeat markers for typing of *Brucella* spp. *J. Clin. Microbiol.* 44, 1982-1993.
 280
 281
 282
 283
 284
 285
 286
 287
 288
 289
 290

Figure legends

Figure 1. Example of fragment length polymorphism at the ADNg 21/22 locus in 18 *Sodalis glossinidius* samples.

The alleles have been separated by electrophoresis on a 10% polyacrylamide gel on the basis of the size of their corresponding DNA sequence. Lane 1 corresponds to the "no template" negative control. Lanes 2-19 correspond to *S. glossinidius* samples. Here three alleles have been separated into three sizes: 163 (4 strains), 172 (13 strains), and 178 (1 strain) base pairs.

1 **Table 1. *S. glossinidius* microsatellite markers, PCR primers and allelic polymorphism**

2

3	Marker	Repeat	Primer sequences	Location at bp ^b	Number	Size of the
4		sequence ^a	(5'-3')		of alleles	alleles (bp)
5						
6	ADNg 5/2	(AC) _{x 5, 7}	GGCCGGTATTCTAACCGAC	4115043-4115222	2	176/180
7			AACTGCCAGGCATCCATTAC			
8	ADNg 21/22	(GCC) _{x 6, 7, 9, 11}	GAGCAAATCTCCCAGCACAT	1450588-1450759	4	163/166/172/178
9			TTCTTGTCCCTCAACCCATC			
10	ADNg 15/16	(AGG) _{x 5, 7}	ATACGGCGAAGCAATGAGAC	3250160-3250283	2	106/112
11			CAGCCTCTAAGCGCTCAACTC			
12						

13 ^a and number of repeats recorded at each locus (5 or 7 at ADNg 5/2, etc); ^bGenBank accession number AP008232

14

Table 2. Combinations of three alleles, one from each of the three loci, recorded among the 40 *S. glossinidius* strains sampled; number and percentage of strains carrying each combination

ADNg 2/5	ADNg 21/22	ADNg 15/16	Number and (%) of strains
180*	172	112	25 (62.5%)
176	163	112	9 (22.5%)
180	178	112	2 (5%)
180	166	112	2 (5%)
176	163	106	2 (5%)

*alleles identified by their size (number of base pairs) – see also Table 1 for details

Figure 1