Evidence of persistence and multiple genetic modifications of LPAI H7N7 in wild mallards in Poland is supported by phylogenetic studies

Krzysztof Śmietanka, Anna Pikula, Zenon Minta, Włodzimierz Meissner

► To cite this version:

Krzysztof Śmietanka, Anna Pikula, Zenon Minta, Włodzimierz Meissner. Evidence of persistence and multiple genetic modifications of LPAI H7N7 in wild mallards in Poland is supported by phylogenetic studies. *Avian Pathology*, 2011, 40 (02), pp.131-138. 10.1080/03079457.2010.537304 . hal-00687799

HAL Id: hal-00687799

<https://hal.science/hal-00687799>

Submitted on 15 Apr 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Evidence of persistence and multiple genetic modifications of LPAI H7N7 in wild mallards in Poland is supported by phylogenetic studies

Journal:	<i>Avian Pathology</i>
Manuscript ID:	CAVP-2010-0111.R1
Manuscript Type:	Original Research Paper
Date Submitted by the Author:	08-Sep-2010
Complete List of Authors:	Śmietanka, Krzysztof; National Veterinary Research Institute, Department of Poultry Diseases Pikula, Anna; National Veterinary Research Institute, Department of Poultry Diseases Minta, Zenon; National Veterinary Research Institute, Department of Poultry Diseases Meissner, Włodzimierz; University of Gdansk, Department of Vertebrate Ecology & Zoology, Avian Ecophysiology Unit
Keywords:	avian influenza virus, H7N7, mallards, epidemiology

SCHOLARONE™
Manuscripts

Cavp-2010-0111.R1

Evidence of persistence and multiple genetic modifications of H7N7low pathogenic avian influenza virus in wild mallards in Poland provided by phylogenetic studies

K. Śmietanka¹, A. Pikuła¹, Z. Minta¹, W. Meissner²

¹National Veterinary Research Institute, Department of Poultry Diseases, Al. Partyzantów 57, 24-100 Pulawy, Poland, and ²University of Gdańsk, Department of Vertebrate Ecology & Zoology, Avian Ecophysiology Unit, Al. Legionów 9, 80-441 Gdańsk, Poland

Running title: Low pathogenic AIV H7N7 in mallards in Poland

Corresponding author: ksmiet@piwet.pulawy.pl

Received: 6 July 2010

Abstract

Genetic characterization of the whole genome of four avian influenza viruses (AIV) H7N7 isolated in 3 successive winter seasons (2007-2009) from wild mallards in 3 cities in Poland was performed. All of the tested strains were of low pathogenicity and no molecular marker associated with an increased adaptation to poultry, mammals or resistance to antiviral drugs was found. The major outcome of the phylogenetic studies was that the isolate A/mallard/Poland/446/09 (detected in December 2009) shared a recent common ancestor with A/mallard/Poland/41/09 (isolated in February 2009) in relation to HA & PB1 genes, with A/mallard/Poland/16/09 (found in January 2009) regarding NA & NS genes, and with A/mallard/Poland/01/08 (recovered in December 2007) as regards NS gene. Interestingly, A/mallard/Poland/16/09 and A/mallard/Poland/446/09 were isolated at the same sampling site almost exactly one year apart and that points to resident population of mallards (and other resident waterfowl) as responsible for the perpetuation of AIV in the given area between successive winters. On the other hand, the ornithological data discussed in detail in the paper strongly suggest that the virus transmission between close sites but located in different urban areas is most likely achieved by migratory birds, a fact additionally supported by a close relatedness between different gene segments of Polish H7N7 and AIV detected in wild birds in Europe. A high heterogeneity of the gene pool found in the study is indicative of frequent reassortment events. Additionally, two H7N7 isolates were shown to possess selected genes closely related to AIV detected in domestic poultry in Italy and Czech Republic. This study corroborates the importance of active surveillance in wild birds as a valuable tool of early warning of AI in poultry.

Introduction

Avian influenza A virus (AIV) is an orthomyxovirus with a genome consisting of eight single-stranded RNA segments encoding for at least 10 different viral proteins: PB2, PB1, PA, HA, NP, NA, M1, M2, NS1, and NS2 (Suarez, 2008). The current classification into subtypes is based on the surface glycoproteins: haemagglutinin (HA) and neuraminidase (NA) and so far 16 HA subtypes and 9 NA subtypes have been described for which wild aquatic birds are considered the natural reservoir (Webster *et al.*, 1992; Swayne & Suarez, 2000; Fouchier *et al.*, 2005). In wild birds AI viruses usually persist in the low pathogenic form but some of the H5 and H7 subtypes have the capacity to become highly pathogenic (HP). The emergence of HP from LP AIV is believed to take place in poultry through a number of processes of which acquisition of multiple amino acids at HA2/HA1 cleavage site plays a crucial role (Swayne & Suarez, 2000).

In recent years HP outbreaks caused by viruses of H7 subtypes, even though overshadowed by H5N1 epidemic (Sims & Brown, 2008), have also had devastating effects on the poultry industry, e.g. resulting in 13 million of birds dead or culled in Italy in 1999-2000 (HPAI H7N1) and 25.6 million birds stamped out in the Netherlands in 2003 (HPAI H7N7) (Capua *et al.*, 2000; Alexander *et al.*, 2008). In 2008 and 2009 outbreaks of HPAI H7N7 were recorded in the UK and Spain but they were contained rapidly and did not spread (OIE online reports, 2010).

Despite the great importance, there are few reports describing phylogenetic analysis of H7N7 strains isolated from their natural reservoir (Campitelli *et al.*, 2008; Jahangir *et al.*, 2010; Metreveli *et al.*, 2010). Furthermore, there is still little information dealing with virus-host ecology.

In the present study we have analyzed the molecular features and phylogenetic relationship of AIV H7N7 isolated from wild mallards in Poland in 2007-2009 based on partial or full-length sequencing of all 8 segments of the viral genome and compared the results to AIV sequences available in the public domain.

Material and methods

Viruses. In the study four avian influenza H7N7 viruses isolated from cloacal swabs collected from wild mallards (*Anas platyrhynchos*) were used (Table 1). The viruses were isolated in 3 successive winters: 2007/2008 (isolate A/mallard/Poland/01/08), 2008/2009 (isolates A/mallard/Poland/16/09 & A/mallard/Poland/41/09), and 2009/2010 (isolate A/mallard/Poland/446/09). The geographical location of sampled mallards is presented in Fig. 1. Noteworthy is the fact that the isolates A/mallard/Poland/16/09 and A/mallard/Poland/446/09 were detected on the same pond (~3500 m²) almost exactly one year apart (January 2009-December 2009). All the viruses were isolated in embryonated chicken specific pathogen free (SPF) eggs followed by identification in haemagglutination inhibition test according to standard procedures (OIE, 2008). Allantoic fluids from inoculated eggs of the 1st passage exhibiting haemagglutinating activity were used directly for RNA isolation or stored at -80°C until use.

RNA extraction, RT-PCR and gene sequencing. The RNA was extracted from infected allantoic fluid using Qiagen RNeasy Mini Kit according to manufacturer's instruction. Reverse transcription and PCR reaction were performed in one tube using Qiagen RT-PCR kit using specific primers (sequences and cycling conditions kindly provided by Dr. W. Dundon,

IZSVe, Legnaro, Italy, available upon request). The complete coding sequences or large portions of all 8 genes were generated using BigDye Terminator v3.1 (Applied Biosystems). The products of the sequencing reactions were sequenced in the 3730xl DNA Analyzer (Applied Biosystems) in Genomed, Warsaw.

Phylogenetic analyses. The nucleotide regions investigated in the phylogenetic analyses were as follows: PB2: 27-1203, PB1: 1-1920, PA: 983-2135, HA: 26-1637, NA: 20-1264, NP: 46-1442, M: 1-937, NS: 1-812. For sequence editing and analysis, modules SeqMan (assembly of the consensus sequence from electrophoregrams) and EditSeq (translation from nucleotide into amino acid sequences) of DNASTar LaserGene (version 8.1.3) were used. Sequences were compared with published sequences of AIV of Eurasian and North American lineages using BLAST local alignment search tool available from the National Center for Biotechnology Information, Bethesda, Maryland, USA (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>). Multiple sequence alignments were established and phylogenies reconstructed with MEGA 4 software (Tamura *et al.*, 2007) with the use of Clustal W and neighbor-joining tree inference method with the Kimura two-parameter substitution model with 1000 bootstrap replicates. Potential N-glycosylation sites were predicted with the use of the NetNGlyc 1.0 server (<http://www.cbs.dtu.dk/services/NetNGlyc/>). Gene sequences were submitted to the EpiFLU nucleotide sequence database (platform.gisaid.org) with the following isolate ID: EPI_ISL_26339, EPI_ISL_60707, EPI_ISL_60708, EPI_ISL_74018.

Results

Molecular characterization of AIV H7N7. All of the investigated isolates had the monobasic amino-acid cleavage site motif at the HA cleavage site: PEIPKGR-GLF (3 isolates) and PELPKGR-GLF (1 isolate) indicating low pathogenicity (Tab.1). The amino-acid residues 226Q and 228G (H3 amino acid numbering) at the receptor binding sites, associated with preferential binding to $\alpha(2,3)$ -linked sialic acid (Vines *et al.*, 1998) were also found in all of the isolates. The H7 HA amino acid sequence revealed five possible glycosylation sequons but not at positions 123 or 149 (H7 numbering, a feature shared by H7 strains isolated from poultry) (Banks *et al.*, 2000; Banks *et al.* 2001). The PB2 E627K mutation, related to an increased adaptation to mammals (Subbarao *et al.*, 1993; Shinya *et al.*, 2004) was not observed. The S31N amino acid substitution in the M2 protein, responsible for conferring resistance to amantadanes (Scholtissek *et al.*, 1998), was not detected in any of the tested H7N7. The NS1 protein did not show any deletion and the D92E substitution, associated with viral resistance to cytokines in pigs (Seo *et al.*, 2002), was not observed.

Phylogenetic analysis. All of the gene segments of the viruses analyzed in this study were of avian origin and clearly belonged to Eurasian lineage (Fig. 2-5 and Supplementary Fig. S1-S4). With regard to the HA gene, all of the Polish H7 isolates fell into the same major clade of phylogenetic tree (Fig. 2) with the sequence homology 96.2 to 99.3%, and the highest similarity was observed between A/mallard/Poland/41/09 and A/mallard/Poland/446/09 (99.3%). The A/mallard/Poland/16/09 was most closely related to A/goose/Czech Republic/1848/2009(H7N9) (99.8%), isolated from poultry.

The NA genes of the tested strains were 90.3% to 98.7% homologous to each other, and A/mallard/Poland/16/09 and A/mallard/Poland/446/09 (isolated at the same sampling site almost exactly 1 year apart) were the most closely related (Fig. 3). Both isolates were also related to A/mute swan/Hungary/5973/2007(H7N7) with sequence identity up to 97.9%.

Interestingly, in contrast to HA gene analysis, the nucleotide homology percentage of the NA gene of A/mallard/Poland/41/09 and A/mallard/Poland/446/09 was very low (90.3%) and both isolates were located on two distant branches of the phylogenetic tree.

In reference to the PB2 gene, two Polish isolates A/mallard/Poland/01/08 and A/mallard/Poland/41/09 clustered in the same subgroup (sequence homology 97.3%), A/mallard/Poland/16/09 was located within a sister subgroup of the same group with up to 96.0% of sequence similarity. The A/mallard/Poland/446/09 showed no significant similarity in relation to the other Polish H7N7 (93.5 to 94.2%) or sequences available at GenBank (Fig. 4).

Regarding the PB1 gene, H7N7 viruses isolated in Poland clustered in 3 different subgroups of the phylogenetic tree. The phylogenetic position of A/mallard/Poland/41/09 and A/mallard/Poland/446/09 on the same branch (sequence homology 99.1%) resembled their location on the phylogenetic tree constructed for the HA gene. Both isolates were similar to A/mute swan/Hungary/5973/2007(H7N7) (nucleotide sequence similarity up to 98.0%, Supplementary Fig. S1). The A/mallard/Poland/01/08 shared a very similar PB1 gene with A/turnstone/Netherlands/1/2007 (nucleotide identity 99.1%).

As for the PA gene, a high diversity was found among Polish isolates, which clustered within 3 different clades with sequence similarity ranging from 94.0 to 96.6% (Supplementary Fig. S2). Some of them, however, showed a very high degree of relationship with PA gene detected in wild waterfowl in Europe, eg. A/mallard/Poland/01/08 to A/mallard/Netherlands/22/2007(H7N1) (99.2%), A/mallard/Italy/299/05(H7N7) (99.0%) or A/common teal/Netherlands/1/2005(H8N4) (99.0%).

A high heterogeneity was also found in relation to NP gene because all of the H7N7 detected in Poland clustered into 4 different subgroups (homology of nucleotide sequences ranged from 93.9 to 96.0%, Supplementary Fig. S3). Similarly to the HA analysis, the highest

sequence similarity (99.4%) was found between A/mallard/Poland/16/09 and A/goose/Czech Republic/1848/2009(H7N9). The A/mallard/Poland/01/08 grouped closely to AIV H7 detected in poultry in Italy: A/turkey/Italy/2685/2003(H7N3) and A/turkey/Italy/220185/2003(H7N3), with similarity of sequences 98.5%.

In the case of the M gene, 97.0 – 98.8 % of nucleotide sequence identity was found but none of the Polish isolates were directly related to one another. On the other hand, a high degree of relationship was observed between A/mallard/Poland/01/08 and A/turnstone/Netherlands/1/2007 (99.5% nucleotide sequence similarity, Fig. 5).

The analysis of NS gene demonstrated that Polish H7N7 belonged to allele A (Ludwig *et al.*, 1991; Zohari *et al.*, 2008). The isolates A/mallard/Poland/01/08, A/mallard/Poland/16/09, and A/mallard/Poland/446/09 showed a high percentage of nucleotide homology (99.0 – 99.1%) and their position on the phylogenetic tree clearly suggest their common origin (Supplementary Fig. S4). The highest nucleotide homology with reference to sequences available in GenBank was observed between A/mallard//Poland/01/08, A/mallard/Netherlands/1/2006 (99.8%) and A/mallard/Altai/1208/2007 (99.6%).

Discussion

Mallards constitute an important host for AIV. According to Munster *et al.* (2007), an average prevalence of AIV infections in this species established in a large-scale surveillance study was 7.3%. However, the frequency of AIV isolations in mallards is dependant on season and geographical location. At the sampling sites in Northern Europe AIV isolation was about 3-fold higher as compared to wintering sites in the West, with peak in the early fall migration (Munster *et al.*, 2007). Among mallards breeding in Poland, like in other European

countries and in North America, two groups can be distinguished (Cramp & Simmons, 1977; Heusmann, 1981). Mallards resident in urban areas do not migrate while most of the birds breeding outside of urban regions show regular movements between breeding and wintering sites. However, in mild climate areas of Western Europe breeding populations of this species are sedentary making only local movements during severe weather conditions (Scott & Rose, 1996). Mallards ringed in Poland during breeding season were recovered in Western and South-Western Europe. On the other hand, Poland is the wintering site for mallards breeding in Northern and North-Eastern Europe (e.g. Scandinavia, Russia, Baltic states) (Fransson & Petterson, 2001; unpublished data of the Polish Ringing Center). According to Stallknecht & Brown (2008) the avian influenza virus maintenance cycle is based on virus transmission on the breeding and wintering grounds between migratory and resident birds but the mechanism is still poorly understood. It is known that in urban areas in Poland both sedentary and migratory mallards form mixed populations in winter. Pair formation in mallards takes place in autumn and winter (Cramp & Simmons, 1977). Females reveal very high philopatry to the breeding grounds, higher than males, and in spring males follow females to the breeding sites (Cramp & Simmons, 1977; Rohwer & Anderson, 1988; Evrard, 1999). Therefore some males of the sedentary populations may join migratory populations whereas some migratory males can stay put on the wintering ground. Such behavioral features create a perfect environment for the transmission of AIV between birds from different populations. Due to the fact that mallards are the most numerous waterfowl in Europe (population estimated at ~7.5 million, Wetlands International, 2006), this species may play a crucial role in the long-lasting perpetuation of AIV, as suggested by Munster & Fouchier (2009).

In our study we analyzed molecular and phylogenetic characteristics of four H7N7 strains isolated from wild mallards in Northern and North-eastern Poland in 3 successive winters (2007/2008, 2008/2009, and 2009/2010). Even though the molecular markers revealed

a profile typical of AIV of wild bird origin (low pathogenicity, no markers related to increased adaptation to poultry and mammals or resistance to some antiviral drugs), these features needed to be analyzed even in AIV isolated from wild birds. There have been sporadic cases of HPAI H7 in wild birds (Swayne, 2008). Moreover, some amino acid substitutions in the binding receptor site of HA and mutations associated with resistance to amantadine in a mallard-derived H7 in Italy have also been found (Campitelli *et al.*, 2008).

The phylogenetic study demonstrated a high degree of inter-relationship between Polish H7N7 but only in relation to selected genes. The major conclusion is that the isolate A/mallard/Poland/446/09 is a reassortant virus that shares a very recent common ancestor with A/mallard/Poland/41/09 (in relation to HA&PB1 gene), A/mallard/Poland/16/09 (with respect to NA & NS gene) and A/mallard/Poland/01/08 (as regards NS gene). The spatial and temporal distribution of the cases was significant: the sampling site of birds carrying A/mallard/Poland/16/09 & A/mallard/Poland/446/09 isolates was located on the same pond in Gdańsk (January-December 2009), approximately 100 km away from the city of Słupsk, in which A/mallard/Poland/41/09 had been detected in March 2009. Both sites were located 330-430 km from the sampling location of the mallard from which A/mallard/Poland/01/08 had been recovered at the end of 2007. The ornithological observations resulted in several resightings of the male (ring number SN 11463), the host of A/mallard/Poland/446/09 isolate. The bird was ringed and sampled on a pond in Gdansk on the 27th of December 2009 and it was recovered at the same site on the 23rd of February, 1st of May and early June of 2010. Additionally, it was seen on the 7th of January and 7th of March in the area within a radius of 2 km from the sampling site (observation by W. Meissner). These findings confirm that an individual carrying A/mallard/Poland/446/09 strain was a resident bird and it reinforces our hypothesis that resident mallards contribute to the long-term circulation of AIV in the given area between successive winters. We suggest that the NA and NS segments of

A/mallard/Poland/16/09 & A/mallard/Poland/446/09 detected in mallards at the same sampling site 1 year apart were derived from avian influenza virus precursor strain(s) that persisted in the Gdańsk area for a long period of time and emerged in January 2009 and then in December 2009. Interestingly, Jahangir *et al.* (2009) found a high isolation rate of AIV H7N7 in northern pintails (*Anas acuta*) in wintering season 2007/2008 in Japan, and similarly to our studies, on one occasion the virus was detected twice at one site, two months apart. A phylogenetic study of H7 and N7 genes revealed that these isolates were very similar or virtually identical (Jahangir *et al.*, 2010). However, the high frequency of H7N7 isolation in that region was noted exclusively in one wintering season as opposed to our study in which we demonstrated the presence of AIV with genetically related genes between two consecutive winter seasons in the same area.

On the other hand, the introduction of new AI viruses and their efflux from the area is caused by migratory waterfowl, a fact supported by ornithological observations and our phylogenetic studies in which we found a high degree of relationship between some of the genome segments of the Polish H7N7 and AIV detected in wild birds in Europe. Despite a very close relationship between HA and PB1 of AIV H7N7 isolated from mallards at close locations in Gdańsk (A/mallard/Poland/446/09) and Słupsk (A/mallard/Poland/41/09), i.e. 102 km away from each other, it is the most likely that the precursor viruses bearing these gene segments were transmitted between these sites via migratory birds since urban mallards show a very high fidelity to their whereabouts and usually move within very restricted area of the city in which they stay (Heusmann, 1981; W. Meissner - unpublished data from studies conducted in three Polish cities).

The AI virus undergoes frequent reassortment events: the H7N7 isolates detected in Poland never shared the same constellation of all genome segments and the close relatedness (>98.5%) was only found in relation to single genes, indicating in most cases different

evolutionary pathways. Of special significance is the fact that despite A/mallard/Poland/41/09 and A/mallard/Poland/446/09 sharing a very closely related HA gene (>99% nucleotide identity) their NA genes were completely different (~90% nucleotide similarity). Other studies have also proved that reassortment events between AIV of various origin are frequent (Hatchette *et al.*, 2004; Macken *et al.*, 2006; Campitelli *et al.*, 2008; Briand *et al.*, 2010).

Interestingly, unlike other genes, H7 gene seems to have established a relatively conserved sublineage for wild mallards in Poland. However, the assumption that this is a result of an adaptive process to this species of birds requires more evidence due to a limited number of available H7 isolates. The fragmentary data of virological surveillance in Poland is on one hand caused by a number and species of birds tested (approximately 3,000 a year, mostly mallards), and on the other hand, possibly insufficient sampling and storage conditions of swabs and faeces, which may have also contributed to the reduction of an overall AIV detection rate (<2% a year). Consequently, it cannot be excluded that there may have been other undetected AIV H7 strains of the same and different sublineages circulating in mallards and other species of waterfowl. This issue will be raised in future after more intensive and targeted surveillance in places with previous records of AIV H7.

Another noteworthy aspect of the phylogenetic studies is a close relationship between 2 genes (HA & NP) of A/mallard/Poland/16/09 and AIV H7N9 isolated from a flock of domestic geese in the Czech Republic. On the other hand, as mentioned above, a high relatedness of its NA & NS to A/mallard/Poland/446/09 was shown. The isolate A/mallard/Poland/01/08 was also found to have an NP gene most closely related to H7N3 viruses detected in poultry in Italy in 2003. There is also strong evidence from other studies that low pathogenic AIV spill over to poultry as phylogenetic studies have shown a direct relationship between AIV genes from wild and domestic birds, even from HPAI outbreaks (Campitelli *et al.*, 2004; Munster *et al.*, 2005; Terregino *et al.*, 2007; Campitelli *et*

al., 2008; Briand *et al.*, 2010; Metreveli *et al.*, 2010). This study corroborates the importance of active surveillance in free living birds as a valuable tool of an early warningsystem of LPAI and HPAI outbreaks. It is very important in relation to H7N7viruses, for which the number of available sequences is still very limited. We also suggest that the survey programmes for AIV should not only be targeted at migratory population at major flyways but that resident waterfowl should also be considered.

Acknowledgements

This work was supported by European Network of Excellence (EPIZONE) WP6.2 “Molecular epidemiology and surveillance of AI and APMV” and EU 6th framework programmes INN-FLU (SSPE-CT-2006-44372) and NEW-FLUBIRD (044490).

References

- Alexander, D.J., Capua, I. & Koch, G. (2008). Highly pathogenic avian influenza outbreaks in Europe, Asia, and Africa since 1959, excluding the Asian H5N1 virus outbreaks. In D.E. Swayne (Eds.). *Avian Influenza*, 1st ed. (pp. 217-237). Ames: Blackwell Publishing.
- Banks, J., Spiedel, E. S., McCauley, J. W. & Alexander, D. J. (2000). Phylogenetic analysis of H7 haemagglutinin subtype influenza A viruses. *Archives of Virology*, 145, 1047–1058.
- Banks, J., Spiedel, E. S., Moore, E., Plowright, L., Piccirillo, A., Capua, I., Cordioli, P., Fioretti, A. & Alexander, D. J. (2001). Changes in the haemagglutinin and the

- neuraminidase genes prior to the emergence of highly pathogenic H7N1 avian influenza viruses in Italy. *Archives of Virology*, 146, 963–973.
- Briand, F.X., Le Gall-Reculé, G., Guillou-Cloarec, C., Ogor, K. & Jestin, V. (2010). Phylogeny and genotyping of recent avian low-pathogenic H5 subtype influenza viruses from French ducks. *Journal of General Virology*, 91, 960-970.
- Campitelli, L., Mogavero, E., De Marco, M. A., Delogu, M., Puzelli, S., Frezza, F., Facchini, M., Chiapponi, C., Foni, E. & other authors (2004). Interspecies transmission of an H7N3 influenza virus from wild birds to intensively reared domestic poultry in Italy. *Virology*, 323, 24–36.
- Campitelli, L., Di Martino, A., Spagnolo, D., Smith, G.J., Di Trani, L., Facchini, M., De Marco, M.A., Foni, E., Chiapponi, C., & other authors (2008). Molecular analysis of avian H7 influenza viruses circulating in Eurasia in 1999-2005: detection of multiple reassortant virus genotypes. *Journal of General Virology*, 89, 48-59.
- Capua, I., Mutinelli, F., Marangon, S. & Alexander, D.J. (2000). H7N1 avian influenza in Italy (1999 to 2000) in intensively reared chickens and turkeys. *Avian Pathology*, 29, 537-543.
- Cramp, S., & Simmons K. E. L. (1977). *The Birds of the Western Palearctic*. Oxford University Press, Oxford.
- Evrard, J. O. (1999). Mallard and Blue-winged Teal philopatry in Northwest Wisconsin. *North American Bird Bander* 24, 38-42.
- Fouchier, R.A., Munster, V., Wallensten, A., Bestebroer, T.M., Herfst, S., Smith, D., Rimmelzwaan, G.F., Olsen, B. & Osterhaus, A.D. (2005). Characterization of a novel influenza A virus hemagglutinin subtype (H16) obtained from black-headed gulls. *Journal of Virology*, 79, 2814-2822.

- Fransson, T., & Pettersson, J. (2001). Swedish Bird Ringing Atlas Volume 1, Divers-raptors. Naturhistoriska riksmuseet, Stockholm.
- Hatchette, T. F., Walker, D., Johnson, C., Baker, A., Pryor, S. P. & Webster, R. G. (2004). Influenza A viruses in feral Canadian ducks: extensive reassortment in nature. *Journal of General Virology*, 85, 2327-2337.
- Heusmann, H. W. (1981). Movements and survival rates of park mallards. *Journal of Field Ornithology*, 52, 214-221.
- Jahangir, A., Ruenphet, S., Ueda, S., Ueno, Y., Shoham, D., Shindo, J., Okamura, M., Nakamura, M. & Takehara, K. (2009). Avian influenza and Newcastle disease viruses from northern pintail in Japan: isolation, characterization and inter-annual comparisons during 2006-2008. *Virus Research*, 143, 44-52.
- Jahangir, A., Ruenphet, S., Shoham, D., Okamura, M., Nakamura, M. & Takehara, K. (2010). Haemagglutinin and neuraminidase characterization of low pathogenic H5 and H7 avian influenza viruses isolated from Northern pintails (*Anas acuta*) in Japan, with special reference to genomic and biogeographical aspects. *Virus Genes*, 40, 94-105.
- Ludwig, S., Schultz, U., Mandler, J., Fitch, W.M. & Scholtissek C. (1991). Phylogenetic relationship of the nonstructural (NS) genes of influenza A viruses. *Virology*, 183, 566-577.
- Macken, C. A., Webby, R. J. & Bruno, W. J. (2006). Genotype turnover by reassortment of replication complex genes from avian influenza A virus. *Journal of General Virology*, 87, 2803-2815.
- Metreveli, G., Zohari, S., Ejdersund, A. & Berg, M. (2010). Phylogenetic analysis of the hemagglutinin gene of low pathogenic avian influenza virus H7N7 strains in mallards in Northern Europe. *Avian Diseases*, 54, 453-456.

- Munster, V.J., Wallensten, A., Baas, C., Rimmelzwaan, G.F., Schutten, M., Olsen, B., Osterhaus, A.D. & Fouchier, R.A. (2005). Mallards and highly pathogenic avian influenza ancestral viruses, northern Europe. *Emerging Infectious Diseases*, 11, 1545-1551.
- Munster, V.J., Baas, C., Lexmond, P., Waldenstrom, J., Wallensten, A., Fransson, T., Rimmelzwaan, G.F., Beyer, W.E.P., Schutten, M., Olsen, B., Osterhaus, A.D. & Fouchier, R.A. (2007). Spatial, temporal, and species variation in prevalence of influenza A viruses in wild migratory birds. *PLoS Pathogens*, 11, e61.
- Munster, V.J., & Fouchier, R.A. (2009). Avian influenza virus: Of virus and bird ecology. *Vaccine*, 27, 6340-6344.
- OIE. World Organisation For Animal Health (2008). Manual of Diagnostic Tests and Vaccines for Terrestrial Animals. http://www.oie.int/eng/normes/en_mmanual.htm.
- Rohwer, F. C. & Anderson, M. G. (1988). Female-biased philopatry, monogamy, and the timing of pair formation in migratory waterfowl. *Current Ornithology* 5, 187–221.
- Scholtissek, C., Rhohe, W., von Hoyningen, V., & Rott, R. (1998). On the origin of the human influenza viruses subtypes H2N2 and H3N2. *Virology*, 87, 13-20.
- Scott, D.A. & Rose, P.M. (1996). Atlas of Anatidae Populations in Africa and Western Eurasia. Wetlands International Publication No.41, Wageningen, The Netherlands. 336pp.
- Seo, S.H., Hoffmann, E. & Webster, R. (2002). Lethal H5N1 influenza viruses escape host anti-viral cytokine responses. *Nature Medicine*, 8, 950–954.
- Shinya, K., Hamm, S., Hatta, M., Ito, H., Ito, T. & Kawaoka, Y. (2004). PB2 amino acid at position 627 affects replicative efficiency, but not cell tropism, of Hong Kong H5N1 influenza A viruses in mice. *Virology*, 320, 258-266.

- Sims L.D., Brown I.H. (2008): Multicontinental epidemic of H5N1 HPAI virus (1996-2007). In D.E. Swayne (Eds.). *Avian Influenza*, 1st ed. (pp. 251-286). Ames: Blackwell Publishing
- Stallknecht, D.E. & Brown, J.D. (2008): Ecology of avian influenza in wild birds. In D.E. Swayne (Eds.). *Avian Influenza*, 1st ed. (pp. 43-58). Ames: Blackwell Publishing.
- Suarez, D.L. (2008). Influenza A virus. In D.E. Swayne (Eds.). *Avian Influenza*, 1st ed. (pp. 3-22). Ames: Blackwell Publishing.
- Subbarao, E.K., London, W. & Murphy B.R. (1993). A single amino acid in the PB2 gene of influenza A virus is a determinant of host range. *Journal of Virology*, 67, 1761-1764.
- Swayne, D.E. & Suarez, D.L. (2000). Highly pathogenic avian influenza. *Revue Scientifique et Technique Office International des Epizooties*, 19, 463-482.
- Swayne, D.E. (2008). The global nature of avian influenza. In D.E. Swayne (Eds.). *Avian Influenza*, 1st ed. (pp. 123-143). Ames: Blackwell Publishing.
- Tamura, K., Dudley, J., Nei, M. & Kumar, S. (2007). MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Molecular Biology and Evolution*, 24, 1596-1599.
- Terregino, C., De Nardi, R., Guberti, V., Scremin, M., Raffini, E., Martin, A.M., Cattoli, G., Bonfanti, L. & Capua, I. (2007). Active surveillance for avian influenza viruses in wild birds and backyard flocks in Northern Italy during 2004 to 2006. *Avian Pathology*, 36, 337-344.
- Vines, A., Wells, K., Matrosowich, M., Castrucci, M.R., Ito, T. & Kawaoka, Y. (1998). The role of influenza A virus hemagglutinin residues 226 and 228 in receptor specificity and host range restriction. *Journal of Virology*, 72, 7626-7631.
- Webster, R.G., Bean, W.J., Gorman, O.T., Chambers, T.M. & Kawaoka, Y. (1992). Evolution and ecology of influenza A viruses. *Microbiological Reviews*, 56, 152-179.

Wetlands International. (2006). Waterbird Population Estimates – Fourth Edition. Wetlands International, Wageningen, The Netherlands

Zohari, S., Gyarmati, P., Ejdersund, A., Berglöf, U., Thorén, P., Ehrenberg, M., Czifra, G., Belák, S., Waldenström, J., Olsen, B. & Berg, M. (2008). Phylogenetic analysis of the non-structural (NS) gene of influenza A viruses isolated from mallards in Northern Europe in 2005. *Virology Journal*, 12, 5:147.

Table 1 *Molecular characteristics of the H7N7 Polish isolates in this study*

Isolate	Location	Sample collection date (ring number of bird)	HA cleavage site (pathogenicity)	PB2 sequence at amino acid 627	NS1 sequence at amino acid 92	M2 sequence at amino acid 31
A/mallard/Poland/01/08	Białystok	28.12.2007 (SN8884)	PEIPKGR-GLF (low pathogenic)	E	D	S
A/mallard/Poland/16/09	Gdańsk	12.01.2009 (SN 10243)	PELPKGR-GLF (low pathogenic)	E	D	S
A/mallard/Poland/41/09	Słupsk	16.02.2009 (SN 10266)	PEIPKGR-GLF (low pathogenic)	E	D	S
A/mallard/Poland/446/09	Gdańsk	27.12.2009 (SN 11463)	PEIPKGR-GLF (low pathogenic)	E	D	S

Figure 1. *Geographical location of the LPAIV H7N7 cases in wild mallards: A - A/mallard/Poland/01/08, B - A/mallard/Poland/16/09, C - A/mallard/Poland/41/09, D - A/mallard/Poland/446/09; isolates B&D detected at the same sampling location*

Figure 2. Phylogenetic tree for the HA gene (nucleotides 26-1637) constructed by the neighbour-joining method. The evolutionary distances were computed using the Maximum Composite Likelihood method and are in the units of the number of base substitutions per site. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (1000 replicates) are shown next to the branches

Figure 3. Phylogenetic tree for the NA gene (nucleotides 20-1264) constructed by the neighbour-joining method. The evolutionary distances were computed using the Maximum Composite Likelihood method and are in the units of the number of base substitutions per site. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (1000 replicates) are shown next to the branches

Figure 4. Phylogenetic tree for the PB2 gene (nucleotides 27-1203) constructed by the neighbour-joining method. The evolutionary distances were computed using the Maximum Composite Likelihood method and are in the units of the number of base substitutions per site. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (1000 replicates) are shown next to the branches

Figure 5. Phylogenetic tree for the M gene (nucleotides 1-937) constructed by the neighbour-joining method. The evolutionary distances were computed using the Maximum Composite Likelihood method and are in the units of the number of base substitutions per site. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (1000 replicates) are shown next to the branches

Supplementary Figure S1. Phylogenetic tree for the PB1 gene (nucleotides 1-1920) constructed by the neighbour-joining method. The evolutionary distances were computed using the Maximum Composite Likelihood method and are in the units of the number of base substitutions per site. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (1000 replicates) are shown next to the branches

Supplementary Figure S2. Phylogenetic tree for the PA gene (nucleotides 983-2135) constructed by the neighbour-joining method. The evolutionary distances were computed using the Maximum Composite Likelihood method and are in the units of the number of base substitutions per site. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (1000 replicates) are shown next to the branches

Supplementary Figure S3. Phylogenetic tree for the NP gene (nucleotides 46-1442) constructed by the neighbour-joining method. The evolutionary distances were computed using the Maximum Composite Likelihood method and are in the units of the number of base substitutions per site. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (1000 replicates) are shown next to the branches

Supplementary Figure S4. Phylogenetic tree for the NS gene (nucleotides 1-812) constructed by the neighbour-joining method. The evolutionary distances were computed using the Maximum Composite Likelihood method and are in the units of the number of base substitutions per site. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (1000 replicates) are shown next to the branches