An optimized hair trap for non-invasive genetic studies of small cryptic mammals

Tobias Erik Reiners, Jorge A. Encarnação, Volkmar Wolters

► To cite this version:

Tobias Erik Reiners, Jorge A. Encarnação, Volkmar Wolters. An optimized hair trap for non-invasive genetic studies of small cryptic mammals. *European Journal of Wildlife Research*, 2011, 57 (4), pp.991-995. 10.1007/s10344-011-0543-9 . hal-00697239

HAL Id: hal-00697239

<https://hal.science/hal-00697239>

Submitted on 15 May 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

An optimized hair trap for non-invasive genetic studies of small cryptic mammals

Tobias Erik Reiners · Jorge A. Encarnação ·
Volkmar Wolters

Received: 18 November 2010 / Revised: 26 April 2011 / Accepted: 2 May 2011 / Published online: 15 May 2011
© Springer-Verlag 2011

Abstract As sample quality and quantity is a crucial factor in non-invasive genetics, we focused on the improvement of sampling efficiency of glue hair traps. We invented an optimized hair trap with moveable parts which enhanced sampling of high-quality genetic material. With the aid of the optimized hair trap, we were able to remotely pluck a sufficient amount of hair bulbs from our study animal the common hamster (*Cricetus cricetus*) with a trapping success of 49.3% after one survey night. The number of collected hairs with bulbs ranged between 1 and 50, with an average of 20.7 ± 14.8 . Subsequently, the use of the hair trap in combination with a simplified laboratory routine allowed us to amplify species-specific microsatellites with an amplification success of 96.2% and ADO of 4.6%. This optimized trap may find usage for species identification or could be used as an instrument for long-term genetic monitoring of mammal populations.

Keywords *Cricetus cricetus* · Remote collecting · Microsatellites · Non-invasive genetic sampling

Communicated by C. Gortázar

T. E. Reiners (✉) · J. A. Encarnação
Department of Animal Ecology, Mammalian Ecology Group,
Justus-Liebig-University of Giessen,
Heinrich-Buff-Ring 26,
35392 Giessen, Germany
e-mail: tobias.reiners@bio.uni-giessen.de

V. Wolters
Department of Animal Ecology,
Justus-Liebig-University of Giessen,
Heinrich-Buff-Ring 26,
35392 Giessen, Germany

Introduction

Non-invasive sampling techniques for genetic studies on mammals become increasingly important due to ethical, logistical, and scientific restrictions. Those methods allow researchers to collect samples without any observer bias or negative consequences for individuals due to handling or stress. Therefore, they are most frequently applied in studies with rare, cryptic, or endangered species (Piggott and Taylor 2003; Waits and Paetkau 2005). The source of sample material together with DNA extraction methods and PCR conditions is among the most crucial sources of error (Beja Pereira et al. 2009; Broquet et al. 2007; Taberlet et al. 1999). Hair samples are a widely used, non-lethal source of DNA, and recent advantages in molecular techniques provide reliable results through the application of strict guidelines and high-performance analysis techniques (Beja Pereira et al. 2009; Mullins et al. 2009). In contrast, very little attention is directed towards the invention, construction, and application of effective traps that facilitate the collection of high-quality sample material. When using hairs, in particular their bulbs, as a source for DNA, these are needed in sufficient amounts containing enough follicular tissue. A minimum of three to ten hair bulbs are needed to reduce genotyping errors significantly (Gagneux et al. 1997). The use of hair traps with mechanical devices (e.g., barbwire) or attached glue-like substances to collect plucked hairs is not ideal as it rarely collects high proportions of hairs with roots (Ebert et al. 2010; Valderrama et al. 1999). Here, we present a new, cost-efficient, optimized hair trap for non-invasive sampling that uses movable parts to collect efficiently hairs with

bulbs. We tested the effectiveness for non-invasive genetic studies by quantifying the amount of hair bulbs, micro-satellite amplification success, and genotyping error rates in the endangered common hamster *Cricetus cricetus* L., 1758.

Materials and methods

Study area

This study was conducted between July and September 2009 in Hesse (Germany). In three different study areas, 223 ha of arable fields were surveyed for the presence of common hamster burrows.

Hair trap

The hair trap is built out of a 5-cm-long PVC pipe with a diameter of 7.5 cm to match the average diameter of common hamster burrows. Through two holes at the upper periphery of the pipe, a 7.5-cm-long steel nail is placed. For trapping hairs, a plastic roll with a diameter of 1.2 cm was attached using the nail as axis, and double-faced adhesive tape was then wrapped around the roll (Fig. 1a and b). When animals went through the pipe, the plastic roll with

the adhesive tape rotates which then results in a vertical extraction of hairs. Hair traps were placed in burrow entrances in the late afternoon and surveyed for hairs early in the next morning (Fig. 1c and d). For each applied trap, the amount of hairs was documented (Fig. 1e and f).

Sample storage

The appropriate storage and preservation of non-invasive samples is crucial (Beja Pereira et al. 2009). As DNA comes in small quantities and low quality, it should be protected against heat stress or sunlight (Taberlet et al. 1999). As freezing and refrigeration result in physical stress, we used dry preservation. We stored samples in 125-ml vials with 5.5-cm height and airtight closure (Sarstaedt®). To dry samples, we attached two packages of Silica Gel (Wisepac®) to each vial. After successful application in the field, the whole glue-plastic roll was moved to a vial.

DNA isolation and genetic analyses

DNA extraction was performed using 20% Chelex following standard protocols (Walsh et al. 1991). This method is used in several other studies with small-sized mammals (Sloane et al. 2000; Moncrief et al. 2008). The roots of ten

Fig. 1 Materials and application of the optimized hair trap. The trap consists of a PVC pipe with two holes, an iron nail, and a plastic roll (a). The plastic roll is wrapped with double-faced adhesive tape and attached inside the pipe by the nail (b). Traps are placed in entrances

of burrows (c), and as an animal passes, hairs are plucked by the rolling tape (d). Most of plucked hair contained bulbs with a sufficient amount of follicular tissue (e and f)

Table 1 Analyzed loci of *C. cricetus* used in the study, listed with GenBank accession no., locus name, repeat motif and primer sequences (sequences in capitals stand for unmodified primers, lowercase sequences for modified primers), assignment to multiplex reaction groups, number of detected alleles, size range of alleles in base pairs (bp), expected heterozygosity (H_E), amplification success, and allelic dropout (ADO)

GenBank accession no.	Locus	Repeat motif	Primer sequence (5'–3')	Multiplex group	Allele numbers	Size range (bp)	H_E	Amplification success (%)	ADO (%)
AM167541	IPK01	(TC) ₂₂ N ₄ (TG) ₁₄	F CTGTGGGCTGATCTCTCTC R gcaagagagacaaagtgig	1	9	99–129	0.425	97.8	2.7
AM167544	IPK05	(TC) ₁₉ (AC) ₇	F gttccatagcccttcactc R cccatttgcaagaatcc	1	6	153–165	0.608	97.8	2.7
AM167545	IPK06	(CT) ₂₄	F TGACCTGAGTTTGATTTC R ATTTGTGCATGTGCATGAC	1	3	80–84	0.505	97.8	0.0
AM167546	IPK07	(CT) ₂₂ N ₂₉ (CT) ₇ N ₁ (TC) ₈	F tccaggtagatttgagtc R tgaacaacatagtagagagagg	1	8	154–170	0.434	97.2	2.0
AM167547	IPK09	(GT) ₂₁	F ctcaacagtcctgaatac R gaggaacaaagfggaaaag	1	5	177–185	0.510	96.1	0.0
AM167548	IPK12	(GT) ₂₂	F gtaggggtccatgagtgag R agaccagcaaatcaaaagag	1	4	100–106	0.522	98.9	0.0
AJ532554	Ccrp4	(GT) ₁₈	F tgagatgatgcatgtgtgc R tticatatgtgtccaagac	2	4	150–164	0.197	95.5	13.5
AJ532556	Ccrp10	(CA) ₂₃	F tatatgcacatcatgtcaeg R ggctcttaagaatcagggtgig	2	5	99–107	0.196	92.9	14.8
AJ532557	Ccrp11	(GT) ₂₂	F geatttgccactatcgttc R tccaactcctcaagttgic	2	4	148–162	0.400	93.9	7.0
AJ532563	Ccrp20	(TC) ₂₃ (CT) ₉ (CT) ₆	F cacttggatctgactttgig R tggggattttggatgattac	2	7	136–142	0.681	94.4	13.8

hairs were incubated in 200 µl of a 20% Chelex solution for 2 h. To test if extracts are sufficient for genetic studies, ten species-specific microsatellites were used (Jakob and Mammen 2006; Neumann and Jansman 2004). Primers were modified, respectively new designed, following the guidelines for non-invasive sampling (Broquet et al. 2007; Taberlet et al. 1999) (Table 1). For PCR, the Qiagen® Multiplex PCR Kit was used, arranging primers into two multiplex reactions (Table 1). All reactions were carried out in a separate laboratory with an Eppendorf® MasterCycler Gradient using a 10-µl reaction volume with 5 µl 2× QIAGEN® Multiplex PCR Master Mix combined with the touchdown-PCR technique to increase annealing specificity (Hecker and Roux 1996).

Assessment of amplification success and error rates

Amplification success was measured by counting the amount of failed amplifications for each locus. To test for allelic dropout (ADO), reactions ($n=77$) and isolations ($n=41$) were repeated randomly. ADO was calculated using the equation presented in Broquet and Petit (2004).

Results

During the study, we applied 199 hair traps to common hamster burrows. At 144 burrows, the trap contained hairs in 98 cases resulting in a 49.3% trapping success after one survey night. The number of collected hairs with bulbs per trap was between 1 and 50, with an average of 20.7 ± 14.8 ($n=98$) bulbs. The proportion of successful amplifications for the species-specific microsatellites over all loci was 96.2%. The rate of ADO over all loci was 4.6% (Table 1). False alleles were not detected in the dataset. Our approach provided enough material to successfully genotype 88 different individuals.

Discussion

Recent developments in molecular techniques and genotyping pronounce the importance of non-invasive sampling (Waits and Paetkau 2005). But most non-invasive techniques are still expensive, because reliable and replicated samples of genetic material are only available using costly methods (Navidi et al. 1992). With our approach, we will reduce costs significantly by improving a basal technique for non-invasive genetic sampling. Our study showed that the newly developed trap is easy to apply and provides a high gain of genetic material. Additionally, the glue-containing part is fast and easily replaceable, and samples can be directly conserved for further analyses.

One of the major drawbacks of non-invasive sampling, the generally small amount and low quality of genetic material (Piggott and Taylor 2003), has been overcome by our current approach. As demonstrated, we were able to amplify species-specific microsatellites with high amplification success (96.2%) and comparably low error rate (ADO, 4.6%). These rates are markedly better than the mean amplification success of 80.2% and mean allelic dropout rate of 18.7% reported from a review of non-invasive studies using hair samples as source for genetic material (Broquet et al. 2007).

The effectiveness of the hair trap with the glue roll was only shown for one burrow-using species. Our approach may also find use in studies with other species as comparable studies using tubes for species identification without genetic sampling were already conducted (Mortelliti and Boitani 2008; Ruiz-Gonzalez et al. 2008; Sanecki and Green 2005). As a conclusion, this trap is well suited for short- and long-term monitoring studies in population genetics of small mammals and provides a suitable technique for highly replicated studies on relevant, large spatial scales.

References

- Beja Pereira A, Oliveira R, Alves PC, Schwartz MK, Luikart G (2009) Advancing ecological understandings through technological transformations in noninvasive genetics. *Mol Ecol Resour* 9 (5):1279–1301
- Broquet T, Petit E (2004) Quantifying genotyping errors in noninvasive population genetics. *Mol Ecol* 13(11):3601–3608
- Broquet T, Ménard N, Petit E (2007) Noninvasive population genetics: a review of sample source, diet, fragment length and microsatellite motif effects on amplification success and genotyping error rates. *Conserv Genet* 8(1):249–260
- Ebert C, Huckschlag D, Schulz HK, Hohmann U (2010) Can hair traps sample wild boar (*Sus scrofa*) randomly for the purpose of non-invasive population estimation? *Eur J Wildl Res* 56(4):583–590
- Gagneux P, Boesch C, Woodruff DS (1997) Microsatellite scoring errors associated with noninvasive genotyping based on nuclear DNA amplified from shed hair. *Mol Ecol* 6(9):861–868
- Hecker KH, Roux KH (1996) High and low annealing temperatures increase both specificity and yield in touchdown and stepdown PCR. *Biotechniques* 20(3):478–485
- Jakob SS, Mammen K (2006) Eight new polymorphic microsatellite loci for genetic analyses in the endangered common hamster (*Cricetus cricetus* L.). *Mol Ecol Notes* 6(2):511–513
- Moncrief ND, Van den Bussche RA, Dueser RD, Loftis D, Cockett NE, Culver M (2008) Diagnostic genetic marker that differentiates eastern fox squirrels from eastern gray squirrels. *J Wildl Manage* 72(1):320–323
- Mortelliti A, Boitani L (2008) Inferring red squirrel (*Sciurus vulgaris*) absence with hair tubes surveys: a sampling protocol. *Eur J Wildl Res* 54(2):353–356
- Mullins J, Statham M, Roche T, Turner P, O'Reilly C (2009) Remotely plucked hair genotyping: a reliable and non-invasive

- method for censusing pine marten (*Martes martes*, L. 1758) populations. Eur J Wildl Res 56(3):443–453
- Navidi W, Arnheim N, Waterman MS (1992) A multiple-tubes approach for accurate genotyping of very small DNA samples by using PCR-statistical considerations. Am J Hum Genet 50(2):347–359
- Neumann K, Jansman H (2004) Polymorphic microsatellites for the analysis of endangered common hamster populations (*Cricetus cricetus* L.). Conserv Genet 5(1):127–130
- Piggott MP, Taylor AC (2003) Remote collection of animal DNA and its applications in conservation management and understanding the population biology of rare and cryptic species. Wildl Res 30(1):1–13
- Ruiz-Gonzalez A, Rubines J, Berdion O, Gomez-Moliner BJ (2008) A non-invasive genetic method to identify the sympatric mustelids pine marten (*Martes martes*) and stone marten (*Martes foina*): preliminary distribution survey on the northern Iberian Peninsula. Eur J Wildl Res 54(2):253–261
- Sanecki GM, Green K (2005) A technique for using hair tubes beneath the snowpack to detect winter-active small mammals in the subnivean space. Eur J Wildl Res 51(1):41–47
- Sloane MA, Sunnucks P, Alpers D, Beheregaray LB, Taylor AC (2000) Highly reliable genetic identification of individual northern hairy-nosed wombats from single remotely collected hairs: a feasible censusing method. Mol Ecol 9(9):1233–1240
- Taberlet P, Waits LP, Luikart G (1999) Noninvasive genetic sampling: look before you leap. Trends Ecol Evol 14(8):323–327
- Valderrama X, Karesh WB, Wildman DE, Melnick DJ (1999) Noninvasive methods for collecting fresh hair tissue. Mol Ecol 8(10):1749–1750
- Waits LP, Paetkau D (2005) Noninvasive genetic sampling tools for wildlife biologists: a review of applications and recommendations for accurate data collection. J Wildl Manage 69(4):1419–1433
- Walsh PS, Metzger DA, Higuchi R (1991) Chelex® 100 as a medium for simple extraction of DNA for PCR-based typing from forensic material. Biotechniques 10(4):506–513