

**Rates of convergence in the strong invariance principle
for non adapted sequences. Application to ergodic
automorphisms of the torus**

Jérôme Dedecker, Florence Merlevède, Françoise Pene

► **To cite this version:**

Jérôme Dedecker, Florence Merlevède, Françoise Pene. Rates of convergence in the strong invariance principle for non adapted sequences. Application to ergodic automorphisms of the torus. High dimensional probability 6, Oct 2011, Banff, Canada. 66, pp.113-138, 2013, High dimensional probability 6, Progr. Probab. <hal-00713797>

HAL Id: hal-00713797

<https://hal.archives-ouvertes.fr/hal-00713797>

Submitted on 2 Jul 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Rates of convergence in the strong invariance principle for non adapted sequences. Application to ergodic automorphisms of the torus

Jérôme Dedecker, Florence Merlevède and Françoise Pène

Abstract. In this paper, we give rates of convergence in the strong invariance principle for non-adapted sequences satisfying projective criteria. The results apply to the iterates of ergodic automorphisms T of the d -dimensional torus \mathbb{T}^d , even in the non hyperbolic case. In this context, we give a large class of unbounded function f from \mathbb{T}^d to \mathbb{R} , for which the partial sum $f \circ T + f \circ T^2 + \dots + f \circ T^n$ satisfies a strong invariance principle with an explicit rate of convergence.

Mathematics Subject Classification (2010). 60F17; 37D30.

Keywords. almost sure invariance principle, strong approximations, non adapted sequences, ergodic automorphisms of the torus.

1. Introduction and notations

Let $(\Omega, \mathcal{A}, \mathbb{P})$ be a probability space, and $T : \Omega \mapsto \Omega$ be a bijective bimeasurable transformation preserving the probability \mathbb{P} . For a σ -algebra \mathcal{F}_0 satisfying $\mathcal{F}_0 \subseteq T^{-1}(\mathcal{F}_0)$, we define the nondecreasing filtration $(\mathcal{F}_i)_{i \in \mathbb{Z}}$ by $\mathcal{F}_i = T^{-i}(\mathcal{F}_0)$. The \mathbb{L}^p norm of a random variable X is denoted by $\|X\|_p = (\mathbb{E}(|X|^p))^{1/p}$.

Let X_0 be a real-valued and square integrable random variable such that $\mathbb{E}(X_0) = 0$, and define the stationary sequence $(X_i)_{i \in \mathbb{Z}}$ by $X_i = X_0 \circ T^i$. Define then the partial sum by $S_n = X_1 + X_2 + \dots + X_n$. According to the Birkhoff-Khinchine theorem, S_n satisfies a strong law of large numbers. One can go further in the study of the statistical properties of S_n . We study here the rate of convergence in the almost sure invariance principle (ASIP). More precisely, we give conditions under which there exists a sequence of

F. Pène is partially supported by the French ANR projects MEMEMO2 and PERTURBATIONS.

independent identically distributed (iid) Gaussian random variables $(Z_i)_{i \geq 1}$ such that

$$\sup_{1 \leq k \leq n} \left| \sum_{i=1}^k (X_i - Z_i) \right| = o(n^{1/p} L(n)) \quad \text{almost surely,} \quad (1.1)$$

for $p \in]2, 4]$ and L an explicit slowly varying function. Let us recall that, in the iid case, Komlós, Major and Tusnády [13] and Major [18] obtained an ASIP with the optimal rate $o(n^{1/p})$ in (1.1) as soon as the random variables admit a moment of order p .

Since the seminal paper by Philipp and Stout [25], many authors have considered this problem in a dependent context, but most of the papers deal with the adapted case, when X_0 is \mathcal{F}_0 measurable (for instance, \mathcal{F}_0 is the past σ -algebra $\sigma(X_i, i \leq 0)$). Unfortunately, it is quite common to encounter dynamical systems for which the natural filtration does not permit the control of any quantity involving terms of the type $\|\mathbb{E}(X_n | \mathcal{F}_0)\|_p$.

In this paper, we shall not assume that X_0 is \mathcal{F}_0 -measurable, and we shall give conditions on the quantities $\|\mathbb{E}(X_n | \mathcal{F}_0)\|_p$, $\|X_{-n} - \mathbb{E}(X_{-n} | \mathcal{F}_0)\|_p$ and $\|\mathbb{E}(S_n^2 | \mathcal{F}_{-n}) - \mathbb{E}(S_n^2)\|_{p/2}$ for (1.1) to hold (see Theorems 3.1 and 3.2 of Section 3). These conditions are in the same spirit as those given by Gordin [7] for $p = 2$ to get the usual central limit theorem. Our proof is based on the approximation

$$\sum_{i=1}^n X_i = M_n + R_n$$

by the martingale $M_n = d_1 + d_2 + \dots + d_n$, where d_i is the martingale difference

$$d_i = \sum_{k \in \mathbb{Z}} (\mathbb{E}(X_k | \mathcal{F}_i) - \mathbb{E}(X_k | \mathcal{F}_{i-1}))$$

introduced by Gordin [7] and Heyde [10]. In the adapted case, similar conditions are given in the recent paper [2], together with a long list of applications.

In the non adapted case, it is easy to see that our results apply to a large class of two-sided functions of iid sequences, or two-sided functions of absolutely regular sequences. But they also apply to very complicated dynamical systems, for which such a representation by functions of absolutely regular sequences is not available. In the next section, we consider the case where T is an ergodic automorphism of the d -dimensional torus \mathbb{T}^d , and \mathbb{P} is the Lebesgue measure on \mathbb{T}^d . In this context, we use the σ -algebras \mathcal{F}_i considered by Le Borgne [14]. As a consequence of Theorem 2.1, we obtain that (1.1) holds for $p = 4$ and $X_i = f \circ T^i$, where $f : \mathbb{T}^d \rightarrow \mathbb{R}$, as soon as the Fourier coefficients $(c_{\mathbf{k}})_{\mathbf{k} \in \mathbb{Z}^d}$ of f are such that

$$|c_{\mathbf{k}}| \leq A \prod_{i=1}^d \frac{1}{(1 + |k_i|)^{3/4} \log^\alpha(2 + |k_i|)} \quad \text{for some } \alpha > 13/8.$$

We also get that there exists a positive ε such that

$$\sup_{1 \leq k \leq n} \left| \sum_{i=1}^k (X_i - Z_i) \right| = o(n^{1/2-\varepsilon}) \quad \text{almost surely,}$$

as soon as

$$|c_{\mathbf{k}}| \leq A \prod_{i=1}^d \frac{1}{(1 + |k_i|)^\delta} \quad \text{for some } \delta > 1/2.$$

These rates of convergence in the almost sure invariance principle complement the results by Leonov [16] and Le Borgne [14] for the central limit theorem and the almost sure invariance principle respectively. Let us mention that Dolgopyat [5] established an ASIP with the rate $o(n^{1/2-\varepsilon})$ (for some $\varepsilon > 0$) valid for ergodic automorphisms of the torus and f a Hölder continuous function. Thanks to the decorrelation estimates obtained in [15], the rate for Hölder observables can be improved by applying the general result of Gouëzel in [8] to get the rate $o(n^{1/4+\varepsilon})$ for every $\varepsilon > 0$, and by applying the results of the present paper to get the rate $o(n^{1/4}L(n))$. To our knowledge, the present work gives the first strong approximation results for such partially hyperbolic transformations T for unbounded (and then non continuous) functions f .

To conclude, let us mention some previous works in the context of dynamical systems: several results have been established with the rate $o(n^{1/2-\varepsilon})$ for some $\varepsilon > 0$ (see [11, 4, 5, 24, 19]). Results giving a rate in $o(n^{1/4+\varepsilon})$ for every $\varepsilon > 0$ can be found in [21, 6, 20, 8]. Most of these results hold for bounded functions f .

Let us reiterate that we can reach the rate $o(n^{1/4}L(n))$ instead of $o(n^{1/4+\varepsilon})$ for every $\varepsilon > 0$. Moreover, our conditions giving the rate $o(n^{1/p}L(n))$ are related to moments of order p of f . Such results are not very common in the context of dynamical systems (let us mention [8] in the particular case of Gibbs-Markov maps, and [3, 23] for generalized Pommeau-Manneville maps).

2. ASIP with rates for ergodic automorphisms of the torus

A probability dynamical system $(\Omega, \mathcal{F}, \mathbb{P}, T)$ is given by a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ and a measurable \mathbb{P} -preserving transformation $T : \Omega \rightarrow \Omega$. Such a dynamical system is said to be ergodic if the only $A \in \mathcal{F}$ such that $T^{-1}A = A$ a.s. are the sets of probability 0 or 1.

If $(\Omega, \mathcal{F}, \mathbb{P}, T)$ is ergodic, the study of the stochastic properties of the stationary sequence $(f \circ T^k)_{k \geq 1}$ starts with the Birkhoff-Khinchine theorem [1, 12]. This theorem ensures that, for every integrable function $f : \Omega \rightarrow \mathbb{R}$, the sequence $n^{-1} \sum_{k=1}^n f \circ T^k$ converges almost surely to $\mathbb{E}(f)$. This means that $(f \circ T^k)_{k \geq 1}$ satisfies a strong law of large number. A natural question is then to investigate further stochastic properties of $(f \circ T^k)_{k \geq 1}$.

We illustrate our general Theorems 3.1 and 3.2 by a concrete example of invertible non hyperbolic dynamical system, which is actually partially hyperbolic. We shall prove a strong invariance principle for a large class of unbounded functions f , with a rate depending on the rate of convergence to

zero of the Fourier coefficients of f . In this context, we use the σ -algebras \mathcal{F}_i considered by Le Borgne [14]. The stationary sequence $(T^i)_{i \in \mathbb{Z}}$ is non adapted to this filtration $(\mathcal{F}_i)_{i \in \mathbb{Z}}$, in the sense that T^i is not \mathcal{F}_i measurable. This is an important difference with the classical probabilistic situation, where the study of stationary sequences can often be done with the help of a natural “past” filtration (think of stationary Markov chains, or of causal linear processes).

Let $d \geq 2$. We consider a group automorphism T of the torus $\mathbb{T}^d = \mathbb{R}^d / \mathbb{Z}^d$. For every $x \in \mathbb{R}^d$, we write \bar{x} its class in \mathbb{T}^d . We recall that T is the quotient map of a linear map $\tilde{T} : \mathbb{R}^d \rightarrow \mathbb{R}^d$ given by $\tilde{T}(x) = S \cdot x$, where S is a $d \times d$ -matrix with integer entries and with determinant 1 or -1. The map $x \mapsto S \cdot x$ preserves the infinite Lebesgue measure λ on \mathbb{R}^d and T preserves the probability Lebesgue measure $\bar{\lambda}$. We suppose that T is ergodic, which is equivalent to the fact that no eigenvalue of S is a root of the unity. In this case, it is known that the spectral radius of S is larger than one (and so S admits at least an eigenvalue of modulus larger than one and at least an eigenvalue of modulus smaller than one). This hypothesis holds true in the case of hyperbolic automorphisms of the torus (i.e. in the case when no eigenvalue of S has modulus one) but is much weaker. Indeed, as mentioned in [14], the following matrix gives an example of an ergodic non hyperbolic automorphism of \mathbb{T}^4 :

$$S := \begin{pmatrix} 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & 2 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 2 \end{pmatrix}.$$

When T is ergodic and non hyperbolic, the dynamical system $(\mathbb{T}^d, T, \bar{\lambda})$ has no Markov partition. However, it is possible to construct some measurable partition [17], to prove a central limit theorem [16]. Moreover, in [14], Le Borgne proved the functional central limit theorem and the Strassen strong invariance principle for $(X_k = f \circ T^k)_k$ under weak hypotheses on f , thanks to Gordin’s method and to the partitions studied by Lind in [17].

We give here rates of convergence in the strong invariance principle for $(X_k = f \circ T^k)_k$ under conditions on the Fourier coefficients of $f : \mathbb{T}^d \rightarrow \mathbb{R}$. In what follows, for $\mathbf{k} \in \mathbb{Z}^d$, we denote by $|\mathbf{k}| = \max_{i \in \{1, \dots, d\}} |k_i|$.

Theorem 2.1. *Let T be an ergodic automorphism of \mathbb{T}^d with the notations as above. Let $p \in]2, 4]$ and q be its conjugate exponent. Let $f : \mathbb{T}^d \rightarrow \mathbb{R}$ be a centered function with Fourier coefficients $(c_{\mathbf{k}})_{\mathbf{k} \in \mathbb{Z}^d}$ satisfying, for any integer $b \geq 2$,*

$$\sum_{|\mathbf{k}| \geq b} |c_{\mathbf{k}}|^q \leq R \log^{-\theta}(b) \quad \text{for some } \theta > \frac{p^2 - 2}{p(p-1)}, \quad (2.2)$$

and

$$\sum_{|\mathbf{k}| \geq b} |c_{\mathbf{k}}|^2 \leq R \log^{-\beta}(b) \quad \text{for some } \beta > \frac{3p-4}{p}. \quad (2.3)$$

Then the series

$$\sigma^2 = \bar{\lambda}((f - \bar{\lambda}(f))^2) + 2 \sum_{k>0} \bar{\lambda}((f - \bar{\lambda}(f))f \circ T^k)$$

converges absolutely and, enlarging \mathbb{T}^d if necessary, there exists a sequence $(Z_i)_{i \geq 1}$ of iid gaussian random variables with zero mean and variance σ^2 such that, for any $t > 2/p$,

$$\sup_{1 \leq k \leq n} \left| \sum_{i=1}^k f \circ T^i - \sum_{i=1}^k Z_i \right| = o(n^{1/p}(\log n)^{(t+1)/2}) \text{ almost surely, as } n \rightarrow \infty. \quad (2.4)$$

Observe that (2.3) follows from (2.2) provided that $\theta > (3p-4)/(2p-2)$. Hence, (2.2) and (2.3) are both satisfied as soon as

$$\sum_{|\mathbf{k}| \geq b} |c_{\mathbf{k}}|^q \leq R \log^{-\theta}(b) \text{ for some } \theta > \frac{3p-4}{2(p-1)}.$$

Example. Let $p \in]2, 4]$. If we assume that the Fourier coefficients of f are such that

$$|c_{\mathbf{k}}| \leq A \prod_{i=1}^d \frac{1}{(1 + |k_i|)^{1/q} \log^\alpha(2 + |k_i|)}, \quad (2.5)$$

for some positive constant A , then the conditions (2.2) and (2.3) are both satisfied provided that $\alpha > (2p^2 - p - 2)/p^2$.

Let us now compare our hypotheses on Fourier coefficients with those appearing in other works. In [16], Leonov proved a central limit theorem (possibly degenerated) when

$$|c_{\mathbf{k}}| \leq A \prod_{i=1}^d \frac{1}{(1 + |k_i|)^{1/2} \log^\alpha(2 + |k_i|)} \text{ for some } \alpha > 3/2. \quad (2.6)$$

In [14], Le Borgne proved the functional central limit theorem and the Strassen strong invariance principle when (2.3) holds true with $\beta > 2$ (and when f is not a coboundary), which is a weaker condition than (2.6). Observe that, as p converges to 2, $(p^2 - 2)/(p(p - 1))$ and $(3p - 4)/p$ both converge to 1.

3. Probabilistic results

In the rest of the paper, we shall use the following notations: $\mathbb{E}_k(X) = \mathbb{E}(X|\mathcal{F}_k)$, and $a_n \ll b_n$ means that there exists a numerical constant C not depending on n such that $a_n \leq Cb_n$, for all positive integers n .

In this section, we give rates of convergence in the strong invariance principle under projective criteria for stationary sequences that are non necessarily adapted to \mathcal{F}_i .

Theorem 3.1. *Let $2 < p < 4$ and $t > 2/p$. Assume that X_0 belongs to \mathbb{L}^p , that*

$$\sum_{n \geq 2} \frac{n^{p-1}}{n^{2/p}(\log n)^{(t-1)p/2}} (\|\mathbb{E}_0(X_n)\|_p^p + \|X_{-n} - \mathbb{E}_0(X_{-n})\|_p^p) < \infty, \quad (3.1)$$

and that

$$\sum_{n \geq 2} \frac{n^{3p/4}}{n^2(\log n)^{(t-1)p/2}} (\|\mathbb{E}_0(X_n)\|_2^{p/2} + \|X_{-n} - \mathbb{E}_0(X_{-n})\|_2^{p/2}) < \infty. \quad (3.2)$$

Assume in addition that there exists a positive integer m such that

$$\sum_{n \geq 2} \frac{1}{n^2(\log n)^{(t-1)p/2}} \|\mathbb{E}_{-nm}(S_n^2) - \mathbb{E}(S_n^2)\|_{p/2}^{p/2} < \infty. \quad (3.3)$$

Then $n^{-1}\mathbb{E}(S_n^2)$ converges to $\sigma^2 = \sum_{k \in \mathbb{Z}} \text{Cov}(X_0, X_k)$ and, enlarging Ω if necessary, there exists a sequence $(Z_i)_{i \geq 1}$ of iid Gaussian random variables with zero mean and variance σ^2 such that

$$\sup_{1 \leq k \leq n} \left| S_k - \sum_{i=1}^k Z_i \right| = o(n^{1/p}(\log n)^{(t+1)/2}) \quad \text{almost surely, as } n \rightarrow \infty. \quad (3.4)$$

Theorem 3.2. *Let $t > 1/2$. Assume that X_0 belongs to \mathbb{L}^4 and that the conditions (3.1) and (3.3) hold with $p = 4$. Assume in addition that*

$$\sum_{n \geq 2} n(\log n)^{4-2t} (\|\mathbb{E}_0(X_n)\|_2^2 + \|X_{-n} - \mathbb{E}_0(X_{-n})\|_2^2) < \infty. \quad (3.5)$$

Then the conclusion of Theorem 3.1 holds with $p = 4$.

Of course, if $(X_i)_{i \in \mathbb{Z}}$ is a sequence of iid random variables in \mathbb{L}^p , then, taking $\mathcal{F}_i = \sigma(X_k, k \leq i)$, all the conditions (3.1), (3.2), (3.3) and (3.5) are satisfied. In that particular case, we obtain an extra power of $\log(n)$ compared to the optimal rate $n^{1/p}$.

The conditions (3.1), (3.2) and (3.5) are similar to the conditions given by Gordin [7] when $p = 2$ for the central limit theorem, and hence Theorem 3.1 and 3.2 have the same range of applicability as Gordin's result.

For the proof of Theorems 3.1 and 3.2, we shall use the approximating martingale M_n introduced by Gordin [7], and we shall give an appropriate upper bound on $R_n = S_n - M_n$. The next step is to get a strong approximation result for the martingale M_n . This will be done by applying Proposition 5.1 in [2], which itself is based on the Skorohod embedding for martingales, as in [26].

3.1. Proofs of Theorems 3.1 and 3.2

Proof. We first notice that since $p > 2$, (3.1) implies that

$$\sum_{n > 0} n^{-1/p} \|\mathbb{E}_0(X_n)\|_p < \infty \quad \text{and} \quad \sum_{n > 0} n^{-1/p} \|X_{-n} - \mathbb{E}_0(X_{-n})\|_p < \infty$$

(apply Hölder's inequality to see this). Let $P_k(X) = \mathbb{E}_k(X) - \mathbb{E}_{k-1}(X)$. Using Lemma 5.1 of the appendix with $q = 1$, we infer that

$$\sum_{k \in \mathbb{Z}} \|P_0(X_k)\|_p < \infty. \quad (3.6)$$

In addition the condition (3.6) implies that $n^{-1}\mathbb{E}(S_n^2)$ converges to the quantity $\sigma^2 = \sum_{k \in \mathbb{Z}} \text{Cov}(X_0, X_k)$.

Let now $d_0 := \sum_{j \in \mathbb{Z}} P_0(X_j)$. Then d_0 belongs to \mathbb{L}^p and $\mathbb{E}(d_0 | \mathcal{F}_{-1}) = 0$. Let $d_i := d_0 \circ T^i$ for all $i \in \mathbb{Z}$. Then $(d_i)_{i \in \mathbb{Z}}$ is a stationary sequence of martingale differences in \mathbb{L}^p . Let

$$M_n := \sum_{i=1}^n d_i \quad \text{and} \quad R_n := S_n - M_n.$$

The theorems will be proven if we can show that

$$R_n = o(n^{1/p}(\log n)^{(t+1)/2}) \quad \text{almost surely as } n \rightarrow \infty, \quad (3.7)$$

and that (3.4) holds true with M_k replacing S_k . Since $\mathbb{E}(d_0^2) = \sigma^2$ and $t > p/2$, according to Proposition 5.1 in [2] (applied with $\psi(n) := n^{2/p}(\log n)^t$), to prove that (3.4) holds true with M_k replacing S_k , it suffices to prove that

$$\sum_{n \geq 2} \frac{1}{n^2 (\log n)^{(t-1)p/2}} \|\mathbb{E}_0(M_n^2) - \mathbb{E}(M_n^2)\|_{p/2}^{p/2} < \infty. \quad (3.8)$$

By standard arguments, (3.7) will be satisfied if we can show that

$$\sum_{r > 0} \frac{\|\max_{1 \leq \ell \leq 2^r} |R_\ell|\|_p^p}{2^r r^{(t+1)p/2}} < \infty. \quad (3.9)$$

Now, by stationarity, $\|\max_{1 \leq \ell \leq 2^r} |R_\ell|\|_p \ll 2^{r/p} \sum_{k=0}^r 2^{-k/p} \|R_{2^k}\|_p$ (see for instance inequality (6) in [27]) and for all $i, j \geq 0$, $\|R_{i+j}\|_q \leq \|R_i\|_q + \|R_j\|_q$. Applying then Item 1 of Lemma 37 in [22], we derive that for any integer n in $[2^r, 2^{r+1}[$,

$$\left\| \max_{1 \leq \ell \leq 2^r} |R_\ell| \right\|_p \ll n^{1/p} \sum_{k=1}^n k^{-(1+1/p)} \|R_k\|_p. \quad (3.10)$$

Therefore using (3.10) followed by an application of Hölder's inequality, we get that for any $\alpha < 1$,

$$\begin{aligned} \sum_{r > 0} \frac{\|\max_{1 \leq \ell \leq 2^r} |R_\ell|\|_p^p}{2^r r^{(t+1)p/2}} &\ll \sum_{n \geq 2} \frac{1}{n (\log n)^{(t+1)p/2}} \left(\sum_{k=1}^n k^{-(1+1/p)} \|R_k\|_p \right)^p \\ &\ll \sum_{n \geq 2} \frac{(\log n)^{(p-1)(1-\alpha)}}{n (\log n)^{(t+1)p/2}} \sum_{k=1}^n k^{-2} (\log k)^{\alpha(p-1)} \|R_k\|_p^p. \end{aligned}$$

Hence taking $\alpha \in]1 - p/(2(p-1)), 1[$ and changing the order of summation, we infer that (3.9) and then (3.7) hold provided that

$$\sum_{n \geq 1} \frac{\|R_n\|_p^p}{n^2(\log n)^{(t-1)p/2}} < \infty. \quad (3.11)$$

On an other hand, we shall prove that condition (3.8) is implied by: there exists a positive finite integer m such that

$$\sum_{n \geq 2} \frac{1}{n^2(\log n)^{(t-1)p/2}} \|\mathbb{E}_{-nm}(M_n^2) - \mathbb{E}(M_n^2)\|_{p/2}^{p/2} < \infty. \quad (3.12)$$

For any nonnegative integer i , we set $V_i := \|\mathbb{E}_0(M_i^2) - \mathbb{E}(M_i^2)\|_{p/2}$. Using that M_n is a martingale, we infer that, for any nonnegative integers i and j ,

$$V_{i+j} \leq V_i + V_j. \quad (3.13)$$

Let now $n \in [2^k, 2^{k+1} - 1] \cap \mathbb{N}$, and write its binary expansion:

$$n = \sum_{\ell=0}^k 2^\ell b_\ell \quad \text{where } b_k = 1 \text{ and } b_j \in \{0, 1\} \text{ for } j = 0, \dots, k-1.$$

Inequality (3.13) combined with Hölder's inequality implies that, for any $\eta > 0$,

$$V_n^{p/2} \leq \left(\sum_{\ell=0}^k V_{2^\ell} \right)^{p/2} \ll 2^{\eta p(k+1)/2} \sum_{\ell=0}^k \left(\frac{V_{2^\ell}}{2^{\eta \ell}} \right)^{p/2}. \quad (3.14)$$

Therefore

$$\sum_{n \geq 2} \frac{1}{n^2(\log n)^{(t-1)p/2}} V_n^{p/2} \ll \sum_{k > 0} \frac{2^{\eta p(k+1)/2}}{2^k k^{(t-1)p/2}} \sum_{\ell=0}^k \left(\frac{V_{2^\ell}}{2^{\eta \ell}} \right)^{p/2}.$$

Changing the order of summation and taking $\eta \in]0, 2/p[$, it follows that (3.8) is implied by

$$\sum_{k \geq 1} \frac{1}{2^k k^{(t-1)p/2}} \|\mathbb{E}_0(M_{2^k}^2) - \mathbb{E}(M_{2^k}^2)\|_{p/2}^{p/2} < \infty \quad (3.15)$$

(actually due to the subadditivity of the sequence (V_i) both conditions are equivalent, see the proof of item 1 of Lemma 37 in [22] to prove that (3.8) entails (3.15)). Now, since (M_n) is a martingale,

$$\begin{aligned} \mathbb{E}_0(M_{2^k}^2) - \mathbb{E}(M_{2^k}^2) &= \sum_{j=1}^k (\mathbb{E}_0((M_{2^j} - M_{2^{j-1}})^2) - \mathbb{E}((M_{2^j} - M_{2^{j-1}})^2)) \\ &\quad + \mathbb{E}_0(d_1^2) - \mathbb{E}(d_1^2), \end{aligned}$$

which implies by stationarity that

$$\begin{aligned} \|\mathbb{E}_0(M_{2^k}^2) - \mathbb{E}(M_{2^k}^2)\|_{p/2} &\leq \sum_{j=0}^{k-1} \|\mathbb{E}_{-2^j}(M_{2^j}^2) - \mathbb{E}(M_{2^j}^2)\|_{p/2} \\ &\quad + \|\mathbb{E}_0(d_1^2) - \mathbb{E}(d_1^2)\|_{p/2}. \end{aligned}$$

Therefore by using Hölder's inequality as done in (3.14) with $\eta \in]0, 2/p[$, we infer that (3.15) is implied by

$$\sum_{k \geq 1} \frac{1}{2^k k^{(t-1)p/2}} \|\mathbb{E}_{-2^k}(M_{2^k}^2) - \mathbb{E}(M_{2^k}^2)\|_{p/2}^{p/2} < \infty. \quad (3.16)$$

Notice now that the sequence $(W_n)_{n>0}$ defined by

$$W_n := \|\mathbb{E}_{-n}(M_n^2) - \mathbb{E}(M_n^2)\|_{p/2}$$

is subadditive. Indeed, for any non negative integers i and j , using that M_n is a martingale together with the stationarity, we derive that

$$\begin{aligned} W_{i+j} &= \|\mathbb{E}_{-(i+j)}(M_i^2) - \mathbb{E}(M_i^2) + \mathbb{E}_{-(i+j)}((M_{i+j} - M_i)^2) \\ &\quad - \mathbb{E}((M_{i+j} - M_i)^2)\|_{p/2} \\ &\leq \|\mathbb{E}_{-i}(M_i^2) - \mathbb{E}(M_i^2)\|_{p/2} + \|\mathbb{E}_{-j}(M_j^2) - \mathbb{E}(M_j^2)\|_{p/2} \\ &\leq W_i + W_j. \end{aligned}$$

Therefore $W_{i+j}^{p/2} \leq 2^{p/2} W_i^{p/2} + 2^{p/2} W_j^{p/2}$. This implies that, for any integer ℓ and any integer $0 \leq j \leq \ell$, $W_\ell^{p/2} \leq 2^{p/2}(W_j^{p/2} + W_{\ell-j}^{p/2})$ in such a way that

$$(\ell + 1)W_\ell^{p/2} \leq 2^{1+p/2} \sum_{j=1}^{\ell} W_j^{p/2}. \quad (3.17)$$

Therefore using (3.17) with $\ell = 2^k$, we infer that condition (3.16) is implied by

$$\sum_{n \geq 2} \frac{1}{n^2 (\log n)^{(t-1)p/2}} \|\mathbb{E}_{-n}(M_n^2) - \mathbb{E}(M_n^2)\|_{p/2}^{p/2} < \infty. \quad (3.18)$$

It remains to prove that (3.12) implies (3.18). With this aim, we have, for any positive integer m ,

$$M_n = \sum_{k=1}^m (M_{k[nm^{-1}]} - M_{(k-1)[nm^{-1}]}) + M_n - M_{m[nm^{-1}]}.$$

Using that M_n is a martingale together with the stationarity, we then infer that

$$\begin{aligned} \|\mathbb{E}_{-n}(M_n^2) - \mathbb{E}(M_n^2)\|_{p/2}^{p/2} &\leq 2^{p/2} m^{p/2} \|\mathbb{E}_{-n}(M_{[nm^{-1}]}^2) - \mathbb{E}(M_{[nm^{-1}]}^2)\|_{p/2}^{p/2} \\ &\quad + 2^{p/2} \|\mathbb{E}_{-n}(M_{n-m[nm^{-1}]}^2) - \mathbb{E}(M_{n-m[nm^{-1}]}^2)\|_{p/2}^{p/2}, \end{aligned}$$

which, together with the fact that $n - m[nm^{-1}] < m$, implies that

$$\begin{aligned} & \left\| \mathbb{E}_{-n}(M_n^2) - \mathbb{E}(M_n^2) \right\|_{p/2}^{p/2} \\ & \leq 2^{p/2} m^{p/2} \left(2^{p/2} \|d_0\|_p^p + \left\| \mathbb{E}_{-n}(M_{[nm^{-1}]}^2) - \mathbb{E}(M_{[nm^{-1}]}^2) \right\|_{p/2}^{p/2} \right) \\ & \leq 2^{p/2} m^{p/2} \left(2^{p/2} \|d_0\|_p^p + \left\| \mathbb{E}_{-m[nm^{-1}]}(M_{[nm^{-1}]}^2) - \mathbb{E}(M_{[nm^{-1}]}^2) \right\|_{p/2}^{p/2} \right), \end{aligned} \quad (3.19)$$

where for the last line we have used the fact that $n \geq m[nm^{-1}]$. We notice now that due to the martingale property of (M_n) and to stationarity, the sequence $(U_i)_{i \geq 0}$ defined for any non negative integer i by

$$U_i := \left\| \mathbb{E}_{-mi}(M_i^2) - \mathbb{E}(M_i^2) \right\|_{p/2}^{p/2}$$

satisfies, for any positive integers i and j ,

$$\begin{aligned} U_{i+j} & \leq \left(\left\| \mathbb{E}_{-m(i+j)}(M_{i+j}^2) - \mathbb{E}(M_{i+j}^2) \right\|_{p/2} \right. \\ & \quad \left. + \left\| \mathbb{E}_{-m(i+j)}((M_{i+j} - M_i)^2) - \mathbb{E}((M_{i+j} - M_i)^2) \right\|_{p/2} \right)^{p/2} \\ & \leq 2^{p/2} U_i + 2^{p/2} U_j. \end{aligned}$$

Hence by (3.17) applied with $W_i^{p/2} = U_i$,

$$U_{[nm^{-1}]} \leq 2^{1+p/2} ([nm^{-1}] + 1)^{-1} \sum_{k=1}^{[nm^{-1}]} U_k \leq 2^{1+p/2} \sum_{k=1}^{[nm^{-1}]} \frac{U_k}{k}. \quad (3.20)$$

Therefore starting from (3.19), considering (3.20) and changing the order of summation, we infer that (3.18) (and so (3.8)) holds provided that (3.12) does. To end the proof, it remains to show that under the conditions of Theorems 3.1 and 3.2, the conditions (3.11) and (3.12) are satisfied. This is achieved by using the two following lemmas. \square

Lemma 3.1. *Let $p \in [2, 4]$. Assume that (3.1) holds. Then*

$$\sum_{n \geq 1} \frac{\max_{1 \leq \ell \leq n} \|R_\ell\|_p^p}{n^2 (\log n)^{(t-1)p/2}} < \infty,$$

and (3.11) holds.

Proof. Since (3.1) implies (3.6), Item 2 of Proposition 5.1 given in the appendix implies that, for any positive integers ℓ and N ,

$$\|R_\ell\|_p \ll \max_{k=\ell, N} \|\mathbb{E}_0(S_k)\|_p + \max_{k=\ell, N} \|S_k - \mathbb{E}_k(S_k)\|_p + \ell^{1/2} \sum_{|j| \geq N} \|P_0(X_j)\|_p.$$

Next, applying Lemma 5.1 given in the appendix with $q = 1$, and using the fact that by stationarity, for any positive integer k ,

$$\|\mathbb{E}_0(S_k)\|_p \leq \sum_{\ell=1}^k \|\mathbb{E}_0(X_\ell)\|_p \quad \text{and} \quad \|S_k - \mathbb{E}_k(S_k)\|_p \leq \sum_{\ell=0}^{k-1} \|X_{-\ell} - \mathbb{E}_0(X_{-\ell})\|_p, \quad (3.21)$$

we derive that for, any positive integers $N \geq n$,

$$\begin{aligned} \max_{1 \leq \ell \leq n} \|R_\ell\|_p &\ll \sum_{k=1}^N \|\mathbb{E}_0(X_k)\|_p + \sum_{k=0}^{N-1} \|X_{-k} - \mathbb{E}_0(X_{-k})\|_p \\ &+ n^{1/2} \sum_{k \geq [N/2]} \frac{\|\mathbb{E}_0(X_k)\|_p}{k^{1/p}} + n^{1/2} \sum_{k \geq [N/2]} \frac{\|X_{-k} - \mathbb{E}_0(X_{-k})\|_p}{k^{1/p}}. \end{aligned} \quad (3.22)$$

The lemma follows from (3.22) with $N = \lceil n^{p/2} \rceil$ by using Hölder's inequality (see the computations in the proof of Proposition 2.2 in [2]). \square

Lemma 3.2. *Let $p \in [2, 4]$ and assume that (3.1) and (3.3) are satisfied. Assume in addition that (3.2) holds when $2 < p < 4$ and (3.5) does when $p = 4$. Then (3.12) is satisfied.*

Proof. Let m be a positive integer such that (3.3) is satisfied. We first write that

$$\begin{aligned} \|\mathbb{E}_{-nm}(M_n^2) - \mathbb{E}(M_n^2)\|_{p/2} &\leq \|\mathbb{E}_{-nm}(S_n^2) - \mathbb{E}(S_n^2)\|_{p/2} \\ &+ 2\|\mathbb{E}_{-nm}(S_n R_n) - \mathbb{E}(S_n R_n)\|_{p/2} + 2\|R_n\|_p^2. \end{aligned}$$

By using Lemma 3.1, and since (3.3) holds, Lemma 3.2 will follow if we can prove that

$$\sum_{n \geq 1} \frac{1}{n^2 (\log n)^{(t-1)p/2}} \|\mathbb{E}_{-nm}(S_n R_n)\|_{p/2}^{p/2} < \infty. \quad (3.23)$$

With this aim we shall prove the following inequality. For any non negative integer r and any positive integer u_n such that $u_n \leq n$, we have that

$$\begin{aligned} \|\mathbb{E}_{-r}(S_n R_n)\|_{p/2} &\ll \sqrt{u_n} (\|\mathbb{E}_0(S_n)\|_2 + \|S_n - \mathbb{E}_n(S_n)\|_2) \\ &+ \max_{k=\{n, n-u_n\}} \|R_k\|_p^2 + \sqrt{n} (\|\mathbb{E}_{-u_n}(S_n)\|_2 + \|S_n - \mathbb{E}_{n+u_n}(S_n)\|_2) \\ &+ \max_{k=\{n, u_n\}} \|\mathbb{E}_{-r}(S_k^2) - \mathbb{E}(S_k^2)\|_{p/2} + \sqrt{n} \left(\sum_{k=1}^n \left\| \sum_{|j| \geq k+n} P_0(X_j) \right\|_2^2 \right)^{1/2}. \end{aligned} \quad (3.24)$$

Let us show how, thanks to (3.24), the convergence (3.23) can be proven. Let us first consider the case where $2 < p < 4$. Notice that the following elementary claim is valid:

Claim 3.1. *If \mathcal{F} and \mathcal{G} are two σ -algebras such that $\mathcal{G} \subset \mathcal{F}$, then for any random variable X in \mathbb{L}^q for $q \geq 1$, $\|X - \mathbb{E}(X|\mathcal{F})\|_q \leq 2\|X - \mathbb{E}(X|\mathcal{G})\|_q$.*

Starting from (3.24) with $r = nm$ and $u_n = n$, and using Claim 3.1, we derive that

$$\begin{aligned} \|\mathbb{E}_{-nm}(S_n R_n)\|_{p/2} &\ll \|\mathbb{E}_{-nm}(S_n^2) - \mathbb{E}(S_n^2)\|_{p/2} \\ &+ \sqrt{n}(\|\mathbb{E}_0(S_n)\|_2 + \|S_n - \mathbb{E}_n(S_n)\|_2) + \|R_n\|_p^2 + n \sum_{|j| \geq n} \|P_0(X_j)\|_2. \end{aligned}$$

This last inequality combined with condition (3.3) and Lemma 3.1 shows that (3.23) will be satisfied if we can prove that

$$\sum_{n \geq 1} \frac{n^{p/4}}{n^2(\log n)^{(t-1)p/2}} (\|\mathbb{E}_0(S_n)\|_2 + \|S_n - \mathbb{E}_n(S_n)\|_2)^{p/2} < \infty, \quad (3.25)$$

and

$$\sum_{n \geq 1} \frac{n^{p/2}}{n^2(\log n)^{(t-1)p/2}} \left(\sum_{|j| \geq n} \|P_0(X_j)\|_2 \right)^{p/2} < \infty. \quad (3.26)$$

To prove (3.25), we use the inequalities (3.21) with $p = 2$. Hence setting

$$a_\ell = \|\mathbb{E}_0(X_\ell)\|_2 + \|X_{-\ell+1} - \mathbb{E}_0(X_{-\ell+1})\|_2, \quad (3.27)$$

and using Hölder's inequality, we derive that for any $\alpha < 1$,

$$\begin{aligned} \sum_{n \geq 1} \frac{n^{p/4}}{n^2(\log n)^{(t-1)p/2}} (\|\mathbb{E}_0(S_n)\|_2 + \|S_n - \mathbb{E}_n(S_n)\|_2)^{p/2} \\ \ll \sum_{n \geq 1} \frac{n^{p/4}}{n^2(\log n)^{(t-1)p/2}} \left(\sum_{\ell=1}^n a_\ell \right)^{p/2} \\ \ll \sum_{n \geq 1} \frac{n^{p/4} n^{(1-\alpha)(p/2-1)}}{n^2(\log n)^{(t-1)p/2}} \sum_{\ell=1}^n \ell^{\alpha(p/2-1)} a_\ell^{p/2}. \end{aligned}$$

Taking $\alpha \in](3p-8)/(2p-4), 1[$ (this is possible since $p < 4$) and changing the order of summation, we infer that (3.25) holds provided that (3.2) does. It remains to show that (3.26) is satisfied. Using Lemma 5.1 and the notation (3.27), we first observe that

$$\begin{aligned} \sum_{n \geq 1} \frac{n^{p/2}}{n^2(\log n)^{(t-1)p/2}} \left(\sum_{|j| \geq n} \|P_0(X_j)\|_2 \right)^{p/2} \\ \ll \sum_{n \geq 1} \frac{n^{p/2}}{n^2(\log n)^{(t-1)p/2}} \left(\sum_{\ell \geq [n/2]} \ell^{-1/2} a_\ell \right)^{p/2}. \end{aligned}$$

Therefore by Hölder's inequality, it follows that for any $\alpha < 1$,

$$\begin{aligned} \sum_{n \geq 1} \frac{n^{p/2}}{n^2(\log n)^{(t-1)p/2}} \left(\sum_{|j| \geq n} \|P_0(X_j)\|_2 \right)^{p/2} \\ \ll \sum_{n \geq 1} \frac{n^{p/2} n^{(1-\alpha)(p/2-1)}}{n^2(\log n)^{(t-1)p/2}} \sum_{\ell \geq [n/2]} \ell^{\alpha(p/2-1)} \ell^{-p/4} a_\ell^{p/2}. \end{aligned}$$

Therefore taking $\alpha \in]1, 2[$ and changing the order of summation, we infer that (3.25) holds provided that (3.2) does. This ends the proof of (3.23) when $p \in]2, 4[$.

Now, we prove (3.23) when $p = 4$. With this aim we start from (3.24) with $r = nm$ and $u_n = \lfloor \sqrt{n} \rfloor$. This inequality combined with condition (3.3), Lemma 3.1 and the arguments developed to prove (3.25) and (3.26) shows that (3.23) will be satisfied for $p = 4$ if we can prove that

$$\sum_{n \geq 1} \frac{1}{n(\log n)^{2(t-1)}} (\|\mathbb{E}_{-\lfloor \sqrt{n} \rfloor}(S_n)\|_2 + \|S_n - \mathbb{E}_{n+\lfloor \sqrt{n} \rfloor}(S_n)\|_2)^2 < \infty, \quad (3.28)$$

and

$$\sum_{n \geq 2} \frac{1}{n^2(\log n)^{2(t-1)}} \|\mathbb{E}_{-nm}(S_{\lfloor \sqrt{n} \rfloor}^2) - \mathbb{E}(S_{\lfloor \sqrt{n} \rfloor}^2)\|_2^2 < \infty. \quad (3.29)$$

We start by proving (3.28). With this aim, using the notation (3.27), we first write that

$$\|\mathbb{E}_{-\lfloor \sqrt{n} \rfloor}(S_n)\|_2 + \|S_n - \mathbb{E}_{n+\lfloor \sqrt{n} \rfloor}(S_n)\|_2 \leq \sum_{k=\lfloor \sqrt{n} \rfloor+1}^{n+\lfloor \sqrt{n} \rfloor} a_k.$$

Therefore by Cauchy-Schwarz's inequality

$$\begin{aligned} \sum_{n \geq 1} \frac{1}{n(\log n)^{2(t-1)}} (\|\mathbb{E}_{-\lfloor \sqrt{n} \rfloor}(S_n)\|_2 + \|S_n - \mathbb{E}_{n+\lfloor \sqrt{n} \rfloor}(S_n)\|_2)^2 \\ \ll \sum_{n \geq 1} \frac{\log n}{n(\log n)^{2(t-1)}} \sum_{k=\lfloor \sqrt{n} \rfloor+1}^{n+\lfloor \sqrt{n} \rfloor} k a_k^2 \\ \ll \sum_{n \geq 1} \frac{1}{n} \sum_{k=\lfloor \sqrt{n} \rfloor+1}^{n+\lfloor \sqrt{n} \rfloor} \frac{k \log k}{(\log k)^{2(t-1)}} a_k^2. \end{aligned}$$

Changing the order of summation, this proves that (3.28) holds provided that (3.5) does. It remains to prove (3.29). With this aim, we set for any positive real x ,

$$h(\lfloor x \rfloor) = \|\mathbb{E}_{-m_{\lfloor x \rfloor}}(S_{\lfloor x \rfloor}^2) - \mathbb{E}(S_{\lfloor x \rfloor}^2)\|_2^2,$$

and we notice that, for any integer $n \geq 0$,

$$\|\mathbb{E}_{-nm}(S_{\lfloor \sqrt{n} \rfloor}^2) - \mathbb{E}(S_{\lfloor \sqrt{n} \rfloor}^2)\|_2^2 \leq h(\lfloor \sqrt{n} \rfloor).$$

In addition, if $x \in [n, n+1[$, then $[\sqrt{n}] = [\sqrt{x}]$ or $[\sqrt{n}] = [\sqrt{x}] - 1$. Therefore

$$\begin{aligned} \sum_{n \geq 3} \frac{1}{n^2(\log n)^{(t-1)p/2}} h([\sqrt{n}]) &\ll \sum_{n \geq 3} h([\sqrt{n}]) \int_{[n, n+1[} \frac{1}{x^2(\log x)^{(t-1)p/2}} dx \\ &\ll \int_3^\infty \frac{1}{x^2(\log x)^{(t-1)p/2}} h([\sqrt{x}]) dx + \int_3^\infty \frac{1}{x^2(\log x)^{(t-1)p/2}} h([\sqrt{x}] - 1) dx \\ &\ll \int_2^\infty \frac{1}{y^3(\log y)^{(t-1)p/2}} h([y]) dy \ll \sum_{n \geq 2} \frac{1}{n^3(\log n)^{(t-1)p/2}} h(n) dy. \end{aligned}$$

For the last inequality, we have used that if $y \in [n, n+1[$, then $[y] = n$. Therefore condition (3.3) implies (3.29). This ends the proof of (3.23) when $p = 4$.

It remains to prove (3.24). With this aim, we start with the decomposition of R_n given in Proposition 5.1 of the appendix with $N = n$. Therefore setting

$$A_n := \sum_{k=1}^n \sum_{j \geq 2n+1} P_k(X_j) + \sum_{k=1}^n \sum_{j \leq n} P_k(X_{-j}),$$

we write that

$$\begin{aligned} R_n = \mathbb{E}_0(S_n) - \mathbb{E}_0(S_n) \circ T^n + \mathbb{E}_{-n}(S_n) \circ T^n + S_n - \mathbb{E}_n(S_n) \\ - (\mathbb{E}_{2n}(S_n) - \mathbb{E}_n(S_n)) \circ T^{-n} - A_n. \end{aligned} \quad (3.30)$$

Starting from (3.30) and noticing that

$$\begin{aligned} \|\mathbb{E}_{-r}(S_n(\mathbb{E}_{-n}(S_n) \circ T^n))\|_{p/2} &\leq \|\mathbb{E}_0(S_n(\mathbb{E}_{-n}(S_n) \circ T^n))\|_{p/2} \\ &\leq \|\mathbb{E}_0(S_n)\|_p \|\mathbb{E}_0(S_{2n} - S_n)\|_p, \end{aligned}$$

and that $\mathbb{E}_{-r}(S_n(S_n - \mathbb{E}_n(S_n))) = \mathbb{E}_{-r}((S_n - \mathbb{E}_n(S_n))^2)$, we first get

$$\begin{aligned} \|\mathbb{E}_{-r}(S_n R_n)\|_{p/2} &\leq 2\|\mathbb{E}_0(S_n)\|_p^2 + \|S_n - \mathbb{E}_n(S_n)\|_p^2 \\ &\quad + \|\mathbb{E}_{-r}(S_n \mathbb{E}_n(S_{2n} - S_n))\|_{p/2} \\ &\quad + \|\mathbb{E}_{-r}(S_n \mathbb{E}_n(S_n \circ T^{-n} - \mathbb{E}_0(S_n \circ T^{-n})))\|_{p/2} \\ &\quad + \|\mathbb{E}_{-r}(S_n A_n)\|_{p/2}. \end{aligned} \quad (3.31)$$

Next, we use the following fact: if X and Y are two variables in \mathbb{L}^p with $p \in [2, 4]$, then for any integer u ,

$$\|\mathbb{E}_u(XY)\|_{p/2} \leq \|\mathbb{E}_u(X^2) - \mathbb{E}(X^2)\|_{p/2} + \|Y\|_p^2 + \sqrt{\mathbb{E}(X^2)} \|Y\|_2. \quad (3.32)$$

Indeed, it suffices to write that

$$\begin{aligned} \|\mathbb{E}_u(XY)\|_{p/2} &\leq \|\mathbb{E}_u^{1/2}(X^2) \mathbb{E}_u^{1/2}(Y^2)\|_{p/2} \\ &\leq \| |\mathbb{E}_u(X^2) - \mathbb{E}(X^2)|^{1/2} \mathbb{E}_u^{1/2}(Y^2) \|_{p/2} + (\mathbb{E}(X^2))^{1/2} \|\mathbb{E}_u^{1/2}(Y^2)\|_{p/2} \\ &\leq \|\mathbb{E}_u(X^2) - \mathbb{E}(X^2)\|_{p/2} + \|Y\|_p^2 + (\mathbb{E}(X^2))^{1/2} \|\mathbb{E}_u^{1/2}(Y^2)\|_{p/2}, \end{aligned}$$

and to notice that, since $p \in [2, 4]$, $\|\mathbb{E}_u^{1/2}(Y^2)\|_{p/2} \leq \|\mathbb{E}_u^{1/2}(Y^2)\|_2 = \|Y\|_2$. Therefore, starting from (3.31) and using (3.32) together with $\mathbb{E}(S_n^2) \ll n$, we infer that

$$\begin{aligned} \|\mathbb{E}_{-r}(S_n R_n)\|_{p/2} &\ll \|\mathbb{E}_0(S_n)\|_p^2 + \|S_n - \mathbb{E}_n(S_n)\|_p^2 \\ &\quad + \|\mathbb{E}_{-r}(S_n \mathbb{E}_n(S_{2n} - S_n))\|_{p/2} \\ &\quad + \|\mathbb{E}_{-r}(S_n \mathbb{E}_n(S_n \circ T^{-n} - \mathbb{E}_0(S_n \circ T^{-n})))\|_{p/2} \\ &\quad + \|\mathbb{E}_{-r}(S_n^2) - \mathbb{E}(S_n^2)\|_{p/2} + \|A_n\|_p^2 + n^{1/2} \|A_n\|_2, \end{aligned}$$

and since $\|\mathbb{E}_0(S_n)\|_p \leq \|R_n\|_p$, $\|S_n - \mathbb{E}_n(S_n)\|_p \leq 2\|R_n\|_p$ and $\|A_n\|_p \leq 8\|R_n\|_p$, we have overall that

$$\begin{aligned} \|\mathbb{E}_{-r}(S_n R_n)\|_{p/2} &\ll \|R_n\|_p^2 + \|\mathbb{E}_{-r}(S_n \mathbb{E}_n(S_{2n} - S_n))\|_{p/2} \\ &\quad + \|\mathbb{E}_{-r}(S_n \mathbb{E}_n(S_n \circ T^{-n} - \mathbb{E}_0(S_n \circ T^{-n})))\|_{p/2} \\ &\quad + \|\mathbb{E}_{-r}(S_n^2) - \mathbb{E}(S_n^2)\|_{p/2} + n^{1/2} \|A_n\|_2. \end{aligned} \quad (3.33)$$

By orthogonality and by stationarity,

$$\begin{aligned} \|A_n\|_2 &\leq \left(\sum_{k=1}^n \left\| \sum_{j \geq 2n+1} P_k(X_j) \right\|_2^2 \right)^{1/2} + \left(\sum_{k=1}^n \left\| \sum_{j \geq n} P_k(X_{-j}) \right\|_2^2 \right)^{1/2} \\ &\leq \left(\sum_{k=1}^n \left\| \sum_{\ell \geq k+n} P_0(X_\ell) \right\|_2^2 \right)^{1/2} + \left(\sum_{k=1}^n \left\| \sum_{\ell \geq k+n} P_0(X_{-\ell}) \right\|_2^2 \right)^{1/2}. \end{aligned} \quad (3.34)$$

Now for any integer u_n such that $u_n \leq n$,

$$\begin{aligned} \|\mathbb{E}_{-r}(S_n \mathbb{E}_n(S_{2n} - S_n))\|_{p/2} &\leq \|\mathbb{E}_{-r}((S_n - S_{n-u_n}) \mathbb{E}_n(S_{2n} - S_n))\|_{p/2} \\ &\quad + \|\mathbb{E}_{-r}(S_{n-u_n} \mathbb{E}_n(S_{2n} - S_n))\|_{p/2} \\ &\ll \|\mathbb{E}_{-r}(S_{u_n}^2) - \mathbb{E}(S_{u_n}^2)\|_{p/2} \\ &\quad + \|\mathbb{E}_0(S_n)\|_p^2 + \sqrt{u_n} \|\mathbb{E}_0(S_n)\|_2 \\ &\quad + \|\mathbb{E}_{-r}(S_{n-u_n} \mathbb{E}_n(S_{2n} - S_n))\|_{p/2}, \end{aligned} \quad (3.35)$$

where for the last inequality we have used (3.32) together with $\mathbb{E}(S_{u_n}^2) \ll u_n$. Next, we write that

$$\begin{aligned} &\|\mathbb{E}_{-r}(S_{n-u_n} \mathbb{E}_n(S_{2n} - S_n))\|_{p/2} \\ &\leq \|\mathbb{E}_{-r}((S_{n-u_n} - \mathbb{E}_{n-u_n}(S_{n-u_n})) \mathbb{E}_n(S_{2n} - S_n))\|_{p/2} \\ &\quad + \|\mathbb{E}_{-r}(\mathbb{E}_{n-u_n}(S_{n-u_n}) \mathbb{E}_n(S_{2n} - S_n))\|_{p/2} \\ &\leq \|S_{n-u_n} - \mathbb{E}_{n-u_n}(S_{n-u_n})\|_p \|\mathbb{E}_0(S_n)\|_p \\ &\quad + \|\mathbb{E}_{-r}(\mathbb{E}_{n-u_n}(S_{n-u_n}) \mathbb{E}_{n-u_n}(S_{2n} - S_n))\|_{p/2} \\ &\leq \|S_{n-u_n} - \mathbb{E}_{n-u_n}(S_{n-u_n})\|_p^2 + \|\mathbb{E}_0(S_n)\|_p^2 \\ &\quad + \|\mathbb{E}_{-r}(S_n \mathbb{E}_{n-u_n}(S_{2n} - S_n))\|_{p/2} \\ &\quad + \|\mathbb{E}_{-r}((S_n - S_{n-u_n}) \mathbb{E}_{n-u_n}(S_{2n} - S_n))\|_{p/2}. \end{aligned}$$

Therefore using (3.32), we infer that

$$\begin{aligned} \|\mathbb{E}_{-r}(S_{n-u_n}\mathbb{E}_n(S_{2n}-S_n))\|_{p/2} &\ll \max_{k=\{n, n-u_n\}} \|R_k\|_p^2 + \sqrt{n}\|\mathbb{E}_{-u_n}(S_n)\|_2 \\ &+ \max_{k=\{n, u_n\}} \|\mathbb{E}_{-r}(S_k^2) - \mathbb{E}(S_k^2)\|_{p/2}. \end{aligned} \quad (3.36)$$

We deal now with the third term in the right-hand side of (3.33). With this aim, we first write that

$$\begin{aligned} &\|\mathbb{E}_{-r}(S_n\mathbb{E}_n(S_n \circ T^{-n} - \mathbb{E}_0(S_n \circ T^{-n})))\|_{p/2} \\ &\leq \|\mathbb{E}_{-r}(S_n\mathbb{E}_n(S_n \circ T^{-n} - \mathbb{E}_{u_n}(S_n \circ T^{-n})))\|_{p/2} \\ &\quad + \|\mathbb{E}_{-r}(S_n\mathbb{E}_{u_n}(S_n \circ T^{-n} - \mathbb{E}_0(S_n \circ T^{-n})))\|_{p/2}. \end{aligned} \quad (3.37)$$

By using (3.32) together with $\mathbb{E}(S_{u_n}^2) \ll n$, stationarity and the fact that $\|S_n - \mathbb{E}_{n+u_n}(S_n)\|_2 \leq 2\|R_n\|_p$, we infer that

$$\begin{aligned} \|\mathbb{E}_{-r}(S_n\mathbb{E}_n(S_n \circ T^{-n} - \mathbb{E}_{u_n}(S_n \circ T^{-n})))\|_{p/2} &\ll \|\mathbb{E}_{-r}(S_n^2) - \mathbb{E}(S_n^2)\|_{p/2} \\ &+ \|R_n\|_p^2 + \sqrt{n}\|S_n - \mathbb{E}_{n+u_n}(S_n)\|_2. \end{aligned} \quad (3.38)$$

On the other hand,

$$\begin{aligned} &\|\mathbb{E}_{-r}(S_n\mathbb{E}_{u_n}(S_n \circ T^{-n} - \mathbb{E}_0(S_n \circ T^{-n})))\|_{p/2} \\ &\leq \|\mathbb{E}_{-r}(S_{u_n}\mathbb{E}_{u_n}(S_n \circ T^{-n} - \mathbb{E}_0(S_n \circ T^{-n})))\|_{p/2} \\ &\quad + \|\mathbb{E}_{-r}(\mathbb{E}_{u_n}(S_n - S_{u_n})\mathbb{E}_{u_n}(S_n \circ T^{-n} - \mathbb{E}_0(S_n \circ T^{-n})))\|_{p/2}. \end{aligned}$$

We apply (3.32) to the first term of the right hand side together with the fact that $\mathbb{E}(S_{u_n}^2) \ll n$. Hence by stationarity and since $\|S_n - \mathbb{E}_n(S_n)\|_p \leq 2\|R_n\|_p$, we derive that

$$\begin{aligned} \|\mathbb{E}_{-r}(S_{u_n}\mathbb{E}_{u_n}(S_n \circ T^{-n} - \mathbb{E}_0(S_n \circ T^{-n})))\|_{p/2} &\ll \|\mathbb{E}_{-r}(S_{u_n}^2) - \mathbb{E}(S_{u_n}^2)\|_{p/2} \\ &+ \|R_n\|_p^2 + \sqrt{u_n}\|S_n - \mathbb{E}_n(S_n)\|_2. \end{aligned}$$

On the other hand, by stationarity,

$$\begin{aligned} &\|\mathbb{E}_{-r}(\mathbb{E}_{u_n}(S_n - S_{u_n})\mathbb{E}_{u_n}(S_n \circ T^{-n} - \mathbb{E}_0(S_n \circ T^{-n})))\|_{p/2} \\ &\leq \|\mathbb{E}_{u_n}(S_n - S_{u_n})\|_p \|\mathbb{E}_{u_n}(S_n \circ T^{-n} - \mathbb{E}_0(S_n \circ T^{-n}))\|_p \\ &\leq \|\mathbb{E}_0(S_{n-u_n})\|_p \|S_n - \mathbb{E}_n(S_n)\|_p \\ &\leq \|\mathbb{E}_0(S_{n-u_n})\|_p^2 + \|R_n\|_p^2. \end{aligned}$$

Therefore we get overall that

$$\begin{aligned} &\|\mathbb{E}_{-r}(S_n\mathbb{E}_{u_n}(S_n \circ T^{-n} - \mathbb{E}_0(S_n \circ T^{-n})))\|_{p/2} \ll \|R_n\|_p^2 + \|\mathbb{E}_0(S_{n-u_n})\|_p^2 \\ &\quad + \|\mathbb{E}_{-r}(S_{u_n}^2) - \mathbb{E}(S_{u_n}^2)\|_{p/2} + \sqrt{u_n}\|S_n - \mathbb{E}_n(S_n)\|_2. \end{aligned} \quad (3.39)$$

Starting from (3.37) and taking into account (3.38) and (3.39), we get that

$$\begin{aligned} & \|\mathbb{E}_{-r}(S_n \mathbb{E}_n(S_n \circ T^{-n} - \mathbb{E}_0(S_n \circ T^{-n})))\|_{p/2} \\ & \ll \sqrt{u_n} \|S_n - \mathbb{E}_n(S_n)\|_2 + \sqrt{n} \|S_n - \mathbb{E}_{n+u_n}(S_n)\|_2 \\ & + \max_{k=\{n, u_n\}} \|\mathbb{E}_{-r}(S_k^2) - \mathbb{E}(S_k^2)\|_{p/2} + \max_{k=\{n, n-u_n\}} \|R_k\|_p^2. \end{aligned} \quad (3.40)$$

Finally, starting from (3.33) and considering (3.34), (3.35), (3.36) and (3.40), we conclude that (3.24) holds. \square

4. Proof of Theorem 2.1

4.1. Preparatory material

Let us denote by E_u , E_e and E_s the S -stable vector spaces associated to the eigenvalues of S of modulus respectively larger than one, equal to one and smaller than one. Let d_u , d_e and d_s be their respective dimensions. Let v_1, \dots, v_d be a basis of \mathbb{R}^d in which S is represented by a real Jordan matrix. Suppose that v_1, \dots, v_{d_u} are in E_u , $v_{d_u+1}, \dots, v_{d_u+d_e}$ are in E_e and $v_{d_u+d_e+1}, \dots, v_d$ are in E_s . We suppose moreover that $\det(v_1|v_2|\dots|v_d) = 1$. Let us write $\|\cdot\|$ the norm on \mathbb{R}^d given by

$$\left\| \sum_{i=1}^d x_i v_i \right\| = \max_{i=1, \dots, d} |x_i|$$

and $d_0(\cdot, \cdot)$ the metric induced by $\|\cdot\|$ on \mathbb{R}^d . Let also d_1 be the metric induced by d_0 on \mathbb{T}^d . We define now $B_u(\delta) := \{y \in E_u : \|y\| \leq \delta\}$, $B_e(\delta) := \{y \in E_e : \|y\| \leq \delta\}$ and $B_s(\delta) = \{y \in E_s : \|y\| \leq \delta\}$. Let $|\cdot|$ be the usual euclidean norm on \mathbb{R}^d .

Let r_u be the spectral radius of $S_{E_u}^{-1}$. For every $\rho_u \in (r_u, 1)$, there exists $K > 0$ such that, for every integer $n \geq 0$, we have

$$\forall h_u \in E_u, \quad \|S^n h_u\| \geq K \rho_u^{-n} \|h_u\| \quad (4.41)$$

and

$$\forall (h_e, h_s) \in E_e \times E_s, \quad \|S^n (h_e + h_s)\| \leq K(1+n)^{d_e} \|h_e + h_s\|. \quad (4.42)$$

Let $\rho_u \in (r_u, 1)$ and K satisfying (4.41) and (4.42). Let us denote by m_u , m_e , m_s the Lebesgue measure on E_u (in the basis v_1, \dots, v_{d_u}), E_e (in the basis $v_{d_u+1}, \dots, v_{d_u+d_e}$) and E_s (in the basis $v_{d_u+d_e+1}, \dots, v_d$) respectively. Observe that $d\lambda(h_u + h_e + h_s) = dm_u(h_u) dm_e(h_e) dm_s(h_s)$.

The properties satisfied by the filtration considered in [17, 14] and enabling the use of a martingale approximation method à la Gordin will be crucial here. Given a finite partition \mathcal{P} of \mathbb{T}^d , we define the measurable partition \mathcal{P}_0^∞ by :

$$\forall \bar{x} \in \mathbb{T}^d, \quad \mathcal{P}_0^\infty(\bar{x}) := \bigcap_{k \geq 0} T^k \mathcal{P}(T^{-k}(\bar{x}))$$

and, for every integer n , the σ -algebra \mathcal{F}_n generated by

$$\forall \bar{x} \in \mathbb{T}^d, \quad \mathcal{P}_{-n}^\infty(\bar{x}) := \bigcap_{k \geq -n} T^k \mathcal{P}(T^{-k}(\bar{x})) = T^{-n}(\mathcal{P}_0^\infty(T^n(\bar{x})).$$

These definitions coincide with the ones of [14] applied to the ergodic torus automorphism T^{-1} . We obviously have $\mathcal{F}_n \subseteq \mathcal{F}_{n+1} = T^{-1}\mathcal{F}_n$. Let $r_0 > 0$ be such that $(h_u, h_e, h_s) \mapsto \overline{h_u + h_e + h_s}$ defines a diffeomorphism from $B_u(r_0) \times B_e(r_0) \times B_s(r_0)$ on its image in \mathbb{T}^d . Observe that, for every $\bar{x} \in \mathbb{T}^d$, on the set $\bar{x} + B_u(r_0) + B_e(r_0) + B_s(r_0)$, we have $d\bar{\lambda}(\bar{x} + \overline{h_u} + \overline{h_e} + \overline{h_s}) = dm_u(h_u)dm_e(h_e)dm_s(h_s)$.

Proposition 4.1 ([17, 14] applied to T^{-1}). *There exist some $Q > 0$, $K_0 > 0$, $\alpha \in (0, 1)$ and some finite partition \mathcal{P} of \mathbb{T}^d whose elements are of the form $\sum_{i=1}^d I_i \bar{v}_i$ where the I_i are intervals with diameter smaller than $\min(r_0, K)$ such that, for almost every $\bar{x} \in \mathbb{T}^d$,*

1. *the local leaf $\mathcal{P}_0^\infty(\bar{x})$ of \mathcal{P}_0^∞ containing \bar{x} is a bounded convex set $\bar{x} + \overline{F(\bar{x})}$, with $0 \in F(\bar{x}) \subseteq E_u$, $F(\bar{x})$ having non-empty interior in E_u ,*
2. *we have*

$$\mathbb{E}_n(f)(\bar{x}) = \frac{1}{m_u(S^{-n}F(T^n\bar{x}))} \int_{S^{-n}F(T^n\bar{x})} f(\bar{x} + \overline{h_u}) dm_u(h_u), \quad (4.43)$$

3. *for every $\gamma > 0$, we have*

$$m_u(\partial(F(\bar{x}))(\gamma)) \leq Q\gamma, \quad (4.44)$$

where

$$\partial F(\beta) := \{y \in F : d(y, \partial F) \leq \beta\},$$

4. *for every $\mathbf{k} \in \mathbb{Z}^d \setminus \{0\}$, for every integer $n \geq 0$,*

$$\left| \mathbb{E}_{-n}(e^{2i\pi(\mathbf{k}, \cdot)})(\bar{x}) \right| \leq \frac{K_0}{m_u(F(T^{-n}(x)))} |\mathbf{k}|^{d_e + d_s} \alpha^n, \quad (4.45)$$

5. *for every $\beta \in (0, 1)$,*

$$\exists L > 0, \forall n \geq 0, \bar{\lambda}(m_u(F(\cdot))) < \beta^n \leq L\beta^{n/d_u}. \quad (4.46)$$

Proof. The first item comes from Proposition II.1 of [14]. Item 2 comes from the formula given after Lemma II.2 of [14]. Item 3 follows from Lemma III.1 of [14] and from the fact that the numbers $a(\mathcal{P}_0^\infty(\cdot))$ considered in [14] are uniformly bounded. Item 4 comes from Proposition III.3 of [14] and from the uniform boundedness of $a(\mathcal{P}_0^\infty(\cdot))$. Item 5 comes from the proof of Proposition II.1 of [14]. \square

According to the first item of Proposition 4.1 and to (4.41), there exists $c_u > 0$ such that, for almost every $\bar{x} \in \mathbb{T}^d$ and every $n \geq 1$, we have

$$\sup_{h_u \in S^{-n}F(T^n(\bar{x}))} |h_u| \leq c_u \rho_u^n. \quad (4.47)$$

Proposition 4.2. *Let $p \geq 2$ and q be its conjugate exponent. Let $\theta > 0$ and $f : \mathbb{T}^d \rightarrow \mathbb{R}$ be a centered function with Fourier coefficients $(c_{\mathbf{k}})_{\mathbf{k} \in \mathbb{Z}^d}$ satisfying*

$$\sum_{|\mathbf{k}| \geq b} |c_{\mathbf{k}}|^q \leq R \log^{-\theta}(b). \quad (4.48)$$

Then

$$\|\mathbb{E}_0(f \circ T^n)\|_p = \|\mathbb{E}_{-n}(f)\|_p = O(n^{-\theta(p-1)/p}).$$

Proof. Recall first that $\mathbb{E}_0(f \circ T^n) = \mathbb{E}_{-n}(f) \circ T^n$. Let us consider α satisfying (4.45). Let $\beta := \alpha^{1/2}$, $\gamma := \max(\alpha^{p/2}, \beta^{1/d_u})$ and

$$\mathcal{V}_n := \{\bar{x} \in \mathbb{T}^d : m_u(F(T^{-n}(\bar{x}))) \geq \beta^n\}.$$

Let $b(n) := \lceil \gamma^{-n/(2p(d+e+d_s))} \rceil$. Let us write

$$f = f_{1,n} + f_{2,n} \text{ where } f_{1,n} := \sum_{|\mathbf{k}| < b(n)} c_{\mathbf{k}} e^{2i\pi(\mathbf{k}, \cdot)} \text{ and } f_{2,n} := \sum_{|\mathbf{k}| \geq b(n)} c_{\mathbf{k}} e^{2i\pi(\mathbf{k}, \cdot)}. \quad (4.49)$$

We have

$$\begin{aligned} \int_{\mathcal{V}_n} |\mathbb{E}_{-n}(f_{1,n})|^p d\bar{\lambda} &\leq \text{esssup}_{\bar{x} \in \mathcal{V}_n} \left(\sum_{|\mathbf{k}| \leq b(n)} |c_{\mathbf{k}}| |\mathbb{E}_{-n}(e^{2i\pi(\mathbf{k}, \cdot)})(\bar{x})| \right)^p \\ &\leq \left(\sum_{|\mathbf{k}| \leq b(n)} |c_{\mathbf{k}}| K_0 \beta^{-n} |\mathbf{k}|^{d_e+d_s} \alpha^n \right)^p, \end{aligned}$$

according to (4.45) and thanks to the definition of \mathcal{V}_n . Now, since $\beta = \alpha^{1/2}$, we get

$$\int_{\mathcal{V}_n} |\mathbb{E}_{-n}(f_{1,n})|^p d\bar{\lambda} \leq 3^{2p} \|f\|_1^p K_0^p \alpha^{\frac{np}{2}} (b(n))^{p(d+e+d_s)}.$$

Hence

$$\int_{\mathcal{V}_n} |\mathbb{E}_{-n}(f_{1,n})|^p d\bar{\lambda} = O(\gamma^n (b(n))^{p(d+e+d_s)}) = O(\gamma^{n/2}). \quad (4.50)$$

Moreover, thanks to (4.46), we have

$$\begin{aligned} \int_{\mathcal{V}_n^c} |\mathbb{E}_{-n}(f_{1,n})|^p d\bar{\lambda} &\leq \bar{\lambda}(\mathcal{V}_n^c) \left(\sum_{|\mathbf{k}| \leq b(n)} |c_{\mathbf{k}}| \right)^p \\ &= O((b(n))^{dp} \beta^{n/d_u}) = O((b(n))^{dp} \gamma^n) = O(\gamma^{n/2}). \end{aligned} \quad (4.51)$$

Since $p \geq 2$ and since $p/q = p - 1$, thanks to (4.48), we have

$$\begin{aligned} \|\mathbb{E}_{-n}(f_{2,n})\|_p^p &\leq \|f_{2,n}\|_p^p \leq \left(\sum_{|\mathbf{k}| \geq b(n)} |c_{\mathbf{k}}|^q \right)^{p/q} \\ &\leq R^{p-1} (\log(b(n)))^{-\theta(p-1)} \ll n^{-\theta(p-1)}. \end{aligned} \quad (4.52)$$

Combining (4.50), (4.51) and (4.52), the proposition follows. \square

Proposition 4.3. *Under the assumptions of Proposition 4.2,*

$$\|\mathbb{E}_0(f \circ T^{-n}) - f\|_p = \|\mathbb{E}_n(f) - f\|_p = O(n^{-\theta(p-1)/p}).$$

Proof. We consider the decomposition (4.49) with $b(n)$ defined by $b(n) = \lceil \rho_u^{-n/(2(d+1))} \rceil$. We have

$$\begin{aligned} \|\mathbb{E}_n(f_{1,n}) - f_{1,n}\|_p &\leq \|\mathbb{E}_n(f_{1,n}) - f_{1,n}\|_\infty \\ &\leq \sum_{|\mathbf{k}| \leq b(n)} |c_{\mathbf{k}}| \|\mathbb{E}_n(e^{2i\pi \langle \mathbf{k}, \cdot \rangle}) - e^{2i\pi \langle \mathbf{k}, \cdot \rangle}\|_\infty \\ &\leq \sum_{|\mathbf{k}| \leq b(n)} |c_{\mathbf{k}}| 2\pi |\mathbf{k}| c_u \rho_u^n, \end{aligned}$$

according to (4.43) and to (4.47). Therefore

$$\|\mathbb{E}_n(f_{1,n}) - f_{1,n}\|_p \ll (b(n))^{d+1} \rho_u^n \ll \rho_u^{n/2}. \quad (4.53)$$

Moreover, thanks to (4.48), we have

$$\begin{aligned} \|\mathbb{E}_n(f_{2,n}) - f_{2,n}\|_p^p &\leq 2^p \|f_{2,n}\|_p^p \leq 2^p \left(\sum_{|\mathbf{k}| \geq b(n)} |c_{\mathbf{k}}|^q \right)^{p/q} \\ &\leq 2^p R^{p-1} (\log(b(n)))^{-\theta(p-1)} \ll n^{-\theta(p-1)}. \end{aligned} \quad (4.54)$$

Considering (4.53) and (4.54), the proposition follows. \square

Proposition 4.4. *Let $p \in [2, 4]$ and set $S_n(f) := \sum_{k=1}^n f \circ T^k$ with $f : \mathbb{T}^d \rightarrow \mathbb{R}$ be a centered function with Fourier coefficients satisfying (4.48) with $\theta > 0$ and*

$$\sum_{|\mathbf{k}| \geq b} |c_{\mathbf{k}}|^2 \leq R \log^{-\eta}(b) \quad \text{for some } \eta > 1. \quad (4.55)$$

Set

$$m := \left\lceil -\frac{4(d_e + d_s) \log(r)}{\log(\alpha)} \right\rceil + 1. \quad (4.56)$$

where r is the spectral radius of S . Then

$$\|\mathbb{E}_{-nm}(S_n^2(f)) - \mathbb{E}(S_n^2(f))\|_{p/2} \ll n^{2-2\theta(p-1)/p} + n^{(3-\eta)/2}.$$

Proof. Let $\beta := \alpha^{1/2}$, $\mathcal{V}_{nm} := \{\bar{x} \in \mathbb{T}^d : m_u(F(T^{-nm}(\bar{x}))) \geq \beta^{nm}\}$, $\gamma := \max(\alpha^{p/8}, \beta^{1/d_u})$ and

$$b(n) := \left\lceil \gamma^{nm/(p(2d+d_e+d_s))} \right\rceil. \quad (4.57)$$

We consider the decomposition (4.49) with $b(n)$ defined by (4.57) and we set

$$S_{1,n}(f) := \sum_{k=1}^n f_{1,n} \circ T^k \quad \text{and} \quad S_{2,n}(f) := \sum_{k=1}^n f_{2,n} \circ T^k.$$

First, we note that

$$\begin{aligned}
& \|\mathbb{E}_{-nm}(S_n^2(f)) - \mathbb{E}(S_n^2(f))\|_{p/2} \\
& \leq \|\mathbb{E}_{-nm}(S_{1,n}^2(f)) - \mathbb{E}(S_{1,n}^2(f))\|_{p/2} \\
& \quad + \|\mathbb{E}_{-nm}(S_{2,n}^2(f)) - \mathbb{E}(S_{2,n}^2(f))\|_{p/2} \\
& \quad + 2\|\mathbb{E}_{-nm}(S_{1,n}(f)S_{2,n}(f)) - \mathbb{E}(S_{1,n}(f)S_{2,n}(f))\|_{p/2} \\
& \leq \|\mathbb{E}_{-nm}(S_{1,n}^2(f)) - \mathbb{E}(S_{1,n}^2(f))\|_{p/2} \\
& \quad + 2\|S_{2,n}(f)\|_p^2 + 4\|\mathbb{E}_{-nm}(S_{1,n}(f)S_{2,n}(f))\|_{p/2}.
\end{aligned}$$

Next using (3.32), we get that

$$\begin{aligned}
\|\mathbb{E}_{-nm}(S_{1,n}(f)S_{2,n}(f))\|_{p/2} & \leq \|\mathbb{E}_{-nm}(S_{1,n}^2(f)) - \mathbb{E}(S_{1,n}^2(f))\|_{p/2} \\
& \quad + \|S_{2,n}(f)\|_p^2 + \|S_{1,n}(f)\|_2 \|S_{2,n}(f)\|_2 \\
& \leq \|\mathbb{E}_{-nm}(S_{1,n}^2(f)) - \mathbb{E}(S_{1,n}^2(f))\|_{p/2} \\
& \quad + 2\|S_{2,n}(f)\|_p^2 + \|S_n(f)\|_2 \|S_{2,n}(f)\|_2.
\end{aligned}$$

By Propositions 4.2 and 4.3, (4.55) implies that

$$\sum_{n>0} \frac{\|\mathbb{E}_{-n}(f)\|_2}{n^{1/2}} < \infty \quad \text{and} \quad \sum_{n>0} \frac{\|f - \mathbb{E}_n(f)\|_2}{n^{1/2}} < \infty,$$

which yields (3.6) with $p = 2$, and then $\|S_n(f)\|_2 \ll \sqrt{n}$. Therefore, we get overall that

$$\begin{aligned}
\|\mathbb{E}_{-nm}(S_n^2(f)) - \mathbb{E}(S_n^2(f))\|_{p/2} & \ll \|\mathbb{E}_{-nm}(S_{1,n}^2(f)) - \mathbb{E}(S_{1,n}^2(f))\|_{p/2} \\
& \quad + \|S_{2,n}(f)\|_p^2 + \sqrt{n}\|S_{2,n}(f)\|_2. \quad (4.58)
\end{aligned}$$

Since $p \geq 2$ and $p/q = p - 1$, (4.48) implies that

$$\begin{aligned}
\|S_{2,n}(f)\|_p & \leq n\|f_{2,n}\|_p \leq n \left(\sum_{|\mathbf{k}| \geq b(n)} |c_{\mathbf{k}}|^q \right)^{1/q} \\
& \leq nR^{(p-1)/p} (\log(b(n)))^{-\theta(p-1)/p} \ll n^{1-\theta(p-1)/p}. \quad (4.59)
\end{aligned}$$

Similarly using (4.55), we get that

$$\|S_{2,n}(f)\|_2 \leq n\|f_{2,n}\|_2 \ll n^{1-\eta/2}. \quad (4.60)$$

We deal now with the first term in the right hand side of (4.58). With this aim, we first observe that, for any non negative integer ℓ , $e^{2i\pi\langle \mathbf{k}, T^\ell(\cdot) \rangle} = e^{2i\pi\langle {}^t S^\ell \mathbf{k}, \cdot \rangle}$,

where ${}^t S^\ell$ is the transposed matrix of S^ℓ . Therefore,

$$\begin{aligned} & \int_{\mathcal{V}_{nm}} \left| \mathbb{E}_{-nm}(f_{1,n} \cdot f_{1,n} \circ T^\ell) - \mathbb{E}(f_{1,n} \cdot f_{1,n} \circ T^\ell) \right|^{p/2} d\bar{\lambda} \\ & \leq \operatorname{esssup}_{\bar{x} \in \mathcal{V}_{nm}} \left(\sum_{|\mathbf{k}|, |\mathbf{m}| \leq b(n): \mathbf{k} + {}^t S^\ell \mathbf{m} \neq 0} |c_{\mathbf{k}}| |c_{\mathbf{m}}| \left| \mathbb{E}_{-nm}(e^{2i\pi(\mathbf{k} + {}^t S^\ell \mathbf{m}, \cdot)})(\bar{x}) \right| \right)^{p/2} \\ & \leq \left(\sum_{|\mathbf{k}|, |\mathbf{m}| \leq b(n)} |c_{\mathbf{k}}| |c_{\mathbf{m}}| K_0 \beta^{-nm} |\mathbf{k} + {}^t S^\ell \mathbf{m}|^{d_e + d_s} \alpha^{nm} \right)^{p/2}, \end{aligned}$$

according to (4.45) and to the definition of \mathcal{V}_{nm} . It follows that

$$\begin{aligned} & \int_{\mathcal{V}_{nm}} \left| \mathbb{E}_{-nm}(f_{1,n} \cdot f_{1,n} \circ T^\ell) - \mathbb{E}(f_{1,n} \cdot f_{1,n} \circ T^\ell) \right|^{p/2} d\bar{\lambda} \\ & \leq \left(\sum_{|\mathbf{k}|, |\mathbf{m}| \leq b(n)} \|f\|_1^2 K_0 \beta^{-nm} (|\mathbf{k}| + r^\ell |\mathbf{m}|)^{d_e + d_s} \alpha^{nm} \right)^{p/2} \\ & \ll \alpha^{\frac{npm}{4}} r^{p\ell(d_e + d_s)/2} (b(n))^{p(2d_e + d_s)/2}. \end{aligned}$$

Hence, since $\gamma \geq \alpha^{p/8}$, $m \geq 4(d_e + d_s) \log(r)/\log(1/\alpha)$, and according to the definition of $b(n)$, we have

$$\sup_{\ell \in \{0, \dots, n\}} \int_{\mathcal{V}_{nm}} \left| \mathbb{E}_{-nm}(f_{1,n} \cdot f_{1,n} \circ T^\ell) - \mathbb{E}(f_{1,n} \cdot f_{1,n} \circ T^\ell) \right|^{p/2} d\bar{\lambda} \ll \alpha^{3nmp/16} r^{pn(d_e + d_s)/2} \ll \gamma^{nm/2}. \quad (4.61)$$

Moreover, for any non negative integer ℓ ,

$$\begin{aligned} & \int_{\mathcal{V}_{nm}^c} \left| \mathbb{E}_{-nm}(f_{1,n} \cdot f_{1,n} \circ T^\ell) \right|^{p/2} d\bar{\lambda} \leq \bar{\lambda}(\mathcal{V}_{nm}^c) \left(\sum_{|\mathbf{k}|, |\mathbf{m}| \leq b(n)} |c_{\mathbf{k}}| |c_{\mathbf{m}}| \right)^{p/2} \\ & \ll (b(n))^{dp} \beta^{nm/d_u} \ll (b(n))^{dp} \gamma^{nm} \ll \gamma^{nm/2}, \quad (4.62) \end{aligned}$$

according to (4.46) and to the definition of $b(n)$ and of γ . Combining (4.61) and (4.62), we then derive that

$$\begin{aligned} & \|\mathbb{E}_{-nm}(S_{1,n}^2(f)) - \mathbb{E}(S_{1,n}^2(f))\|_{p/2} \\ & \leq 2 \sum_{i=1}^n \sum_{j=0}^{n-i} \|\mathbb{E}_{-nm}(f_{1,n} \circ T^i f_{1,n} \circ T^{i+j}) - \mathbb{E}(f_{1,n} \circ T^i f_{1,n} \circ T^{i+j})\|_{p/2} \\ & \leq n^2 \sup_{\ell \in \{0, \dots, n\}} \|\mathbb{E}_{-nm}(f_{1,n} f_{1,n} \circ T^\ell) - \mathbb{E}(f_{1,n} f_{1,n} \circ T^\ell)\|_{p/2} \ll n^2 \gamma^{nm/p}. \end{aligned} \quad (4.63)$$

Considering (4.59), (4.60) and (4.63) in (4.58), the proposition follows. \square

4.2. End of the proof of Theorem 2.1

Proof. Propositions 4.2 and 4.3 give (3.1) provided (2.2) is satisfied. Propositions 4.2 and 4.3 give (3.2) (when $p \in]2, 4[$) and (3.5) (when $p = 4$), provided (2.3) is satisfied. Finally, Proposition 4.4 gives (3.3) provided (2.2) and (2.3) are satisfied. The proof follows now from Theorem 3.1 when $p \in]2, 4[$ and from Theorem 3.2 when $p = 4$. \square

5. Appendix

As in Section 3, let $P_k(X) = \mathbb{E}_k(X) - \mathbb{E}_{k-1}(X)$.

Lemma 5.1. *Let $p \in [2, \infty[$. Then, for any real $1 \leq q \leq p$ and any positive integer n ,*

$$\sum_{k \geq 2n} \|P_0(X_k)\|_p^q \ll \sum_{k \geq n} \frac{\|\mathbb{E}_0(X_k)\|_p^q}{k^{q/p}}$$

and
$$\sum_{k \geq 2n} \|P_0(X_{-k})\|_p^q \ll \sum_{k \geq n} \frac{\|X_{-k} - \mathbb{E}_0(X_{-k})\|_p^q}{k^{q/p}}.$$

Proof. The first inequality is Lemma 5.1 in [2]. To prove the second one, we first consider the case $p > q$ and we follow the lines of the proof Lemma 5.1 in [2] with $P_k(X_0)$ replacing $P_{-k}(X_0)$. We get that

$$\sum_{k \geq 2n} \|P_0(X_{-k})\|_p^q \ll \sum_{k \geq n+1} k^{-\frac{q}{p}} \left(\sum_{\ell \geq k} \|P_0(X_{-\ell})\|_p^p \right)^{q/p}.$$

Now, we notice that, by the Rosenthal's inequality given in Theorem 2.12 of [9], there exists a constant c_p depending only on p such that

$$\begin{aligned} \sum_{\ell \geq k} \|P_0(X_{-\ell})\|_p^p &= \sum_{\ell \geq k} \|P_\ell(X_0)\|_p^p \\ &\leq c_p \left\| \sum_{\ell \geq k} P_\ell(X_0) \right\|_p^p = c_p \|X_0 - \mathbb{E}_k(X_0)\|_p^p = c_p \|X_{-k} - \mathbb{E}_0(X_{-k})\|_p^p. \end{aligned} \quad (5.1)$$

Now when $p = q$, inequality (5.1) together with the fact that by Claim 3.1, for any integer k in $[n+1, 2n]$, $\|X_0 - \mathbb{E}_{2n}(X_0)\|_p^p \leq 2^p \|X_0 - \mathbb{E}_k(X_0)\|_p^p$ imply the result. Indeed we have

$$\sum_{k \geq 2n} \|P_0(X_{-\ell})\|_p^p \leq c_p \|X_0 - \mathbb{E}_{2n}(X_0)\|_p^p \ll \sum_{k=n+1}^{2n} k^{-1} \|X_0 - \mathbb{E}_k(X_0)\|_p^p.$$

The proof is complete. \square

Proposition 5.1. *Let $p \in [1, \infty[$ and assume that*

$$\text{the series } d_0 = \sum_{i \in \mathbb{Z}} P_0(X_i) \text{ converges in } \mathbb{L}^p. \quad (5.2)$$

Let $M_n := \sum_{i=1}^n d_0 \circ T^i$ and $R_n := S_n - M_n$. Then, for any positive integers n and N ,

$$\begin{aligned} R_n &= \mathbb{E}_0(S_n) - \mathbb{E}_0(S_N) \circ T^n + \mathbb{E}_{-n}(S_N) \circ T^n - \sum_{k=1}^n \sum_{j \geq n+N+1} P_k(X_j) \\ &\quad + S_n - \mathbb{E}_n(S_n) - (\mathbb{E}_{n+N}(S_N - \mathbb{E}_N(S_N))) \circ T^{-N} - \sum_{k=1}^n \sum_{j \geq N} P_k(X_{-j}), \end{aligned}$$

and

$$\begin{aligned} \|R_n\|_p^{p'} &\ll \| \mathbb{E}_0(S_n) \|_p^{p'} + \| \mathbb{E}_0(S_N) \|_p^{p'} + \| S_n - \mathbb{E}_n(S_n) \|_p^{p'} + \| S_N - \mathbb{E}_N(S_N) \|_p^{p'} \\ &\quad + \sum_{k=1}^n \left\| \sum_{j \geq k+N} P_0(X_j) \right\|_p^{p'} + \sum_{k=1}^n \left\| \sum_{j \geq k+N} P_0(X_{-j}) \right\|_p^{p'}, \end{aligned}$$

where $p' = \min(2, p)$.

Proof. Notice first that the following decomposition is valid: for any positive integer n ,

$$\begin{aligned} R_n &= \sum_{k=1}^n \left(X_k - \sum_{j=1}^n P_j(X_k) \right) - \sum_{k=1}^n \sum_{j \geq n+1} P_k(X_j) - \sum_{k=1}^n \sum_{j=0}^{\infty} P_k(X_{-j}) \\ &= R_{n,1} + R_{n,2}, \end{aligned} \tag{5.3}$$

where

$$R_{n,1} := \mathbb{E}_0(S_n) - \sum_{k=1}^n \sum_{j \geq n+1} P_k(X_j), \quad R_{n,2} := S_n - \mathbb{E}_n(S_n) - \sum_{k=1}^n \sum_{j=0}^{\infty} P_k(X_{-j}). \tag{5.4}$$

Let N be a positive integer. According to item 1 of Proposition 2.1 in [2],

$$\begin{aligned} R_{n,1} &= \mathbb{E}_0(S_n) - \mathbb{E}_n(S_{n+N} - S_n) \\ &\quad + \mathbb{E}_0(S_{n+N} - S_n) - \sum_{k=1}^n \sum_{j \geq n+N+1} P_k(X_j). \end{aligned} \tag{5.5}$$

On an other hand, we write that

$$\sum_{j=0}^{\infty} P_k(X_{-j}) = \sum_{j=0}^{N-1} P_k(X_{-j}) + \sum_{j \geq N} P_k(X_{-j}).$$

Therefore

$$R_{n,2} = S_n - \mathbb{E}_n(S_n) - (\mathbb{E}_{n+N}(S_N - \mathbb{E}_N(S_N))) \circ T^{-N} - \sum_{k=1}^n \sum_{j \geq N} P_k(X_{-j}). \tag{5.6}$$

Starting from (5.3) and considering (5.5) and (5.6), the first part follows. We turn now to the second part of the proposition. Applying Burkholder's inequality and using stationarity, we obtain that there exists a positive constant

c_p such that, for any positive integer n ,

$$\begin{aligned} \left\| \sum_{k=1}^n \sum_{j \geq n+N+1} P_k(X_j) \right\|_p^{p'} &\leq c_p \sum_{k=1}^n \left\| \sum_{j \geq n+N+1} P_k(X_j) \right\|_p^{p'} \\ &= c_p \sum_{k=1}^n \left\| \sum_{j \geq N+k} P_0(X_j) \right\|_p^{p'}, \end{aligned} \quad (5.7)$$

and

$$\begin{aligned} \left\| \sum_{k=1}^n \sum_{j \geq N} P_k(X_{-j}) \right\|_p^{p'} &\leq c_p \sum_{k=1}^n \left\| \sum_{j \geq N} P_k(X_{-j}) \right\|_p^{p'} \\ &= c_p \sum_{k=1}^n \left\| \sum_{j \geq N+k} P_0(X_{-j}) \right\|_p^{p'}. \end{aligned} \quad (5.8)$$

The second part of the proposition follows from item 1 by taking into account the stationarity and by considering the bounds (5.7) and (5.8). \square

References

- [1] G. D. Birkhoff, *Proof of the ergodic theorem*. Proc. Nat. Acad. Sci. USA. **17** (1931), 656–660.
- [2] J. Dedecker, P. Doukhan and F. Merlevède, *Rates of convergence in the strong invariance principle under projective criteria*. Electron. J. Probab. **17** (2012), 1–31.
- [3] J. Dedecker, S. Gouëzel and F. Merlevède, *The almost sure invariance principle for unbounded functions of expanding maps*. ALEA, Lat. Am. J. Probab. Math. Stat. **9** (2012), 141–163.
- [4] M. Denker and W. Philipp, *Approximation by Brownian motion for Gibbs measures and flows under a function*. Ergodic Theory Dynam. Systems **4** (1984), 541–552.
- [5] D. Dolgopyat, *Limit theorems for partially hyperbolic systems*. Trans. Amer. Math. Soc. **356** (2004), 1637–1689.
- [6] M. J. Field, I. Melbourne I. and A. Török, *Decay of correlations, central limit theorems and approximation by Brownian motion for compact Lie group extensions*, Ergodic Theory Dynam. Systems **23** (2003), 87–110.
- [7] M. I. Gordin, *The central limit theorem for stationary processes*. Dokl. Akad. Nauk SSSR **188** (1969), 739–741.
- [8] S. Gouëzel, *Almost sure invariance principle for dynamical systems by spectral methods*. Ann. Probab. **38** (2010), 1639–1671.
- [9] P. Hall P. and C. C. Heyde, *Martingale limit theory and its application*. Probab. Math. Stat. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York-London 1980.
- [10] C. C. Heyde, *On the central limit theorem for stationary processes*. Z. Wahrscheinlichkeitstheorie und Verw. Gebiete **30** (1974), 315–320

- [11] F. Hofbauer and G. Keller, *Ergodic properties of invariant measures for piecewise monotonic transformations*. Math. Z. **180** (1982), 119–140.
- [12] A. Khintchine, *Zu Birkhoffs Lösung des Ergodenproblems*. Math. Ann. **107** (1933) 485–488.
- [13] J. Komlós, P. Major G. and Tusnády, *An approximation of partial sums of independent RV's, and the sample DF. II*. Z. Wahrscheinlichkeitstheorie und Verw. Gebiete **34** (1976), 33–58.
- [14] S. Le Borgne, *Limit theorems for non-hyperbolic automorphisms of the torus*. Israel J. Math. **109** (1999), 61–73.
- [15] S. Le Borgne S. and F. Pène F, *Vitesse dans le théorème limite central pour certains systèmes dynamiques quasi-hyperboliques*. Bull. Soc. Math. France **133** (2005), 395–417.
- [16] V. P. Leonov, *Central limit theorem for ergodic endomorphisms of compact commutative groups*. Dokl. Acad. Sci. USSR **135** (1960), 258–261.
- [17] D. A. Lind, *Dynamical properties of quasihyperbolic toral automorphisms*. Ergodic Theory Dynamical Systems **2** (1982), 49–68.
- [18] P. Major, *The approximation of partial sums of independent RV's*. Z. Wahrscheinlichkeitstheorie und Verw. Gebiete **35** (1976), 213–220.
- [19] I. Melbourne and M. Nicol, *Almost sure invariance principle for nonuniformly hyperbolic systems*. Commun. Math. Phys. **260** (2005), 131–146.
- [20] I. Melbourne I. and M. Nicol, *A vector-valued almost sure invariance principle for hyperbolic dynamical systems*. Ann. Probab. **37** (2009), 478–505.
- [21] I. Melbourne and A. Török, *Central limit theorems and invariance principles for timeone maps of hyperbolic flows*. Commun. Math. Phys. **229** (2002), 57–71.
- [22] F. Merlevède and M. Peligrad, *Rosenthal inequalities for martingales and stationary sequences and examples*, to appear in Ann. Probab. (2012). arXiv:1103.3242.
- [23] F. Merlevède and E. Rio, *Strong approximation of partial sums under dependence conditions with application to dynamical systems*. Stochastic Process. Appl. **122** (2012), 386–417.
- [24] N. Nagayama, *Almost sure invariance principle for dynamical systems with stretched exponential mixing rates*. Hiroshima Math. J. **34** (2004), 371–411.
- [25] W. Philipp and W. F. Stout, *Almost sure invariance principle for partial sums of weakly dependent random variables*. Mem. of the Amer. Math. Soc. **161** (1975), Providence, RI: Amer. Math. Soc.
- [26] Q. M. Shao, *Almost sure invariance principles for mixing sequences of random variables*. Stochastic Process. Appl. **48** (1993), 319–334.
- [27] W. B. Wu, *Strong invariance principles for dependent random variables*. Ann. Probab. **35** (2007), 2294–2320.

Jérôme Dedecker

Université Paris Descartes, Sorbonne Paris Cité,

Laboratoire MAP5 and CNRS UMR 8145

e-mail: jerome.dedecker@parisdescartes.fr

Florence Merlevède
Université Paris Est,
LAMA and CNRS UMR 8050
e-mail: florence.merlevede@univ-mlv.fr

Françoise Pène
Université de Brest,
Laboratoire de Mathématiques de Bretagne Atlantique and CNRS UMR 6205
e-mail: francoise.pene@univ-brest.fr