Methodological characteristics of the national dietary surveys carried out in the European Union as included in the EFSA Comprehensive European food consumption database

Journal:	<i>Food Additives and Contaminants</i>
Manuscript ID:	TFAC-2010-502.R1
Manuscript Type:	Original Research Paper
Date Submitted by the Author:	14-Mar-2011
Complete List of Authors:	Merten, Caroline; EFSA, DATEX Ferrari, Pietro; IARC Bakker, Martine; RIVM Boss, Alexander Hearty, Aine; University College Dublin, Institute of Food & Health, University College Dublin, Leclercq, Catherine; INRAN Lindtner, Oliver; BfR Tlustos, Christina; Food Safety Authority of Ireland Verger, Philippe; WHO Volatier, Jean-Luc Arcella, Davide; EFSA
Methods/Techniques:	Exposure assessment
Additives/Contaminants:	Nutrition
Food Types:	
Abstract:	In 2009 competent organisations in the European Union (EU) provided the European Food Safety Authority (EFSA) with data from the most recent national dietary survey, at the level of individuals' consumption. Twenty different Member States (MS) provided EFSA with data from 22 different national dietary surveys, with consumption figures for adults and, when available, for children. MSs' dietary data were assembled into the EFSA Comprehensive European Food Consumption Database (Comprehensive Database).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

	<p>In this paper an overview of the methodologies and protocols employed in the different national dietary surveys is provided. Specifically, details about dietary assessment methods, interview administration, sampling design, portion size estimation, dietary software, evaluation of under-reporting and non dietary information collected are described. This information is crucial to evaluate the level of accuracy of food consumption data and to anticipate and acknowledge utmost important sources of heterogeneity of national databases included in the Comprehensive database. The Comprehensive database constitutes a unique resource for the estimation of consumption figures across the EU and represents a useful tool to assess dietary exposure to hazardous substances and nutrient intake in Europe. Nevertheless, the many substantial methodological differences that characterise the Comprehensive database are acknowledged and critically discussed.</p>

SCHOLARONE™
Manuscripts

Peer Review Only

Methodological characteristics of the national dietary surveys carried out in the EU as included in the EFSA Comprehensive European Food Consumption Database.

Merten¹ C., Ferrari¹ P., Bakker² M., Boss³ A., Hearty⁴ Á., Leclercq⁵ C., Lindtner⁶ O., Tlustos⁷ C., Verger⁸ P., Volatier⁹ J.L. and Arcella¹ D.

¹ European Food Safety Authority (EFSA)

² Rijksinstituut voor Volksgezondheid en Milieu (RIVM)

³ Food Standards Agency (FSA)

⁴ University College Dublin (UCD)

⁵ Istituto Nazionale di Ricerca per gli Alimenti e la Nutrizione (INRAN)

⁶ Federal Institute for Risk Assessment (BfR)

⁷ Food Safety Authority of Ireland (FSAI)

⁸ World Health Organization (WHO)

⁹ Agence Nationale de Sécurité Sanitaire de l'Alimentation, de l'Environnement et du Travail (ANSES)

AUTHOR FOR CORRESPONDENCE:

Davide Arcella

Data Collection and Exposure Unit

European Food Safety Authority (EFSA)

Largo Palli 5/a, I-43121 Parma

Tel +39 0521 036828

Fax +39 0521 0360828

E-mail: davide.arcella@efsa.europa.eu

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

KEYWORDS: Dietary survey, exposure assessment, food consumption, 24-hour recall, food records.

RUNNING TITLE: Comprehensive European Food Consumption Database.

LIST OF ABBREVIATIONS

BMR:	Basal Metabolic Rate
Comprehensive Database:	EFSA Comprehensive European Food Consumption Database
DCF:	Data Collection Framework
EGFCD:	Expert Group on Food Consumption Data
EI:	Energy Intake
EU:	European Union
FFQ:	Food Frequency Questionnaire
FPQ:	Food Propensity Questionnaire
HBS:	Household Budget Surveys
MS:	Member State
PSMAs:	Portion Size Measurement Aids
WHO:	World Health Organization

ABSTRACT

In 2009 competent organisations in the EU provided the European Food Safety Authority (EFSA) with data from the most recent national dietary survey, at the level of individuals' consumption. Twenty different Member States (MS) provided EFSA with data from 22 different national dietary surveys, with consumption figures for adults and, when available, for children. MSs' dietary data were assembled into the EFSA Comprehensive European Food Consumption Database (Comprehensive Database). In this paper an overview of the methodologies and protocols employed in the different national dietary surveys is provided. Specifically, details about dietary assessment methods, interview administration, sampling design, portion size estimation, dietary software, evaluation of under-reporting and non dietary information collected are described. This information is crucial to evaluate the level of accuracy of food consumption data and to anticipate and acknowledge utmost important sources of heterogeneity of national databases included in the Comprehensive database. The Comprehensive database constitutes a unique resource for the estimation of consumption figures across the EU and represents a useful tool to assess dietary exposure to hazardous substances and nutrient intake in Europe. Nevertheless, the many substantial methodological differences that characterise the Comprehensive database are acknowledged and critically discussed.

Introduction

Food consumption data reflect what individuals or population groups consume in terms of foods, beverages, including drinking water, and supplements. Food consumption in a population can be estimated through surveys at individual level (Individual dietary surveys) or household level (Household budget surveys). Alternatively, consumption figures might be approximated through food supply data derived from food balance sheets. Individual dietary surveys are the only surveys that provide information on the distribution of food consumption in well-defined groups of individuals and are therefore preferred for the assessment of dietary exposure within the risk assessment process. Data from individual dietary surveys are also assumed to more closely reflect actual consumption (Kroes et al., 2002). National dietary surveys are presently carried out in many European countries and provide valuable information to be used in national policy and in nutritional surveillance.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

In 2007, the European Food Safety Authority (EFSA) established the “Expert Group On Food Consumption Data” (EGFCD), an EFSA network with representatives from each EU Member State (MS) to provide a platform for discussion to ultimately define practical guideline steps for the collection and collation of food consumption data. At the end of 2008, EFSA started a project aimed at establishing the EFSA Comprehensive European Food Consumption Database (Comprehensive Database). Within this project, competent organisations, nominated in each MS by the Permanent Representative to the European Union, were requested to provide EFSA with data from the most recent national dietary survey in their country.

In October 2009, the EGFCD endorsed the Guideline of EFSA on “Methods and protocols for the collection of national food consumption data in the view of a pan-European dietary survey” (EFSA, 2009). The main objective of this Guideline is to recommend general principles for the collection of dietary information to estimate the intake of foods and nutrients to perform risk assessment for a variety of biological agents and chemical substances evaluated by EFSA Scientific Panels.

The aim of this paper is to give an overview of the different methodologies used for the collection of food consumption and related data included in the Comprehensive Database. This information is crucial to evaluate the level of accuracy of food consumption data and to anticipate and acknowledge utmost important sources of heterogeneity for the comparison of this data at European level. Such knowledge will assist with realising estimates of uncertainty when assessing exposure to hazardous substances.

Materials and methods

Twenty MSs signed a collaboration agreement with EFSA for the provision and processing of dietary data collected through national surveys. The information provided made the establishment of the Comprehensive database possible. In order to be included in the database, the dietary data had to be collected at individual level by means of (replicates of) 24-hour dietary recalls or dietary records. Dietary surveys were requested to be representative at national level, at least for the adult population. The consumption data should be provided at the most disaggregated level recorded. All participating institutions provided EFSA with a database schema describing their food consumption and related data tables. Based on this information a common data model was developed for the transmission of the food

consumption data. Data providers coded all food descriptors present in the food consumption database according to a unique classification system developed by EFSA (EFSA, 2010). Data providers were also asked to disaggregate industrially produced composite foods or home-made dishes, such as a ready-made frozen pizza or a home cooked beef stew, into their main raw ingredients. The transmission of food consumption data was accomplished through an application designed by EFSA, called Data Collection Framework (DCF).

Data providers systematically compiled a report describing in detail the methodology employed in the dietary survey, according to an agreed protocol with the objective of providing detailed information on each of these sections: dietary method, administration of the interview, sampling design, portion size estimation, dietary software, food coding, evaluation of under-reporting and non dietary information.

All information contained in the reports was checked for completeness and consistency. When necessary, clarifications were requested to the data providers. Where applicable, information reported was verified against the related food consumption data provided to EFSA.

Results

Twenty different MSs provided food consumption data to EFSA at individual level collected in 22 different national dietary surveys. Table 1 shows the MS institutions that implemented the food consumption survey in their respective country. The methodological characteristics of the surveys are presented in Table 2. Five different types of survey methodologies were conducted: 7-day food records were carried out in five surveys, 3-day food records in three surveys, one-day 24-hour dietary recalls in six surveys and 2-day 24-hours dietary recalls in seven surveys. In Finland a 48 hours dietary recall method was used.

All countries which used 2-day 24-hour dietary recalls conducted the interview on two non-consecutive days with the exception of the most recent survey in Spain (Spain II) where about 35% of the interviews were conducted on consecutive days. The average length between non-consecutive days ranged from 3 (Spain II) to 79 days (Czech Republic). The food record surveys were consistently conducted on consecutive days.

The methodology used within the studies conducted in Denmark (Groth and Fagt, 1997) and France (Lafay et al., 2002) have been reported as validated. In the United Kingdom, a doubly

151 labelled water validation study was carried out as part of the feasibility study prior to the
152 main survey.

153
154 Fourteen surveys also included the administration of a Food Frequency Questionnaire (FFQ)
155 or of a Food Propensity Questionnaire (FPQ) whereas in Germany a diet history was
156 administered to the same study subjects that completed 24-hour dietary recalls. In addition,
157 Germany and Finland collected further dietary information, by means of food records, in a
158 sub-sample of the study population, whereas an additional 24-hour recall was administered to
159 all subjects within the Spain I survey. This information was not transmitted to EFSA.

160 The methods used for the administration of the interview are presented in Table 3. The
161 number of face-to-face meetings between interviewer and subjects to collect food
162 consumption information varied from none up to four meetings. In 15 surveys, at least one
163 face-to face interview took place at the study subject's home. In Ireland, subjects were
164 interviewed either at home or at their working place. In Slovakia, Bulgaria II and Finland the
165 interviews were conducted at a medical centre, whereas in Spain II subjects were interviewed
166 either at the university campus, health centres and pharmacies where the subjects were also
167 recruited for the study. In the Netherlands the interviews were conducted by phone without
168 any prior physical contact. In the German study there was only one face-to-face meeting with
169 the subject which aimed at collecting information on the socio demographic status, along
170 with information on nutritional and purchase behaviour, health aspects and anthropometric
171 measures. The 24-hours recalls were both conducted by phone a few weeks later. In Austria
172 no verbal contact with the study subjects was established. The subjects completed a 24-hour
173 dietary recall according to documented guidelines and returned it by mail to the study centre
174 (postal survey method). In thirteen surveys interviewers had a background in nutrition and in
175 the remaining surveys interviewers were trained staff without a nutrition related background.
176 The survey period, sampling strategy and response rate are shown in Table 4. In three surveys
177 (Estonia, Ireland and Sweden) the food consumption data were collected prior to the year
178 2000. In 16 surveys the study population was sampled at individual level whereas in the
179 remaining 6 surveys, it was sampled at household level. Sample units were selected randomly
180 in all surveys but different sampling frames were used. The national population register was
181 the most used sampling frame (in 8 surveys). In Spain, participants were randomly selected
182 from universities, health centres and pharmacies, whereas in Slovakia subjects were sampled
183 from lists of employees of confectionary and bakery manufactures. The response rate varied
184 from 27% (Hungary) to 96 % (Slovakia and Poland).

In six surveys the target population included children below 10 years old and in about half of the surveys the target population included adolescents from 10 to 17 years (Table 5). In all surveys, with the exception of Bulgaria II, which focused on children, surveys included the age range of 18 to 64 years. In nine surveys elderly people from 64 to 74 years were also included. In the Dutch dietary survey only young adults, from 19 to 30 years of age, were included.

The sample size, the stratification variables and the distribution by gender are shown in Table 6. The sample size across the surveys varied between 410 (Slovenia) and 13,926 (Germany) subjects. The vast majority of the surveys were nationally representative for gender (in 20 surveys), age groups (in 19) and geographical areas (in 15). Six surveys were also representative for urban and rural areas. In addition, the Dutch and Irish data were representative for education level. Austria was only representative according to the employment status. In six countries weighting factors were used to make the sample representative at national level for at least age groups, gender and regions.

The weekday and seasonal representativeness of the surveys are shown in Table 7. In six surveys record or recall days did not evenly cover week and weekend days. For example, in Slovakia only 5% of the records for which the consumption date was known related to weekend days. Twelve surveys captured consumption figures across all seasons. In the remaining surveys the seasonality was not fully covered, with only one season represented in Bulgaria I (Spring), Estonia (Summer), Hungary (Winter) and The Netherlands (Fall).

The applied exclusion criteria are shown in Table 8. The majority of MSs (in 17 surveys) excluded institutionalised persons, such as elderly in retirement homes or people residing in hospitals, prisons or military barracks. In seven surveys pregnant and breastfeeding women were excluded. Information on the diet of pregnant and breastfeeding women were available only from nine different surveys. In three surveys, specific population groups were purposely over represented, notably children aged 3–17 years in France, teenagers aged 15–18 years and people aged >75 years in Belgium, and subjects with lower education level in The Netherlands.

The methods used to estimate portion size are shown in Table 9. Three surveys were conducted using the weighing method, either as the sole method (United Kingdom for food consumed inside the home) or combined with other measurement tools (Ireland and Spain I),

1
2
3 219 to estimate the amount of food consumed. In the British survey, for food eaten outside of
4
5 220 home, a ruler and information on household measures and known packaging size were used.
6
7 221 In the majority of surveys (19) a combination of 2 or more measurement tools were used and
8
9 222 in 16 studies the picture book was used as one of these tools. In all but four (Ireland, Latvia,
10
11 223 Slovenia, Estonia) of these 16 surveys a validated or tested picture book was used. EPIC-soft
12
13 224 picture book was used in four surveys (Belgium, Germany, The Netherlands and Italy). Out
14
15 225 of the six surveys in which no picture book was used, two were weighed surveys (United
16
17 226 Kingdom and Spain I), Austria relied on household measurements only, Spain II was
18
19 227 conducted using household measurements and packaging size, while in the Slovakian survey
20
21 228 the interviewer estimated portion sizes without any tool but relied only on the subject's
22
23 229 description. In Hungary, subjects used "reference tables" to estimate and fill in the portion
24
25 230 sizes in the record. Three out of the six dietary surveys including children <10 years of age
26
27 231 (Bulgaria II, Denmark and Italy) reported the use of a picture book with small portion sizes
28
29 232 appropriate for children. The remaining three (Poland, Latvia and France) did not use specific
30
31 233 tools for children.
32
33 234
34
35 235 Details of the dietary software and related databases used in the different surveys are
36
37 236 presented in Table 10. In Belgium, Germany, and the Netherlands EPIC SOFT (Slimani et
38
39 237 al., 1999) was used, in Austria and France an *ad hoc* software using MS Access was
40
41 238 developed and used. In all other surveys different software were used. Most countries used
42
43 239 integrated portion size databases (in 17 surveys) and integrated standard recipe databases (in
44
45 240 20 surveys). In 10 surveys no yield factors were used when breaking down recipes and/or
46
47 241 composite dishes into the main raw ingredients. Only Sweden did not disaggregate recipes
48
49 242 and/or composite dishes into the main ingredients. In 19 surveys, the software links the food
50
51 243 consumption databases to food composition databases. With this respect, the EPIC SOFT
52
53 244 program is an exception since it does include food composition information only on
54
55 245 macronutrients. The availability of brand name, household processing and packaging
56
57 246 information in each survey per food record are presented in Table 11. Brand information was
58
59 247 available only in nine surveys for a percentage of records varying from 1% to 29%. In
60
248 Germany only a flag variable indicating the availability of the brand name was provided
249 because, at the time of the data transmission, data concerning brand name were still in the
250 cleaning phase. The description of the food incorporated information on brand, household
251 processing and packaging in Ireland and, for some of the food items, in UK. In these
252 circumstances it was not possible to calculate the percentage of food records including this

information. In Finland the brand information is also incorporated directly in the food name. Household processing information was available in 14 surveys for a percentage of records varying from 1% to 45% with the number of different processing methods reported in the various surveys varying from 1 to 24. Packaging information is available from three surveys for a percentage of records varying from 1% to 24%. In Belgium and Spain II the packaging information differentiates between physical characteristic such as carton or metal, whereas in Germany packaging information is available only for two food groups (fat and sauces).

The technique used in the various surveys to identify under-reporters is described in table 12. In eleven surveys, individual level under-reporters were identified among adults, by comparing the ratio of Energy Intake (EI) to Basal Metabolic Rate (BMR) to Goldberg cut off points (Goldberg 1991), showing frequency of under-reporters ranging from 11.7% (in Hungary) to 37.3% (in Finland). The cut-off was corrected on the basis of the physical activity level, as suggested by Black (2000a and 2000b) only in the survey from France. In Italy and in the Netherlands under-reporting among adults was assessed at the population level. Under reporting among children was only assessed in France, by comparing the logarithm of energy intake with the mean minus three standard deviations, and in Italy, by examining the means of the ratio of energy intake on estimated energy expenditure distributions. Under-reporters were excluded before transmitting data to EFSA only in the Austrian survey where subjects with a ratio of EI/BMR below 0.65 were excluded.

Information on specific study subjects' long term dietary pattern (e.g. vegetarian, health related or slimming) had been collected in half of the surveys (Table 13). In Germany information on further special diets like Halal was collected. Non dietary information collected within the survey is presented in Table 14. Data on body weight and height were consistently available. In nine surveys direct measurements were taken, while in the remaining, self reported measures were used. In 12 and eight surveys information on study subjects' physical activity level, self-reported by means of questionnaires, and ethnicity were collected, respectively.

Discussion

Dietary method

Only data collected through dietary records and 24-hours dietary recalls have been accepted to be included in the Comprehensive Database. The EGFCDC considered both methods

1
2
3 287 suitable for the requirements of providing data at a sufficiently detailed level for the purposes
4
5 288 of exposure assessment (EFSA, 2009) but differences exist between these two methods. The
6
7 289 dietary record method might be a more suitable tool for the elicitation of a greater level of
8
9 290 detail but, among others, a potential drawback might be the selection bias due to high
10
11 291 illiteracy levels in some minority groups across Europe (Biró et al., 2002). On the other hand,
12
13 292 an accurate memorisation of food consumed the preceding day is required for the 24-hours
14
15 293 dietary recalls method. This might represent a difficult cognitive task for the respondent, in
16
17 294 particular, for the very young or very old study subjects, and could affect the precision in the
18
19 295 quantification of the foods consumed (Thompson and Subar, 2001). Six of the countries
20
21 296 providing data into the Comprehensive Database collected dietary data on children younger
22
23 297 than 10 years. Half of these countries used the 24-hour recall method and the other half used
24
25 298 the dietary record method.
26
27 299
28 300 Food consumption data collected on more than one day per subject are required to assess
29
30 301 chronic exposure. At least two independent short-term assessments days are needed to apply
31
32 302 statistical modelling to estimate habitual intake (Dodd et al., 2006). The EGFCF (EFSA,
33
34 303 2009) recommends the collection of dietary information for two non-consecutive days for
35
36 304 both the 24-hour recall and the dietary record methods. Recording days are here considered
37
38 305 as non-consecutive if there is an interval of at least two weeks between them. Collecting data
39
40 306 on non-consecutive days has the potential advantage of making dietary measurements less
41
42 307 prone to correlation between errors in dietary assessments (Kipnis et al., 2003, Day et al.,
43
44 308 2004). However, in half of the surveys where 24-hour dietary recalls were used, information
45
46 309 on only one single day was collected per subject. Such data can nevertheless be useful to
47
48 310 estimate acute exposure. The seven surveys that used 2-day 24-hour recall method were
49
50 311 conducted on non-consecutive days, with the exception of Finland where a 48-hour dietary
51
52 312 recall method was used. In three of these seven surveys the average distance between two
53
54 313 interview days is below the recommended two weeks interval.
55
56 314
57 315 It has been argued that, accuracy of self-reported dietary information may decrease as the
58
59 316 number of days increases (Biro et al., 2002, Gersovitz et al., 1978, Moreno et al. 2005,
60
317 Whybrow et al., 2008). On the other hand, increasing the number of assessment days of a
318 survey affects the distribution of consumption, particularly at the upper tails (EFSA, 2006). In
319 particular, increasing the number of survey days (for both recalls and records) has the

advantage of reducing the effect of study subjects' day-to-day variation, thus leading to an improved estimation of consumption variability (Willett, 1989). As survey duration increases, also the observed percentage of subjects reporting non zero consumption for commonly and rarely eaten foods becomes larger (Nusser et al., 1995), whereas the observed mean and high percentiles consumption, in consumers only, decreases, as also illustrated by Lambe et al. (2000). Half of the surveys using the record methodology in the Comprehensive Database (Denmark, France, Ireland and United Kingdom) collected information for more than three days per subject.

Some episodically consumed foods may become of public health concern, as a result of high level contamination with a given hazardous substance. In order to accurately estimate the usual intake of these foods, additional information on their frequency of consumption could complement 24-hour dietary recalls or dietary records (Tooze et al., 2006; Kipnis et al., 2009), as emphasised in the EGFC (2009) guideline. Twelve countries reported using a FFQ or a FPQ along with a 24 h recall or dietary record method. In one country (Ireland), the FFQ focused solely on the intake of meat. Data from FFQ or FPQ may be used in acute exposure assessments to assess the proportion of consumers of a given food, and in chronic exposure assessments to determine the frequency distribution of the consumption of rarely consumed foods. Data from FFQ or FPQ was, however, not transmitted to EFSA.

Administration of the interview

The vast majority of surveys in the Comprehensive Database used the face-to-face interview method, mainly at the respondent's home. There were also two surveys conducting 24-hour dietary recalls by phone (Germany and the Netherlands), while in Austria one self-administered 24-hour dietary recall was collected via postal mail. Numerous studies (Derr et al., 1992; Lyu et al., 1998; Casey et al., 1999; Tran et al., 2000; Bogle et al., 2001) concluded that the telephone 24-hour dietary recall interview method provides a reliable and cost-effective alternative to a face-to-face method, with similar response rates being observed for both methods whereas there are not as many studies in the literature that have examined postal questionnaires as a method of obtaining dietary information. In general it seems that both face-to-face and telephone interviews are more suitable for collecting dietary and health data compared to postal questionnaires (Sibbaldi et al., 1994). The effect of the number of face-to-face meetings is likely to improve the accuracy of dietary estimates as the number of

1
2
3 353 contacts with the study subjects increases and especially if conducted by personnel with a
4 354 nutritional background.
5
6 355
7
8 356 The home visit is considered the most appropriate location to conduct dietary interviews
9 357 since it offers the advantage of collecting additional information on foods consumed, such as
10 358 brand level data and food packaging, and could allow for direct weighing of some dietary
11 359 items (De Henauw et al., 2002). On the other hand, other locations like medical centres are
12 360 likely to offer better conditions for the collection of additional information, such as biological
13 361 samples (Riboli et al., 1997).
14
15 362
16
17 363 Dietary interviews should be ideally conducted by nutritionists or dieticians. In alternative
18 364 adequately trained staff could be employed. In the Comprehensive Database, over half of the
19 365 surveys were conducted by an interviewer with a nutritional background.
20 366
21
22 367 *Sampling design*
23
24 368 The Comprehensive Database includes data from the most recent national dietary surveys
25 369 carried out in 20 MSs of the European Union. Some surveys (Estonia, Ireland, Spain and
26 370 Sweden) started before 2000, thus limiting the accuracy of consumption estimates of dietary
27 371 items recently introduced into the market.
28
29 372
30
31 373 Sample representativeness is a crucial aspect for the evaluation of the food consumption data
32 374 gathered in the Comprehensive Database. Representativeness is a qualitative parameter
33 375 related to the sampling design of a dietary survey. Significant biases can arise from a survey
34 376 sample that does not represent the population at national level.
35
36 377
37
38 378 Sample units (individuals or households) were selected randomly in all dietary surveys, but
39 379 differences in the sampling design are observed. Concerning the sampling frames used,
40 380 National population registers, address registers, postcode address files, general population
41 381 census, consumer panels and electoral lists (only for the adult population) can all be
42 382 considered as optimal sampling frames as far as these lists are regularly updated and each
43 383 subject within the national population register is likely to be selected. The national register of
44 384 the general practitioner's practices used in Bulgaria seems to be a good sampling frame for
45 385 children since, due to the national legislation, all children must be listed there. In Italy the use
46 386 of telephone books has the potential of introducing unwanted bias, as non-telephone

households, telephone numbers issued after the publication of the telephone directory, and unlisted numbers will be automatically excluded. In Spain, the use of universities, health centres and pharmacies to randomly recruit subjects is likely to constitute a potential source of bias. In Slovakia, the study population cannot be considered representative of the general population since subjects were only selected among employees of confectionary and bakery manufactures.

On the other hand, the use of the household as a sampling unit seems to be a convenient choice since an interviewer could collect information from more subjects during the same visit. However, food consumption estimates are likely to be mutually dependent when subjects from the same household are interviewed, thus leading to a reduced variability in terms of dietary pattern observed. In France and UK, only one respondent per household was included in the study. The random selection of subjects from households sampled within the national household budget survey in Poland seems to offer a good choice since it allows the comparability between the two sets of data. A cost reduction of the survey can be obtained by the use of a multistage sampling design where, for example, a random choice of cities, as primary units, of households, as secondary units, and finally individuals within households is adopted. This method can be combined with a stratification for instance on regions or rural/urban areas to reduce sampling bias because the part of the variability of the studied indicator explained by the strata is controlled (Cochran, 1977). All surveys in the Comprehensive Database were stratified for gender and age groups with the exception of Austria. The number of subjects participating in the dietary survey varied sizeably from survey to survey. The adult population group (from 18 to 64 years of age) was the only represented age group in all 20 MSs with a number of study subjects varying from 400 (Slovenia) to 10,419 (Germany) where the study population was sampled at a very fine geographical level (Federal State level). A large sample gives the opportunity to collect more information on rarely consumed foods.

The target population of a national dietary survey includes all people living in the country at the time of the study. However, institutionalised subjects, such as the elderly in retirement homes or people residing in hospitals, prisons or military barracks, have not been included in the large majority of dietary surveys. Their exclusion is often considered necessary to simplify sampling procedures, as outlined in the recommendation of the EGFCF (EFSA, 2009) for a future pan-European dietary survey.

1
2
3 421 Despite pregnant and breastfeeding women being specifically excluded only in 7 surveys,
4 422 where included their number is overall rather low, i.e. from 0 (Spain I) to 59 (Denmark)
5 423 breastfeeding women and from 0 (Spain II) to 52 (Germany) pregnant women. Therefore
6 424 dietary estimates of these important subgroups should be treated cautiously.
7
8
9
10
11 425

12 426 One important aspect of food consumption data is their representativeness over the different
13 427 seasons. Ideally, 24-hours dietary recalls and dietary records should be collected uniformly
14 428 over the four seasons. This was achieved in the majority of surveys in the Comprehensive
15 429 Database, with the exception of Bulgaria I, Bulgaria II, Estonia, Finland, Hungary, Latvia,
16 430 Netherlands, Poland and Spain II where an uneven distribution of recording days over
17 431 seasons was reported. This issue is particularly relevant when using food consumption data to
18 432 assess exposure to hazardous chemicals mainly present in seasonal foods. In addition, it has
19 433 been suggested that a proportionate collection of dietary estimates in week versus weekend
20 434 days should be adopted, at population level, in the sampling phase (Lyhne et al., 2005).
21 435 Weekend days were particularly under represented in Austria, Bulgaria and Slovakia and
22 436 over represented in Bulgaria II. The effects of uneven sampling fractions over days of the
23 437 week are potentially relevant for foods that exhibit specific consumption patterns related to
24 438 weekend consumption, e.g. alcoholic drinks.
25
26
27
28
29
30
31
32
33
34
35
36

37 440 It has been argued that the response rates of dietary surveys should be high enough to ensure
38 441 that individual dietary estimates can be generalised to the general population, thus avoiding
39 442 under-sampling of specific population subgroups. Response rates were relatively low in
40 443 Hungary (27%), Spain II (28%), Italy (33%), Belgium (41%), Netherlands (42%) and
41 444 Germany (42% for the total study). The high response rate in Slovakia (98%) is likely due to
42 445 the very convenient sampling frame used. No information was collected on type of incentives
43 446 for the study subjects. The use of incentives is a common method for increasing survey
44 447 response but can also introduce bias in dietary estimates by inadvertently drawing individuals
45 448 from selected population subgroups or by influencing respondents' responses (Singer &
46 449 Kulka, 2002).
47
48
49
50
51
52
53
54
55

56 451 *Portion size estimation*

57 452 Systematic bias and large random error may occur while quantifying foods and no gold
58 453 standard exists for estimation of portion size (Wrieden et al., 2009). The use of different aids
59 454 will depend on survey methodology, target population (Foster et al., 2009) and the level of
60

accuracy required. Weighing is considered to be the most precise method for measuring food intake, however it is time-consuming, costly and disruptive (Wolper et al., 1995). It was used, alone or in combination with other methods, in the United Kingdom, Ireland and Spain only. As an alternative to weighing all food items eaten, consumption can be measured based on the subjects' estimates of portion size. The parallel use of different Portion Size Measurement Aids (PSMAs), such as photographs, household measures, rulers etc, was considered the most convenient option by the EGFCD (EFSA, 2009) in order to obtain best estimates for different foods. Indeed the majority of surveys (in 19) used a combination of two or more measurement tools, with 16 using the picture book as one of the tools and all but four of these used a validated or tested picture book. Measuring children's dietary intake is challenging. Foster et al. (2009) investigated whether the estimate of portion sizes for children would improve if they were provided with age appropriate tools. They concluded that providing children with food photographs depicting age-appropriate portion sizes increased accuracy of estimates compared with estimates using photographs designed for use with adults. The EGFCD also stressed the importance of using age-appropriate tools and portion size aids which are representative of the food available on the market and of the food portions actually consumed. Six of the 22 surveys included children <10 years of age. Bulgaria II, Denmark and Italy reported the use of the same picture book as for the adults but with special sets of smaller portion sizes for children.

It might be advisable to more closely examine estimated food portion quantities in those surveys' data using only household measurement tools (Austria), household measurement tools in combination with packaging size (Spain II) and, in particular, in those two countries reporting no use of any PSMAs to quantify portion sizes. This latest case is related to the dietary surveys carried out in Slovakia, where the interviewer estimated portions relying only on the subject's description, and in Hungary, where "reference tables" were used by all the study subjects.

Dietary software and database

Different software (and associated databases) for the collection and/or processing of food consumption data were used in the 22 surveys, thus introducing extra complexity for the comparability of dietary information across surveys. Only three surveys (Belgium, Germany and the Netherlands) used the EPIC SOFT program (Slimani et al., 1999), thus making these three surveys more comparable.

489 Cautious interpretation of consumption data is advisable in almost half of the surveys which
490 did not use yield factors to disaggregate composite foods from cooked into their raw
491 ingredients. Hence, the cooked foods are considered as consumed as such and consumption
492 estimates of these particular food groups might differ from those in countries where yield
493 factors were used to disaggregate the recipes into the ingredients as purchased.
494 Overestimation of exposure might result from these five surveys in certain foods such as
495 pasta or rice (the cooked weight of one portion is far higher than its raw weight) whereas
496 underestimation may result in other foods such as meat or fish (their weight decrease when
497 cooked since they lose water).

498
499 In Sweden recipes were not disaggregated into ingredients. As a result, an underestimation of
500 the foods used regularly as ingredients in recipes e.g. cheese or tomato can be expected in the
501 Swedish data. For some exposure assessments within the focus of EFSA, additional
502 information to the food name might be needed. For example, the concentration of heat-
503 generated food toxicants, food additives and substances migrating from packaging materials
504 in foods can differ by household processing method, by brand names, or kind of packaging,
505 respectively. It was requested that national data providers provide available information on
506 brand, household processing and packaging, additionally to the food name information. The
507 available information on brand was very limited at country level. Concerning the information
508 on household processing, in about two third of surveys different information on household
509 processing for varying percentages of food records was collected. Information on baking,
510 frying and grilling of food was available in the vast majority of the surveys where household
511 processing information was collected. Information on packaging material was only available
512 in Belgium (24% of food records), Germany (5%) and Spain II (1%) and its use for exposure
513 assessment purpose is likely to be very limited.

514
515 Available brand and household processing information might differ throughout the surveys
516 since different survey objectives might result in the collection of different information on the
517 foods consumed. Similar to the brand names, information on household processing is
518 sometimes only available as a part of the food description, instead of than as a separate
519 variable, and this cannot easily be extracted. It should be noted that for the brand names, and
520 partly also for the packaging information, the level of detail related to all food items might
521 not be sufficient for a complete exposure assessment. If brand level information is available,
522 it will further need to be linked to chemical occurrence data specific for that branded food

(e.g. food additives). The use of this information for exposure assessment must be decided case by case, based on the availability of information for the food categories of interest. Information on brand name, food packaging and household processing is available for some of the countries that have used a 24 hour recall as well as for some countries that have used food records. This confirmed that this information can be collected by both protocols. However, the three surveys using the 24 hour dietary recall by means of EPIC SOFT program provided a higher number of different descriptors of household processing information. In summary, the use of brand information is limited to certain foods and only available in a restricted number of surveys, information on household processing might be useful in a larger number of countries whereas the available packaging information appears to be of poor use for exposure assessment. In this respect, the EGFC (EFSA, 2009) recommended the collection of information on brand name, physical characteristics of the packaging and household cooking procedures. However, other data sources, like market share data from marketing research studies, could be linked to the food consumption data in order to fill the information gap for exposure assessments.

Under-reporting

Under-reporting was assessed in a number, but not all, surveys by comparison of individual energy intake estimates with Goldberg cut-off points. In the evaluated surveys a varying proportion of under-reporters was identified, i.e. between 12% to 37%. It was assessed that the Goldberg's cut-offs have moderately low sensitivity in identifying under-reporters (Black, 2000b), thus suggesting that, although most study participants identified as extreme under-reporters are likely to have truly underestimated energy intake, a proportion of study-subjects identified as normal-reporters are likely to be under-reporters. One effective way to improve the identification of under-reporters is to use complementary information on individuals' physical activity levels. This strategy was followed in France, where the cut-off points were corrected on the basis of physical activity levels (Black, 2000b). However, this approach finds a natural limitation in the fact that the assessments of physical activity are also prone to measurement errors (Ferrari et al., 2007).

In the guidance document of the EGFC (EFSA, 2009) it was suggested that individual measurements should not be excluded on the basis of estimated under- or over-reporting assessed with short-term dietary assessment methods. Rather, dietary measurements considered unreliable by the interviewers should be removed from the database.

1
2
3 557 The inclusion in the database of surveys with high frequency of under-reporters may lead to
4
5 558 an underestimation of mean dietary exposure and of the percentage of consumers of some
6
7 559 particular foods, e.g. foods with high fat or sugar content, whose estimation is more prone to
8
9 560 under-reporting (Becker, 1999). On the other hand, under-reporting is likely to have less
10
11 561 impact on the assessment of high percentiles of consumption.
12
13 562
14 563 An extensive review from Gorber et al. (2007), evaluating the relationship between self-
15
16 564 reported and directly measured height and weight, showed trends of under-reporting for
17
18 565 weight (ranging from -0.1 kg to -6.5 kg in women and from -0.1 kg to -3.2 kg in men) and
19
20 566 over-reporting for height (ranging from +0.6 cm to +7.5 cm. A slight over-estimation of
21
22 567 exposure expressed per kg body weight is therefore expected when using data from the 13
23
24 568 dietary surveys that collected the weight information through self reporting.
25
26 569
27 570 Uncertainty analysis
28
29 571 The uncertainty related to exposure estimates based on the Comprehensive Database is
30
31 572 presented and discussed for a number of methodological parameters characterising each
32
33 573 dietary survey. It has been highlighted that different survey methodologies have a potential to
34
35 574 induce uncertainties on exposure, especially for the comparison of results between countries.
36
37 575 Although in some circumstances the effect and direction of potential bias in exposure
38
39 576 estimates can be, to a certain extent, anticipated, this is not consistently possible. For
40
41 577 example, it is recognised that estimating high percentiles of habitual consumption, when only
42
43 578 one record or recall day is available per subject, leads to overestimation of intake. In dietary
44
45 579 surveys the degree of such overestimation is often an unknown quantity in the absence of
46
47 580 accurate information on true variability of a given food. Similarly, the effect of a non-random
48
49 581 selection of survey participants might be difficult to predict in magnitude and direction. The
50
51 582 effects of an uneven sampling over days of the week can be, for example, particularly
52
53 583 relevant for the consumption of alcoholic drinks and not for bread, whereas the reported
54
55 584 consumption of soft drinks and certain types of fruits and vegetables can be significantly
56
57 585 affected if the survey design does not capture consumption patterns evenly across the four
58
59 586 seasons. With these examples it is intended to stresses the fact that uncertainty is likely to be
60
587 specific (i) to food or food groups, (ii) to the characteristic of populations in terms of dietary
588 habits, and possibly (iii) to subgroups within a population.
589 Existing guidance documents on uncertainty refer to the uncertainty in exposure assessment
590 (EFSA, 2006 and WHO, 2008) but not to uncertainty of data inputs. Because the uncertainty

of a database can be evaluated only by considering the objective of the assessment, it is therefore deemed extremely difficult to quantitatively anticipate an overall level of uncertainty for each dietary survey. The description of the surveys given in the present paper will enable the risk assessor to perform a qualitative uncertainty analysis and, to a limited extend, also a quantitative analysis when using the Comprehensive Database to assess exposure. The qualitative evaluation of uncertainty represents a valuable option by considering, if possible, the effect of a given parameter on the estimation of the consumption of specific food or food groups. This qualitative approach is in accordance with an EFSA guidance related to uncertainty in dietary exposure assessment (EFSA, 2006). Nevertheless it is recognized that in some circumstances qualitative evaluations can be complemented by methods for quantitative uncertainty analysis of inputs to models for exposure, such as intervals and probability bounds, fuzzy methods, probabilistic methods and sensitivity analysis, as recently summarised in a WHO (2008) monograph. These methods, requiring careful consideration, could be used to evaluate the uncertainty related to specific methodological parameters characterising the Comprehensive Database.

Conclusions

The food consumption data gathered at EFSA to be part of the Comprehensive European food consumption database offers a unique resource for risk assessment activities routinely carried out by EFSA. However, dietary data collected at national level cannot be directly compared due to the different study designs, methodologies and protocols adopted in the different MSs. In particular, differences existed with respect to a number of parameters affecting the level of detail and the accuracy of the collected data, such as: the dietary assessment method used, the description and codification of the food consumed, the number of days per subject, the sampling design, the quantification of portion sizes, the software applications used and non-dietary information collected. Furthermore, in three countries, data provided to EFSA came from national dietary surveys carried out more than ten years ago.

The collection of accurate food consumption data at a European level following a harmonized methodology and protocols is a primary long term objective for EFSA, and it has been recognised as a top priority for collaboration with the EU MSs.

1
2
3 624 **ACKNOWLEDGMENTS:**
4

5 625 Authors are extremely grateful to Verena Nowak and Melanie Fröhler (Austria), Stefanie
6 626 Vandevijvere (Belgium), Stefka Petrova (Bulgaria), Marcela Dofkova and Jiří Ruprich
7 627 (Czech Republic), Tue Christensen (Danmark), Liis Kambek (Estonia), Harri Sinkko, Heli
8 628 Reinivuo, and Liisa Valsta (Finland), Lionel Lafay (France), Carolin Krems and Thorsten
9 629 Heuer (Germany), Maria Szeitzne Szabo (Hungary), Christina Tlustos (Ireland), Laura
10 630 D’Addezio and Aida Turrini (Italy), Gatis Ozolins (Latvia), Evelien de Boer (Netherlands),
11 631 Włodzimierz Sekuła and Maciej Oltarzewski (Poland), Anna Turzova (Slovak Republic),
12 632 Matej Gregorič (Slovenia), Ana M^a López-Sobaler and Rosa Ortega Anta (Spain), Helene
13 633 Enghardt Barbieri (Sweden) and Robin Clifford (United Kingdom) for the information
14 634 provided.
15
16
17
18
19
20
21
22
23
24
25

26 636 **References**
27
28

29 637 Agence française de sécurité sanitaire des aliments (AFSSA). 2009. Report of the
30 638 2006/2007 Individual and National Study on Food Consumption 2 (INCA 2).Synthèse de
31 639 l’étude individuelle nationale des consommations alimentaires 2 (INCA 2) 2006-2007. 1-44.
32
33 640
34
35 641 Becker W, Lennernäs M, Gustafsson IB, Haraldsdottir J, Nydahl M, Vessby B, Ytterfors A.
36 642 1998. Precoded food records compared with weighted food record for measuring dietary
37 643 habits in a population of Swedish adults. Scand J Nutr. 42 : 145-149.
38
39 644
40
41 645 Becker W, Pearson M. 2002. Riksmaten 1997-98. Kostvanor och näringsintag I Sverige.
42 646 Metod-och resultatanalys.(Dietary habits and nutrient intake in Sweden 1997-98)
43 647 Livsmedelsverket (National FoodAdministration). 1-201. Available from:
44 648 <http://www.slv.se/upload/dokument/rapporter/kostundersokningar/riksmat.pdf>.
45
46 649
47
48 650 Becker W, Foley S, Shelley E, Gibney M. 1999. Energy under-reporting in Swedish and Irish
49 651 dietary surveys: Implications for food-based dietary guidelines. Br J Nutr. 81 (2): 127-131.
50
51 652
52
53 653 Biró G, Hulshof KF, Ovesen L, Amorim Cruz JA. 2002. EFCOSUM Group.Selection of
54 654 methodology to assess food intake. Eur J Clin Nutr.56 (2):25-32.
55
56
57
58
59
60

- 655 Black AE a. 2000. Critical evaluation of energy intake using the Goldberg cut-off for energy
656 intake: basal metabolic rate. A practical guide to its calculation, use and limitations. *Int J*
657 *Obes.* 24: 1119-1130.
- 658
- 659 Black AE b. 2000. The sensitivity and specificity of the Goldberg cut-off for EI: BMR for
660 identifying diet reports of poor validity. *Eur J Clin Nutr.* 54: 395-404.
- 661
- 662 Bogle M, Stuff J, Davis L, Forrester I, Strickland E, Casey PH, Ryan D, Champagne C,
663 McGee B, Mellad K, Neal E, Zaghoul S, Yadrick K, Horton J. 2001. Validity of a telephone-
664 administered 24-hour dietary recall in telephone and non-telephone households in the rural
665 Lower Mississippi Delta region. *J Am Diet Assoc.* 101(2):216-22.
- 666
- 667 Casey PH, Goolsby SL, Lensing SY, Perloff BP, Bogle ML. 1999. The use of telephone
668 interview methodology to obtain 24-hour dietary recalls. *J Am Diet Assoc.* 99(11): 1406-11.
- 669 Cochran W. 1977. *Sampling techniques* (3rd edition). Wiley, New York.
- 670
- 671 Day NE, Wong MY, Bingham S, Khaw KT, Luben R, Michels KB, Welch A, Wareham NJ.
672 2004. Correlated measurement error—implications for nutritional epidemiology. *Int J*
673 *Epidemiol.* 33:1373–81.
- 674
- 675 De Henauw S, Brants HA, Becker W. 2002. Operationalization of food consumption surveys
676 in Europe: recommendations from the European Food Consumption Survey Methods
677 (EFCOSUM) Project. *Eur J Clin Nutr.* 56 (2): 75-88.
- 678
- 679 De Vriese S, Huybrecht I, Moreau M, De Henauw S, De Backer G, Kornlitzer M, Leveque A,
680 Van Oyen H. 2005. The Belgian food consumption survey: aim, design and methods. *Arch*
681 *Public Health.* 63: 1-16.
- 682
- 683 Derr JA, Mitchell DC, Brannon D, Smiciklas-Wright H, Dixon LB, Shannon BM. 1992. Time
684 and cost analysis of a computer-assisted telephone interview system to collect dietary recalls.
685 *Am J Epidemiol.* 136(11): 1386-92.
- 686

- 687 Dodd K, Guenther P, Freedman LS, Subar AF, Kipnis V, Midthune D, Tooze JA, Krebs-Smith
688 SM. 2006. Statistical methods for estimating usual intake of nutrients and foods: a review of
689 the theory. *J Am Diet Assoc.* 106:1640-1650.
- 690
- 691 Dubuisson C, Lioret S, Touvier M, Dufour A, Calamassi-Tran G, Volatier JL, Lafay L. 2010.
692 Trends in food and nutritional intakes of French adults from 1999 to 2007: results from the
693 INCA surveys. *Brit J Nutr.* 103: 1035-1048.
- 694
- 695 European Food Safety Authority (EFSA). 2006. Guidance of the Scientific Committee on a
696 request from EFSA related to uncertainties in dietary exposure assessment (adopted on 14
697 December 2006). *EFSA Journal* 438: 1-54. Available from:
698 http://www.efsa.europa.eu/EFSA/Scientific_Opinion/sc_op_uncertainty%20exp_en.5.pdf
- 699
- 700 European Food Safety Authority (EFSA). 2009. General principles for the collection of
701 national food consumption data in the view of a pan-European dietary survey. *EFSA Journal.*
702 27(12):1435-1486. Available from: http://www.rivm.nl/vcp/Images/1435_tcm68-68599.pdf
- 703
- 704 European Food Safety Authority (EFSA). 2010. Technical report of EFSA on the Evaluation
705 of the FoodEx, the food classification system applied in the development of the EFSA
706 Comprehensive European Food Consumption Database. In press.
- 707
- 708 Elmadfa I, Freisling H, Nowak V, Hofstätter D. 2009. Austrian Nutrition Report 2008
709 (Österreichischer Ernährungsbericht 2008). 1. Edition. Vienna.
- 710
- 711 Ferrari P, Friedenreich C, Matthews CE. The role of measurement error in estimating levels of
712 physical activity. *Am J Epidemiol.* 2007 Oct 1;166(7):832-40.
- 713
- 714 Foster E, Adamson AJ, Anderson AS, Barton KL, Wrieden WL. 2009. Estimation of portion
715 size in children's dietary assessment: lessons learnt. *Eur J Clin Nutr.* 63: 45-49.
- 716
- 717 Gabrijelčič Blenkuš M, Gregorič M, Tivadar B, Koch V, Kostanjevec S, Turk Fajdiga V, Žalar
718 A, Lavtar D, Kuhar D, Rozman U. 2009. Prehrambene navade odraslih prebivalcev Slovenije
719 z vidika varovanja zdravja. Pedagoška fakulteta, Inštitut za varovanje zdravja RS. Ljubljana.

- 720 Gersovitz M, Madden JP, Smiciklas-Wright H. 1978. Validity of the 24-dietary recall and
721 seven day record for group comparisons. *J Am Diet Assoc.* 73:48-55.
722
- 723 Goldberg GR, Black AE, Jebb SA, Cole TJ, Murgatroyd PR, Coward WA, Prentice AM. 1991.
724 Critical evaluation of energy intake data using fundamental principles of energy physiology:
725 Derivation of cut-off limits to identify under-recording. *Eur J Clin Nutr.* 45: 569-81.
726
- 727 Gorber CS, Tremblay M, Moher D, Gorber B. 2007. A comparison of direct measures for
728 assessing height, weight and body mass index: a systematic review. *The international*
729 *association for the study of obesity. Obes Rev.* 8: 307-326.
730
- 731 Groth M, Fagt S. 1997. Teknisk rapport 1 (Technical report 1). Copenhagen: National Food
732 Agency (in Danish).
733
- 734 Harrington KE, Robson PJ, Kiely M, Livingstone MBE, Lambe J, Gibney MJ. 2001. The
735 north/south Ireland food consumption survey: survey design and methodology. *Pub Health*
736 *Nutr.* 4: 1037-1042.
737
- 738 Henderson L, Gregory J, Swan G. 2002. National Diet and Nutrition Survey: adults aged 19 to
739 64 years. Volume 1: Types and quantities of foods consumed. TSO. London: 2002.
740 Available from:
741 www.food.gov.uk/science/dietarysurveys/ndnsdocuments/ndnspreviousurveyreports
742
- 743 Kiely M, Flynn A, Harrington KE, Robson PJ, Cran G. 2001. Sampling description and
744 procedures used to conduct the North/South Ireland food consumption survey. *Pub Health*
745 *Nutr.* 4:1029-1035.
746
- 747 Kipnis V, Subar AF, Midthune D, Freedman LS, Ballard-Barbash R, Troiano RP, Bingham S,
748 Schoeller DA, Schatzkin A, Carroll RJ. 2003. Structure of dietary measurement error: results
749 of the OPEN biomarker study. *Am J Epidemiol.* 158:14-21.
750
- 751 Kipnis V, Midthune D, Buckman DW, Dodd KW, Guenther PM, Krebs-Smith SM, Subar AF,
752 Tooze JA. 2009. Modelling data with excess zeros and measurement error: application to

- 753 evaluating relationships between episodically consumed foods and health outcomes. *Biometr.*
754 65: 1003-1010.
755
- 756 Krems C, Bauch A, Götz A, Heuer T, Hild A, Möseneder J, Brombach C. 2006. Methoden der
757 Nationalen Verzehrstudie II. *Ernährungsumschau*. 53 (2): 44-50.
758
- 759 Kroes R, Muller D, Lambe J, Lowik MR, van Klaveren J, Kleiner J, Massey R, Mayer S,
760 Urieta I, Verger P, Visconti A. 2002. Assessment of intake from the diet. *Food Chem Toxicol.*
761 40 (2-3): 327-85.
762
- 763 Lafay L, Mennen L, Six M, Calamassi-Tran G, Hercberg S, Volatier JL. 2002. Etude de
764 validation d'un carnet de consommation alimentaire de 7 jours pour l'enquête INCA 2 -
765 ENNS. In: INSEE, editor. *Journées de Méthodologie Statistique*.
766
- 767 Lambe J, Kearney J, Leclercq C, Zunft HFJ, De Henauw S, Lamberg-Allardt CJE, Dunne A,
768 Gibney MJ. 2000. The influence of survey duration on estimates of food intakes and its
769 relevance for public health nutrition and food safety issues. *Eur J Clin Nutr.* 54: 166-173.
770
- 771 Le Moullec N, Deheeger M, Preziosi P, Monteiro P, Valeix P, Rolland-Cachera MF, Potiers de
772 Courcy G, Christides JP, Cherouvrier F, Galan P, Hercberg S. 1996. Validation du manuel-
773 photos utilisé pour l'enquête alimentaire de l'étude SU.VI.MAX. *Cah Nutr Diét.* 31 :158-163.
774
- 775 Leclercq C, Arcella D, Piccinelli R, Sette S, Le Donne C, Turrini A. 2009. The Italian national
776 food consumption survey INRAN-SCAI 2005-06: main results in terms of food consumption.
777 *Publ Health Nutr.* 12(12): 2504 –2532.
778
- 779 Lioret S, Dubuisson C, Dufour A, Touvier M, Calamassi-Tran G, Maire B, Volatier JL, Lafay
780 L. 2010. Trends in food intake in French children from 1999 to 2007: results from the INCA
781 (Etude Individuelle Nationale des Consommations Alimentaires) dietary surveys. *Brit J Nutr.*
782 103:585-601.
783
- 784 Lyhne N, Christensen T, Velsing Groth, Fagt S, Biloft-Jensen A, Hartkopp H, Hinsch HJ,
785 Matthiesen J, Moller A, Saxholt E, Trolle E. 2005. *Danskernes kostvaner 2000-2002.*
786 *Hovedresultater. Danmarks Fodevareforskning.* 1-168.

- 787 Lyu LC, Hankin JH, Liu LQ, Wilkens LR, Lee JH, Goodman MT, Kolonel LN. 1998.
788 Telephone vs face-to-face interviews for quantitative food frequency assessment. J Am Diet
789 Assoc. 98(1): 44-8.
790
- 791 Max Rubner-Institut (MRI). 2008. Ergebnisbericht der Nationalen Verzehrsstudie II - Teil 1
792 (in German: Report of the National Nutrition Survey II - Part 1). Karlsruhe (Germany), page
793 144. Available from: [http://www.was-esse-](http://www.was-esse-ich.de/uploads/media/NVS_II_Abschlussbericht_Teil_1.pdf)
794 [ich.de/uploads/media/NVS_II_Abschlussbericht_Teil_1.pdf](http://www.was-esse-ich.de/uploads/media/NVS_II_Abschlussbericht_Teil_1.pdf)
795
- 796 Moreno LA, Kersting M, De Henauw S, González-Gross M, Sichert-Hellert W, Matthys C,
797 Mesena MI, Ross N. 2005. How to measure dietary intake and food habits in adolescents: the
798 European perspective. Int J Obes. 29: 66-77.
799
- 800 Nusser SM, Fuller WA, Guenther PM. 1995. Estimating usual dietary intake distributions:
801 adjusting for measurement error and nonnormality in 24-hour food intake data. Dietary
802 assessment research series report 76. Staff report 95-SR 80.
803
- 804 Ocké MC, Hulshof KFAM, Van Rossum CTM. 2005. The Dutch national food consumption
805 survey 2003. Methodological issues. Arch Public Health. 63: 227-241.
806
- 807 Ortega RM, López-Sobaler AM, Ballesteros-Arribas JM, Pérez-Farinós N, Rodríguez-
808 Rodríguez E, Aparicio A, Perea JM, Andrés P. 2010. Estimation of salt intake by 24-hour
809 urinary sodium excretion in a representative sample of Spanish adults. Br J Nutr. In press.
810
- 811 Ovaskainen ML, Paturi M, Reinivuo H, Hannila ML, Sinkko H, Lehtisalo J, Pynnönen-Polari,
812 Männistö S. 2008. Accuracy in the estimation of food servings against the portions in food
813 photographs. Eur J Clin Nutr. 62: 674-681.
814
- 815 Paturi M, Tapanainen H, Reinivuo H. 2008. The National FINDIET 2007 Survey. (In Finnish,
816 summary, figures and tables in English) No. B23. Helsinki: National Public Health Institute.
817 Available from :
818 http://www.ktl.fi/attachments/suomi/julkaisut/julkaisusarja_b/2008/2008b23.pdf.
819

- 820 Petrova S, Angelova K. 2006. Food-Based Dietary Guidelines for Bulgarian adults – scientific
821 background for development and formulation. *Advances in Bulgarian Science* . 4: 19 – 33.
822
- 823 Petrova S, Ovcharova D, Rangelova L, Duleva V, Angelova K, Kalinov K, Dimitrov P,
824 Bojilova D, Baikova D, Vatrlova K, Popivanova A, Marinova M, Antonova Z, Duneva Z.
825 2009. National survey on nutrition of infants and children under 5 years and family child
826 rearing. A Report for UNICEF Bulgaria. NCPHP, pp 1 – 361 (In Bulgarian). Available from:
827 <http://www.ncphp.government.bg>
828
- 829 Pietinen P, Paturi M, Reinivuo H, Tapanainen H, Valsta LM. 2010. FINDIET 2007. Survey:
830 energy and nutrient intakes. *Publ Health Nutr*. 13(6A):920-4.
831
- 832 Pomerleau J, McKee M, Robertson A, Vaask S, Pudule I, Grinberga D, Kadziauskiene K,
833 Abaravicius A, Bartkeviciute R. 1999. Nutrition and lifestyle in the Baltic Republics.
834 European centre on health of societies in transition and world health organization regional
835 office for Europe, 1-61.
836
- 837 Reinivuo H, Hirvonen T, Ovaskainen M-L, Korhonen T, Valsta LM. 2010. Dietary survey
838 methodology of FINDIET 2007 with a risk assessment perspective. *Publ Health*
839 *Nutr*.13(6A):915-919.
840
- 841 Requejo AM, Ortega AM, Lopez Sobaler AM, Navia B, Andres P, Jodral M, Quintas E,
842 Redondo R, Menendez L, Perea JM, Aparicio A, Lozano MC, Bermejo L, Mena C, Faci M,
843 Lolo JM, Rodriguez N, Cocho M, Diez C, Alvarez C. 2002. Estudio sobre dietas y habitos
844 alimentarios en la poblacion Española. Consejo de Seguridad Nuclear. 1-303.
845
- 846 Riboli E, Kaaks R. 1997. The EPIC Project: rationale and study design. *European Prospective*
847 *Investigation into Cancer and Nutrition*. *Int J Epidemiol*. 26(1):6-14.
848
- 849 Rodler I, Biró L, Greiner E, Zajkás G, Szórád I, Varga A, Domonkos A, Agoston H, Balazs A,
850 Mozsary E, Vitrai J, Hermann D, Boros J, Nemeth R, Zsuzsanna K. 2005. Taplalkozasi
851 vizsgalat Magyarországon, 2003-2004. (Dietary survey in Hungary, 2003-2004) *Orvosi*
852 *Hetilap*, 146. Evfolyam, 34, 1781-1789.
853

- 854 Ruprich J, Dofkova M, Rehurkova I, Slamenikova E, Resova D. 2006. Individual food
855 consumption - the national study SISP04. CHFCH NIPH in Prague. Available from:
856 <http://www.chpr.szu.cz/spotrebapotravin.htm>
- 857 Šantare D, Ozoliņš G, Joffe R. 2008. Latvijas iedzīvotāju pārtikas patēriņa pētījums: mērķi,
858 norise, metodes. LU Raksti, Jelgava, iesniegta publicēšanai.
- 859 Sekula W, Nelson M, Figurska K, Oltarzewski M, Weisell R, Szponar L 2004. Comparison
860 between household budget survey and 24 hour recall data in a nationally representative
861 sample of Polish households. *Publ Health Nutr.* 8:430-439.
- 862 Sibbaldi B, Addington-Hall J, Brenneman D, Freeling P. 1994. Telephone versus postal
863 surveys of general practitioners: methodological considerations. *Br J Gen Pract.* 44: 297-300.
- 864 Singer E, Kulka RA. 2002. Paying Respondents for Survey Participation. In *Studies of Welfare*
865 *Populations: Data Collection and Research Issues*. Panel on Data and Methods for Measuring
866 the Effects of Changes in Social Welfare Programs, edited by Michele Ver Ploeg, Robert A.
867 Moffitt, and Constance F. Citro. Committee on National Statistics, Division of Behavioral
868 and Social Sciences and Education. Washington, DC: National Academy Press. 105-128.
- 869 Slimani N, Deharveng G, Charrondiere RU. 1999. Structure of the standardized computerized
870 24- hour diet recall interview used as reference method in the 22 centres participating in the
871 EPIC project. *Comput Methods Programs Biomed.* 8: 251-256.
- 872 Szponar L, Sekuła W, Nelson M, Weisell RC. 2001. The Household Food Consumption and
873 Anthropometric Survey in Poland. *Publ Health Nutr.* 4 (5B): 1183 – 1186.
- 874 Szponar L, Sekuła W, Rychlik E, Oltarzewski M, Figurska K. 2003. Badania indywidualnego
875 spożycia żywności i stanu odżywienia w gospodarstwach domowych (Household Food
876 Consumption and Anthropometric Survey). *NFNI Studies* 101, Warszawa (text in Polish,
877 summary in English).
- 878 Thompson FE, Subar AF. 2001. Dietary assessment methodology. In: *Nutrition in the*
879 *prevention and treatment of disease*. Coulson A, Boushey C (ed). Elsevier, 2nd edition.
- 880 Tooze JA, Midthune D, Dodd KW, Freedman LS, Krebs-Smith SM, Subar AF, Carroll RJ,
881 Kipnis V. 2006. A new statistical method for estimating the usual intake of episodically
882 consumed foods with application to their distribution. *J Am Diet Assoc.* 106(10):1575-87.
- 883 Tran KM, Johnson RK, Soultanakis RP, Matthews DE. 2000. In-person vs telephone-
884 administered multiple-pass 24-hour recalls in women: validation with doubly labelled water.
885 *J Am Diet Assoc.* 100(7):777-83.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

886 WHO (2008) Guidance on Characterizing and Communicating Uncertainty in Exposure
887 Assessment and Guidance on Data Quality in Chemical Exposure Assessment have been
888 published as a two-part publication (Harmonization Project Document No 6).
889 Whybrow S, Horgan G, Stubbs RJ. 2008. Low energy reporting and duration of recording
890 period. Eur J Clin Nutr. 62: 1148-1150.
891 Willet W. 1998. Nutritional Epidemiology. United Kingdom. Oxford University Press.
892 Wolper C, Heshka S, Heymsfield SB. 1995. Measuring food intake: An overview. Thousand
893 Oaks: Sage Publishing.
894 Wrieden WL, Momen NC. 2009. Workshop 3: Novel approaches for estimating portion sizes.
895 Eur J Clin Nutr. 63 (1): 80-81.

Table 1 – Basic information on the dietary surveys included in the “Comprehensive European Food Consumption Database”

Country	Name of the dietary survey (Acronym)	Institution providing the data	Reference publication
Austria	Austrian Study On Nutritional Status (ASNS)	Institute of Nutritional Sciences - University of Vienna	Elmadfa <i>et al.</i> , 2009.
Belgium	Diet National 2004	Scientific Institute of Public Health	De Vriese <i>et al.</i> , 2005.
Bulgaria I	National Survey Of Food Intake And Nutritional Status	National Centre of Public Health Protection	Petrova & Angelova, 2006.
Bulgaria II	NUTRICHILD	National Centre of Public Health Protection	Petrova <i>et al.</i> , 2009.
Czech Republic	SISP04	National Institute of Public Health	Ruprich <i>et al.</i> , 2006..
Denmark	Danish National Survey of Dietary Habits and Physical Activity	National Food Institute, Technical University of Denmark	Lyhne <i>et al.</i> 2005.
Estonia	NDS 1997	National Institute for Health Development	Pomerleau <i>et al.</i> , 1999.
Finland	FINDIET 2007	National Public Health Institute - Nutrition Unit [§]	Paturi <i>et al.</i> , 2008. Reinivuo <i>et al.</i> , 2010. Pietinen <i>et al.</i> , 2010.
France	INCA2	French Food Safety Authority (AFSSA) ^{§§}	AFSSA, 2009, Lioret <i>et al.</i> 2010. Dubuisson <i>et al.</i> 2010.
Germany	German National Nutrition Survey II (NVS II)	Max Rubner-Institut (Bundesforschungsinstitut für Ernährung und Lebensmittel)	MRI, 2008; Krems <i>et al.</i> , 2006.
Hungary	National Repr Surv	Hungarian Food Safety Office	Rodler <i>et al.</i> , 2005.
Ireland	NSIFCS	Food Safety Authority of Ireland	Kiely <i>et al.</i> , 2001. Harrington <i>et al.</i> , 2001.
Italy	INRAN-SCAI 2005–06	National Research Institute for Food and Nutrition (INRAN)	Leclercq <i>et al.</i> , 2009.
Latvia	EFSA_TEST	Food Centre Food and Veterinary Service of Latvia	Šantare <i>et al.</i> , 2008.
Netherlands	VCP2003	National institute of public health and the environment, TNO Quality of Life	Ocké <i>et al.</i> , 2005.
Poland	IZZ-FAO-2000	National Food and Nutrition Institute	Sekula <i>et al.</i> , 2004. Szponar <i>et al.</i> , 2001 and 2003.
Slovakia	SK MON 2008	Food Research Institute	Not available.
Slovenia	CRP-2008	National Institute of Public Health of Slovenia	Gabrijelčič Blenkuš <i>et al.</i> 2009.
Spain I	AESAN -FIAB	Universidad Complutense de Madrid	Requejo <i>et al.</i> , 2002.
Spain II	AESAN	Universidad Complutense de Madrid	Ortega <i>et al.</i> , 2010.
Sweden	RIKSMATEN 1997-98	Swedish National Food Administration	Becker and Pearson, 2002.
United Kingdom	National Diet & Nutrition Survey (NDNS)	Food Standards Agency (FSA)	.Henderson <i>et al.</i> 2002.

[§] currently National Institute for Health and Welfare, ^{§§} currently French agency for food, environmental and occupational health safety

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 2 – Information on the dietary method used within the dietary surveys

Country	Method	Number of replicates	Days between non consecutive replicates* Mean (5 th – 95 th percentile)	Additional food frequency (FFQ) or propensity (FPQ) questionnaire^
Austria	24 h dietary recall	1	Not applicable	No
Belgium	24 h dietary recall	2	23 (12 – 42)	Yes
Bulgaria I	24 h dietary recall	1	Not applicable	Yes
Bulgaria II	24 h dietary recall	2	3 (2 – 5)	Yes
Czech Republic	24 h dietary recall	2	79 (43 – 141)	Yes
Denmark	Food record	7	Consecutive days	No
Estonia	24 h dietary recall	1	Not applicable	Yes
Finland	48 h dietary recall	1	Not applicable	Yes
France	Food record	7	Consecutive days	No
Germany	24 h dietary recall	2	16 (7 – 39)	Diet history
Hungary	Food record	3	2 consecutive days and 1 non consecutive day [§]	No
Ireland	Food record	7	Consecutive days	Yes, only focused on meat
Italy	Food record	3	Consecutive days	No
Latvia	24 h dietary recall	2	68 (27 – 106)	Yes
Netherlands	24 h dietary recall	2	11 (8 – 17)	Yes
Poland	24 h dietary recall	1	Not applicable	No
Slovakia	24 h dietary recall	1	Not applicable	No
Slovenia	24 h dietary recall	1	Not applicable	Yes
Spain I	Food record	3	Consecutive days	Yes
Spain II	24 h dietary recall	2	3 (1 – 16)	Yes
Sweden	Food record	7	Consecutive days	No
United Kingdom	Food record	7	Consecutive days	Yes

* Information extracted from the “Comprehensive European Food Consumption Database”.

^ Information collected by means of FFQ or FPQ was not transmitted to EFSA

§ 2 consecutive weekdays and one weekend day

Table 3 – Administration of the interview

Country	Number of meetings	Method of administration	Place of interview	Interviewer with nutritional background
Austria	0	Post	Not applicable	Not applicable
Belgium	2	Face to face	Home	Yes
Bulgaria I	1	Face to face	Home	Yes
Bulgaria II	2	Face to face	Medical centre	Yes
Czech Republic	2	Face to face	Home	No
Denmark	1	Face to face	Home	No
Estonia	1	Face to face	Home	No
Finland	1	Face to face	Study centre	Yes
France	2	Face to face	Home	Yes
Germany	1 [§]	Telephone	Not applicable	No
Hungary	1	Face to face	Home	Yes
Ireland	4	Face to face	Home or work place	Yes
Italy	3	Face to face	Home	No
Latvia	2	Face to face	Home	No
Netherlands	0	Telephone	Not applicable	Yes
Poland	1	Face to face	Home	No
Slovakia	1	Face to face	Medical centre	Yes
Slovenia	1	Face to face	Home	Yes
Spain I	2	Face to face	Universities, health centre, pharmacies	Yes
Spain II	3	Face to face	Home, universities, health centre, pharmacies	Yes
Sweden	1	Face to face	Home	No
United Kingdom	4	Face to face	Home	No

[§] The meeting was not used for 24h dietary recall administration

Table 4 – Survey period, sampling design and response rate

Country	Survey period	Sampling method and sampling frame	Sample unit	Response rate (%)
Austria	May '05 - February '06	Random from telephone book, Job centres, gynaecologists, university	Individual	48
Belgium	February '04 - February '05	Random from the national population register	Individual	41
Bulgaria I	March '04 - August '04	Random from the national population register	Individual	85
Bulgaria II	April '07 - August '07	Random from the register of general practitioner's practices	Individual	78
Czech Republic	November '03 - November '04	Random from the address register	Household	54
Denmark	June '00 - December '02	Random from the national population register	Individual	53
Estonia	July 1997 - August 1997	Random from the national population register	Individual	67
Finland	January '07 - March '07	Random from the national population register	Individual	62
France	December '05 - April '07	Random from the general population census	Household	60
Germany	November '05 - January '07	Random from the national population register	Individual	42 [§]
Hungary	October '03 - December '03	Random from the general population census	Individual	27
Ireland	October 1997 - October 1999	Random from the electoral list	Individual	63
Italy	October '05 - December '06	Random from the telephone book	Household	33
Latvia	June '08 - November '08	Random from a consumer panel	Individual	56
Netherlands	October '03 - December '03	Random from a consumer panel	Individual	42
Poland	September '00 - November '00	Random from the sample of the household budget survey	Household	96
Slovakia	January '08 - December '08	Random among employees of confectionary and bakery manufactures and canteen	Individual	96
Slovenia	September '07 - April '08	Random from the national population register	Individual	52
Spain I	January 1999 - November '01	Random from the university, health centre, pharmacies	Individual	71
Spain II	January '09 - September '09	Random from the university, health centre, pharmacies	Individual	28
Sweden	January 1997 - January 1998	Random from the national population register	Household	60
United Kingdom	July '00 - June '01	Random from the postcode address file	Household	47

[§] For the total study

Table 5 – Number of subjects according to age group and gender

Country	From 0* to 1 [#] years old [§]		From 1* to 3 [#] years old		From 3* to 6 [#] years old [§]		From 6* to 10 [#] years old [§]		From 10* to 18 [#] years old		From 18* to 64 [#] years old		From 64* to 74 [#]		Older than 74* years	
	Total	% Males	Total	% Males	Total	% Males	Total	% Males	Total	% Males	Total	% Mal	Total	% Male	Total	% Males
Austria	.	.	.	.	.	.	.	.	.	.	2123	37	.	.	.	.
Belgium	.	.	.	.	.	.	.	.	611	48	1356	50	502	49	776	52
Bulgaria I	.	.	.	.	.	.	.	.	162	50	691	48	130	55	221	46
Bulgaria II	861	50	428	51	434	51	.	.	.	.	.	.	.	.	.	.
Czech Republic	.	.	.	.	.	.	.	.	85	51	1666	48	.	.	.	.
Denmark	.	.	.	.	159	52	331	53	479	48	2821	46	283	50	45	51
Estonia	.	.	.	.	.	.	.	.	.	.	1858	45	8	13	.	.
Finland	.	.	.	.	.	.	.	.	.	.	1575	46	463	49	.	.
France	.	.	.	.	163	49	319	50	973	46	2276	41	240	42	108	46
Germany	.	.	.	.	.	.	.	.	1011	49	10419	44	190	47	588	45
Hungary	.	.	.	.	.	.	.	.	.	.	1074	41	188	39	98	35
Ireland	.	.	.	.	.	.	.	.	.	.	952	50	6	50	.	.
Italy	16	56	36	56	67	49	126	48	247	44	2277	46	302	46	252	31
Latvia	.	.	.	.	.	.	190	50	496	50	1382	50	2	0	.	.
Netherlands	.	.	.	.	.	.	.	.	.	.	750	47	.	.	.	.
Poland	.	.	79	59	137	48	272	51	666	50	2527	45	316	40	137	37
Slovakia^{&}	.	.	.	.	.	.	.	.	1	100	2756	50	2	50	.	.
Slovenia^{&}	.	.	.	.	.	.	.	.	.	.	400	49	7	57	.	.
Spain I	.	.	.	.	.	.	.	.	86	55	982	48	.	.	.	.
Spain II	.	.	.	.	.	.	.	.	.	.	418	47	.	.	.	.
Sweden	.	.	.	.	.	.	.	.	7	57	1081	48	112	54	10	50
United Kingdom	.	.	.	.	.	.	.	.	.	.	1724	44	.	.	.	.

[§] Information extracted from the Comprehensive European Food Consumption Database;

*included in the age range;

[#] excluded from the age range

[&] subjects missing because no gender or age information available

Table 6 – Stratification of the sample and gender distribution

Country	Age range* (years)	Number of subjects*	Sample stratification variables				Weighting factors to normalise the sample	Gender**§	
			Gender	Age groups	Geographic al areas	Others		Females (%)	Males (%)
Austria	19 - 64	2123	No	No	No	Employment status	Yes	63	37
Belgium	14 - 105	3245	Yes	Yes	Yes		Yes	50	50
Bulgaria I	16 - 95	1204	Yes	Yes	Yes	Urban vs. rural residence	No	51	49
Bulgaria II	0.1 - 5	1723	Yes	Yes	Yes	Urban vs. rural residence	No	49	51
Czech Republic	16 - 64	1751	Yes	Yes	Yes	Urban vs. rural residence	No	52	48
Denmark	4 - 75	4118	Yes	Yes	No		No	53	47
Estonia	19 - 65	1866	Yes	Yes	No	Urban vs. rural residence	No	55	45
Finland	25 - 74	2038	Yes	Yes	Yes		No	53	47
France	3 - 79	4079	Yes	Yes	Yes	Size of urban area	Yes	56	44
Germany	14 - 80	13926	Yes	Yes	Yes		Yes	55	45
Hungary	18 - 96	1360	Yes	Yes	No		No	60	40
Ireland	20 - 65	958	Yes	Yes	Yes	Education level Urban vs. rural residence, Social status, Employment status	No	50	50
Italy	0.1 - 98	3323	No	No	Yes	Household structure	No	45	55
Latvia	7 - 66	2070	Yes	No	Yes		No	50	50
Netherlands	19 - 30	750	Yes	Yes	Yes	Education level	Yes	53	47
Poland	1 - 96	4134	Yes	Yes	No		No	54	46
Slovakia	17 - 68	2761	Yes	Yes	Yes		No	50	50
Slovenia	18 - 65	410	Yes	Yes	No		No	51	49
Spain I	17 - 60	1068	Yes	Yes	Yes		No	51	49
Spain II	18 - 60	418	Yes	Yes	Yes	Urban vs. rural residence	No	53	47
Sweden	17 - 79	1210	Yes	Yes	Yes		No	52	48
United Kingdom	19 - 64	1724	No	No	Yes	Population density and socio- economic status	Yes	56	44

* Information extracted from the “Comprehensive European Food Consumption Database”.

§ Weighting factors to normalise the sample not used to calculate these percentages.

Deleted: Region, p

Deleted: No

Table 7 – Percentage of record or recall days according to the day of the week and season

Country	% of record or recall days according to the day of the week*			% of record or recall days according to the season*				
	Week days	Week end days	Unclassified	Spring 21.03-21.06	Summer 22.06-22.09	Fall 23.09-22.12	Winter 23.12-20.03	Unclassified
Austria	49	14	37	21	26	25	27	1
Belgium	76	24	0	26	25	27	23	0
Bulgaria I	92	8	0	100	0	0	0	0
Bulgaria II	54	46	0	60	40	0	0	0
Czech Republic	74	26	0	34	23	12	31	0
Denmark	72	28	0	25	26	39	10	0
Estonia	73	27	0	0	100	0	0	0
Finland	67	33	0	9	0	0	91	0
France	71	29	0	20	17	24	39	0
Germany	75	25	0	20	27	40	13	0
Hungary	67 [§]	33 [§]	0	0	0	0	100	0
Ireland	71	29	0	26	28	27	18	0
Italy	78	22	0	26	24	25	25	0
Latvia	72	28	0	0	49	50	0	0
Netherlands	71	29	0	0	0	100	0	0
Poland	77	23	0	0	31	69	0	0
Slovakia	78	5	17	23	19	29	7	23
Slovenia	76	24	0	11	14	56	19	0
Spain I	43	30	27	28	7	25	22	17
Spain II	73	26	0	75	19	0	6	0
Sweden	71	29	0	0	0	0	0	100
United Kingdom	71	29	0	31	24	22	23	0

* Information extracted from the "Comprehensive European Food Consumption Database".

[§] Percentages reported by the national data provider.

Table 8 – Exclusion criteria

Country	Institutionalised subjects excluded	Pregnant and breastfeeding women excluded	Number of breastfeeding women*	Number of pregnant women*	Over sampled population groups
Austria	Yes	No	Not available	Not available	
Belgium	Yes	No	7	9	Subjects from 15 to 18 and older than 75 years
Bulgaria I	Yes	Yes	Not applicable	Not applicable	No
Bulgaria II	Yes	NA	Not applicable	Not applicable	No
Czech Republic	Yes	No	Not available	Not available	No
Denmark	Yes	No	59	50	No
Estonia	No	Yes	Not applicable	Not applicable	No
Finland	No	No	Not available	22	No
France	Yes	No	20	27	Children
Germany	Yes	No	36	52	No
Hungary	No	No	Not available	Not available	No
Ireland	Yes	Yes	Not applicable	3	No
Italy	Yes	No	10	19	No
Latvia	Yes	Yes	Not applicable	Not applicable	No
Netherlands	Yes	Yes	Not applicable	Not applicable	Males with low education level group
Poland	Yes	No	26	23	No
Slovakia	No	No	Not available	Not available	No
Slovenia	Yes	Yes	Not applicable	Not applicable	No
Spain I	Yes	No	0	3	No
Spain II	Yes	No	Not available	0	No
Sweden	Yes	No	16	11	No
United Kingdom	Yes	Yes	Not applicable	Not applicable	No

* Information extracted from the "Comprehensive European Food Consumption Database".

Table 9 – Portion sizes estimation

Country	Portion sizes estimated by				
	Weighing	Picture book	Household measures	Known packaging size	Ruler
Austria	No	No	Yes	No	No
Belgium	No	Yes, based on EPIC-soft	Yes	No	No
Bulgaria I	No	Yes, validated	Yes	Yes	No
Bulgaria II	No	Yes, validated	Yes	Yes	No
Czech Republic	No	Yes, tested in a convenient sample	Yes	No	Yes
Denmark	No	Yes, validated	Yes	No	No
Estonia	No	Yes, not validated	Yes	No	No
Finland	No	Yes, validated (Ovaskainen et al., 2008)	Yes	Yes	Yes
France	No	Yes, validated (Le Moullec et al., 1996)	Yes	Yes	No
Germany	No	Yes, based on EPIC-soft	Yes	No	No
Hungary	No	No	No	No	No
Ireland	Yes	Yes, not validated	Yes	Yes	No
Italy	No	Yes, based on EPIC-soft	Yes	Yes	No
Latvia	No	Yes, not validated	Yes	No	No
Netherlands	No	Yes, based on EPIC-soft	Yes	No	Yes
Poland	No	Yes, tested in a convenient sample	Yes	Yes	No
Slovakia	No	No	No	No	No
Slovenia	No	Yes, not validated	Yes	No	No
Spain I	Yes	No	Yes	Yes	No
Spain II	No	No	Yes	Yes	No
Sweden	No	Yes, validated (Becker et al., 1998)	Yes	No	No
United Kingdom	Yes	No	<u>Yes</u>	Yes	Yes

Deleted: No

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 10 – Dietary software and database used

Country	Dietary software	Database used before providing the data to EFSA:			
		Portion size	Standard recipe	Yield factors	Food composition
Austria	MS Access 2003 based on German nutrient data base BLS II.3	Yes	Yes	Yes, from raw to cooked foods	Yes
Belgium	EPIC SOFT Program	Yes	Yes	Yes, from raw to cooked foods*	No
Bulgaria I	NUTRICALC	Yes	Yes	Yes, from cooked to raw foods*	Yes
Bulgaria II	NUTRICALC	Yes	Yes	Yes, from cooked to raw foods*	Yes
Czech Republic	Paradox for Windows	Yes	Yes	Yes, from cooked to raw foods*	Yes
Denmark	GIES	Yes	Yes	Yes, from cooked to raw foods*	Yes
Estonia	Finnish micro nutrica nutritional analysis program	Yes	Yes	Yes, from cooked to raw foods*§	Yes
Finland	Finessi	Yes	Yes	Yes, from cooked to raw foods*§	Yes
France	MS Access	Yes	Yes	No	Yes
Germany	EPIC SOFT Program	Yes	Yes	Yes, from raw to cooked foods*	Yes
Hungary	NutriCompEtrend	Yes	Yes	Yes, from cooked to raw foods	Yes
Ireland	WISP-DES	Yes	Yes	Yes, from cooked to raw foods*	Yes
Italy	INRAN-DIARIO 3.1	Yes	Yes	Yes, from cooked to raw foods*§	Yes
Latvia	PGAIS	No	Yes	Yes, from raw to cooked foods	Yes
Netherlands	EPIC SOFT Program	Yes	Yes	Yes, from cooked to raw foods*	No
Poland	Dieta FAO	Yes	Yes	Yes, from cooked to raw foods*	Yes
Slovakia	Alimenta 4.3	No	Yes	Yes, from cooked to raw foods§	Yes
Slovenia	Blaise 4.7	No	Yes	Yes, from raw to cooked foods§	Yes
Spain I	DIAL software	Yes	Yes	No	Yes
Spain II	DIAL software	Yes	Yes	No	Yes
Sweden^	MATs version 4.03	No	No	No	Yes
United Kingdom	Intake 2	No	No	No	Yes

* Yield factors included in the recipe database.

§ only for some foods

^ Sweden did not disaggregate their recipes

Table 11 – Additional information on food coding

Country	Specific information collected on:		
	Brand* (% of food records)	Household processing* (% of food records)	Packaging* (% of food records)
Austria	3	1	0
Belgium	19	22	24
Bulgaria I	0	25	0
Bulgaria II	0	57	0
Czech Republic	0	0	0
Denmark	0	32	0
Estonia	0	0	0
Finland	0	75	0
France	29	18	0
Germany	13	14	5
Hungary	0	40	0
Ireland ^o	0	0	0
Italy	27	0	0
Latvia	12	26	0
Netherlands	21	20	0
Poland	0	45	0
Slovakia	0	0	0
Slovenia	1	1	0
Spain I	0	41	0
Spain II	4	18	1
Sweden	0	0	0
United Kingdom	0	0	0

* Information extracted from the "Comprehensive European Food Consumption Database".

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 12 – Method of identification of under-reporters, cut-off values and percentage of under-reporters

Country	Method	Cut off value	Under-reporters excluded	Under-reporters (%)
Austria	Individual level	0.65 (Goldberg et al., 1991) for adults	Yes	12%
Belgium	Individual level	0.965 (Black, 2000a)) for adults	No	NR^
Bulgaria I	Individual level	0.9 (Goldberg et al., 1991) for adults	No	NR^
Bulgaria II	None	-	-	-
Czech Republic	Individual level	0.96 (Black, 2000a) for adults	No	12.8
Denmark	Individual level	1.1 (Goldberg et al., 1991) for adults	No	NR^
Estonia	None	-	-	-
Finland	Individual level	1 (Goldberg et al., 1991) for adults	No	37.3
France	Individual level	1.01-1.27 (Black, 2000b)* for adults	No	26.9
Germany	Individual level	-	No	-
Hungary	Individual level	1.1 (Goldberg et al., 1991) for adults	Yes	11.7
Ireland	Individual level	1.1 (Goldberg et al., 1991) for adults	No	18.0
Italy	Population level	1.55 (Goldberg et al., 1991) for adults	No	-
Latvia	None	-	-	-
Netherlands	Population level	1.53 (Goldberg et al., 1991) for adults	No	-
Poland	None	-	-	-
Slovakia	None	-	-	-
Slovenia	None	-	-	-
Spain I	None	-	-	-
Spain II	None	-	-	-
Sweden	Individual level	1.1 (Goldberg et al., 1991) for adults	No	24.7
United Kingdom	None	-	-	-

* Cut-off values defined by Black (2000a) vary according to age and sex, taking into account the specific intra and inter individual variability of Physical Activity Levels (PAL)

^ NR: Not reported;

Table 13 – Number of subjects according to the different [self-reported](#) eating patterns.

Deleted: identified

Country	Normal diet*	Vegetarian diet*	Diet related to health condition*	Slimming diet*	Vegetarian and slimming diet*	Unspecified*
Austria	2123					0
Belgium	2642	1	331			271
Bulgaria I	863	1	70	116		154
Bulgaria II						1723
Czech Republic	1572	9	86	66		18
Denmark						4118
Estonia						1866
Finland	1377	29	584	26	22	0
France	3167	19	314	181	1	397
Germany	10839	287	2106	141	1	552**
Hungary	1360					0
Ireland	764	9	77	70		38
Italy	3124		80	76		43
Latvia						2070
Netherlands	691	12	8	24		15
Poland						4134
Slovakia	83					2678
Slovenia	410					0
Spain I	1051		10	1		6
Spain II	398		4	16		0
Sweden	1188	18	2			2
United Kingdom	1333	66		314	11	0

* Information extracted from the "Comprehensive European Food Consumption Database"

** In Germany information on further special diets, like halal, was also collected but here these are considered as unspecified.

Table 14 – Non dietary information

Country	Body weight and height	Information available on		
		Physical activity* [§] (% of subjects)	Education level* (% of subjects)	Ethnicity* (% of subjects)
Austria	Estimated	100	100	14
Belgium	Estimated	100	100	0
Bulgaria I	Measured	0	100	100
Bulgaria II	Measured	0	100	100
Czech Republic	Estimated	100	100	0
Denmark	Estimated	0	0	0
Estonia	Estimated	100	100	100
Finland	Measured	0	99	0
France	Measured	74 [^]	100	0
Germany	Measured	0	100	0
Hungary	Estimated	0	0	0
Ireland	Measured	100	100	100
Italy	Estimated	0**	83	0
Latvia	Estimated	0	100	0
Netherlands	Estimated	100	100	100
Poland	Measured	0	100	0
Slovakia	Estimated	3	3	6
Slovenia	Estimated	100	99	0
Spain I	Estimated	80	28	0
Spain II	Measured	100	99	0
Sweden	Estimated	0	100	0
United Kingdom	Measured	96	100	100

* Information extracted from the "Comprehensive European Food Consumption Database"

** Data not transmitted to EFSA

[^] Questionnaire administered only to subjects between 15 and 79 years old

[§] Information on physical activity collected by means of questionnaires within all surveys.

Formatted: Superscript

Formatted: Superscript