USUTU VIRUS IN ITALY, AN EMERGENCE OR A SILENT INFECTION?

Giovanni Savini, Federica Monaco, Calogero Terregino, Annapia Di Gennaro, Luca Bano, Chiara Pinoni, Roberta de Nardi, Paolo Bonilauri, Monica Pecorari, Luigina Di Gialleonardo, et al.

► To cite this version:

Giovanni Savini, Federica Monaco, Calogero Terregino, Annapia Di Gennaro, Luca Bano, et al.. USUTU VIRUS IN ITALY, AN EMERGENCE OR A SILENT INFECTION?. *Veterinary Microbiology*, 2011, 151 (3-4), pp.264. 10.1016/j.vetmic.2011.03.036 . hal-00717083

HAL Id: hal-00717083

<https://hal.science/hal-00717083>

Submitted on 12 Jul 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Title: USUTU VIRUS IN ITALY, AN EMERGENCE OR A SILENT INFECTION?

Authors: Giovanni Savini, Federica Monaco, Calogero Terregino, Annapia Di Gennaro, Luca Bano, Chiara Pinoni, Roberta De Nardi, Paolo Bonilauri, Monica Pecorari, Luigina Di Gialleonardo, Lebara Bonfanti, Andrea Polci, Paolo Calistri, Rossella Lelli

PII: S0378-1135(11)00210-0
DOI: doi:10.1016/j.vetmic.2011.03.036
Reference: VETMIC 5259

To appear in: *VETMIC*

Received date: 12-10-2010
Revised date: 15-3-2011
Accepted date: 31-3-2011

Please cite this article as: Savini, G., Monaco, F., Terregino, C., Di Gennaro, A., Bano, L., Pinoni, C., De Nardi, R., Bonilauri, P., Pecorari, M., Di Gialleonardo, L., Bonfanti, L., Polci, A., Calistri, P., Lelli, R., USUTU VIRUS IN ITALY, AN EMERGENCE OR A SILENT INFECTION?, *Veterinary Microbiology* (2010), doi:10.1016/j.vetmic.2011.03.036

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

USUTU VIRUS IN ITALY, AN EMERGENCE OR A SILENT INFECTION?

Giovanni Savini¹, Federica Monaco¹, Calogero Terregino², Annapia Di Gennaro¹, Luca Bano², Chiara Pinoni¹, Roberta De Nardi², Paolo Bonilauri³, Monica Pecorari⁴, Luigina Di Gialleonardo¹, Lebara Bonfanti², Andrea Polci¹, Paolo Calistri¹, Rossella Lelli¹

¹ Istituto G. Caporale, Teramo, Italy

² Istituto Zooprofilattico Sperimentale delle Venezie, Legnaro (PD), Italy

³ Istituto Zooprofilattico Sperimentale della Lombardia ed Emilia Romagna, Reggio Emilia, Italy

⁴ Department of Diagnostic and Laboratory Services and Legal Medicine, University of Modena and Reggio Emilia, Azienda Ospedaliera Policlinico, Modena, Italy

Corresponding author: Giovanni Savini
Dept. of Virology
National Reference Center for West Nile disease
OIE Reference laboratory for Bluetongue
Istituto G. Caporale Teramo
Via Campo Boario, 64100 - Teramo - Italy
Telephone: +390861 332440
Fax No. +390861 332251
E-mail: g.savini@izs.it

ABSTRACT

A two year study (2008-2009) was carried out to monitor the Usutu virus (USUV) circulation in Italy. Sentinel horses and chickens, wild birds and mosquitoes were sampled and tested for the presence of USUV and USUV antibodies within the WND National Surveillance plan. Seroconversion evidenced in sentinel animals proved that in these two years the virus has circulated in Tuscany, Emilia Romagna, Veneto and Friuli Venezia Giulia regions. In Veneto USUV caused a severe blackbird die-off disease involving at least a thousand birds. Eleven viral strains were detected in organs of 9 blackbirds (52.9%) and two magpies (0.5%) originating from Veneto and Emilia Romagna regions. USUV was also detected in a pool of *C. pipiens* caught in Tuscany. According to the alignment of the NS5 partial sequences, no differences between the Italian USUV strains isolated from Veneto, Friuli and Emilia Romagna regions were observed. The Italian North Eastern strain sequences were identical to those of the strain detected in the brain of a human patient and shared a high similarity with the isolates from Vienna and Budapest. Conversely, there were few differences between the Italian strains which circulated in the North Eastern regions and the USUV strain detected in a pool of *C. pipiens* caught in Tuscany. A high degree of similarity at both nucleotide and amino acid level was also found when the full genome sequence of the Italian North Eastern isolate was compared with that of the strains circulating in Europe. The North Eastern Italian strain sequence exhibited 97% identity to the South African reference strain SAAR-1776. The deduced amino acid sequences of the Italian strain differed by 10 and 11 amino-acids from the Budapest and Vienna strains, respectively, and by 28 from the SAAR-1776 strain. According to this study two strains of USUVs are likely to have circulated in Italy between 2008 and 2009. They have developed strategies of adaptation and evolution to spread into new areas and to become established.

KEY WORDS: Chickens, genome sequencing, horse, Italy, NS5, Usutu virus, wild birds

INTRODUCTION

Usutu virus (USUV) is a single strand RNA virus included in the mosquito-borne cluster of the genus *Flavivirus*, within the Family *Flaviviridae*. Together with human pathogens like Japanese encephalitis virus (JEV), Murray Valley encephalitis virus (MVEV), Saint Louis encephalitis virus (SLEV) and West Nile virus (WNV) (Kuno et al., 1998; Bakonyi et al., 2004), it is included in the Japanese encephalitis virus serogroup (Heinz et al., 2000). As all members of this group, the virus is maintained in the environment through a bird-mosquito life cycle. Even if grouped with important human pathogens, USUV was considered for many years as an exotic virus of minor importance confined to a small area of sub-Saharan Africa and characterised by a very low pathogenicity for humans and animals.

In recent years, coinciding with its occurrence in Europe, the virus appears to have changed its behaviour, and fatalities associated with USUV infection were reported in European wild birds (Weissenböck et al., 2002; Chvala et al., 2007; Bakonyi et al., 2007; Steinmetz et al., 2010) and, in 2009, two cases of human encephalitis have been associated to the USUV infection (Pecorari et al., 2009, Cavrini et al., 2009). All these events have greatly changed the way of looking at these infections. Information on USUV biologic features, host spectrum, and pathogenesis is becoming of high priority. In Italy, before the clinical cases reported in humans, USUV was evidenced in sentinel chickens in the Trentino region in 2005 (Rizzoli et al., 2007) and in the Emilia Romagna region in 2007 (Lelli et al., 2008). Few fatal cases of USUV were also described in owls and blackbirds near Milan in Lombardia between 2006 and 2008 (Manarolla et al., 2010). As main theatre of these events, Italy definitely plays an important role for this infection at both epidemiological and public health level. This paper describes two years of outbreaks and USUV circulation in Italy as reported by the surveillance plan and provides information to better understand the epidemiological as well as the virological characteristic of the USUV Italian isolates.

MATERIALS AND METHODS

With the aim of monitoring flavivirus circulation in Italy, an extensive surveillance plan has been implemented involving horses as well as wild and domestic birds. Insect traps were also placed in some

horse stables and mosquitoes were collected and identified. The surveillance activities included: i) regular serological testing of sentinel-animals located in the proximity of fourteen wetland areas which were selected because of the presence of migratory birds. These areas could represent a risk of virus introduction and spread. The sentinel animals consisted of: horses sampled twice, in spring and in late autumn, and chickens sampled as described by Monaco et al. (2008); ii) the virological examination of dead wild birds and serological monitoring of backyard poultry flocks; in 2009 more emphasis was placed on sampling horses, birds and mosquitoes of areas of Northern Italy in which WNV and USUV have circulated in 2008 (Figure 1).

Wild birds

In 2009, brains, hearts, and spleens of 1045 Italian wild bird carcasses belonging to a least 22 species were examined for the presence of USUV (Table 4). Between 2008 and 2009, 440 serum samples belonging to 7 species were further examined for the presence of USUV antibodies.

Sentinel animals

According to the National surveillance plan implemented for West Nile virus since 2001, 1797 and 2567 chicken samples were examined serologically for USUV in 2008 and 2009, respectively. The samples were from animals located in the proximity of the 14 wetland areas distributed all around Italy (Figure 1). Equally, from the same locations, 643 and 522 serum samples from sentinel horses were tested for the presence of USUV neutralising antibodies in 2008 and 2009, respectively (Figure 1). They were tested at least twice a year, before and after the mosquito season. In total, 1349 and 1275 sentinel horse serum samples were examined in 2008 and 2009, respectively.

Horse sera

Besides the sentinel animals, additional 800 and 4130 serum samples of horses originating from all Italian regions except Abruzzo and Aosta Valley, were examined for the presence of USUV antibodies in 2008 and 2009, respectively.

Mosquito pools

Twenty pools of 50 mosquitoes each were tested for the presence of the USUV. They were from the Tuscany region and comprised 9 pools of *C. pipiens*, 9 of *Aedes albopictus* and 2 of other species.

Laboratory testing and diagnostic protocol

Samples (blood, sera, mosquitoes and tissue) were sent for serological and virological analyses to the OIE and National Reference laboratory for West Nile and Usutu viruses, Istituto G. Caporale Teramo, Italy (ICT).

To detect the presence of USUV and WNV specific antibody response, serum samples were tested by ELISA and virus neutralisation test in microtitre plates (SN) (Lelli et al., 2008). To rule out possible cross-reactions, positive samples (titre $\geq 1:10$) were also tested for TBEV and WNV antibodies by PRNT (OIE, 2008a; Lelli et al., 2008).

To detect USUV in organs, EDTA blood samples and mosquitoes, a specific RT-PCR targeting the NS5 was used (Lelli et al., 2008; Bakonyi et al., 2007). Virus isolation was attempted in the same samples by inoculating VERO E6 cell cultures. Samples negative at the first passage were sub-cultured twice in the same cells (Bernard et al., 2001; Lelli et al., 2008). On blackbird carcasses collected during the outbreak, bacteriological, parasitological and toxicological tests as well as virological tests other than those described were also performed to exclude the presence of other pathogens. In particular, samples of brain, heart, liver, kidney, spleen, lung and intestine were processed for viral isolation attempts in 9-10 day old SPF chicken embryonated eggs (OIE, 2008b), chicken embryo fibroblast (CEF) and chicken embryo liver (CEL). Virus isolation attempts in cells were carried out with homogenized tissue samples suspended in phosphate-buffered saline with antibiotics. After centrifugation ($1,000 \times g$, 10 min), the supernatants were inoculated onto confluent CEF, CEL and Vero cell cultures. Cell cultures were incubated at 37°C for 7 days and were checked daily for the occurrence of cytopathic effects (CPE). Three blind passages were made in absence of cytopathic effect. At the end of the third passage and in case of CPE, the presence of USUV in the supernatants was confirmed or excluded

either by electron microscopy or by Immunofluorescence (IFA) using a monoclonal antibody (produced by ICT, Italy) against USUV.

To confirm the identification of the virus, a second USUV specific RT-PCR targeting the Pre-M region of viral RNA was performed on the positive samples. The amplified products of both the Pre-M and NS5 were sequenced (602 bp of preM and 583 bp of NS5) and aligned with the corresponding sequences deposited in Genbank (gb|AY453411.1; gb|EF206350.1) using ClustalW (table 1). NS5 datasets were analyzed using the PHYLIP program (Phylogeny Interference Program Package version 3.6a) (Felsenstein, 1993). For phylogenetic analysis the appropriate nucleotide substitution model selected was the Kimura two-parameter model (Kimura, 1980) with transition/transversion of 2.0. Statistical support at the internodes was assessed by 100 bootstrap replication.

Total RNA was extracted from samples with the High Pure Viral Nucleic Acid kit (Roche, Germany), eluted in 50µl of Elution buffer according to the manufacturer instruction and used for the RT-PCR reaction performed by the RT-PCR Onestep kit (Qiagen). Concisely, for each sample the mixture contained 21.8 µl of Rnase free water, 10 µl of Q-solution, 2 µl of enzyme mix, 1.6 µl of each primer (50 pmoli/µl), 10 µl of One-step buffer 5X, 2 µl of dNTPs (10mM) and 5 µl of denatured RNA in a total volume of 50 µl. The reverse transcription was performed at 45°C for 30' followed by the activation phase at 95°C for 5'. Forty cycles at 95° for 30", 55°C for 30" and 72°C for 2' were used for the amplification followed by 5' at 72°C for the final elongation.

Full genome sequencing

Total RNA was also extracted from 200µl of cell culture supernatant from the USUV as described below. Viral RNA was reverse transcribed and amplified with a set of 66 primer pairs targeting overlapping sequences of nearly 1000 bp of the viral genome. The primer sequences are available upon request. Amplification and sequencing were repeated twice to avoid introduction of artificial substitutions. Raw sequence data were assembled using Contig Express (Vector NTI suite 9.1, Invitrogen, USA) and the consensus sequence was aligned with the Central European and South African strains (AY453411,

AY453412 and EF206350) to evaluate the variation within the strains. Both, the nucleotidic and the deduced amino acid sequences were compared using Vector NTI suite 9.1 (Invitrogen, USA).

Outbreak data

Clinical signs, mortality, morbidity and macroscopic lesions observed during the outbreaks were reported and fully described.

Serological and virological data obtained in this study were investigated and each proportion was compared by using the beta distribution (Sivia, 1996).

RESULTS

Case history

Between August and October 2009 widespread anomalous death episodes of blackbirds (*Turdus merula*) were reported in at least six different municipalities of the Treviso province (Veneto region, North-Eastern Italy) (Figure 2). It was estimated that the mortality in the different outbreaks concerned almost a thousand of birds altogether. From each outbreak, groups of 5-10 subjects, for a total of 35 blackbirds, have been examined at the Istituto Zooprofilattico Sperimentale delle Venezie and comprised some dead or very ill subjects spontaneously dead few minutes after submission to the laboratory.

At necropsy the main part of the subjects was lean and the birds showed no remarkable gross changes such as hepato-splenomegaly, hepatic congestion, enlarged gallbladder, enlarged and discoloured kidneys and hyperaemic meninges and brain.

Bacteriological and toxicological tests gave negative results. Virological tests on embryonated eggs gave negative results as well. After about 3 days of incubation, focal cell rounding and shrinkage were observed in VERO cell cultures inoculated with brain, heart and kidney homogenates. Within 2 days, the majority of the cells showed CPEs. In detail, most of the inoculated cells lost their adherence to the

surface of the culturing flask and floated in the supernatant. The CPE was more evident in the subsequent two passages. No CPEs or cell degeneration was observed in CEF and CEL cells.

The supernatants of the flasks with CPE when examined by electron microscopy showed the presence of spherical viruses of about 50 nm diameter. To confirm the presence of USUV in the reported outbreaks, tissue culture lysates and/or original organs from birds (no=6) representative of each outbreak municipality were sent to the ICT. At the National Reference Centre USUV was then detected and identified. Parasitological test highlighted the concomitant presence of haemosporidian parasites in 18 of the 35 subjects examined and found positive to USUV.

Surveillance plan

Sentinel animals

In 2008, 9 sentinel chickens seroconverted to the USUV in the period September -October. Seroconversions were observed in Tuscany, 8 in the province of Pisa and 1 in the province of Pistoia. In 2009, 2 sentinel chickens seroconverted in the province of Gorizia (Friuli Venezia Giulia) in the month of October (Figure 2).

In 2008, no seroconversion was reported in horses while in 2009, 22 horses from Veneto, Emilia Romagna, Tuscany and Friuli Venezia Giulia regions seroconverted (Table 2 and Figure 2).

Horses

When tested by SN, some sera reacted to USUV only, some others have titers for both USUV and WNV. No cross reactions between the USUV and WNV were found. A higher ($P>0.05$) prevalence of positive horses were found in 2008 (89.2%) compared to 2009 (7.8%). Similar results were obtained when the data were analysed according to provinces and regions (Tables 3).

Wild birds

Virological investigation

One thousand and forty five tissue samples belonging to more than 22 wild bird species were tested for USUV. Species and origin are listed on Table 4. Presence of USUV was detected in 11 samples, 9 (52.9%) from blackbirds and 2 (0.5%) from magpies of Emilia Romagna and Veneto. Blackbird was the species in which USUV was more frequently ($P>0.05$) detected (Figure 2).

Serological investigation

Of the 440 serum samples tested, USUV neutralising antibodies were detected in 20 rock pigeon, 1 blackbird and 2 magpie sera. The positive animals were from Veneto and Emilia Romagna (table 4).

Characterisation of the isolates

Sequence comparison

Results of the alignment of NS5 are shown in figure 3. No differences were observed among the Italian USUV strains isolated from Veneto, Friuli and Emilia Romagna regions when the NS5 partial sequences were aligned. Sequences were identical to those of the strain detected in the brain of the human patient and shared a high similarity with those of the Vienna, Budapest and PI-09-Cp isolates. The NS5 partial sequence of the Italian strain detected in *C. pipiens* caught in Tuscany was more similar to Budapest and Vienna isolates than to the other Italian isolates.

High degree of similarity at both nucleotide and aminoacid level was also found when the full genome sequence of the Italian isolate (RA-09-Tm) was compared with that of the strains circulating in Europe, (Table 5). The Italian USUV strain sequences exhibited 97% identity to the South African reference strain SAAR-1776. The nucleotide sequences were translated to putative amino acid sequences. The deduced amino acid sequences of the Italian strain differed by 10 and 11 amino-acids from the Budapest and Vienna strains, respectively (Table 6). The Italian USUV showed 99% amino acid identity to the SAAR-1776 strain. The amino acid differences are summarized in table 6.

DISCUSSION

1 In the last decades the emergence of new diseases and/or the recrudescence of old disease into new
 2 areas exhorted the scientific community to propose a new category of pathogens: the so called
 3 emerging and re-emerging pathogens. Among the emerging pathogens, arboviruses, which by
 4 definition are viruses that are transmitted to vertebrates by arthropods (mosquitoes, ticks, sandflies,
 5 midges ...), are nowadays definitely playing an important role. USUV is an arbovirus and, in view of the
 6 recent events, it represents the typical example of an emerging pathogen. It appeared in Austria in
 7 2001 causing widespread mortality among the population of Eurasian blackbirds and some other bird
 8 species (Weissenböck et al., 2002). In the following years presence of USUV was associated to mortality
 9 of birds south east (Hungary) and west (Switzerland) of Austria (Bakonyi et al., 2007; Steinmetz et al.,
 10 2010). In Italy a silent virus circulation was detected since 2007 (Lelli et al., 2008). In the last years
 11 presence of this virus has been associated with fatal cases in birds. More recently the Italian strain was
 12 detected in the brain of immune deficient patients with neuroinvasive infection (Pecorari et al., 2009;
 13 Cavrini et al., 2009). The data of this study prove that, after being evidenced in 2007, USUV has
 14 circulated widely in Italy at least in Tuscany, Veneto, Emilia Romagna and Friuli regions.
 15 Seroconversions of either sentinel horses or chickens in both 2008 and 2009 and the detection of virus in
 16 wild birds, support the hypothesis of an endemic cycle in local bird and mosquito species able to
 17 maintain and disperse USUV in Italy. A rather complex relationship between virus, hosts and vectors is
 18 needed to maintain an arbovirus in the environment. Vector competence for a particular virus and
 19 the contemporary presence of susceptible vertebrate host species with viraemia levels permitting
 20 vector transmission, are key factors for the spread of the virus in a given environment. USUV is thought
 21 to be maintained in nature by a mosquito-bird transmission cycle. In Africa ornithophilic mosquitoes of
 22 the genera *Culex*, *Coquillettidia* and *Mansonia* are thought to be the main vectors (Pfeffer and
 23 Dobler, 2010). In Austria *Culex* spp may play a major role (Pfeffer and Dobler, 2010), although USUV has
 24 not been isolated from mosquitoes so far but it has been only detected in overwintering *Culex pipiens*
 25 pools by real time RT-PCR. In this study USUV was detected in a pool of *C. pipiens* and was isolated
 26 from organs of dead blackbirds suggesting that blackbirds and *C. pipiens* might play an important role
 27 in the Italian endemic cycle. The fact that many horses often had neutralising titers to both WNV and

USUV seems to imply that both viruses might share the same vector/s. It is now well documented that *C. pipiens* represents the main vector of WNV in Italy and its involvement as a vector has also been suggested for USUV (Tamba et al., 2010). *Culex pipiens* is commonly regarded as ornithophilic (Hamers et al., 2008; Marra et al., 2004; Turell et al., 2005; Simpson et al., 2009). In the United States of America it has been observed that it displayed an innate host preference for *Turdus* genus (Simpson et al., 2009). Preferential feeding by a vector on virus-competent hosts may influence transmission dynamics as contact rates are shifted away from other abundant, but less competent hosts (Simpson et al., 2009). In USUV infection, the blackbird (*Turdus merula*), a species of the *Turdus* genus, has been the one most commonly involved. Other studies showed that also certain owl species and house sparrows are susceptible to USUV infection (Manarolla et al., 2009; Steinmetz et al., 2010). A considerable number of different bird species were examined in this study, USUV however was detected in magpies and blackbirds only. As in Central Europe (Chvala et al., 2007; Bakonyi et al., 2007; Steinmetz et al., 2010), the blackbird was the species found more significantly affected (81.9%). In this species as reported in the province of Treviso, USUV was able to determine clinical signs and high mortality rate with symptoms and lesions resembling those observed in Austria (Weissenböck et al., 2002; Chvala et al., 2004).

USUV has been detected in 9 out of 11 carcasses of birds which died because of nervous symptoms. Six of them were from the outbreaks reported in the province of Treviso.

Detection of USUV in dead birds does not prove that it is the cause of the death of the animal. Excess deaths occurring in a restricted area and the detection of USUV in all dead birds tested as occurred in the Treviso area might indicate that the bird deaths, are likely to be due to USUV. Almost 50% of the blackbirds examined in this study were also parasitised by hemosporidia. Presence of parasites in free-ranging wild birds are a common finding (Cole and Friend, 1999). They debilitate the hosts, and thus might facilitate USUV infections. Although other species like pigeons and magpies have been found serologically and/or virologically infected, no reports of mass die-offs of species other than blackbirds were recorded. This indicates that several species of birds are susceptible to USUV infection, but only a few get fatal disease. In view of the fact that Usutu virus infects wild birds and causes high death rates,

especially in blackbirds, the finding that the chicken embryos, CEF, CEL were apparently resistant to Usutu virus infection could appear unexpected. Other authors, however, have found a similar result when infecting chicken embryo and CEF with USUV (Bakonyi et al., 2005).

Dramatic bird mortality could be due to a particular feeding preference of the vector, to a particular susceptibility of the host species, being completely naive in the new endemic areas or, alternatively, to the high pathogenicity of some USUV strains for certain bird species because of genetic mutations. Like the Central European strains, the Italian North Eastern strains were pathogenic and capable of causing mortality in wild birds, particularly in blackbirds. The Italian North eastern strains however were also able to infect humans and cause encephalitis. Considering that *C. pipiens*, the likely vector of USUV, has been demonstrated to serve both as an enzootic and a bridge vector for humans (Hamer et al., 2008), USUV infection in humans was not surprising. What was surprising, was the encephalitis associated to the infection.

It has been shown that neurovirulence and neuroinvasiveness of Flaviviruses are mainly associated with sequence variation of the protein E and to a lesser extent to other viral proteins such as NS1 and NS2b (Beasley et al., 2005; Brault et al., 2004; Botha et al., 2008). The RA-09-Tm, a strain isolated from a blackbird in the same area where the human infection occurred, was fully sequenced and sequences compared with those of the Budapest, Vienna and South African isolates. The full genome sequence of the USUV isolated in North Eastern Italy grouped with the Budapest and Vienna isolates sharing 99% of similarity differing from both European isolates in 10-11 amino-acids only. These amino-acid substitutions were distributed throughout the entire genome. No differences were noticed in the amino-acid sequences of the protein E, the most commonly referred as potential site of pathogen determinants. It seems then that other host related variables such as susceptibility, age, immune and general health status might have affected the USUV pathogenicity for humans.

The genetic analysis also confirmed that as a whole the European USUV isolate cluster is closely related to the South African USUV. Their nucleotidic and amino acidic sequences shared 97% and 99% similarity, respectively. The minor differences observed in the amino acidic sequence were the same reported by Bakonyi et al. (2004). These authors, assumed that the Vienna strain evolved

1 independently from the South African USUV. A different picture however emerged when the NS5
 2 partial sequences of the Italian, South African and Central European isolates were compared. Three
 3 genetically different clusters could be evidenced. One included all strains isolated in the North Eastern
 4 regions of Italy, a second which grouped the Central European and the Tuscany strains and a third
 5 which comprised the South African strain. The NS5 583 bp partial sequences of the North Eastern and
 6 Tuscany isolates differ in 5 nucleotides. With all caution induced by the limited NS5 fragment
 7 sequenced, one might also hypothesize that not all the Italian strains clustered together. The Tuscany
 8 strain might indeed differ from the other Italian isolates, being closer to the Central European isolates
 9 and for it a further different origin could be hypothesised. According to the seroconversion observed
 10 either in horses or in chicken, USUV has circulated in Tuscany during 2008 and 2009. Unlike the other
 11 Italian USUV strains, the Tuscany strain circulation as shown by the horse and chicken seroconversions
 12 did not coincide with excess bird mortality. In the same area a massive blackbird mortality was
 13 however described in 1996 (Mani et al., 1998). At that time, a JEV-like virus (Mani et al., 2009) was
 14 identified as the virus responsible of the outbreak. It could be supposed that the Tuscany USUV strain
 15 might have been circulating in Italy at least since 1996 and might be the cause of excess mortality in
 16 naive bird population. It might be further hypothesised that it might have established subsequently an
 17 efficient silent transmission cycle between local bird and mosquito species. It has been shown that
 18 after an initial severe bird mortality, the virus might become endemic. The high seroprevalence
 19 observed in these situations in the susceptible hosts might be an indication of the establishment of a
 20 stable population immunity which, however, does not preclude virus activity with new infections and
 21 seroconversions without causing further excess mortality (Meister et al., 2007). The seroconversion in
 22 sentinel animals in both 2008 and 2009 implied the presence of active virus circulation in Tuscany. In
 23 accordance with the horse seroprevalence data, the virus activity in this region (31.58%) was probably
 24 much lower than that occurring in the same year in the North Eastern regions (>90%). In 2008
 25 incidence of USUV in horses in the North eastern regions was high and decreased markedly in the
 26 following year. In birds it has been observed that even after natural infection flaviviral antibody titres
 27 are generally low and subject to considerable variations within short times (Meister et al., 2007). A

similar phenomenon could have occurred in the horse population tested. In this species USUV infection generally does not induce high antibody titers. Horses are not susceptible to Usutu and do not show a strong immune response, however they appear to be good sentinel animals to monitor USUV circulation.

CONCLUSIONS

This study suggests the possibility that two strains of USUVs have circulated in Italy between 2008 and 2009. They have developed strategies of adaptation and evolution to spread into new areas and become established. Their circulation has been associated to high blackbird mortality and, although sharing 100% similarity at amino acidic level with the other European strains, at least the Italian North eastern strain showed a distinctive zoonotic potential. Keeping in mind that dealing with a member of the genus *Flavivirus* the unexpected is often the most frequent event particularly with respect to disease severity, unusual clinical manifestations, and unpredicted methods of transmission, further investigations involving different bird and mammal species are still necessary to show the most important host species, natural reservoirs, and vectors of Usutu virus and to estimate its epidemiologic impact and possible threat to domesticated animals and to the human population. This information will allow us to better understand the diffusion routes and dynamics of potential zoonotic arboviruses and know better the different elements of their complex biological cycle. Therefore it will be possible to design more efficient surveillance, control and prevention activities for these emerging viral zoonoses with great benefit for human and veterinary public health.

REFERENCES

- Bakonyi, T., Gould, E.A., Kolodziejek, J., Weissenböck H., Nowotny, N., 2004. Complete genome analysis and molecular characterization of **Usutu** virus that emerged in Austria in 2001: comparison with the South African strain SAAR-1776 and other flaviviruses. *Virology* 328, 301–310.
- Bakonyi, T., Lussy, H., Weissenböck, H., Hornyák, Á., Nowotny, N., 2005. In Vitro Host-Cell susceptibility to **Usutu** Virus. *Emerging Infectious Diseases* 11, 298-301.

- 1 Bakonyi, T., Erdélyi, K., Ursu, K., Ferenczi, E., Csörgo, T., Lussy, H., Chvala, S., Bukovsky, C., Meister, T.,
2 Weissenböck H., Nowotny, N., 2007. Emergence of **Usutu** Virus in Hungary. Journal of Clinical
3 Microbiology 45, 3870-3874.
- 4 Beasley, D.W., Whiteman, M.C., Zhang, S., Huang, C.Y., Schneider, B.S., Smoth, D.R., Gromowsky, G.D.,
5 Higgs, S., Kinney, R.M., Barrett, A.D., 2005. Envelope protein glycosilation status influence mouse
6 neuroinvasion phenotype of genetic lineage 1 West Nile Virus strains. J Virol. 79, 8339-47.
- 7 Bernard, K.A., Maffei, J.G., Jones, S.A., Kauffman, E.B., Ebel, G.D., Dupuis II, A.P., Ngo, K.A., Nicholas,
8 D.C., Young, D.M., Shi, P.Y., Kulasekera, V.L., Eidson, M., White, D.J., Stone, W.B., NY State West Nile Virus
9 Surveillance Team, Kramer, L.D., 2001. West Nile Virus Infection in Birds and Mosquitoes, New York State,
10 2000. Emerging Infectious Diseases 7, 679-685.
- 11 Botha, E.M., Markotter, W., Wolfaardt, M., Paweska, J.T., Swanepoel, R., Palacios, G., Nel, L.H., Venter,
12 M., 2008. Genetic determinants of virulence in pathogenic lineage 2 West Nile virus strains. Emerg
13 Infect Dis.; Feb14(2), 222-230.
- 14 Brault, A.C., Langevin, S.A., Bowen, R.A., Panella, N.A., Biggerstaff, B.J., Miller, B.R., Komar, N., 2004.
15 Differential virulence of West Nile strains for American crows. Emerg Infect Dis. Dec 10(12), 2161-68.
- 16 Cavrini, F., Gaibani, P., Longo, G., Pierro, A.M., Rossini, G., Bonilauri, P., Gerundi, G.E., Di Benedetto, F.,
17 Pasetto, A., Girardis, M., Dottori, M., Landini, M.P., Sambri, V., 2009. Usutu virus infection in a patient who
18 underwent orthotropic liver transplantation, Italy, August-September 2009. Euro Surveill. 17, 17-19.
- 19 Chvala S., Kolodziejek J., Nowotny N., Weissenböck H., 2004. Pathology and viral distribution in fatal
20 Usutu virus infections of birds from the 2001 and 2002 outbreaks in Austria. J. Comp. Pathol., 131, 176-
21 185.
- 22 Chvala S., Bakonyi T., Bukovsky C., Meister T., Brugger K., Rubel F., Nowotny N., Weissenböck H., 2010.
23 Monitoring of Usutu virus activity and spread by using dead bird surveillance in Austria, 2003-2005. Vet.
24 Microbiol., 122, 237-245.
- 25 Felsenstein, J., 1993. PHYLIP (Phylogeny Inference Package)3.5c Manual (computer program).
26 University of Washington, Dept. of Genetics SK-50, Seattle, WA 98195, USA.

- 1 Hamer, G.L., Kitron, U.D., Brawn, J.D., Loss, S.R., Ruiz, M.O., Goldberg, T.L., Walker, E.D., 2008. *Culex*
2 *pipiens* (Diptera: Culicidae): A bridge vector of West Nile virus to humans. *J. Med. Entomol.* 45, 125–128.
- 3 Heinz, F.X., M.S. Collett, R.H. Purcell, E.A. Gould, C.R. Howard, M. Houghton, R.J.M., Moormann, Rice,
4 C.M., Thiel, H.J., 2000. Family *Flaviviridae*. In *Virus Taxonomy. Seventh Report on International Committe*
5 *on Taxonomy of Viruses*, Eds M.H.V., Van Regenmortel, C.M., Fauquet, D.H.L., Bishop, E.B., Carstens,
6 M.K., Estes, S.M., Lemon, J., Maniloff, M.A., Mayo, D.J., McGeoch, C.R., Pringle, R.B., Wickner. San
7 Diego, San Francisco, New York, Boston, London, Sydney, Tokyo, Academic press. pp 859-878.
- 8 Hubálek, Z., Halouzka, J., Juricová, Z., Sikutová, S., Rudolf, I., Honza, M., Janková, J., Chytil, J., Marec,
9 F., Sitko, J., 2008a. Serologic Survey of Birds for West Nile Flavivirus in Southern Moravia (Czech Republic)
10 *Vector-Borne and Zoonotic Diseases* 8(5), 659-666.
- 11 Hubálek, Z., Wegner, E., Halouzka, J., Tryjanowski, P., Jerzak, L., Sikutová, S., Rudolf, I., Kruszcwicz, A.G.,
12 Jaworski, Z., Wlodarczyk, R., 2008. Serologic Survey of Potential Vertebrate Hosts for West Nile Virus in
13 Poland. *Viral Immunology* 21, DOI: 10.1089/vim.2007.0111.
- 14 Kimura, M., 1980. A simple model for estimating evolutionary rates of base substitutions through
15 comparative studies of nucleotide sequences. *J. Mol. Evol.* 16, 111-120.
- 16 Kuno, G., G.J. Chang, K.R. Tsuchiya, N. Karabatsos and C.B. Cropp, 1998. Phylogeny of the genus
17 *Flavivirus*. *J. Virol.* 72, 73–83.
- 18 Lelli, R., Savini, G., Teodori, L., Filipponi, G., Di Gennaro, A., Leone, A., Di Gialleonardo, L., Venturi, L.,
19 Caporale V., 2008. Serological evidence of *Usutu* virus occurrence in north-eastern Italy. *Zoonoses*
20 *Public Health.* 55, 361-67.
- 21 Mani, P., Rossi, G., Perrucci, S., Bertini, S., 1998. Mortality of *Turdus merula* in Tuscany. *La Selezione*
22 *Veterinaria* 8-9, 749-753.
- 23 Mani, P., Legrottagnie, R., Bertelloni, F., Fratini, F., Filogari D., Rossi G., 2009. I virus dell'Encefalite
24 giapponese (JEV) in uccelli selvatici sinantropici (*Passer italiae*, *Turdus merula*, *Sturnus vulgaris*) e da
25 richiamo (*Turdus musicae*) in Toscana. *Proceeding of International Conference "Passeri in crisi?"*
26 *Ecologia Urbana*, 21,101-102.

- 1 Manarolla, G., Bakonyi, T., Gallazzi, D., Crosta, L., Weissenböck, H., Dorrestein, G.M., Nowotny, N., 2010,
- 2 Usutu virus in wild birds in northern Italy. *Vet. Microbiol.* 141, 59-63.
- 3 Marra, P. P., S. Griffing, C. Caffrey, A. M. Kilpatrick, R. McLean, C. Brand, E. Saito, A. P. Dupuis, L. Kramer,
- 4 and R. Novak. 2004. West Nile virus and wildlife. *Bioscience* 54, 393-402.
- 5 Meister, T., H. Lussy, T. Bakonyi, S. Sikutova', I. Rudolf, W. Vogl, H. Winkler, H. Frey, Z. Hubalek, N.
- 6 Nowotny and H. Weissenböck, 2007. Serological evidence of continuing high USUTU virus (Flaviviridae)
- 7 activity and establishment of herd immunity in wild birds in Austria, *Vet. Microbiol.* 2008, 127:237-248.
- 8 Monaco F, Lelli R, Teodori L, Pinoni C, Di Gennaro AP, Polci A, Calistri P, Savini G. *Re-emergence of West*
- 9 *Nile virus in Italy.* *Zoonoses and Public Health.* 2009; Jul 23. (Epub ahead of print) doi:10.1111/j.1863-
- 10 2378.2009.01245.x.
- 11 Pecorari, M., Longo, G., Gennari, W., Grottola, A., Sabbatini, A.M.T., Tagliazucchi, S., Savini, G.,
- 12 Monaco, F., Simone, M.L., Lelli, R., Rumpianesi, F., 2009. First human case of USUTU virus (USUV)
- 13 neuroinvasive infection. *Eurosurveillance*, August-September 2009. *Euro Surveill.* 17, 15-17.
- 14 Pfeffer, M. and Dobler, G., 2010 Emergence of zoonotic arboviruses by animal trade and migration.
- 15 *Parasites & Vectors* 2010, 3, 35-49
- 16 Rizzoli, A., R. Rosà, F. Rosso, A. Buckley and E. Gould, 2007. West Nile Virus Circulation Detected in
- 17 Northern Italy in Sentinel Chickens, *Vector-Borne and Zoonotic Dis.* 7, 411–417.
- 18 Sivia, DS, 1996. *Data analysis – a Bayesian tutorial.* Oxford University Press.
- 19 Simpson, J.E., Folsom-O'Keefe, C.M., Childs, J.E., Simons, L.E., Andreadis, T.G., Diuk-Wasser, M.A., 2009.
- 20 Avian Host-Selection by *Culex pipiens* in Experimental Trials. *PloS ONE* 4, e7861.
- 21 doi:10.1371/journal.pone.0007861
- 22 Steinmetz H.W., Bakonyi T., Weissenböck H., Hatt J.M., Eulenberger U., Robert N., Hoop R., Nowotny N.,
- 23 2010. Emergence and establishment of Usutu virus infection in wild and captive avian species in and
- 24 around Zurich, Switzerland-Genomic and pathologic comparison to other central European outbreaks.
- 25 *Vet. Microbiol.* doi:10.1016/j.vetmic.2010.09.018.Tamba, M., Bonilauri, P., Bellini, R., Calzolari, M., Albieri,
- 26 A., Sambri, V., Dottori, M., Angelini, P., 2010. Detection of Usutu Virus Within a West Nile Virus
- 27 Surveillance Program in Northern Italy. *Vector-Borne and Zoonotic Diseases*, DOI: 10.1089/vbz.2010.0055

- 1 Turell, M. J., D. J. Dohm, M. R. Sardelis, M.L.O. Guinn, T. G. Andreadis, and J. A. Blow. 2005. An update
2 on the potential of North American mosquitoes (Diptera : Culicidae) to transmit West Nile virus. *J. Med.*
3 *Entomol.* 42, 57-62.
- 4 Weissenböck, H., J. Kolodziejek, A. Url, H. Lussy, B. Rebel-Bauder and N. Nowotny, 2002. Emergence of
5 Usutu virus, an African mosquito-borne flavivirus of the Japanese encephalitis virus group, central
6 Europe. *Emerg. Infect. Dis.* 8, 652–656.
- 7 World Organisation for Animal Health (Office International des Épizooties: OIE), 2008a. West Nile fever,
8 Chapter 2.1.20. In *Manual of diagnostic tests and vaccines for terrestrial animals*. OIE, Paris, France,
9 377-385.
- 10 World Organisation for Animal Health (Office International des Épizooties: OIE), 2008b. Avian influenza,
11 Chapter 2.3.4. In *Manual of diagnostic tests and vaccines for terrestrial animals*. OIE, Paris, France, 465-
12 481.

Table 1

Strains of Usutu viruses included in NS5 and PrM partial sequencing

<i>Date</i>	<i>ID</i>	<i>Bankit/GenBank</i>	<i>Biological matrix</i>	<i>Species</i>	<i>Geographical Origin</i>	<i>NS5</i>	<i>PrM</i>
01/09/09	TV1-09-Tm	1430426	organs	<i>Blackbird (T. merula)</i>	Treviso (Veneto)	x	x
16/09/09	TV2-09-Tm	1430425	organs	<i>Blackbird (T. merula)</i>	Treviso (Veneto)	x	
18/09/09	UD-09-Tm	1430431	organs	<i>Blackbird (T. merula)</i>	Udine (Friuli)	x	
23/10/09	MO-09-Hu	1430429	cerebral spinal fluid	<i>Human</i>	Modena (Emilia Rom.)	x	x
20/10/09	RA-09-Tm	1430432	organs	<i>Blackbird (T. merula)</i>	Ravenna (Emilia Rom.)	x	x
29/09/09	TV3-09-Tm	1430433	organs	<i>Blackbird (T. merula)</i>	Treviso (Veneto)	x	x
07/10/09	VE-09-Tm	1430428	organs	<i>Blackbird (T. merula)</i>	Venezia (Veneto)	x	
2009	PI-09-Cp	1430405	mosquitoes	<i>Culex pipiens</i>	Tuscany	x	
2001	Vienna'01	AY453411	organs	<i>Blackbird (T. merula)</i>	Austria	x	
1959	SAAR-1776	AY453412	mosquitoes	<i>Culex neavei</i>	South Africa	x	
2005	Budapest'05	EF206350	organs	<i>Blackbird (T. merula)</i>	Hungary	x	

Table 2

Seroconversions to Usutu virus in horses in 2009

Region	Province	Number of seroconversions
EMILIA ROMAGNA	FERRARA	5
		1
	REGGIO EMILIA	5
FRIULI VENEZIA GIULIA	UDINE	1
TUSCANY	FIRENZE	1
	LUCCA	1
	PISA	1
	PISTOIA	4
VENETO	VERONA	2
	VICENZA	1

Table 3

Usutu virus seroprevalence in Italian horses in 2008-2009 according to regions

Year	Region	No. of horses tested	No. of positives	% of positives
2008	CAMPANIA	2	0	0,00
	EMILIA ROMAGNA	371	366	98,65
	FRIULI VENEZIA GIULIA	4	1	25,00
	LAZIO	13	0	0,00
	LOMBARDY	112	109	97,32
	MARCHE	12	0	0,00
	SICILY	12	0	0,00
	TUSCANY	19	6	31,58
	VENETO	255	227	89,02
	Total	800	709	88,6
2009	BASILICATA	207	0	0,00
	CALABRIA	71	0	0,00
	CAMPANIA	84	0	0,00
	EMILIA ROMAGNA	739	100	13,53
	FRIULI VENEZIA GIULIA	207	6	2,90
	LAZIO	619	0	0,00
	LIGURIA	3	0	0,00
	LOMBARDY	195	20	10,26
	MARCHE	162	0	0,00
	MOLISE	96	0	0,00
	PIEDMONT	1	0	0,00
	APULIA	59	0	0,00
	SARDINIA	86	0	0,00
	SICILY	280	0	0,00
	TUSCANY	404	22	5,45
	TRENTINO ALTO ADIGE	1	0	0,00
	UMBRIA	65	0	0,00
	VENETO	851	203	23,85
	Total	4130	351	8,5
Total 2008 and 2009		4930	1060	21,5

Table 4

Virological and serological findings in Italian wild birds examined for USUTU virus in the years 2008-2009

Accepted Manuscript

Region	Species	Common name	RT-PCR		Serum-neutralisation	
			No. of animals tested	No. of positives	No. of animals tested	No. of positives
BASILICATA	<i>Columba livia</i>	Rock pigeon	1	0	0	0
	<i>Gallus gallus domesticus</i>	Chicken	0	0	2	0
CAMPANIA	<i>Gallus gallus domesticus</i>	Chicken	0	0	1	0
	-	Wild birds	1	0	0	0
EMILIA ROMAGNA	<i>Otus scops</i>	Eurasian scops owl	1	0	0	0
	<i>Accipiter gentilis</i>	Goshawk	1	0	0	0
	<i>Tyto alba</i>	Barn owl	1	0	0	0
	<i>Athena noctua</i>	Little owl	2	0	0	0
	<i>Columba livia</i>	Rock pigeon	1	0	17	17
	<i>Corvus corone corvix</i>	Carrion crow	147	0	0	0
	<i>Corvus corax</i>	Common raven	2	0	0	0
	<i>Larus occidentalis</i>	Seagull	8	0	0	0
	<i>Pica pica</i>	Magpie	223	2	0	0
	<i>Anas platyrhynchos</i>	Mallard	1	0	0	0
	<i>Garrulus glandarius</i>	European jay	21	0	0	0
	<i>Bubo bubus</i>	Eurasian eagle owl	1	0	0	0
	<i>Turdus merula</i>	Blackbird	8	2	0	0
	<i>Erithacus rubecula</i>	European robin	1	0	0	0
	<i>Columba livia</i>	Rock pigeon	1	0	0	0
	<i>Accipiter nisus</i>	Eurasian sparrow	2	0	0	0
	<i>Sturnus vulgaris</i>	Common starling	112	0	0	0
	<i>Streptopelia turtur</i>	Eurasian collared dove	1	0	0	0
FRIULI VENEZIA GIULIA	<i>Corvus corone corvix</i>	Carrion crow	3	0	6	0
	<i>Corvus corax</i>	Common raven	0	0	1	0
	<i>Pica pica</i>	Magpie	39	0	80	0
	<i>Gallus gallus domesticus</i>	Chicken	0	0	5	2
LAZIO	<i>Anas platyrhynchos</i>	Mallard	0	0	1	0
	<i>Corvus corone corvix</i>	Carrion crow	38	0	0	0
MARCHE	<i>Corvus corax</i>	Common raven	5	0	0	0
	<i>Pica pica</i>	Magpie	20	0	0	0
	<i>Sturnus vulgaris</i>	Common starling	1	0	0	0
MOLISE	<i>Gallus gallus domesticus</i>	Chicken	2	0	8	0
SARDINIA	<i>Corvus corone corvix</i>	Carrion crow	8	0	0	0
	<i>Phoenicopterus ruber roseus</i>	Greater flamingo	0	0	1	0
	<i>Gallus gallus domesticus</i>	Chicken	0	0	3	0
SICILY	<i>Gallus gallus domesticus</i>	Chicken	0	0	14	0
TUSCANY	<i>Anas platyrhynchos</i>	Mallard	0	0	6	6
	<i>Columba livia</i>	Rock pigeon	159	0	0	0
	<i>Pica pica</i>	Magpie	4	0	0	0
	<i>Gallus gallus domesticus</i>	Chicken	0	0	5	5
	<i>Turdus philomelos</i>	Song thrush	7	0	0	0
	<i>Streptopelia turtur</i>	Eurasian collared dove	12	0	0	0
TRENTINO	<i>Tetrao urogallus</i>	Capercaille	1	0	0	0
UMBRIA	<i>Corvus corax</i>	Common raven	0	0	73	0
	<i>Pica pica</i>	Magpie	0	0	57	0
	<i>Gallus gallus domesticus</i>	Chicken	0	0	1	0
VENETO	<i>Columba palumbus</i>	Wood pigeon	0	0	1	0
	<i>Columba livia</i>	Rock pigeon	0	0	135	3
	<i>Corvus corone corvix</i>	Carrion crow	55	0	3	0
	<i>Fringilla coelebs</i>	Chaffinch	1	0	0	0
	<i>Larus occidentalis</i>	Seagull	6	0	0	0
	<i>Pica pica</i>	Magpie	113	0	17	2
	<i>Turdus merula</i>	Blackbird	9	7	1	1
	<i>Erithacus rubecula</i>	European robin	1	0	0	0
	<i>Sturnus vulgaris</i>	Common starling	3	0	0	0
	<i>Turdus philomelos</i>	Song thrush	22	0	0	0
	<i>Streptopelia turtur</i>	Eurasian collared dove	6	0	0	0
	-	Wild birds	5	0	0	0
Total			1045	11	404	36

Table 5

Percentage of nucleotidic and amino-acid sequence similarity between European and South African USUTU strains

Legend:

Vienna'01: Vienna strain 2001

SAAR-1776: South African strain SAAR-1776

Budapest'05: Budapest strain 05

RA-09-Tm: Italian strain Ravenna 2009

	Vienna'01	SAAR-1776	Budapest'05	RA-09-Tm
Vienna'01		97	100	99
SAAR-1776	99		97	97
Budapest'05	100	99		99
Italy'09	100	99	100	

aa

bp

Table 6

Comparison between the amino-acid deduced sequences of the European and South African Usutu strains

Position bp	Vienna'01	SAAR-1776	Budapest'05	RA-09-Tm	Position aa	Gene	Vienna'01	SAAR-1776	Budapest'05	RA-09-Tm
454-56	GTT	GCC	GTT	GTT	120	C	val	ala	val	val
622-24	ACT	ACT	ATT	ACT	176	M	thr	thr	ila	thr
637-39	AAA	AAA	AAA	ACT	181		lys	lys	lys	arg
1351-53	ATG	GTG	ATG	ATG	419	E	met	val	met	met
1801-03	GGC	AGC	GGC	GGC	569		gly	ser	gly	gly
1933-35	GGC	AGC	GGC	GGC	613		gly	ser	gly	gly
2242-44	ACA	CCA	ACA	ACA	716		thr	pro	thr	thr
2464-66	AAC	AGC	AAC	AAC	790		asn	ser	asn	asn
2584-86	GAA	GAG	GAA	GGA	830	NS1	glu	glu	glu	gly
2911-13	GCA	GTA	GCA	GTA	939		asn	val	asn	val
3445-47	AGG	AAG	AGG	AGG	1117		arg	lys	arg	arg
3532-34	CAT	TAC	CAT	TAT	1146	NS2a	his	tyr	his	tyr
3685-87	ACT	ACT	ATT	ACT	1197		thr	thr	ile	thr
3895-97	AAC	GAC	AAC	AAC	1267		asn	asp	asn	asn
3898-900	TTT	CTC	TTT	TTT	1268		phe	leu	phe	phe
4066-68	ATC	ATT	ACC	ACC	1324		ile	ile	thr	thr
4900-02	ATA	ATA	ATA	GTA	1602	NS3	ile	ile	ile	val
4948-50	ATC	GTC	ATC	ATC	1618		ile	val	ile	ile
5179-81	CTG	ATG	CTG	CTG	1695		leu	met	leu	leu
5431-33	GCT	GTT	GTT	GTT	1779		ala	val	val	val
6184-86	AGA	ATA	AGA	AGA	2030		arg	ile	arg	arg
6190-02	GAG	CAG	GAG	GAG	2032		glu	gln	glu	glu
6412-14	GCT	TCC	GCT	GCT	2106		ala	ser	ala	ala
6592-94	CTT	TTT	CTT	CTT	2166	NS4a	leu	phe	leu	leu
6955-57	ATA	ATG	ATA	ATA	2287	NS4b	ile	met	ile	ile
6964-66	AGC	GGC	AGC	AGC	2290		ser	gly	ser	ser
7006-08	AAC	AAC	AAC	AGC	2304		asn	asn	asn	ser
7195-97	TTA	TTG	TTG	CTG	2367		phe	leu	leu	leu
7744-46	AAG	AGG	AAG	AAG	2550	NS5	lys	arg	lys	lys
8029-31	ATG	ATG	ATG	ATA	2645		met	met	met	ile
8503-05	ACC	GCC	ACC	ACC	2803		thr	ala	thr	thr
8641-43	AGC	GGC	AGC	AGC	2849		ser	gly	ser	ser
10183-85	CCA	CCA	CCA	CTA	3363		pro	pro	pro	leu
10375-77	AAT	GAT	AAT	AAT	3427		asn	asp	asn	asn

Figure 1

Areas surveyed for the presence of Usutu virus and Usutu virus antibodies in 2008 and 2009

Figure 2

Usutu virus circulation in Italy in 2008 and 2009

Figure 3

Neighbor joining tree compiled from aligned nucleic acid sequences of the partial NS5 segments of Usutu virus isolates. The tree is rooted with the same sequence of the Eg101 WNV strain.

The most significant support value from the bootstrap analysis is reported.

Legend:

Eg101: West Nile strain Egypt 101

PI 09 Cp: Usutu strain detected in *C. pipiens* from Tuscany in 2009

Vienna01: Usutu strain Vienna 2001

Budapest05: Usutu strain Budapest 2005

SAAR-1776: Usutu strain from South African 1959

VE 09 Tm: Usutu strain isolated in 2009 in the province of Venice from blackbird (*T. merula*)

TV1 09 Tm: Usutu strain isolated in 2009 in the province of Treviso from blackbird (*T. merula*)

TV2 09 Tm: Usutu strain isolated in 2009 in the province of Treviso from blackbird (*T. merula*)

TV3 09 Tm: Usutu strain isolated in 2009 in the province of Treviso from blackbird (*T. merula*)

MO 09 Tm: Usutu strain isolated in 2009 in the province of Modena from blackbird (*T. merula*)

RA 09 Tm: Usutu strain isolated in 2009 in the province of Ravenna from blackbird (*T. merula*)

UD 09 Tm: Usutu strain isolated in 2009 in the province of Udine from blackbird (*T. merula*)