Transmission between chickens of an H7N1 Low Pathogenic Avian Influenza virus isolated during the epidemic of 1999 in Italy

J.L. Gonzales, J.A. van Der Goot, J.A. Stegeman, A.R.W. Elbers, G. Koch

► To cite this version:

J.L. Gonzales, J.A. van Der Goot, J.A. Stegeman, A.R.W. Elbers, G. Koch. Transmission between chickens of an H7N1 Low Pathogenic Avian Influenza virus isolated during the epidemic of 1999 in Italy. *Veterinary Microbiology*, 2011, 152 (1-2), pp.187. 10.1016/j.vetmic.2011.04.022 . hal-00719082

HAL Id: hal-00719082

<https://hal.science/hal-00719082>

Submitted on 19 Jul 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Title: Transmission between chickens of an H7N1 Low Pathogenic Avian Influenza virus isolated during the epidemic of 1999 in Italy

Authors: J.L. Gonzales, J.A. van der Goot, J.A. Stegeman, A.R.W. Elbers, G. Koch

PII: S0378-1135(11)00236-7
DOI: doi:10.1016/j.vetmic.2011.04.022
Reference: VETMIC 5285

To appear in: *VETMIC*

Received date: 4-1-2011
Revised date: 6-4-2011
Accepted date: 14-4-2011

Please cite this article as: Gonzales, J.L., van der Goot, J.A., Stegeman, J.A., Elbers, A.R.W., Koch, G., Transmission between chickens of an H7N1 Low Pathogenic Avian Influenza virus isolated during the epidemic of 1999 in Italy, *Veterinary Microbiology* (2010), doi:10.1016/j.vetmic.2011.04.022

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Transmission between chickens of an H7N1 Low Pathogenic
Avian Influenza virus isolated during the epidemic of 1999
in Italy**

J.L. Gonzales ^{a,b}, J.A. van der Goot ^{a,c}, J.A. Stegeman ^b, A.R.W. Elbers ^a, G. Koch ^c.

^a Department of Epidemiology, Crisis organization and Diagnostics,

Central Veterinary Institute (CVI), part of Wageningen UR, P.O. Box 65, 8200 AB

Lelystad, The Netherlands.

^b Department of Farm Animal Health, Faculty of Veterinary Medicine, Utrecht

University P.O. Box 80, 151, 3508 TD Utrecht, The Netherlands.

^c Department of Virology, Central Veterinary Institute (CVI), part of Wageningen UR,

P.O. Box 65, 8200 AB Lelystad, The Netherlands.

Correspondence: Jose L Gonzales, Central Veterinary Institute (CVI), part of

Wageningen UR, P.O. Box 65, 8200 AB Lelystad, The Netherlands. Email:

jose.gonzales@wur.nl

25 **Abstract**

26 The transmissibility of an H7N1 Low Pathogenic Avian Influenza (LPAI) virus
 27 isolated from a turkey flock during the large epidemic in Italy in 1999, was
 28 experimentally studied in chickens. Four group transmission experiments were
 29 performed. Infection and transmission were monitored by means of virus isolation on
 30 swab samples and antibody detection in serum samples. From the results of these
 31 groups, we estimated the mean infectious period at 7.7 (6.7 – 8.7) days, the
 32 transmission rate parameter at 0.49 (0.30 – 0.75) infections per infectious chicken per
 33 day and the basic reproduction ratio at 3.8 (1.3 – 6.3). These estimates can be used for
 34 the development of surveillance and control programmes of LPAI in poultry.

35
 36 **Keywords:** Avian Influenza, LPAI, H7N1, Transmission, Transmission parameters,
 37 Reproduction ratio

39 **1. Introduction**

40 Highly pathogenic Avian influenza (HPAI) is caused by AI viruses of H7 or
 41 H5 subtypes, that have evolved from preceding Low Pathogenic Avian Influenza
 42 viruses (LPAIv) (Banks et al., 2001; Capua and Maragon, 2000; DEFRA, 2008;
 43 Garcia et al., 1996). The H7N1 LPAI epidemic in Italy in 1999, which preceded the
 44 HPAI (1999 – 2000) epidemic, is one of the largest LPAI epidemics reported in a
 45 western country. In the absence of control measures – no compulsory regulations for
 46 control of LPAI outbreaks were in place – a total of 199 LPAI infected poultry farms
 47 were detected before the HPAI virus (HPAIv) took over (Busani et al., 2009; Capua
 48 and Maragon, 2000). Current knowledge does not allow to predict when an H5 or H7
 49 LPAIv strain will mutate into a HPAIv strain. However, assuming that the molecular

changes necessary for the virulence transformation occur at random, the probability that a LPAIv strain will mutate into a HPAIv strain will depend on the number of virus replicates, which in turn is associated with the number of birds acquiring infection. Hence, besides knowledge of the molecular biology of these viruses, quantitative knowledge of the transmission of LPAIv is important for the development of surveillance programmes that target early detection (see for example Graat et al., (2001)), hence reducing the probability of both undesired mutations and/or large epidemics.

To our knowledge, until now only one study quantifying LPAI virus transmission between chickens has been published in the peer reviewed literature – a chicken-derived H5N2 LPAI precursor virus of the 1983 HPAI epidemic in Pennsylvania (Van der Goot et al., 2003) – and there is no information about the transmissibility between chickens of H7 LPAIv subtypes. The objective of this study was to characterize – by means of transmission experiments – the between-chicken transmission of an H7N1 LPAIv isolated from turkey during the 1999 epidemic in Italy.

2. Material and methods

2.1 Virus and inoculation

The LPAIv used for this study was the H7N1 LPAI A/turkey/Italy/1067/99 virus isolate. Its intravenous pathogenicity index and amino acid sequence of the hemagglutinin gene (GenBank accession number AF364134) have been published elsewhere (Banks et al., 2001; Capua and Maragon, 2000). Birds were inoculated both intranasally and intratracheally with 0.1 ml/route of inoculum containing 10^6 EID₅₀ (50% egg infectious dose)/ml.

75

76 *2.2 Birds and experimental procedure*

77 A total of two transmission experiments were performed and each experiment
 78 consisted of two replicates (here referred to as trials). A trial comprised 10 six-week-
 79 old specific pathogen free (SPF) white leghorn chickens (Charles Rivers, The
 80 Netherlands). Five of the 10 birds were inoculated and the remaining five kept as
 81 contacts. The experiments were carried out in the High Containment Unit at the
 82 Central Veterinary Institute of Wageningen UR in Lelystad using biosafety level 3
 83 procedures. Birds were housed in different pens in the same experimental room. Next
 84 to each experimental pen, 2 pairs of birds, housed in separate smaller pens, were
 85 placed as sentinels to detect whether between-pen (indirect) transmission should be
 86 considered and monitor independence of the trials. Chickens were housed 1 week
 87 before challenge to allow adaptation and during this period they were tested for
 88 presence of antibodies against influenza A virus using an in-house competitive NP
 89 ELISA (de Boer et al., 1990) and for virus by virus isolation. The day of inoculation
 90 (day = 0), contacts were removed from their pen and placed back 24 hours later.
 91 Trachea and cloaca swabs were collected daily from 1 to 10 days post inoculation
 92 (dpi) and then on 14, 17 and 21 dpi. Serum samples were taken at 7, 14 and 21 dpi.
 93 All surviving birds were euthanized at 21dpi. All experiments complied with the
 94 Dutch Law on Animal Experiments and were reviewed by an ethical committee.

95 Trachea and cloaca swabs were used for virus isolation using SPF
 96 embryonated chicken eggs that were incubated for 9 days. Serum samples were tested
 97 for antibodies using the Hemagglutinin Inhibition test (HI) using 8 Hemagglutination
 98 units (HAU) of H7N1 as antigen. Diagnostic procedures were similar as those
 99 described by van der Goot et al. (2003). A bird was considered infected when it

100 scored positive in virus isolation or serology. A bird was considered infectious for the
 101 time it remained positive in virus isolation.

102

103 *2.3 Data analysis*

104 The basic reproduction ratio (R_0), which is defined as the average number of
 105 new infections caused by one infectious individual in a susceptible population, was
 106 estimated by the final size (FS) and the generalized linear model (GLM) methods as
 107 described by van der Goot et al. (2003; 2005). The latent period was estimated as the
 108 time (in days) from inoculation to the first day the inoculated chicken was positive to
 109 virus isolation. The infectious period was estimated by fitting a parametric survival
 110 regression model with a Weibull distribution. In some cases, infected birds were
 111 detected positive by virus isolation, late, for example at 10 dpi and were tested again
 112 at 14 dpi when they became negative and assumed to have recovered (Table 1).
 113 These cases were treated as interval censoring in the survival analysis. Survival
 114 analysis and the GLM method were carried out with the statistical software R (R
 115 Development Core Team, 2005).

116

117 **3. Results**

118 In the first experiment, all inoculated birds in both trials became infected and
 119 transmitted virus to 4 and 5 contact birds in the first and second trial respectively
 120 (Tables 1a and 1b). In the second experiment (trials 3 and 4) all inoculated and
 121 contact birds became infected (Tables 1c and 1d). No apparent clinical signs were
 122 observed in infected birds. The sentinels remained negative during the course of the
 123 experiment, which confirmed the independence of the trials.

Inoculated birds were positive to virus isolation at 1 dpi. The estimated mean length of the infectious period (IP) was 7.7 (95% Confidence Interval (CI): 6.7 – 8.7) days and the mean estimate of the transmission parameter β was equal to 0.49 (95% CI: 0.30 – 0.75) infections/day per chicken. R_0 was estimated to be 3.8 (95% CI: 1.3 – 6.3) by the GLM method and 4.0 (95% CI: 1.7 – 11.0) by the FS method (Table 2).

4. Discussion

Herein the transmissibility in chickens of the H7N1 LPAIv, which caused a severe epidemic in poultry in Italy in 1999 and later evolved into a HPAI epidemic, was characterized. We have, in particular, estimated β and the IP. These parameters give an indication of how fast the virus would spread to another susceptible bird and how long an infected chicken would remain infectious. Due to, mainly, the subclinical presentation of LPAI infections, it is difficult to accurately follow the course of the infection in field conditions, and characterization of transmission in the field would be limited to estimates of the R_0 (Comin et al., 2010). Our experimental approach allowed us to follow the course of the infection under controlled conditions and have objective estimates of the transmission parameters.

The mean estimates of the IP, β and R_0 in chickens for this H7N1 LPAIv are higher than those observed for an H5N2 LPAIv (van der Goot et al., 2003). This is remarkable because the latter strain was isolated from a chicken whereas our strain originated from turkey. Upon introduction, the virus here studied would reach its peak (highest prevalence of infectious animals) faster and at a higher prevalence than the H5N2 virus. Thus, it could be hypothesized that the infectivity of the flock would be higher, resulting in a higher probability of transmission (assuming similar contact rate) to other flocks. In this scenario, surveillance programmes would need to sample

149 at a higher frequency if the objective is to detect in an early stage infection and to
150 reduce the probability of between- flock transmissions.

151 Of the two methods used to estimate R_0 , the GLM method yielded a more
152 precise estimate (Table 2). This is because the GLM method uses all the detailed (day
153 to day) information about the time course of the infection chain, while the FS method
154 uses only the total number of animals that got infected throughout the experiment. We
155 also used the FS method, because with that method there is no dependency on whether
156 or not there is a latency period, as intrinsically assumed in the GLM method.

157 In conclusion, this study revealed valuable information on transmission
158 parameters for LPAI virus in chickens. Such information can be used to improve or
159 develop surveillance programmes and control measures.

160

161 **Acknowledgement**

162 The research was financially supported by EU contract QLK2-CT-2002-01454

163 AVIFLU and the Dutch Ministry of Economic Affairs, Agriculture and Innovation.

164 We thank Giovanni Cattoli (Istituto Zooprofilattico Sperimentale delle Venezie) for
165 providing the virus for this study.

166

167

168

168 **References**

- 169 Banks, J., Speidel, E.S., Moore, E., Plowright, L., Piccirillo, A., Capua, I., Cordioli,
170 P., Fioretti, A., Alexander, D.J., 2001, Changes in the haemagglutinin and the
171 neuraminidase genes prior to the emergence of highly pathogenic H7N1 avian
172 influenza viruses in Italy. *Archives of Virology* 146, 963-973.
- 173 Busani, L., Valsecchi, M.G., Rossi, E., Toson, M., Ferrè, N., Pozza, M.D., Marangon,
174 S., 2009, Risk factors for highly pathogenic H7N1 avian influenza virus
175 infection in poultry during the 1999-2000 epidemic in Italy. *The Veterinary*
176 *Journal* 181, 171-177.
- 177 Capua, I., Maragon, S., 2000, The avian influenza epidemic in Italy, 1999-2000.
178 *Avian Pathol* 29, 289-294.
- 179 Comin, A., Klinkenberg, D., Stegeman, A., Busani, L., Marangon, S., 2010. Estimate
180 of basic reproduction number (R_0) of low pathogenicity avian influenza
181 outbreaks using a bayesian approach. In: *Meeting of the Society for*
182 *Veterinary Epidemiology and Preventive Medicine*, Nantes, France, pp. 145 -
183 153.
- 184 de Boer, G.F., Back, W., Osterhaus, A.D.M.E., 1990, An ELISA for detection of
185 antibodies against influenza A nucleoprotein in humans and various animal
186 species. *Archives of Virology* 115, 47-61.
- 187 DEFRA 2008. Highly pathogenic avian influenza – H7N7, Oxfordshire, June 2008.
188 Situation at 12.30pm Wednesday 2nd July.
- 189 Garcia, M., Crawford, J.M., Latimer, J.W., Rivera-Cruz, E., Perdue, M.L., 1996,
190 Heterogeneity in the haemagglutinin gene and emergence of the highly
191 pathogenic phenotype among recent H5N2 avian influenza viruses from
192 Mexico. *J Gen Virol* 77, 1493-1504.

- 193 Graat, E.A.M., de Jong, M.C.M., Frankena, K., Franken, P., 2001, Modelling the
194 effect of surveillance programmes on spread of bovine herpesvirus 1 between
195 certified cattle herds. *Veterinary Microbiology* 79, 193-208.
- 196 R Development Core Team 2005. R: A Language and Environment for Statistical
197 Computing. (Vienna, Austria: Foundation for Statistical Computing, 2005).
- 198 van der Goot, J.A., De Jong, M.C.M., Koch, G., van Boven, M., 2003, Comparison of
199 the transmission characteristics of low and high pathogenicity avian influenza
200 A virus (H5N2). *Epidemiology and Infection* 131, 1003-1013.
- 201 van der Goot, J.A., Koch, G., de Jong, M.C.M., van Boven, M., 2005, Quantification
202 of the effect of vaccination on transmission of avian influenza (H7N7) in
203 chickens *PNAS* 102 18141-18146.
204

Table 1. Overview of the transmission process in the four group-transmission trials

Table 1a		Days post – inoculation											Serology
Trial 1	1	2	3	4	5	6	7	10	14	17	21		
Inoculated	+ ^a	+	+	++	++	+	-	-	nd	nd	nd	+ (7) ^c	
Inoculated	+	++	++	++	++	+	+	-	nd	nd	nd	+ (7)	
Inoculated	++	++	+	+	+	+	+	-	nd	nd	nd	+ (6)	
Inoculated	+	+	++	++	+	+	+	-	nd	nd	nd	+ (7)	
Inoculated	+	+	+	+	-	-	-	-	nd	nd	nd	+ (4)	
Contact	nd ^b	-	+	+	++	++	++	-	-	-	-	+ (7)	
Contact	nd	+	-	-	-	+	+	-	-	-	-	- (2)	
Contact	nd	-	-	-	-	-	-	+	-	-	-	+ (5)	
Contact	nd	-	-	-	-	-	-	-	+	-	-	+ (6)	
Contact	nd	-	-	-	-	-	-	-	-	-	-	-	

Table 1b		Days post – inoculation											Serology
Trial 2	1	2	3	4	5	6	7	10	14	17	21		
Inoculated	+	+	+	+	-	-	-	-	nd	nd	nd	+ (8)	
Inoculated	+	++	++	++	++	+	-	-	nd	nd	nd	+ (6)	
Inoculated	+	+	+	+	+	+	-	-	nd	nd	nd	+ (5)	
Inoculated	++	++	++	++	++	+	+	-	nd	nd	nd	+ (6)	
Inoculated	+	+	+	+	++	++	-	-	nd	nd	nd	+ (6)	
Contact	nd	-	-	-	-	+	++	-	-	-	-	-	
Contact	nd	-	-	+	++	++	++	++	-	+	-	+ (6)	
Contact	nd	-	-	-	-	+	+	+	-	-	-	+ (5)	
Contact	nd	-	-	-	-	-	++	+	+	-	-	+ (5)	
Contact	nd	-	-	-	-	++	++	+	-	-	-	+ (6)	

Table 1c		Days post – inoculation												Serology
Trial 3	1	2	3	4	5	6	7	8	9	10	14	17	21	
Inoculated	+	+	+	+	-	+	-	-	-	-	-	-	-	+ (6)
Inoculated	+	+	-	+	+	+	-	-	-	-	-	-	-	+ (3)
Inoculated	+	+	+	+	-	++	-	-	-	-	-	-	-	+ (5)
Inoculated	+	++	++	++	+	+	+	+	-	-	-	-	-	+ (7)
Inoculated	+	+	+	+	+	++	+	+	+	+	-	-	-	+ (5)
Contact	nd	-	-	+	+	-	-	+	-	-	-	-	-	+ (5)
Contact	nd	-	+	+	+	++	+	-	+	-	-	-	-	+ (4)
Contact	nd	-	+	++	+	+	++	+	+	+	-	-	-	+ (6)
Contact	nd	-	-	+	+	+	+	+	+	+	+	-	-	+ (7)
Contact	nd	-	-	+	-	+	-	+	-	-	-	-	-	- (2)

Table 1d		Days post – inoculation												Serology
Trial 4	1	2	3	4	5	6	7	8	9	10	14	17	21	
Inoculated	+	+	+	++	++	+	-	-	-	-	-	-	-	+ (7)
Inoculated	+	++	++	++	++	++	+	-	-	-	-	-	-	+ (7)
Inoculated	+	+	+	+	+	-	-	-	-	-	-	-	-	+ (6)
Inoculated	+	++	++	++	++	+	+	-	-	-	-	-	-	+ (7)
Inoculated	+	+	+	+	+	+	+	-	+	-	-	-	-	+ (6)
Contact	nd	-	-	-	+	++	++	+	+	+	+	+	-	+ (8)
Contact	nd	-	+	+	+	-	++	+	+	+	-	+	-	+ (6)
Contact	nd	-	-	-	-	+	+	+	+	+	-	-	-	+ (6)
Contact	nd	-	+	+	-	-	+	++	+	+	-	-	-	+ (6)
Contact	nd	-	-	++	+	+	+	+	+	+	+	-	-	+ (6)

^a ++, +|, -|, - : virus isolation results of “trachea | cloaca” swab samples. The single “-” means that both type of samples were negative.

^b nd: test not done

^c HI results at day 21 post-inoculation. Number in parenthesis are log₂ HI titers. Titers higher than 2 were considered positive.

Accepted Manuscript

Table 2 Estimates (95% confidence interval) of transmission parameters of H7N1 LPAI

Parameter	Estimates
Number of contacts infected per trial	4, 5, 5, 5
Latent period ^b	≤1
Mean length Infectious period	7.7 (6.7 – 8.7)
Transmission rate β (day ⁻¹)	0.49 (0.30 – 0.75)
Reproduction ratio R_0 (GLM) ^a	3.8 (1.3 – 6.3)
Reproduction ratio R_0 (FS) ^a	4.0 (1.7 – 11.0)

^a R_0 estimates using the generalized lineal model (GLM) or the final size (FS) methods.

^b Inoculated birds were swab-sampled 24 hours post inoculation.