Source and profile of paralytic shellfish poisoning toxins in shellfish in Daya Bay, South China Sea

Wang Zhao-Hui, Nie Xiang-Ping, Jiang Shi-Jun, Zhang Jian-Gang, Cao Yu,
Zhang Yu-Juan, Wang Da-Zhi

► To cite this version:

Wang Zhao-Hui, Nie Xiang-Ping, Jiang Shi-Jun, Zhang Jian-Gang, Cao Yu, et al.. Source and profile of paralytic shellfish poisoning toxins in shellfish in Daya Bay, South China Sea. *Marine Environmental Research*, 2011, 72 (1-2), pp.53. 10.1016/j.marenvres.2011.04.007 . hal-00720180

HAL Id: hal-00720180

<https://hal.science/hal-00720180>

Submitted on 24 Jul 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Accepted Manuscript

Title: Source and profile of paralytic shellfish poisoning toxins in shellfish in Daya Bay, South China Sea

Authors: Wang Zhao-Hui, Nie Xiang-Ping, Jiang Shi-Jun, Zhang Jian-Gang, Cao Yu, Zhang Yu-Juan, Wang Da-Zhi

PII: S0141-1136(11)00050-X

DOI: [10.1016/j.marenvres.2011.04.007](https://doi.org/10.1016/j.marenvres.2011.04.007)

Reference: MERE 3523

To appear in: *Marine Environmental Research*

Received Date: 12 July 2010

Revised Date: 18 March 2011

Accepted Date: 28 April 2011

Please cite this article as: Zhao-Hui, W., Xiang-Ping, N., Shi-Jun, J., Jian-Gang, Z., Yu, C., Yu-Juan, Z., Da-Zhi, W. Source and profile of paralytic shellfish poisoning toxins in shellfish in Daya Bay, South China Sea, *Marine Environmental Research* (2011), doi: [10.1016/j.marenvres.2011.04.007](https://doi.org/10.1016/j.marenvres.2011.04.007)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Source and profile of paralytic shellfish poisoning toxins in shellfish in Daya Bay, South China Sea

Wang Zhao-Hui ^{a, b*}, Nie Xiang-Ping ^{a, b}, Jiang Shi-Jun ^{a, b}, Zhang Jian-Gang ^a, Cao Yu ^a,

Zhang Yu-Juan ^a, Wang Da-Zhi ^c

^a College of Life Science and Technology, Jinan University, Guangzhou, 510632, China

^b Key Laboratory of Eutrophication and Red Tide Prevention, Education Department of Guangdong Province, Guangzhou 510632, China

^c Key Laboratory for Marine Environmental Science of Ministry of Education, Xiamen University, Xiamen 361005, PR China

Corresponding author: Wang Zhao-Hui

Telephone: 8620-85225100

Fax: 8620-85220239

E-mail: twzh@jnu.edu.cn

Correspondence address: Institute of Hydrobiology, College of Life Science and Technology,
Jinan University, Guangzhou, 510632, China

Abstract

Changes in cell density and cyst flux of *Alexandrium tamarense*, paralytic shellfish poisoning (PSP) toxin contents in shellfishes, and environmental parameters were measured in two stations in Daya Bay, South China Sea from March 2005 to July 2006. Vegetative cells of *A. tamarense* occurred sporadically; however, they presented abundantly during the winter months. Meanwhile, cyst flux reached its maximum level just following the peak abundance of motile cells. The PSP contents in shellfish were generally low, but higher in winter with the maximum of 14 015 µg STX equiv./kg. The majority of toxins were found in digestive glands, with a maximum of 66 227 µg STX equiv./kg. There were significant positive relationships between toxin level and vegetative cell density and cyst flux. This indicates that vegetative cells and cysts of *Alexandrium* significantly influenced PSP level, and could be an important source of PSP toxins in shellfish during winter.

Key words: Paralytic shellfish poisoning (PSP) toxins; *Alexandrium tamarense*; Cyst; Phytoplankton; Algal toxins; Toxicity; South China Sea

1. Introduction

Paralytic shellfish poisoning (PSP) toxins are biotoxins produced by micro-phytoplankton that can accumulate in filter-feeding shellfish (Shumway, 1990; Hallegraeff, 1995). Subsequent ingestion of contaminated shellfish by humans can result in serious illness or even death. The incidence of human poisoning has increased markedly since the early 1970s (Hallegraeff, 1995). Species within the dinoflagellate genus *Alexandrium*, such as *A. tamarense* and *A. catenella*, have been known as important producers of PSP toxins. Resting cysts of these toxic dinoflagellates have been shown to contain toxins (Bravo et al., 1998; Persson et al., 2006) and sometimes have been reported to contain six (Oshima et al., 1992) to ten folds (Yentsch and Mague, 1979) more toxin than vegetative cells. Dinoflagellate cysts are one of the food sources of bivalves (Tsujino and Uchida, 2004; Persson et al., 2006). Many studies have suggested that consumption of cysts may be the direct cause of toxicity in bivalves, especially during the winter months (Shumway et al., 1988; Schwinghamer et al., 1994; Anderson, 1997; Bravo et al., 1998).

Daya Bay is a site of recurrent PSP episodes (Anderson et al., 1996; Lin et al., 1999) with the highest PSP toxin levels among the Chinese coastal areas (Lin et al., 1999; Jiang et al., 2000a). Human mortalities and hospitalizations have occurred from time to time in nearby areas following consumption of toxic mussels. Five persons were poisoned due to consumption of the contaminated shellfish in 1989 (Jian et al., 1991), and three persons died after shellfish consumption in 1993 (Lin et al., 1999). Bivalves collected from this sea area had PSP toxin concentrations exceeding the regulatory limit of 800 µg STX equiv./kg, and a maximum concentration of 10 000 µg STX equiv./kg was reported (Jiang et al., 2000a; Yang et al., 2005). The highest contents of PSP toxins were detected during the winter months (Jiang et al., 2000a). Despite the long-known occurrence of PSP toxins, PSP causative species, such as *Alexandrium* spp. and *Gymnodinium catenatum*, were seldom recorded in phytoplankton survey (Zhou et al., 1998; Liu et al., 2006). However, abundant *Alexandrium* cyst were recorded in surface sediments from this area, and it was suggested that high PSP toxin levels in bivalves during the winter months were probably due to ingestion of cysts of

Alexandrium (Wang et al., 2004). Therefore, we set up sediment traps at two stations of Daya Bay, bred shellfish synchronously at the same areas, and conducted phytoplankton surveys. For phytoplankton community and its relationships with environmental parameters, see Wang et al. (2009). In this study, we analyzed the annual data of *Alexandrium* vegetative cell and its cyst, and the PSP concentrations in shellfish. The purpose is to investigate the seasonal variation in the population size of *Alexandrium* (cell density and cyst flux) and their relationship with PSP levels in shellfish.

2 Materials and Methods

2.1 Site description

Daya Bay is located in the southeast part of the South China Sea (22°30′-22°50′N, 114°30′-114°50′E). It is a semi-enclosed shallow embayment with depth ranging from 6 m to 15 m. It covers an area of 650 km² and is separated into several small sub-basins. The bay has a mild and wet subtropical climate with an annual mean air temperature of 22 °C. The coldest months are January and February, with a monthly mean air temperature of about 15 °C, and the hottest months are July and August, with a monthly mean air temperature around 28 °C. Salinity usually lies between 22 and 33 with little fluctuation except during the typhoon season from July to September. The Daya Bay Nuclear Power Station (DNPS), the first nuclear power station in China, has been operating since February 1994, and it discharges cooling water at the rate of 2.9×10^7 m³/year (Liu et al., 2006).

Sampling stations (Fig. 1) were located in two cultural areas of the bay. St. 1 was located at 22°33′25.7″N, 114°30′54.1″E, where *Pteria* spp. were bred in suspended cages. St. 2 was at a fish cage area (22°33′51.7″N, 114°31′20.2″E), which was closer to the shore and close to the residential area.

2.2 Nutrients and hydrographic parameters

Temperature (T), Salinity (S) and dissolved oxygen (DO) were recorded *in situ* with an YSI meter (YSI 85, YSI Incorporated, USA) once every three days. Transparency and nutrients were measured every two weeks. Transparency was measured with a Secchi disk.

Water samples for nutrient determination were collected in the subsurface water column (0.5 m), immediately filtered through glass fibre filters (Whatman GF/F, porosity=0.7 μ m), and placed in a freezer. Nutrients, including dissolved inorganic nitrogen (DIN, the sum of ammonium, nitrite and nitrate), dissolved inorganic phosphorus (DIP, orthophosphate) and dissolved silicon (DSi, silicate), were analysed following standard methods of State Oceanic China (SOC, 1991) within 24 hours.

2.3 Phytoplankton sampling and observation

Water samples of 1000 mL were collected in the subsurface water column once every three days for phytoplankton observation and fixed immediately with neutralized formalin to a final concentration of 4%. The fixed water samples were finally concentrated to 20 mL by sedimentation. Phytoplankton species were counted from aliquots of 0.5-1.0 mL sub-samples using a microscope (Leica DMIRB) at magnifications of 100x to 600x. The phytoplankton abundance was presented as cell numbers per millilitre (cells/mL).

The thecal plate morphology of *Alexandrium* cells was examined using phase contrast and UV epifluorescence microscopy after calcofluor staining (Fritz and Triemer, 1985) under a Leica DMIRB microscope. The thecal plate tabulation and the shapes of plates and apical pore complex proved that the dominant *Alexandrium* species was *A. tamarense*.

2.4 Sediment collection and cyst observation

The sediment trap was set to collect settling materials from March 24, 2005 to June 10, 2006. The trap consisted of a cylindrical PVC collector (height 35cm, diameter 7cm, height/diameter ratio 5), flexible tube holder, subsurface buoy, surface buoy marker, and anchor. The trap was moored at 1.0 m from the sea bottom (depth at the deployment location was about 6 m).

Trap samples were collected every two weeks. The settled material was suspended in 2000 mL seawater, and the suspension was divided into two 1000 mL aliquots, one of which was fixed with 4% buffered formalin and subsequently concentrated to 50 mL by sedimentation.

Sediment samples were gently sonicated for 30 seconds. The sonicated material was then

sieved through 125 μm and 20 μm metallic screens to remove larger particles and detritus, and the slurry remaining on the 20 μm screen was washed into a beaker and collected into a plastic tube to a final sample volume of 10 mL with filtered seawater. For observation, 0.1 to 0.5 mL aliquots of sample were placed on a 1 mL counting chamber and diluted with appropriate distilled water.

Observation was under an inverted light microscope (Leica DMIRB). Only living cysts that had not germinated were counted. At least three observations were conducted for each sample until 200 dinoflagellate cysts were observed. Mean daily cyst flux (cysts/cm² d) was calculated. Four seasons during the survey were defined as: spring, March to May; summer, June to August; autumn, September to November; and winter, December to next February.

2.5 Shellfish breeding and PSP analysis

2.5.1 Shellfish breeding and sampling

At the beginning of the survey on March 24, 2005, 600 scallops (*Chlamys nobilis*) were bought from the local shellfish farm. Half of the scallops were placed in lantern nets and hung from long lines 3 m below the surface in each of the two stations. Green mussels (*Perna viridis*) were collected from fishing crossrails and ropes and shellfish cages. About 10 scallops (mean shell length of $9.48 \pm 0.44\text{cm}$) and 20 green mussels (mean shell length of $9.19 \pm 0.95\text{cm}$) were collected in each station for PSP toxin determination every two weeks. The shells were dissected immediately after being washed with a brush and rinsed with DI water. Whole soft tissues were removed from the shells and washed with DI water gently. Digestive glands were carefully removed from the rest of tissues. Both the digestive glands and other tissues were weighed and placed in plastic bags separately and stored in a freezer. During the transportation to the laboratory (within 4 h), the tissue samples were stored at $-20\text{ }^{\circ}\text{C}$.

2.5.2 Mouse bioassay

As the toxicities of most samples were below detection level, the digestive glands were treated and analysed first in order to screen out any contaminated samples. If the digestive glands were toxic to test mice, the remaining tissues of the same sample then treated and assayed. Sample treatment and mouse bioassay were performed according to the standard

AOAC procedure (Association of Official Analytical Chemists, AOAC, 1990) with the exception that samples sizes were smaller (about 20 g digestive gland homogenate, equal volume of 0.1N HCl and digestive gland homogenate). Mouse bioassays were conducted using ICR strains, 19–21 g male mice. The mice (3-6 mice per extract) were observed for 1 h, and the mean survival time was used to calculate the toxicity. When the extracts were too toxic, it was necessary to dilute the shellfish extract until the survival time was between 5 and 7 min. Bioassay results were calculated using a computer-based version of Sommer's table and correction table for mouse weight (Sommer and Meyer, 1937). The mouse bioassay was standardized using pure saxitoxin purchased from the National Research Council of Canada, Institute for Marine Biosciences (Halifax, Canada). The toxicity of the sample equals to the sum of toxicities of the digestive glands and the other tissues per kg total tissue ($\mu\text{g STX equiv./kg}$), and the toxicity of the digestive gland was expressed as $\mu\text{g STX equiv. per kg digestive gland}$.

2.5.3 HPLC analyses

Four digestive gland samples with high toxicity in mouse bioassay were selected for HPLC analysis (Table 1). Each sample was divided into two sub-samples. One sub-sample was extracted in 0.1 N HCl according to the AOAC extraction procedure as described in 2.5.2, and another was extracted in 1% (v/v) acetic acid solution according to Lawrence et al. (2004). Toxin contents and profiles were determined using the post-column oxidation method (Oshima, 1995) modified by Anderson et al. (1996). The HPLC system used was an Agilent Series 1100, fitted with an autosampler and fluorescence detector (Agilent, USA). Three separate isocratic elutions were employed to separate different sets of PSP toxins at a flow rate of 0.8 mL/min: (1) 2 mM tetrabutyl ammonium phosphate solution adjusted to pH 6.0 with acetic acid for C toxins; (2) 2 mM 1-heptanesulphonic acid in 10 mM ammonium phosphate buffer (pH 7.1) for GTX and dcGTX group; and (3) 2 mM 1-heptanesulphonic acid in 30 mM ammonium phosphate buffer (pH 7.1): acetonitrile (100:5) for the STX group. Toxin standards, GTX1-5, neoSTX, STX, dcSTX, dcGTX2/3 and C1, C2, were purchased from the National Research Council of Canada, Institute for Marine Biosciences (Halifax,

Canada). The standards solutions were diluted with 0.03 N acetic acid as required. PSP toxin profiles were calculated from the percentages of individual toxin concentrations of the total PSP toxin concentrations for each sample (mol %). The relative toxicities ($\mu\text{g STX equiv./kg}$) were calculated based on the specific toxicities given by Oshima (1995).

3 Results

3.1 Environmental conditions

Water temperature ranged from 14.0 to 32.8 °C, with the lowest recorded from December to early March and the highest from July to November. Salinity was relatively stable with an annual mean value of 29.3, except for the rainy season from April to May and from July to September.

Nutrient concentrations showed similar seasonal changes at the two stations. Concentrations of DIN ranged from 2.04 $\mu\text{mol/L}$ to 14.94 $\mu\text{mol/L}$, and showed an annual cycle characterized by higher values during spring and summer and lower values during late autumn and winter. Concentrations of DIP remained consistently low, and ranged from 0.044 $\mu\text{mol/L}$ to 1.67 $\mu\text{mol/L}$. DSi concentrations were remarkably high during the flood season between July and September, and low in winter and the early spring dry season. Annual mean values in DIN and DIP showed no significant differences between the two stations; however, DSi level was obviously higher at St. 1 than St. 2, with mean concentrations of 17.8 $\mu\text{mol/L}$ and 12.2 $\mu\text{mol/L}$, respectively.

3.2 Abundance of *A. tamarensis* vegetative cells

The vegetative cells of *A. tamarensis* occurred sporadically in the water column and showed a regular seasonal cycle. Vegetative cells only occurred during spring (March to May) and winter (November to February) (Fig. 2). The abundance during spring was low, ranging from 0.2 to 1.4 cells/mL. The vegetative cells were not detected during summer and autumn (Fig. 2) when water temperature exceeded 25 °C. Cell numbers increased at the end of November with decreasing water temperature and peaked in January and February, when water temperatures were around 20 °C. Cell numbers were markedly higher in St. 1 than in St.

2, and the maximum cell number of 603 cells/mL was observed in St. 1 on February 19, 2006. Maximum number coincided with low water temperature and low nutrient concentrations during the winter months.

The phytoplankton assemblages were dominated by diatoms, which constituted 93.2% of total phytoplankton population. Dinoflagellates occurred in low numbers with a mean abundance of 9.4 cells/mL, and represented only 1.2% of total abundance. The abundance of total phytoplankton was 3 920 cells/mL when the maximum cell number of *A. tamarens* occurred. The most dominant species was *Skeletonema costatum* at that time, with a cell density of 3 230 cells/mL and proportion of 82.5%. *A. tamarens* accounted for 98.3% of dinoflagellate, but only 15.3% of total phytoplankton.

3.3 Cyst flux of *A. tamarens*

Cysts of *A. tamarens* occurred frequently in the sediment traps, but in low numbers (mostly <5 cysts/cm² d). Cysts appeared to have a seasonal cycle similar to that of the vegetative population and occurred more abundantly during spring and winter (Fig. 2, Fig. 3). High cyst fluxes were observed approximately two weeks to one month after the vegetative cell peaks. For example at St. 1, there was a significant increase in cyst flux in January 2006 just following vegetative cell peaks in December and January, and the maximum flux of 42.6 cysts/cm² d was observed on February 11-26, 2006, right after the maximum number of vegetative cells. Meanwhile, a strong positive correlation existed between cyst flux and the cumulative number of vegetative cell ($\text{Cysts} = 6.64 \times 10^{-5} \times \text{Cells} + 0.77$, $r = 0.995$, $n = 27$, $p < 0.0001$).

3.4 PSP toxins in shellfish

The extracts of shellfish tissues in the present study killed mice with typical symptoms associated with PSP toxins. Seasonal variations in toxin concentration are illustrated in Fig. 4. The PSP toxins in the two stations were generally below the regulatory limit, except for the winter months. The concentrations of PSP toxins were lower in spring and early summer, and PSP toxins were not detected between August and October. Toxin concentrations increased significantly in January, and a maximum level of 14 015 µg STX equiv./kg was recorded in *P.*

viridis samples collected from St. 1 on January 3, 2006 (Fig. 4). The majority of toxins were detected in digestive glands, with the maximum of 66 227 µg STX equiv./kg in digestive glands from *C. nobilis* collected at St. 1 on January 3, 2006 (Fig. 5). Toxins in other tissues were very low to undetectable except for five samples collected in January 2006, with a maximum level of 392 µg STX equiv./kg. The results showed that the hepatopancreatic tissues from both *P. viridis* and *C. nobilis* were unsafe for human consumption during the winter months, while the tail muscle would be safe.

Toxin levels in digestive glands from *C. nobilis* (mean value of 3 373 µg STX equiv./kg) were significantly higher than those from *P. viridis* (mean value of 1 531 µg STX equiv./kg). However, the contribution of digestive gland to total tissues is lower in *C. nobilis* than in *P. viridis*. Therefore, toxin concentrations in total tissues of the two shellfish were similar, with mean values of 399 and 394 µg STX equiv./kg in *C. nobilis* and *P. viridis*, respectively.

The sharp increase in concentrations of PSP toxins in January was coincident with the occurrence of *Alexandrium* population (both vegetative cells and cysts), with peak concentrations or abundances between January and March. PSP concentrations were clearly higher in St. 1 than in St. 2, and the averages in digestive glands and total tissues were 3 796 and 632 µg STX equiv./kg in St. 1 and 1 107 and 161 µg STX equiv./kg in St. 2, respectively.

3.5 PSP toxin profiles

Fig. 6 illustrates the toxin profiles in four digestive gland samples treated by HCl and acetic acid, respectively. Of the 12 components analyzed in this study, GTX5 and dcGTX2 were not detected in any of the samples, and only trace amounts of decarbamoyl toxins were detected with a mean proportion of 2.2%. The PSP toxin profiles in HCl extracts were rich in carbamate toxins (mean 63.7% of the total) and dominated by GTX2 (mean 35.1%), GTX3 (mean 26.2%) and GTX4 (mean 15.7%). C toxins accounted for a medium amount of the total (around 0-12% for C1, 1-7% for C2). In addition to these components, a trace amount of decarbamoyl toxins (mean 3.3%) was detected. In contrast, in acetic acid extracts, large amounts of C toxins were detected instead, and C1 (around 38-68% of the total, mean 53.6%) was the most abundant. On the other hand, concentrations of GTX toxins dropped

significantly and accounted for 4-31% of the total, with an average of 17.8%. The relative concentrations of STX toxins (including STX, dcSTX and neoSTX) in acetic acid extracts (mean 22.1%) were similar to those in HCl extracts (mean 18.4%); however, percentages of STX increased from 6.7% in HCl extracts to 13.9% in acetic acid extracts on average.

Though samples collected from St. 1 on 3 January 2006 were most toxic during study period (Fig. 4, 5), the toxin profiles (sample 1 and 1') showed less toxic GTX2 and C1 (Fig. 6). Other less toxic samples (samples 2-4, and 2'-4') had higher proportions of more toxic components such as STX, neoSTX and GTX4. However, toxin concentrations on a molar basis ($\mu\text{mol/kg}$) in sample 1 and 1' were 5-8-fold greater than those in other samples (Table 1).

Toxin concentrations on a molar basis ($\mu\text{mol/kg}$) were comparable between any two parallel samples treated by HCl or acetic acid (Table 1). The toxicities in HCl extracts approximately coincided with those analyzed by mouse assay. However, samples treated by acetic acid showed only one-third to two-thirds of those treated by HCl, especially in samples with rich contents of C1 toxin (Table. 1, Fig. 6).

4 Discussion

High abundance of *A. tamarensis* vegetative cells was observed in this study, with a maximum number of 603 cells/mL. In previous phytoplankton studies, vegetative cells of *Alexandrium* were seldom observed in the water column during seasonal routine observation due to the long interval of sampling (Liu et al., 2006). However, peak abundance of *A. tamarensis* was captured in this survey by highly frequent sampling (once every three days).

Alexandrium tamarensis showed a high dispersal of temperature/cell density couples (Fig. 7). It never occurred at temperatures greater than 30 °C, and peak abundances occurred when temperature lies between 16 and 20 °C. *A. tamarensis* is a stenohaline alga (ca. 15-25 °C) with optimal growth at 17-22 °C (Yan et al., 2002). The blooms of *Alexandrium* usually occurred in temperatures between 13 and 18 °C (Giacobbe et al., 1996; Gracc et al., 2004). After the operation of DNPS, the annual mean water temperature increased by 0.4-1.1 °C (Tang et al.,

2003), and the variation in temperature was more significant in winter months, increasing from 16.04 °C to 18.03 °C (Wang et al., 2008), which seem to favor the growth of *Alexandrium*. Meanwhile during this period, nutrient loadings had increased significantly in the bay (Wang et al., 2008), facilitating the rapid growth of phytoplankton, especially the small diatoms with high growth rates (Liu et al., 2006; Wang et al., 2009). The phytoplankton abundances peaked in spring and summer during the early 1990s (Zhou et al., 1998), but frequent winter peaks have been observed since the late 1990s (Liu et al., 2006; Wang et al., 2009). Generally, Daya Bay is characterized by high levels and rapid compensation of nutrients, however, temporary shortage of nutrients occurs when the diatom bloom exhausts the nutrients, particularly during the dry low-nutrient winter season (Wang et al., 2009). Dinoflagellates such as *Alexandrium* are more competitive in nutrient depletion conditions due to its higher nutrient affinities and capability to utilize organic nutrients (Collos et al., 2007). As a result of higher winter temperatures and increased nutrient loading, *Alexandrium* has occurred more frequently and abundantly during winter months, which contributes to the high PSP concentrations in shellfish.

Cyst fluxes were recorded during summer and autumn (June to October), when no vegetative cells of *A. tamarense* were observed in the water column. The absence of vegetative cells in the subsurface water column during the observation does not necessarily indicate their absence in the entire water column. More likely, the water sample collection did not catch *A. tamarense* cell patches due to random sampling. In addition, this period is also the typhoon season in South China Sea. Strong storms may cause a high resuspension rate, which could result in redeposition of bottom cysts into sediment traps.

Similar to the total phytoplankton abundance, cell density and cyst flux of *A. tamarense* were clearly higher in St. 1, located in a shellfishery area, than those in St. 2, located in a fish cage area. This resulted in higher PSP levels in shellfish collected from St. 1. The low primary production in the fishery area was observed in another study in this area as well (Huang et al., 2005). Meanwhile, higher concentrations of *Alexandrium* cyst were recorded in surface sediments from the shellfishery area in the same sea area (Wang et al., 2004). This

phenomenon may be interpreted as compensatory growth of phytoplankton against predatory pressure by shellfish.

Daya Bay has been reported to have the highest PSP concentrations among the Chinese coastal areas (Lin et al., 1999; Yang et al., 2005). Jiang et al. (2000a) found that the highest concentration of PSP toxins occurred in January during their survey conducted between 1997 and 1999. However, *Alexandrium* was seldom observed in previous phytoplankton surveys (Liu et al., 2006). Our results showed that high cell numbers and cyst fluxes of *Alexandrium* coincided with high PSP levels in shellfish. In addition, content of PSP toxins showed a strong positive relationship with the accumulative abundances of vegetative cells and cysts [$\ln(\text{PSP}) = 0.41 \times \ln(\text{Cells}) + 2.34$, $r = 0.647$, $n = 27$, $p = 0.002$; $\ln(\text{PSP}) = 0.37 \times \text{Cysts} + 5.22$, $r = 0.373$, $n = 27$, $p = 0.04$]. These results suggest that vegetative cells and cysts of *Alexandrium* influenced significantly the PSP levels, and could have contributed to high PSP levels in shellfish during winter.

However, toxin levels were not always consistent with concentrations of vegetative cells and cysts. The inconsistency of PSP contamination in shellfish and concentration of *Alexandrium* cells and cysts might be explained by the variability of *Alexandrium* toxicity, which depends on environmental conditions, among species and strains (Anderson, 1990). Toxicities in *Alexandrium* vegetative cells and cysts seem to be species and strain specific. The average toxicity of *A. tamarense* vegetative cells is approximately 16 pg STX equiv./cell (Persson et al., 2006). Oshima et al. (1992) reported as much as 269 pg STX equiv./cyst in *A. tamarense*, and Bricelj and Shumway (1998) reported 325 pg STX equiv./cell of an *Alexandrium* sp. Beppu et al. (2008) measured the toxicities of six strains of *Alexandrium* from the coastal water of Hiroshima Prefecture, Japan, and found that variation in the toxin level was varied about 30 fold among strains (from 3.7 to 111.5×10^{-6} MU/cell), or 10 fold even between *A. tamarense* strains isolated from the same sea area in the same month of different years.

The toxin profiles of HCl treated samples were similar to those reported by Jiang et al. (2000b) in the same sea area, in which GTX toxins were found to be the main component.

GTX toxins were dominant in contaminated shellfish as determined by the AOAC HCl extraction method in other geographic regions (Lassus et al., 1989; Ichimi et al., 2001; Abouabdellah et al., 2008). Toxin profiles of acetic acid extracts showed good similarity with those of *Alexandrium* strains isolated from the nearby sea areas (Jiang et al., 2000b; Wang et al., 2006), which were predominated by C1/C2 toxins and had only trace amounts of STX toxins. The results suggested PSP toxins in filter-shellfish detected in this study were possibly produced by a local strain. It is well known that the AOAC extraction procedure used in the sample preparation converts the sulphocarbamoyl toxins (C1–4, B1, B2) into corresponding carbamates (Laycock et al., 1995; Anderson et al., 1996), i.e., C3,4 to GTX1,4 and C1,2 to GTX2,3 (Turrell et al., 2007). Our results also showed evident toxin conversion through the HCl procedure, as displayed by the remarkable increase of GTX2, 3 and sharp decrease of C1, and meanwhile a comparatively stable overall toxin concentrations on molar basis (mol/kg). The carbamates are considerably more toxic than the corresponding sulphocarbamoyls, and, consequently, changes in toxin profile after HCl treatment will increase the toxicity of the shellfish extract (Kao, 1993). In our study, toxicities of HCl extracts were 1.5-3.5-fold higher than those of corresponding acetic acid extracts. Despite the increase toxicity through HCl procedure, we suggest using HCl extracts in the mouse bioassay. This procedure better protects human health by considering the potential conversion of C toxins into more poisonous GTX toxins in human body.

5 Conclusion

Our one-year observation found that high PSP toxin levels in shellfish coincided with peak abundances of cell and cyst of *A. tamarense* during winter months, though the toxin levels were generally below the regulatory limit for most seasons of the year. The significant, positive relationship between toxin level, and vegetative cell and cyst abundance reveals possible toxin source from *Alexandrium* thriving during winter months. More specifically, the majority of the toxins were accumulated in the digestive glands.

353 **Acknowledgements**

354 This research was funded by the National Natural Science Foundation of China (41076093,
355 40306020), and the Fundamental Research Funds for the Central Universities. We are very
356 grateful to Miss Na Wang of Xiamen University for HPLC analysis.

References

- Abouabdellah, R., Taleb, H., Bennouna, A., Erler, K., Chafik, A., Moukrim, A., 2008. Paralytic shellfish poisoning toxin profile of mussels *Perna perna* from southern Atlantic coasts of Morocco. *Toxicon* 51, 780–786.
- Anderson, D.M., 1997. Diversity of harmful algal blooms in coastal waters. *Limnol. Oceanogr.* 42, 1009–1022.
- Anderson, D.M., Kulis, D.M., Sullivan, J.J., Hall, S., 1990. Toxin composition variations in one isolate of the dinoflagellate *Alexandrium fundyense*. *Toxicon* 28, 885–893.
- Anderson, D.M., Kulis, D.M., Qi, Y., Zheng, L., Lu, S., Lin, Y., 1996. Paralytic shellfish poisoning in Southern China. *Toxicon* 14, 579–590.
- AOAC, 1990. Paralytic shellfish poison. Biological method. Final action. In: Hellrich, K. (Eds.), *Official Methods of Analysis*, 15th ed., Sec. 959.08. Association of Official Analytical Chemists, Arlington, Virginia, USA, pp. 881–882.
- Beppu, R., Najima, K., Tsuruda, S., Gomez-Dalan, G., Narte-Quilantang M., Taniyama S., Sagara, T., Nishio, S., Takayama, H., Miyazawa, K., Asakawa, M., 2008. Occurrence of PSP-producing dinoflagellate *Alexandrium tamiyavanichii* in Bingo-Nada, the central coastal water of Seto Inland Sea, Hiroshima Prefecture, Japan. *Mar. Pollut. Bull.* 56, 758–763.
- Bravo, I., Franco, J.M., Reyero, M.I., 1998. PSP toxin composition of three life cycle stages of *Gymnodinium catenatum*. In: Reguera, B., Blanco, J., Ferná'ndez, M.L., Wyatt, T. (Eds.), *Harmful Algae*. Xunta de Galicia and IOC of UNESCO, Grafisant, Santiago de Compostela, pp. 356–358.
- Bricelj, V.M., Shumway, S.E., 1998. Paralytic shellfish toxins in bivalve molluscs: occurrence, transfer kinetics, and biotransformation. *Rev. Fish. Sci.* 6, 315–383.
- Collos, Y., Vaquer, A., Laabir M., Abadie, E., Laugier, T., Pastoureaud, A., Souchu, P., 2007. Contribution of several nitrogen sources to growth of *Alexandrium catenella* during blooms in Thau lagoon, southern France. *Harmful Algae*, 6, 781–789.

- 384 Fritz, L., Triemer, R.E., 1985. A rapid simple technique utilizing Calcofluor White M2R for
385 the visualization of dinoflagellate thecal plates. *J. Phycol.* 21, 662–664.
- 386 Giacobbe M, Goliva D, Maimone G. 1996, Environmental factors and seasonal occurrence of
387 the dinoflagellate *Alexandrium minutum*, a PSP potential producer, in a Mediterranean
388 lagoon. *Estuar. Coast. Shelf Sci.* 42, 539-549.
- 389 Gracéc, E., Bravo, I., Vila, M., Figueroa, R.I., Masó, M. and Sampedro, N. 2004. Relationship
390 between vegetative cells and cyst production during *Alexandrium minutum* bloom in
391 Arenys de Mar harbour (NW Mediterranean). *J. Plankton Res.* 26, 637-645.
- 392 Hallegraeff, G. M., 1995. Harmful algal blooms: a global overview. In: Hallegraeff, G.M.,
393 Anderson, D.M., Cembella, A.D. (Eds.), *Manual on Harmful Marine Microalgae*. IOC
394 Manuals and Guides No. 33. UNESCO, pp. 1–22.
- 395 Huang, H., Lin, Q., Wang, W, Jia, X., 2005. Impact of cage fish farming on water
396 environment in Daya Bay. *South China Fish. Sci.* 1 (3), 9-17 (in Chinese, with English
397 abstract).
- 398 Ichimi, K., Suzuki, T., Yamasaki, M., 2001. Non-selective retention of PSP toxins by the
399 mussel *Mytilus galloprovincialis* fed with the toxic dinoflagellate *Alexandrium tamarense*.
400 *Toxicon* 39, 1917–1921.
- 401 Jian, Y., Deng, F., 1991. Analysis of seafood poison caused by clam *Pinna pectinara*. *Res.*
402 *Dev. South China Sea* 4, 42–45 (in Chinese, with English abstract).
- 403 Jiang, T., Yin, Y., Luo, Y., Chen, J., Qi, Y., 2000a. Study on the paralytic poisoning in
404 shellfish of Daya Bay and Dapeng Bay, Guangdong. *China Environ. Sci.* 20, 341-344 (in
405 Chinese, with English abstract).
- 406 Jiang, T., Yin, Y., Chen, Z., Luo, Y., Huang, W., Qi, Y., 2000b. HPLC analysis of paralytic
407 shellfish toxins in shellfish from Daya Bay. *Mar. Environ. Sci.* 19(3), 16–19 (in Chinese,
408 with English abstract).
- 409 Kao, C.Y., 1993. Paralytic shellfish poisons. In: Falconer, I.R. (Eds.), *Algal Toxins in*
410 *Seafood and Drinking Waters*. Academic Press, London, pp. 75–86.

- 411 Lassus, P., Fremy, J.M., Ledoux, M., Bardouil, M., Bohec, M., 1989. Patterns of experimental
412 contamination by *Protogonyaulax tamarens* in some French commercial shellfish.
413 *Toxicon* 27, 1313–1321.
- 414 Lawrence, J.F., Niedzwiadek, B., Menard, C., 2004. Quantitative determination of paralytic
415 shellfish poisoning toxins in shellfish using prechromatographic oxidation and liquid
416 chromatography with fluorescence detection: interlaboratory study. *J. AOAC Int.* 87,
417 83–100.
- 418 Laycock, M.V., Kralovec, J., Richards, R., 1995. Some in vitro chemical interconversions of
419 paralytic shellfish poisoning (PSP) toxins useful in the preparation of analytical standards.
420 *J. Mar. Biotechnol.* 3, 121–125.
- 421 Lin, Y., Jia, X., Yang, M., Zhong, Y., Quan, G., Fan, P., 1999. Paralytic shellfish poison in
422 contaminated shellfish along coast of China. *Trophic Oceanol.* 18, 90-96 (in Chinese,
423 with English abstract).
- 424 Liu, S., Huang, H., Huang, L., Lian, J., Long, A., Li, T., 2006. Ecological response of
425 phytoplankton to the operation of Daya Bay nuclear power station. *Mar. Environ. Sci.* 25
426 (2), 9-13 (in Chinese, with English abstract).
- 427 Oshima, Y., 1995. Post-column derivatization HPLC methods for paralytic shellfish poisons.
428 In: Hallegraeff, G.M., Anderson, D.M., Cembella, A.D. (Eds.), *Manual on Harmful*
429 *Marine Microalgae*. IOC Manuals and Guides No. 33. Intergovernmental Oceanographic
430 Commission (UNESCO), Paris, pp. 81–94.
- 431 Oshima Y., Bolch C.M., Hallegraeff G.M. 1992. Toxin composition of resting cysts of
432 *Alexandrium tamarens* (Dinophyceae). *Toxicon* 30, 1539-1544.
- 433 Persson, A., Smith, B.C., Wikfors, G.H., Quilliam, M., 2006. Grazing on toxic *Alexandrium*
434 *fundyense* resting cysts and vegetative cells by the eastern oyster (*Crassostrea virginica*).
435 *Harmful Algae* 5, 678–684.
- 436 Schwinghamer, P., Hawryluk, M., Powell, C., Mackenzie, C.H., 1994. Resuspended
437 hypnozygotes of *Alexandrium fundyense* associated with winter occurrence of PSP in
438 inshore Newfoundland waters. *Aquaculture* 122(2-3), 171-179.

- 439 Shumway, S.E., 1990. A review of the effects of algal blooms on shellfish and aquaculture. J.
440 World Aquat. Soc. 21, 65–104.
- 441 Shumway, S.E., Sherman-Caswell, S., Hurst, J.W., 1988. Paralytic shellfish poisoning in
442 Maine: monitoring a monster. J. Shellfish Res. 7, 643–652.
- 443 SOC (State Oceanic China). 1991. The criterion of Marine Monitoring. Oceanographical
444 Press, Beijing, HY003. 4-91. pp. 205-282 (in Chinese).
- 445 Sommer, H., Meyer, K.F., 1937. Paralytic shellfish poisoning. Arch. Path. 24, 560–598.
- 446 Tang, D., Kester, D. R., Wang, Z., Li, J., Kawamura, H., 2003. AVHRR satellite remote
447 sensing and shipboard measurements of the thermal plume from the Daya Bay, nuclear
448 power station, China. Remote Sens. Environ. 84, 506–515.
- 449 Tsujino, M., Uchida, T., 2004. Fate of resting cysts of *Alexandrium* spp. ingested by
450 *Perinereis nuntia* (Polychaeta) and *Theola fragilis* (Mollusca). J. of Exp. Mar. Biol. Ecol.
451 303, 1-10.
- 452 Turrell, E.A, Lacaze, J.P., Stobo, L., 2007. Determination of paralytic shellfish poisoning
453 (PSP) toxins in UK shellfish. Harmful Algae 6, 438–448.
- 454 Wang, D., Zhang, S., Gu, H., Chan, L., Hong, H., 2006. Paralytic shellfish toxin profiles and
455 toxin variability of the genus *Alexandrium* (Dinophyceae) isolated from the Southeast
456 China Sea. Toxicon 48, 138–151.
- 457 Wang Y S, Lou Z P, Sun C C, Sun S. 2008. Ecological environment changes in Daya
458 Bay, China, from 1982 to 2004. Marine Pollution Bulletin, 56(11): 1871-1879.
- 459 Wang, Z., Matsuoka, K., Qi, Y., Chen, J., Lu, S., 2004. Dinoflagellate cyst records in recent
460 sediments from Daya Bay, South China Sea. Phycol. Res. 52, 396-407.
- 461 Wang, Z., Zhao, J., Zhang Y., Cao, Y., 2009. Phytoplankton community structure and
462 environmental parameters in aquaculture areas of Daya Bay, South China Sea. J Environ.
463 Sci. China. 21, 1268-1275.

- 464 Yan, T., Zhou, M. J., Qian, P. Y., 2002. Study on the combined effects of temperature,
465 salinity and irradiance on the growth of dinoflagellate *Alexandrium tamarense*. Acta
466 Oceanol. Sin. 24(2), 114-120 (in Chinese, with English abstract).
- 467 Yang, M., Lin, Q., Lu, X., Cai, W., 2005. The paralytic shellfish poison content in the oysters
468 in important maricultural waters along Guangdong Coast. Mar. Environ. Sci. 24(1), 48-50
469 (in Chinese, with English abstract).
- 470 Yentsch, C.M., Mague, F.C., 1979. Motile cells and cysts: two probable mechanisms of
471 intoxication of shellfish in New England waters. In: Taylor, D.L., Seliger, H.H. (Eds.),
472 Toxic Dinoflagellate Blooms. Elsevier/North-Holland, pp. 127–130.
- 473 Zhou, P., Lin, Y., Wang, Z., 1998. A statistical analysis on phytoplankton community features
474 in waters of Daya Bay. Tropic Oceanol. 17 (3), 57-64 (in Chinese, with English abstract).

Tables

Table 1 PSP toxin contents in digestive gland samples by HPLC analysis and mouse bioassay using the AOAC HCl procedure and acetic acid extraction procedure, respectively

	1	1'	2	2'	3	3'	4	4'
Toxin concentration by								
HPLC	191.5	173.4	18.6	15.3	23.5	18.6	22.8	24.4
($\mu\text{mol/kg}$)								
Toxicity by HPLC								
analysis	41 584	11 993	4 477	1 842	5 377	3 424	6 135	2 832
($\mu\text{g STX equiv./kg}$)								
Toxicity by mouse								
bioassay	51 565	-	6 636	-	6 780	-	5 551	-
($\mu\text{g STX equiv./kg}$)								

1 and 1': *P. viridis* samples from St. 1 on January 3, 2006, 2 and 2': *P. viridis* samples from St. 1 on January 22, 2006, 3 and 3': *P. viridis* samples from St. 1 on February 26, 2006, 4 and 4': *C. nobilis* samples from St. 1 on March 16, 2006. Samples 1-4 were treated by HCl, samples 1'-4' were treated by acetic acid. "-" indicates without bioassay data.

Figure captions

Fig. 1 Sampling stations in Daya Bay, South China Sea. Also shown is the location of the Daya Bay Nuclear Power Station (DNPS). St. 1 was located at a shellfish cultural area, and St. 2 was at a fish cage area close to a residential area.

Fig. 2 Seasonal changes in cell density of *Alexandrium tamarense*

Fig. 3 Seasonal changes in cyst flux of *Alexandrium tamarense*

Fig. 4 Concentrations of PSP toxins in shellfish

Fig. 5 Concentrations of PSP toxins in digestive glands of shellfish

Fig. 6 PSP toxin profiles in digestive glands from shellfish samples. 1 and 1': *P. viridis* samples from St. 1 on January 3, 2006, 2 and 2': *P. viridis* samples from St. 1 on January 22, 2006, 3 and 3': *P. viridis* samples from St. 1 on February 26, 2006, 4 and 4': *C. nobilis* samples from St. 1 on March 16, 2006. Samples 1-4 were treated by HCl, samples 1'-4' were treated by acetic acid.

Fig. 7 Relationship between cell density of *Alexandrium tamarense* and water temperature.

Title: Source and profile of paralytic shellfish poisoning toxins in shellfish in Daya Bay, South China Sea

Highlights

> Annual survey on *Alexandrium* population and PSP was conducted. > *Alexandrium* cells and cysts occurred abundantly during the winter months. > PSP contents in shellfish were generally low, but high in winter. > *Alexandrium* thriving during winter months could be possible toxin source in shellfish.

Fig. 1 Sampling stations in Daya Bay, South China Sea. Also shown is the location of the Daya Bay Nuclear Power Station (DNPS). St. 1 was located at a shellfish cultural area, and St. 2 was at a fish cage area close to a residential area.

Fig. 2 Seasonal changes in cell density of *Alexandrium tamarense*

Fig. 3 Seasonal changes in cyst flux of *Alexandrium tamarensis*

Fig. 4 Concentrations of PSP toxins in shellfish

Fig. 5 Concentrations of PSP toxins in digestive glands of shellfish

Fig. 6 PSP toxin profiles in digestive glands from shellfish samples. 1 and 1': *P. viridis* samples from St. 1 on January 3, 2006, 2 and 2': *P. viridis* samples from St. 1 on January 22, 2006, 3 and 3': *P. viridis* samples from St. 1 on February 26, 2006, 4 and 4': *C. nobilis* samples from St. 1 on March 16, 2006. Samples 1-4 were treated by HCl, samples 1'-4' were treated by acetic acid.

Fig. 7 Relationship between cell density of *Alexandrium tamarense* and water temperature.