Accepted Manuscript

Title: Complex phase distribution and seismic velocity structure of the transition zone: convection model predictions for a magnesium-endmember olivine-pyroxene mantle

Authors: Michael H.G. Jacobs, Arie P. van den Berg

PII: S0031-9201(11)00042-2
DOI: doi:10.1016/j.pepi.2011.02.008
Reference: PEPI 5384

To appear in: *Physics of the Earth and Planetary Interiors*

Received date: 27-8-2010
Revised date: 12-2-2011
Accepted date: 24-2-2011

Please cite this article as: Jacobs, M.H.G., van den Berg, A.P., Complex phase distribution and seismic velocity structure of the transition zone: convection model predictions for a magnesium-endmember olivine-pyroxene mantle, *Physics of the Earth and Planetary Interiors* (2008), doi:10.1016/j.pepi.2011.02.008

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 Complex phase distribution and seismic velocity structure of the
2 transition zone: convection model predictions for a
3 magnesium-endmember olivine-pyroxene mantle

4 Michael H.G. Jacobs, Institut für Metallurgie, TU Clausthal, Clausthal-Zellerfeld, Germany
Arie P. van den Berg, Dept. Theoretical Geophysics, Institute of Earth Sciences,
Utrecht University, Utrecht, The Netherlands

Dedicated to Bernard de Jong on the occasion of his seventieth birthday

5 February 12, 2011

6 1

¹Projects/Multiplephasetransitions/pepi1102_revision

Abstract

We have investigated the complex spatial distribution of mineral phases in the Earth's upper mantle, resulting from lateral variations of temperature in a convecting mantle, through numerical modelling of mantle convection. Our convection model includes a self-consistent thermodynamic description for an olivine-pyroxene composition in the SiO_2 , MgO system. The thermodynamic model is based on lattice vibrations and allows for the calculation of thermophysical properties as well as seismic wavespeeds.

Our modelling results show a complex structure in the behavior of physical properties, in particular the seismic shear wavespeed, in a depth range including the mantle transition zone, 400-700 km. We demonstrate that this behavior is related to the distribution of mineral phases in the olivine-pyroxene system. Especially near cold downwelling flows, representing subducting lithospheric plates, our model results show strong lateral variation of mineral phases and associated shear wavespeed. We show that, typically, pockets of contrasting mineral phases smaller than 100 km occur in subduction regions.

In line with current developments in seismic imaging of the mantle transition zone we have computed reflectivity profiles from the shear wavespeed distribution obtained from the convection results. We applied frequency filtering to the raw reflectivity data to investigate the requirements for resolving the heterogeneous structure of the transition zone. Our results show that heterogeneous structure from contrasting mineral phase regions is resolved in the reflectivity profiles, for periods below twenty seconds which may be feasible in seismic imaging applications. This opens a perspective for detailed mapping of mineral phase distribution in the mantle offering new constraints on the thermal conditions of the transition zone region, thus providing valuable constraints for geodynamic models of the upper mantle.

1 Introduction

The mantle transition zone in the depth range 400-700 km is characterized by a large number of mineral phases especially in regions with strong lateral temperature variation, related to subducting lithospheric slabs. Seismological mapping of the existing phase boundaries provides constraints on the geotherm in such regions from identification with phase diagram data for candidate mantle materials. This way Boehler (1996) estimated the temperature at the spinel to post-spinel '660 km' phase boundary at 1900 ± 100 K, based on a pressure of 23.9 GPa from the PREM model (Dziewonski and Anderson, 1981). A recent example of this procedure, extended to lateral variation, is an estimate of temperature distribution in the heterogeneous D'' region in the bottom region of the mantle, from high resolution imaging of seismic reflectors, identified as the post perovskite phase boundary (Van der Hilst et al., 2007).

Lateral variation of the upper mantle transition zone has been investigated seismologically using underside reflections, observed as precursors of SS and PP phases (Shearer et al., 1999, Deuss and Woodhouse, 2001, Deuss et al., 2006), resulting in estimates of lateral variation of the depth of the '410' and '660 km' phase boundaries and regional variation in the visibility of the '520 km' phase boundary, associated with the wadsleyite (β) to ringwoodite (γ) transition in the (main) olivine component of the upper mantle. Recently Cao et al. (2010) applied seismic imaging methods, based on the Generalized Radon Transform (GRT) and related to the approach in (van der Hilst et al., 2007) and (Wang et al., 2008), to SS precursors originating from the transition zone for a tectonically undisturbed region of the Pacific. This study shows clear resolution of the topography of the '410' and '660 km' boundaries and laterally varying intermediate structure interpreted as the '520 km' discontinuity that is shown to resolve locally in two separate reflections in the shorter period ranges of the signal. This could be an indication for additional phase boundaries related to

mineral phases from the pyroxene system such as majorite and akimotoite that would be stable for possible temperature conditions in the transition zone.

The above examples illustrate that seismic visibility of mineral phase boundaries in complex configurations, as predicted for the mantle transition zone, is of great importance for constraining local temperature, which in turn is of great value for dynamical reconstructions of the upper mantle. In this paper we investigate this issue, of the seismic expression of the mineral distribution in the transition zone, by means of numerical modelling of mantle convection, taking into account physically self-consistent thermophysical parameters and mineral phase diagram of the olivine-pyroxene system. To our knowledge no mantle convection studies have addressed the prediction of heterogeneous seismic structure from the distribution of mineral phases in the transition zone, with the exception of (Jacobs et al., 2006) restricted to an olivine mantle. Here our mantle convection model is based on the magnesium endmember of the olivine-pyroxene mineralogy (Jacobs and de Jong, 2007). The same model, further introduced in section 2.2 and the appendix, is also used to compute longitudinal and shear wave speed. Other workers have computed seismic wave speed distributions from mantle convection results without similar consistency between the mineral physics and the convection model (Goes et al., 2004), (Schuberth et al., 2009a,b). We have chosen to apply the mineral physics model of Jacobs and de Jong (2007), in spite of its restricted composition, because of its degree of internal consistency and accuracy in representing available experimental data, that is superior to alternative mineral physics models that include a wider range of compositions, as is further illustrated below in section 2.2. The extension of our mineral physics model for a broader range of compositions, in particular iron, aluminium and calcium, is work in progress (Jacobs and de Jong, 2009).

The magnesium endmember olivine-pyroxene model is a simplified model relative to a more realistic, for instance pyrolitic, mantle composition (Ringwood, 1975), yet it contains the most significant features of the mantle transition zone including the major phase transitions. The monovariant phase transitions of the magnesium-endmember system produce corresponding sharp transitions in physical properties including the seismic wave velocities. We use this endmember model for an exploratory test (1) to investigate the applicability of a self-consistent thermodynamic model in mantle convection computations, (2) to couple the self-consistently calculated sound wave velocities with mineral phase partitioning in the mantle and (3) to investigate the expression of the different mineral phases in seismic reflectivity traces.

In line with the above we have computed model predictions of the complex spatial distribution of mineral phases and corresponding seismic wave velocities resulting from our convection model for the whole mantle, in particular for the transition zone region. We convert the convection results (spatial distribution of pressure and temperature) to shear velocity values for the upper 1000 km of the mantle and correlate the resulting heterogeneous seismic velocity structures with corresponding variations of both temperature and mineral phase distribution. To put the results in perspective with recent developments in seismological reflectivity imaging of mantle structure, we compute 1-D radial reflectivity profiles from our convection results and consider the effect of limited seismic data bandwidth on the representation of the heterogeneous structure by reflectivity time traces. This modelling procedure can help in investigating the potential of seismological resolution of the detailed structure of the transition zone. By focusing on the transition zone we avoid strong compositional heterogeneity of other mantle regions such as the D'' zone at the bottom of the mantle, not accounted for in our model (Xu et al., 2008).

Seismic expression of temperature variations in mantle convection models has been investigated by Goes et al. (2004) who focus on large scale mantle plumes, and Schubert et al. (2009a,b), who focus on lateral variation in 3-D global convection models in comparison with evidence from global

February 12, 2011

4

seismic tomography. In contrast we focus on modelling small scale heterogeneity in the transition zone and its expression in high resolution seismic imaging methods.

In the next sections we first describe the convection model and underlying parameterization of the mineral physics model. This is followed by a presentation of the results from convection modelling and a discussion of the implications of the results and conclusions.

2 Description of the convection and thermodynamical model

2.1 Convection model

We apply a model of a compressible mantle, taking into account the effects of density increase with pressure and solid state phase transitions. The model is defined for a two-dimensional cylindrical domain geometry. The governing equations are,

$$\partial_j(\rho u_j) = 0 \quad (1)$$

expressing mass conservation using a truncated anelastic liquid approximation (Jarvis and McKenzie, 1981, Steinbach et al., 1989, Steinbach, 1991, Ita and King, 1994). Conservation of momentum is expressed in the Stokes equation for a highly viscous fluid of infinite Prandtl number,

$$-\partial_i \Delta P + \partial_j \tau_{ij} + \Delta \rho g(r) e_i = 0, \quad i = 1, 2 \quad (2)$$

where $g(r)$ and e_i are, respectively, the amplitude and direction cosines of the local radial gravity acceleration vector. The density perturbation $\Delta \rho$ in (2) is defined in terms of a reference profile ρ_r ,

$$\rho(P, T) = \rho_r(P) + \Delta \rho(P, T) \quad (3)$$

where $\rho_r(P) = \rho(P, T_r(P))$ and $T_r(P) = \langle T(P) \rangle$ is a reference temperature profile computed from a lateral average. The dynamic pressure gradient in (2) is defined by subtracting $\rho_r(P) g_i$ from the gradient of pressure $\partial_i P$ in the Stokes equation.

Assuming that the effect of the rate of volume change, that scales with a bulk viscosity is negligible on the convective time scale (Jarvis and McKenzie, 1980), the deviatoric shear stress tensor for the compressible fluid is defined as,

$$\tau_{ij} = \eta \left(\partial_i u_j + \partial_j u_i - \frac{2}{3} \partial_k u_k \delta_{ij} \right) \quad (4)$$

where the summation convention is implied for repeated indices and η is the temperature and pressure dependent viscosity further specified in Table 1.

The energy transport equation used is,

$$\rho c_p \frac{DT}{Dt} - \alpha \rho g w T = \partial_j (k \partial_j T) + \tau_{ij} \partial_j u_i + \rho H \quad (5)$$

D/Dt is the material time derivative and the second term describes the effect of adiabatic heating. The terms on the right specify thermal conduction, heating by viscous dissipation and radiogenic heating respectively. Further symbol definitions are given in Table 1.

Symbol	Definition	Value	Unit
h	Depth of the mantle model	$3 \cdot 10^6$	m
z	Depth coordinate aligned with gravity	-	-
P	Thermodynamic pressure	-	-
ΔP	Dynamic pressure	-	-
T	Temperature	-	-
$T_{surface}$	Surface temperature	373	K
ΔT	Temperature scale	3500	K
u_i	Velocity field component	-	-
w	Radial velocity aligned with gravity	-	-
$e_{ij} = \partial_j u_i + \partial_i u_j$	Strain rate tensor	-	-
$e'_{ij} = e_{ij} - \frac{1}{3} e_{kk} \delta_{ij}$	Strain rate deviator	-	-
$e' = (1/2 e'_{ij} e'_{ij})^{1/2}$	Second invariant of deviator	-	-
$\eta(T, z) = \eta_0 \exp(cz - bT)$	P,T dependent viscosity	-	-
	$b = \ln(\Delta\eta_T), c = \ln(\Delta\eta_P)$	-	-
	$\Delta\eta_T = 100, \Delta\eta_P = 10$	-	-
η_0	Viscosity scale value	10^{23}	Pa s
$\tau_{ij} = \eta e'_{ij}$	Viscous stress tensor	-	-
α_0	Thermal expansivity scale value	$3 \cdot 10^{-5}$	K ⁻¹
ρ	Density	-	-
ρ_0	Density scale value	4500	kgm ⁻³
k	Uniform conductivity value	4.7	Wm ⁻¹ K ⁻¹
$\kappa = \frac{k}{\rho c_p}$	Thermal diffusivity	-	-
c_p	Specific heat	-	-
c_{p0}	Specific heat scale value	1250	Jkg ⁻¹ K ⁻¹
g	Gravitational acceleration	9.8	ms ⁻²
H	constant internal heating	$2 \cdot 10^{-12}$	Wkg ⁻¹
$Ra = \frac{\rho_0 \alpha_0 g \Delta T h^3}{\kappa_0 \eta_0}$	Rayleigh number	10^6	-
$Di = \frac{\alpha_0 g h}{c_{p0}}$	dissipation number	0.47	-

Table 1: Physical parameters

The thermal conductivity k is set to a uniform value. The physical properties ρ , c_p and α that appear as coefficients in equation (5) are included in our model through a general interface based on pressure temperature (P, T) domain tabulation of the material property values in the relevant pressure and temperature range for the Earth's mantle, further discussed in sections 2.2 and A.3 (van den Berg et al., 2002; Jacobs et al., 2006). The effect of heat exchange due to solid state phase transitions is not included in the formulation of the energy equation (5). In the extended boussinesq approximation of Christensen and Yuen (1985) this effect is included in terms of parameterizations of the Clapeyron slope and the density contrast across the phase boundary, resulting in modified expressions for effective heat capacity and thermal expansivity in the energy equation. Nakagawa et al., (2009) have extended this approach to a compressible model using tabulated thermophysical properties for a multicomponent model, where effective c_p and α are expressed in the temperature- and pressure derivatives of the enthalpy. In our model calculations for the binary system MgO, SiO₂ we have

not included such extensions that would require differentiation of the enthalpy across discontinuities at the monovariant phase transitions. We assume that this approximation will only have a minor influence on the overall dynamics and temperature structure of our model. This is further discussed in section 4.

The coupled model equations (1), (2) and (5) are solved using finite element methods. Domain discretization, boundary and initial conditions and computational aspects of the coupling between the model equations and the underlying mineral physics model are discussed in Appendix A.3.

2.2 Thermodynamical and mineralogical model

In our convection model the thermophysical properties density ρ , thermal expansivity α and specific heat c_P , are derived from a thermodynamic formalism based on lattice dynamics. This formalism by Jacobs and de Jong (2005, 2006, 2007), further denoted as ‘phonon-thermodynamic’ method, incorporates the vibrational density of states, for the purely magnesian endmember model of the MgO – SiO₂ system. The overall composition is specified by the mole fractions of the olivine and pyroxene components, (Mg₂SiO₄)_{0.56}(MgSiO₃)_{0.44}.

The absence of at least iron in our model system, relative to a natural system, of pyrolitic composition, prevailing in the mantle has three consequences. The first one is that density in our model system is lower than for a pyrolitic mantle, affecting sound wave velocities and the pressure-depth distribution. This is illustrated in Fig. 1 showing adiabatic profiles of seismic velocities, derived from the phonon-thermodynamic model, for an Earth-like mineral composition within the MgO – SiO₂ system. This figure shows that the main features of the mantle transition zone are reproduced by our mineral physics model based on a restricted composition (not including Al, Ca and Fe). The effect, of adding the major component Fe, on the seismic velocities of olivine, amounts to an overall reduction of the velocity by about 1-2%. Secondly, addition of more elements to the model system displaces phase boundaries. For instance the olivine-wadsleyite phase boundary will shift to an approximately 1 GPa lower pressure when Fe content is increased from zero to 10%. The wadsleyite-ringwoodite transition shifts in a similar way to lower pressure for increased iron content. Thirdly, specific phase boundaries in the model system split up by the addition of other elements, enclosing multi-phase regions. For instance the olivine-wadsleyite phase boundary splits up in a field in which olivine and wadsleyite coexist when iron is added in our model. Although the restricted model cannot be used to represent the ‘real Earth’ it is sufficiently realistic for our present purpose of exploring the seismic expression of multiple phase regions in the transition zone generated by the mantle convection process.

Despite the issue of compositional completeness of our present thermodynamic model we emphasize that the approach we follow here, is not restricted to a particular thermodynamic model. The mantle convection computations are set up in a flexible way such that different thermodynamical models, once available, can be applied, as illustrated further in section A.3 below. In future work this will enable us to refine our present calculations with an extended thermodynamic model containing more elements, and moreover to investigate effects of applying alternative thermodynamic models.

A thermodynamic model incorporating iron, which is constrained by phonon data and applicable in the range from zero Kelvin up to the melting point, is currently under development. Following the work of Jacobs and de Jong (2009), we anticipate that this model is particularly important for iron-bearing materials, which are characterized by antiferromagnetic-paramagnetic transitions at low temperatures, e.g. 67 K for fayalite and 12 K for ringwoodite. This requires a thermodynamic

framework, which is adequate to reliably describe thermophysical properties from zero Kelvin up to the melting temperature and to pinpoint inconsistencies in existing laboratory data.

A main advantage of the application of the phonon-thermodynamic method in numerical models of mantle convection is that it enables one to compute the required thermo-physical parameters as well as the phase diagram in a single self-consistent framework. This is important because for many substances crucial experimental data, such as for thermal expansivity and bulk modulus, have been measured at conditions far away from those prevailing in the mantle transition zone. Therefore, to obtain values for thermodynamic properties at mantle conditions, any physical model is forced to make large extrapolations of available thermodynamic data. For instance Jacobs and de Jong (2005b) showed, by using a calphad-type parameterization technique, that thermal expansivity and volume of gamma-Mg₂SiO₄ (ringwoodite) could not be constrained with sufficient accuracy to enable a prediction of bulk sound velocity for gamma-(Mg_{0.91}Fe_{0.09})₂SiO₄ to within 0.8%, twice the tomographic accuracy. Figure 12 of their work demonstrates that bulk sound velocity for this substance modeled by Fabrichnaya (1998), Stixrude and Bukowinsky (1993) and their own work are quite different, calling for a method which more tightly constrains thermodynamic data.

Such accuracy issues are also important because the representation of phase equilibrium boundaries is quite sensitive to variations in the Gibbs energy in pressure-temperature space. For instance a change in the Gibbs energy of akimotoite by about 500 J/mol (0.04 %) leads to a temperature shift of about 300 K of the quadruple point where wadsleyite, ringwoodite, akimotoite and stishovite are in equilibrium. Saxena (1996, 2010) gives a detailed analysis of how uncertainties in thermochemical properties affect the accuracy of the description of phase boundaries.

For all mineral phases present in the model phase diagram of Fig. 5, even for the phases stable at high pressure, experimental data, such as heat capacity, thermal expansivity and bulk modulus, are available at 1 bar pressure. These 1-bar data do not suffer from uncertainties associated with equations of state of pressure reference materials, as is the case for data obtained at high pressure. It is therefore of great importance to constrain thermodynamic analyses of these phases with 1 bar experimental data. Such data are very well fitted by methods based on polynomial expressions for these properties, (Fabrichnaya et al., 2004), (Saxena, 1996), (Holland and Powell, 1998).

However it has been demonstrated by Jacobs and Oonk (2000) that the methods based on polynomial approximations are difficult to use when thermodynamic properties are extrapolated to conditions prevailing in the lower mantle because of physical anomalies that occur in the computed thermal expansivity. Therefore the methods based on polynomials do not perform well, compared to alternative methods, for instance methods based on Mie-Grüneisen-Debye theory, when applied for the full pressure and temperature range of the Earth's mantle.

Significant progress in representing experimental data at lower mantle pressures has been made by Stixrude and Lithgow-Bertelloni (2005). They used the Mie-Grüneisen-Debye method (MGD) to construct a thermodynamic database for the CFMAS system. However due to the simplified construction of the Vibrational Density of State (VDoS) inherent to the Debye model and the use of the quasi-harmonic approximation, experimental data at 1-bar conditions are less well represented by the MGD method compared to the polynomial approach. This is illustrated for forsterite in Fig. 2, where computed values of the specific heat at constant pressure are compared with experimental data.

Jacobs and de Jong (2005, 2007) introduced a method, based on Kieffers (1979) more elaborate model for the VDoS that also includes the effect of anharmonicity. This method, here labeled MGK-a, combines an accurate representation of both 1-bar and high pressure (lower mantle) experimental

data. As a consequence the MGK-a method is better constrained by, both 1-bar and high pressure, experimental data.

Figure 2 shows for forsterite, that including anharmonicity combined with a realistic VDoS improves the representation of 1-bar heat capacity significantly. The effect of anharmonicity on vibrational frequencies has been measured for forsterite by Gillet (1991, 1997). Its effect on the heat capacity of forsterite is about 2.5% at 2000 K and 1 bar pressure and it propagates for about 0.4% in the Gibbs energy. Although this effect seems small at first sight, Jacobs and de Jong (2005) showed that it has a significant effect on location and slope of the phase boundary between forsterite and wadsleyite. Neglecting anharmonicity shifts this phase boundary by 2 GPa to lower pressures, whereas the slope decreases with almost a factor two. Not only forsterite, but also other mineral phases such as perovskite and akimotoite are affected by anharmonicity, impacting the position and slopes of phase boundaries as has been demonstrated by Jacobs and de Jong (2007).

In view of the demonstrated impact on the phase diagram, of both a detailed VDoS representation and of anharmonicity, in the lattice dynamics model, a unique characteristic of the MGK-a method, we consider this the most reliable method available to date for the computation of both the phase diagram and the thermo physical properties of the mantle convection model specified in (2)-(5), for an olivine-pyroxene system.

The thermodynamic formalism (MGK-a) by Jacobs et al. (2007) is based on microscopic properties such as vibrational frequencies and microscopic Grüneisen parameters. Relative to polynomial representations of thermodynamic data, this method is additionally constrained by Raman and Infrared spectroscopic data, resulting in more tight constraints of thermodynamic data. Jacobs et al. (2006) applied this to Mg_2SiO_4 to obtain better constrained thermodynamic properties for ringwoodite. In contrast to many thermodynamic parameterizations available in the literature their vibrational model is tightly constrained by experimental data below room temperature, such as low temperature heat capacity data, resulting in a better partitioning of physical effects at high temperature, such as intrinsic anharmonicity in forsterite and akimotoite and cation disorder in majorite. This also implies better constrained descriptions of entropy and enthalpy, necessary for the determination of Clapeyron slopes. This high degree of physical self-consistency and agreement with experimental data is considered particularly important in modeling complex, heterogeneous, configurations in the mantle transition zone.

Thus the properties ρ , c_P and α , required in model calculations based on equations (2), (5), together with both seismic velocities v_p , v_s , can be computed for the complete (P, T) range of the Earth's mantle, in a single physically self-consistent framework, without relying on separate parameterizations for individual mineral phases, as is common practice in models based on an (extended) Boussinesq approximation (EBA), (Christensen and Yuen, 1985, Steinbach et al., 1989, Zhao et al., 1992). In this respect our method is similar to the methods applied by Ita and King (1994), Piazonni et al. (2009) and Nakagawa et al. (2009), that are based on thermodynamical models, with varying degree of self-consistency in the representation of thermophysical parameters and seismic wave velocities. The phonon-thermodynamic method is further outlined in the appendix.

3 Results of numerical modelling

We have modelled thermal convection in a 2-D cylindrical domain with an opening angle of ninety degrees. The ratio of the core radius and the outer radius R_c/R is 0.54, similar to the value for Earth. The depth of the mantle is set to 3000 km. The model is both bottom and internally heated. Uniform and constant radiogenic heating is applied, set to a value $H = 2 \cdot 10^{-12}$ W/kg, slightly

lower than the chondritic value, in agreement with the smaller surface/volume ratio of the 2-D cylindrical domain, compared to the 3-D spherical case. Temperature is prescribed constant at the outer surface and the core mantle boundary with a temperature contrast $\Delta T = 3500$ K across the mantle. Zero heatflux symmetry conditions are imposed on the radial boundaries.

Computations were started from an initial adiabatic temperature field, with a potential temperature $T_p = 1600$ K, truncated at the top- and bottom thermal boundary layers. The model equations were integrated in time for about 1 Gyr.

A newtonian, temperature and pressure dependent viscosity model is used with an exponential parameterization defined in Table 1.

Fig. 3 shows results for a representative snapshot of the temperature distribution resulting from the convection model, at a model time of 837 Myr. The snapshot shown is characteristic for the model results with narrow cold downwellings, descending all the way to the core mantle boundary. These downwellings migrate along the outer boundary, similar to subducting lithospheric plates subject to trench migration and slab roll-back. The temperature field is shown in the top row of frames. The bottom row displays corresponding plots of the lateral variation of the seismic shear wavespeed. The temperature field is characterized by two colliding cold downwellings and a region of hot upwelling, in the area indicated by 1500 km wide zoom-in windows, outlined in black in frame (a), that correspond to frames (b) and (c). The cold downwellings show only limited dynamic interaction with the endothermic phase boundaries near 700 km depth, with a slight deflection of the right hand downwelling. This is related to the small value of the Clapeyron slope, -1.9 MPa/K, of the post-spinel phase transition at the bottom of the transition zone in our model. The temperature distribution of the zoom-in windows is smooth and without significant features related to the transition zone.

Lateral variation of the seismic shear velocity δv_s is shown in frames (d,e,f) that correspond to the temperature frames (a,b,c) respectively. The lateral variation is defined with respect to a smooth background profile, to bring out more clearly the discontinuities in the velocity, related to the different phase boundaries, $\delta v_s(\mathbf{x}) = (v_s(\mathbf{x}) - \langle v_s(r) \rangle) / \langle v_s(r) \rangle$. The smooth background profile $\langle v_s(r) \rangle$ is computed from the horizontally averaged values by applying a boxcar moving average filter with a 200 km halfwidth.

The velocity and temperature variations shown in Fig.3 are closely correlated in the lower mantle region away from the D'' region, where several lens-like structures of post-perovskite show up in the seismic velocity (Hernlund et al., 2005, van der Hilst et al., 2007).

This indicates that seismic velocity is related in a simple way to temperature and pressure, in this region without phase transitions. This situation is quite different for the transition zone region between about 400 and 700 km depth as illustrated particularly in Fig. 3e,f. The large scale structure of the smooth temperature field associated with the vertical downwellings is dominant in the lateral variation but much of the small scale structure of the velocity anomalies is not reflected in the smooth variation of the temperature field.

A smooth relation between temperature and seismic wavespeed is underlying interpretations of seismic travel time tomography (Goes et al., 2004), (Schuberth et al., 2010b). The poor correlation between small scale lateral variation of shear wave speed and temperature illustrated in Fig. 3 shows that assumption of such relation is not applicable for small scale variation in the depth range of the transition zone. A clear difference in correlation of seismic velocity and temperature, between regions inside and outside the transition zone is illustrated more directly in Fig. 4, showing discrete samples of wavespeed data obtained from several isobaric cross-sections at different sample depths, taken from the convection results.

Only the data samples taken at 797 km depth (27.5 GPa), in the shallow lower mantle occupied by post-spinel, show a continuous relation between shear velocity and temperature.

The velocity-temperature plots of Fig. 4 can be interpreted with the phase-diagram for the magnesium endmember model applied in the convection model shown in Fig. 5. Horizontal dashed lines in Fig. 5 indicate the pressure values of the isobaric velocity-temperature plots in Fig. 4.

The intersection of the 9.4 GPa isobar, at 300 km depth, with the phase boundary curve of the ortho- and clino enstatite transition at about 1750 K corresponds to the transition shown in the bottom velocity-temperature curve in Fig. 4. The loop in the sampling curves for 300 and 636 km depth of Fig. 4 is a result of the spread in pressure in the sample points that are located in a 20 km depth range around the indicated depth.

In a similar way the intersection of the 21 GPa isobar, at 636 km depth, with the wadsleyite-ringwoodite ($\beta \rightarrow \gamma$) phase boundary near 1800 K corresponds to the velocity transition of the middle curve in Fig. 4. The latter phase transition corresponds to the ‘520 km’ discontinuity that has been the target of a growing number of seismological investigations (Shearer et al., Deus et al., 2001, 2006, Cao et al., 2010). In our modelling results this transition occurs around 650 km depth as will be shown below. The fact that this transition occurs at depths greater than 520 km in our model is a consequence of the absence of iron in the magnesium endmember composition model used here. This composition related pressure shift of the $\beta \rightarrow \gamma$ transition has been calculated from the thermodynamic database of Saxena (1996) as 1.25 GPa. This effect is amplified in a shift to smaller depth, due to the increased pressure gradient associated with the density increase resulting from adding 10 % iron. Fig. 4 clearly illustrates that interpreting lateral variation in seismic velocity in terms of temperature variation only, will not produce correct results in a multi-phase environment like the mantle transition zone.

That strong lateral variation of physical properties, related to variation in mineral phase, can occur in the mantle transition zone, especially in regions of large lateral variation in temperature, is illustrated in Figure 5, by three radial temperature profiles. The profiles are labeled by their angular coordinate, positioned in the zoom-in window around the cold downwellings of Fig. 3b,e. Adiabats with a range of potential surface temperatures of 1400-1800 K are drawn for reference in Figure 5. This temperature range also includes the red profile, positioned between the two cold downwellings, that corresponds to a background reference profile, more representative for the mantle outside the cold disturbance of the downwelling flows. From the temperature excursion across several phase boundaries of the two low temperature geotherms, associated with the downwelling flows, it is clear that several mineral phases can exist at a given depth/pressure depending on the laterally varying temperature field. This is more directly illustrated in Fig. 6 showing the mineral phase distribution, frames (a,b) and corresponding lateral variation of the shear velocity δv_s (c,d) for the two zoom-in windows indicated by black boxes in Fig. 3(a,d). Dashed lines in Fig. 6a correspond to the radial temperature profiles shown in the phase diagram of Fig. 5. Individual pockets of different mineral associations intersected by the radial profiles can be identified with the help of the phase diagram of Fig. 5. Fig. 6a shows a complex configuration of regional domains occupied by contrasting mineral phases, indicated by the same phase labels as in Fig. 5. The large negative temperature perturbations, of up to 1000 K, of the two cold downwellings, shift the local geotherms across several stability fields in the phase diagram in Fig. 5. This causes the strong lateral variation in the phase distribution shown in Fig. 6a.

Frame 6b shows similar lateral variation in the mineral phase distribution for a zoom-in window with a smaller, hot, mantle anomaly. The lateral variation of temperature is mainly reflected here in the variation of the thickness of the various mineral phase layers. The forementioned phase

boundary between wadsleyite and ringwoodite, associated with the ‘520 km’ discontinuity, occurs in two different settings here. First as the transition $\beta + ak \rightarrow \gamma + ak$ that is more prominent in the core of the cold downwellings. The local thickening of the $\beta + ak$ layer is a consequence of the diamond shaped stability field of this mineral assemblage shown in Fig.5. Second, in warmer areas, as the transition $\beta + pv \rightarrow \gamma + pv$. This results in an intermittent behavior of the $\beta \rightarrow \gamma$ boundary illustrated in 6a,b, which could be related to lateral variations in seismic observations of the ‘520 km’ boundary reported by (Deuss, 2009) and (Cao et al., 2010). If such detail of the $\beta \rightarrow \gamma$ transition would indeed be observable in the seismic data this would provide strong constraints on lateral variation of the temperature field in the transition zone.

In the modelling results the bottom of the transition zone that represents the boundary between the upper and lower mantle consists of a layer of ringwoodite and perovskite, that transforms into a post-spinel assemblage of periclase and perovskite for pressures higher than the endothermic post-spinel transition. The pyroxene component has transformed to perovskite at shallower level at the nearby strongly endothermic transition of akimotoite to perovskite (Chudinovskikh and Boehler, 2004, Jacobs and de Jong, 2007). The magnitude of the Clapeyron slope in (Jacobs and de Jong, 2007) of 4.1 MPa/K is smaller than the value of 6 MPa/K reported in (Chudinovskikh and Boehler, 2004). The Clapeyron slope of the ringwoodite to postspinel transition of (Jacobs and de Jong, 2007) is -1.9 MPa/K. The relatively small magnitude of the latter transition and the splitting of the phase transition to perovskite of the olivine and pyroxene reduce the dynamic effect of the composite endothermic phase boundary between the upper and lower mantle, in line with the weak interaction of the cold downwellings with these phase boundaries as shown in Fig. 3.

The radial profile at 37° , halfway between the cold downwellings, is representative for the global horizontally averaged mantle. The corresponding geotherm in Fig. 5 is close to an adiabat of 1700K. The transition zone has a five layer configuration here with, from top to bottom 1) $\beta +$ clino-enstatite, 2) $\beta +$ majorite, 3) $\beta +$ akimotoite, 4) $\beta +$ pv 5) $\gamma +$ perovskite. The transition zone is followed by the lower mantle consisting of a post spinel assemblage of perovskite $MgSiO_3 +$ periclase MgO .

Fig. 6c,d show the lateral variation of the shear velocity for the corresponding zoom-in windows of frames (a) and (b). There is a strong correlation between the velocity anomalies and the mineral phase distribution, to the degree that individual pockets of contrasting mineral content can be identified in the anomaly pattern of the shear velocity. This is in contrast with the corresponding smooth temperature field distributions shown in Fig. 3(b,c).

The ‘410 km’ seismic discontinuity is clearly visible in the velocity frames as the boundary between second and third phase layer from the surface. This phase boundary shows the well-known uplift near the core of the two cold downwellings, in agreement with the positive Clapeyron slope of the $\alpha \rightarrow \beta$ phase transition (Turcotte and Schubert, 2002). The phase boundary between layer 1 and 2, related to the transition from ortho-enstatite to clino-enstatite shows an even greater topography related to the high value of the Clapeyron slope, that increases for lower temperatures, as shown in Fig. 5.

Fig. 7 shows a compilation of radial distributions of temperature (a,c) and seismic shear velocity (b,d). The different curves in each frame correspond to the profiles indicated by dotted lines in Fig. 6a,b. The black curves in frames (a) and (c) correspond to the horizontally averaged geotherm that is close to the 1700K adiabat in the transition zone.

The curves in frames (c),(d) correspond to profiles near 70° in a less disturbed and warmer upper mantle region. The geotherms in Fig. 7a show temperature contrasts over 1000 K, illustrating the

wide range of temperatures found in the mantle transition zone, in areas of subducting lithosphere.

Corresponding profiles of the shear wavespeed are shown in Fig. 7b,d. The smoothened global background profile, $\langle v_s(r) \rangle$, used in the calculation of lateral variations of shear velocity, is plotted as a dashed curve. Seismic discontinuities corresponding to the phase boundary crossings of the different temperature profiles can be clearly recognized, showing substantial differences between the different profiles, in particular in Fig.7b, representing the region with stronger lateral variation in temperature.

The direct correlation between mineralogy and lateral variation of seismic velocity shown in these results clearly demonstrates that the temperature dependence of the seismic wave velocities is strongly non-linear because of the mineralogy involved. This is in contrast with assumptions underlying methods of temperature and density estimation from seismic tomographic data.

Detailed seismic velocity structure of the mantle transition zone, including lateral variation, is an important target of seismological investigation, (Shearer et al.,1999, Deus and Woodhouse, 2001, Deuss et al, 2006, Lebedev et al., 2002, Cao et al., 2010). Studies like those by Deuss and co-workers and Cao et al., 2010, using SS precursors originating from underside reflections of the transition zone layering, produce reflectivity-time traces as a measure of radial distribution of velocity contrasts. Here we have computed a similar approximation of the radial distribution of the shear wave reflectivity from the radial derivative of the logarithmic seismic impedance. The computed reflectivity-depth profiles were then converted to time traces by a conversion from depth to two-way traveltime, using the spatially variable shear wave speed, followed by equidistant interpolation of the time traces with a sample frequency $f_s = 2.56$ Hz. Fig. 8 shows a spectral analysis of a representative reflectivity time trace that samples the transition zone in the region of complex phase distribution corresponding to Fig. 6a, near the radial profile labeled 35° . The signal power is concentrated near a frequency of 0.1 Hz that corresponds to wavelength's between about 50 and 60 km in the transition zone. Structures of such wavelength are reasonably well resolved by our finite element mesh with a nodal point spacing of 6 km in the transition zone. We tested the effect of the finite element resolution on our results by comparison with a model computation with half the number of nodal points with respect to the model results shown. This comparison produced practically identical results in particular also in the powerspectrum shown in Fig. 8. The mesh resolution is further specified in section A.3.

The raw reflectivity data traces obtained by the forementioned procedure are shown in Fig. 9a, for the region corresponding to Fig. 6a. To investigate the seismic signal bandwidth required to resolve the underlying velocity structure we applied low-pass frequency filtering to the raw reflectivity traces. The result of second order Butterworth filtering of the raw data traces with corner periods of 20 and 40 seconds are shown in frames b) and c) of Fig. 9 respectively.

The corresponding shear velocity distribution for the region is shown in Fig. 10, for interpretation of the reflectivity traces. This figure shows clearly contrasting velocity values that coincide with the different phase regions shown in Fig. 6. The discontinuities outlining the different phase regions in Fig. 10 are closely correlated with the pattern of reflectivity maxima in Fig. 9. Pockets of anomalous 'low temperature' phases like the ringwoodite regions in the cold downwellings are clearly visible in the reflectivity traces.

Table 2 lists reflectivity maxima corresponding to several phase boundaries, mostly related to the pyroxene component, that can be identified in locations, away from the irregular structures related to the two cold downwellings in the mantle region shown.

Time in seconds	Phase transition
200	α -olivine + clino-enstatite $\rightarrow$ β -olivine + clino-enstatite
225	β -olivine + clino-enstatite $\rightarrow$ β -olivine + majorite
240	β -olivine + majorite $\rightarrow$ β -olivine + akimotoite
not-resolved	β -olivine + akimotoite $\rightarrow$ γ -olivine + akimotoite
270	γ -olivine + akimotoite $\rightarrow$ γ -olivine + perovskite
280	γ -olivine + perovskite $\rightarrow$ periclase + perovskite

Table 2: Approximate two-way travel times and phase transitions related to the reflectivity maxima for the background mantle parts of Fig. 9.

The ‘520 km’ seismic discontinuity, associated with the $\beta \rightarrow \gamma$ transition of the olivine component, is represented in our modelling results by the contrast between the $\beta + ak$ (green in Fig.6ab) and $\gamma + pv$ (pink in Fig.6ab) layers near the boundary between the upper and lower mantle. In the reflectivity traces of Fig.9 this transition is visible as the second peak from the bottom near 270 seconds. The thin layer of $\gamma + ak$, replaced laterally by $pv + \beta$, as shown in Fig.6ab is not resolved in the reflectivity traces.

The ‘520 km’ discontinuity is observed in a peridotite mantle, of pyrolitic composition, roughly halfway between the $\alpha \rightarrow \beta$ (‘410 km’) and $\gamma \rightarrow MgO + pv$ (‘660 km’) phase boundaries (Deuss and Woodhouse, 2001, Cao et al., 2010). Another reason for the relative proximity of the ‘520’ and ‘660 km’ boundaries, besides the forementioned absence of iron in our magnesian endmember model, is the rather high background temperature of the mantle model where the geotherm intersects the spinel to post-spinel boundary ($\gamma + pv \rightarrow MgO + pv$) in Fig. 5 at a temperature of about 2050 K, higher than the experimental value 1900 ± 100 K in (Boehler, 2000). This explanation by a temperature effect is in agreement with the occurrence of two thicker pockets of ringwoodite and akimotoite ($\gamma + ak$), in the phase distribution shown in Fig. 6a, intersected by dashed profile lines, corresponding to the two colder geotherms labeled 35° and 39° in the phase diagram of Fig. 5. From Fig. 9 it can be concluded that the main mineral phase distribution of the transition zone, in the present magnesian endmember model, are clearly expressed in the reflectivity structure, at least for the period range below 20 seconds, well accessible in broadband seismological data, as illustrated in 9b.

4 Discussion and conclusions

Recent progress in mapping sharp contrasts in seismic shear wavespeed in the transition zone (Deuss et al., 2006, Cao et al., 2010) calls for an assessment of phase related heterogeneity generated in mantle dynamical models. In order to investigate such heterogeneous structure of the mantle transition zone, resulting from lateral variation in temperature, we have modelled thermal convection in the Earth’s mantle, applying a mineral physics model from which the complex phase diagram for the olivine-pyroxene system can be derived in a self-consistent manner.

Our mantle convection model includes thermophysical properties such as density, thermal expansivity and specific heat, for the magnesium endmembers of a olivine-pyroxene mantle. We have computed lateral variations in seismic shear wave velocities applying the same lattice vibrational model as used in the computation of thermophysical properties (Jacobs and de Jong, 2007).

Our computations are based on a restricted chemical composition of the model material, in particular without the approximately 10% iron content of a pyrolitic mantle composition (Ringwood,

1975). We applied this model in our calculations because of the available corresponding thermo-physical database (Jacobs and de Jong, 2007) that is unique in accurate prediction of both the phase diagram and material properties specific heat, thermal expansion and density, as discussed in section 2.2. The absence of iron in our model results in a shift to higher pressure of major phase transitions of the model olivine component, olivine-wadsleyite and wadsleyite-ringwoodite, associated with the ‘410 km’ and ‘520 km’ seismic discontinuity respectively (Bina and Helffrich, 1994, Jacobs and de Jong, 2005b). In contrast the spinel-postspinel boundary corresponding to the ‘660 km’ discontinuity is less affected by the model iron content. The restricted composition also implies a lower model density and associated pressure gradient, which also results in a depth increase of the phase boundaries. The monovariant phase transitions of our mineral physics model do not have multi-phase regions characteristic for models containing solid solutions. This implies that phase transitions in our model are associated with discontinuities in physical properties, in contrast to a model containing 10% iron in solid solution that shows smoother transitions between phase domains (Bina and Helffrich, 1994, Stixrude, 1997). In this sense our model results represents an endmember case for the computation of the seismic expression of phase boundaries in the transition zone.

Our model calculations were done with the simplifying assumption that the effect of latent heat exchange due to phase transitions can be neglected to first order in the large scale dynamics and global temperature distribution. The dynamical effect of the phase boundary on the mantle flow operates mainly through the phase related density contrast in the righthand side of the Stokes equation (2). The localized temperature effect from the latent heat exchange produces a shift in the transition pressure that is estimated to be smaller than about 0.4 GPa. We note that this value is similar to the uncertainty in the experimental determination of phase boundaries (Frost, 2003). In view of the forementioned shift of the transition-depth of phase boundaries, resulting from absence of iron in our model, estimated at about 1 to 2 GPa for $\alpha \rightarrow \beta$ and $\beta \rightarrow$ ringwoodite respectively, the effect of latent heat is acceptable.

In spite of these differences with a more realistic mantle composition, our model reproduces thermophysical properties, including seismic wavespeeds, as well as mineral phase boundaries to within experimental uncertainty, for the magnesium endmember of the olivine-pyroxene system. Self-consistency of the underlying mineral physics model and high accuracy in representing experimental data, as discussed in section 2.2, is an important characteristic of our convection model, especially when applied to mantle regions with a large number of mineral phases with contrasting physical properties such as the upper mantle transition zone.

We have analysed the distribution of mineral phase and corresponding shear wavespeed, computed from the pressure-temperature fields obtained in the mantle convection calculations. In relatively undisturbed regions, away from cold downwellings and hot mantle plumes, the modelling results show heterogeneity mainly in the variation of layer thicknesses of the different phase associations. Here we observe interesting similarities with seismological results from SS and precursors from underside reflections of the transition zone (Deuss et al., 2006, Cao et al, 2010). The seismological results show lateral variations in amplitude and depth of seismic reflectors interpreted as topography of the ‘410’ and ‘660 km’ phase boundaries. The ‘520 km’ boundary has a more regional character (Deuss, 2009) and frequency dependent signature that could be related to phase transitions in the pyroxene component (Cao et al., 2010). In our model results ringwoodite occurs in association with stishovite and with akimotoite in the core of a cold downwelling. Away from cold downwellings a thin layer of ringwoodite and akimotoite exists intermittantly replaced by an equally thin layer of wadsleyite and perovskite. This layering can be identified in the reflectivity traces computed from the wavespeed distributions. This configuration is related to the vicinity of the stability fields for

these phase associations, along an isobar of 22 GPa, in the phase domain of Fig. 5. The lateral boundaries between these associations pinpoint the temperature of the ‘quadruple point’ at 22 GPa in Fig. 5 and the middle (black) data set shown in Fig. 4. This example illustrates the possibility of lateral variation in the signature of seismic reflectors due to phase transitions of the pyroxene component.

Other transitions in the pyroxene component that are clearly visible in the reflectivity traces of Fig. 9 are clinoenstatite to majorite, close to the ‘410 km’ discontinuity, and the majorite to akimotoite transition about halfway in the transition zone.

The strongest heterogeneity in our model results is found in regions with cold downwelling flows, simulating lithospheric subduction, where the geotherm deviates by up to 1000 K from the horizontally averaged background value. This heterogeneity shows up as distinct pockets of mineral phase with corresponding, contrasting seismic velocity distribution, where the outlines are clearly visible in the reflectivity traces. It should be noted here that our modelling results have been obtained assuming equilibrium thermodynamics, thereby neglecting possible effects of phase transition kinetics that may become significant in the coldest cores of old subducting lithospheric slabs (Daessler et al., 1996; Vacher et al., 1999).

Our main objective in this paper has been to investigate the heterogeneity due to the complex phase diagram of a magnesium endmember olivine-pyroxene system, for the mantle depth range including the transition zone, and relate the results to recent developments in seismological reflectivity imaging of the mantle.

We have shown that lateral variations of temperature result in a laterally heterogeneous phase distribution, especially near subduction regions. The results show that the distribution of seismic shear wavespeed is clearly dominated by the phase distribution rather than by the temperature field.

The major phase transitions of both the olivine and pyroxene subsystems show up clearly in the reflectivity traces computed from the shear velocity distribution in the signal bandwidth with periods below 20 s, indicating that these phase boundaries can, in principle, be imaged by seismological methods. Geodynamical models that can generate the detailed structure of the transition zone region can therefore be helpful in the interpretation of such seismological data.

Further progress requires that multi-phase regions, in particular related to a finite iron composition will be modelled. This requires the extension of our thermodynamic data base with iron, which is currently in progress (Jacobs and de Jong, 2009).

Acknowledgments

We acknowledge constructive reviews from Takashi Nakagawa and Yuji Higo that were helpful for improving the manuscript. Rob van der Hilst is thanked for making available unpublished results. Computational resources for this work were made available by the Netherlands Research Center for Integrated Solid Earth Science (ISES 3.2.5). Collaboration between A. van den Berg and M. Jacobs has been supported through ISES project ME-2.7.

References

- [1] Ashida T., Kume S., and Ito E. Thermodynamic aspects of phase boundary among alpha-, beta-, and gamma-Mg₂SiO₄. In Manghnani M.H., Syono Y (eds), *High-pressure research in mineral physics*. Terra Scientific Publishing Company (TERRAPUB), Tokyo/American Geophysical Union, Washington DC, pp 269-274, 1987.
- [2] Bina, C.R. and G.R. Helffrich, Phase transition Clapeyron slopes and transition zone seismic discontinuity topography, *J. Geophys. Res.*, **99**, 15,853-15,860, 1994.
- [3] Boehler, R. High-pressure experiments and the phase diagram of lower mantle and core materials, *Reviews of Geophysics*, **38**, 221-245, 2000.
- [4] Cao, Q., Wang, P., Lamm, R., van der Hilst, R.D. and M.V. de Hoop, Imaging the upper mantle transition zone with a generalized Radon transform of SS precursors, *Phys. Earth Planet. Inter.*, **180**, 80-91, 2010.
- [5] Christensen, U.R. and D.A. Yuen, Layered convection induced by phase transitions, *J. Geophys. Res.*, **90**, 10291-10300, 1985.
- [6] Chudinovskikh, L., and R. Boehler, MgSiO₃ phase boundaries measured in the laser-heated diamond cell, *Earth Planet. Sci. Lett.*, **219**, 285-296, 2004.
- [7] Connolly, J.A.D., Computation of phase equilibria by linear programming: A tool for geodynamic modeling and its application to subduction zone decarbonation, *Earth Planet. Sci. Lett.*, **236**, 524-541, 2005.
- [8] Cuvelier, C., Segal, A. and A.A. van Steenhoven, *Finite Element Methods and Navier-Stokes Equations*, D. Reidel Publishing Company, Dordrecht, 1986.
- [9] Daessler, R., Yuen, D.A., Karato, S. and M.R. Riedel, Two-dimensional thermo-kinetic model for the olivine-spinel phase transition in subducting slabs, *Phys. Earth Planet. Inter.*, **94**, 217-239, 1996.
- [10] Deuss, A. and J.H. Woodhouse, Seismic observations of splitting of the mid transition zone in Earth's mantle, *Science*, **294**, 354-357, 2001.
- [11] Deuss, A., Redfern, S.A.T., Chambers, K. and J.H. Woodhouse, The nature of the 660-kilometer discontinuity in Earth's mantle from global seismic observations of PP precursors. *Science*, **311**, 198-201, 2006.
- [12] Deuss, A., Global Observations of Mantle Discontinuities Using SS and PP Precursors, *Surveys in Geophysics*, **30**, 301-326, 2009.
- [13] Dziewonski, A.M. and D.L. Anderson, Preliminary reference earth model (PREM), *Phys. Earth Planet. Int.*, **25**, 297-356, 1981.
- [14] Fabrichnaya, O. The assessment of thermodynamic parameters for solid phases in the Fe-Mg-O and Fe-Mg-Si-O systems. *Calphad*, **22**, 85-125, 1998.
- [15] Frost, D.J., The structure and sharpness of (Mg, Fe)₂SiO₄ phase transformations in the transition zone, *Phys. Earth Planet. Int.*, **216**, 313-328, 2003.

- [16] Gillet, P., Richet, P., Guyot, F. and G. Fiquet, High-temperature thermodynamic properties of forsterite, *J. Geophys. Res.*, **96**, 11 805-11 816, 1991.
- [17] Gillet P, Daniel I, Guyot F. and G. Fiquet, Anharmonic properties of Mg_2SiO_4 -forsterite measured from the volume dependence of the Raman spectrum, *Eur. J. Mineral.*, **9**, 255-262, 1997.
- [18] Goes, S., Cammarano, F. and U. Hansen, Synthetic seismic signature of thermal mantle plumes, *Earth Planet. Sci. Lett.*, **218**, 403-419, 2004.
- [19] Gillet, P., Daniel, I., Guyot, F., Matas, J., and J.-C. Chervin, A thermodynamic model for MgSiO_3 -perovskite derived from pressure, temperature and volume dependence of the Raman mode frequencies, *Phys. Earth Planet. Inter.*, **117**, 361-384, 2000.
- [20] Helffrich, G.R. and C.R. Bina, Frequency dependence of the visibility and depths of mantle seismic discontinuities, *Geophys. Res. Lett.*, **21**, 24, 2613-2616, 1994.
- [21] Hernlund, J.W., Thomas, C. and P.J. Tackley, A doubling of the post-perovskite phase boundary and structure of the Earth's lowermost mantle, *Nature*, **434**, 882-886, 2005.
- [22] Holland, T. and R. Powell, An internally consistent thermodynamic dataset for phases of petrological interest, *J. Metamorphic Geol.*, **16**, 309-343, 1998.
- [23] Ita, J. and S.D. King, Sensitivity of convection with an endothermic phase change to the form of governing equations, initial conditions, boundary conditions, and equation of state, *J. Geophys. Res.*, **99**, 15,919-938, 1994.
- [24] Jacobs, M.H.G. and B.H.W.S. de Jong, Quantum-thermodynamic treatment of intrinsic anharmonicity; Wallace's theorem revisited, *Phys Chem Minerals*, **32**, 614-626, 2005a.
- [25] Jacobs, M.H.G. and B.H.W.S. de Jong, An investigation into thermodynamic consistency of data for the olivine, wadsleyite and ringwoodite form of $(\text{Mg, Fe})_2\text{SiO}_4$. *Geochim. Cosmochim. Acta*, **69**, 4361-4375, 2005b.
- [26] Jacobs, M.H.G., van den Berg, A.P. and B.H.W.S. de Jong, The derivation of thermo-physical properties and phase equilibria of silicate materials from lattice vibrations: applications to convection in the earth's mantle, *Calphad*, **30**, 131-146, 2006.
- [27] Jacobs, M.H.G. and B.H.W.S. de Jong, Placing constraints on phase equilibria and thermo-physical properties in the system MgO-SiO_2 by a thermodynamically consistent vibrational method, *Geochimica et Cosmochimica*, **71**, 3660-3655, 2007.
- [28] Jacobs, M.H.G. and B.H.W.S. de Jong, Thermodynamic mixing properties of olivine derived from lattice vibrations, *Phys. Chem. Minerals*, **36**, 365-389, DOI 10.1007/s00269-008-0284-9, 2009.
- [29] Jarvis, G.T. and D.P. McKenzie, Convection in a compressible fluid with infinite Prandtl number, *J. Fluid Mech.*, **96**, 525-583, 1980.
- [30] Kieffer, S.W., Thermodynamics and lattice vibrations of minerals: 3 Lattice dynamics and an approximation for minerals with application to simple substances and framework silicates, *Rev. Geophys. Space Physics*, **17**, 35-59, 1979.

- [31] Lebedev, S., Chevrot, S. and R.D. van der Hilst, Seismic evidence for olivine phase changes at the 410- and 660-kilometer discontinuities, *Science*, **296**, 1300-1302, 2002.
- [32] Nakagawa, T., Tackley, P.J., Deschamps, F. and J.A.D. Connolly, Incorporating self-consistently calculated mineral physics into thermochemical mantle convection simulations in a 3-D spherical shell and its influence on seismic anomalies in Earth's mantle, *Geochemistry Geophysics Geosystems*, **10**, Q03004, doi:10.1029/2008GC002280, 2009.
- [33] Orr R.L., High temperature heat contents of magnesium orthosilicate and ferrous orthosilicate, *J. Am. Chem. Soc.*, **75**, 528-529, 1953.
- [34] Piazzoni, A.S., Steinle-Neumann, G., Bunge, H-P., D. Dolejš, A mineralogical model for density and elasticity of the Earth's mantle, *Geochem. Geophys. Geosys.*, **8**, 1-23, 2007, Q11010, doi:10.1029/2007GC001697.
- [35] Reynard, B. and D.C. Rubie, High-pressure, high-temperature Raman spectroscopic study of ilmenite-type MgSiO_3 , *Am. Miner.*, **81**, 1092-1096, 1996.
- [36] Ringwood, A.E., *Composition and petrology of the earth's mantle*, McGraw Hill, New York, 1975.
- [37] Robie R.A., Hemingway B.S and H. Takei, Heat capacities and entropies of Mg_2SiO_4 , Mn_2SiO_4 and Co_2SiO_4 between 5 and 380 K. *Am. Mineral.*, **67**, 470-482, 1982.
- [38] Saxena, S.K., Earth mineralogical model: Gibbs free energy minimization in the system $\text{MgO} - \text{FeO} - \text{SiO}_2$, *Geochim. Cosmochim. Acta*, **60**, 2379-2395, 1996.
- [39] Saxena, S.K., Thermodynamic modelling of the Earth's interior, *Elements*, **6**, 321-325, 2010.
- [40] Schuberth, B.S.A., Bunge, H.-P., Steinle-Neumann, G., Moder, C. and J. Oeser, Thermal versus elastic heterogeneity in high-resolution mantle circulation models with pyrolite composition: High plume excess temperatures in the lowermost mantle, *Geochemistry Geophysics Geosystems*, **10**, Q01W01, 2009.
- [41] Schuberth, B.S.A., Bunge and J. Ritsema, Tomographic filtering of high-resolution mantle circulation models: Can seismic heterogeneity be explained by temperature alone?, *Geochemistry Geophysics Geosystems*, **10**, Q05W03, 2009.
- [42] Segal, A., Praagman, N.P., The SEPRAN package, Technical report, <http://ta.twi.tudelft.nl/sepran/sepran.html>.
- [43] Shearer, P.M., Flanagan, M.P. and M.A. Hedlin, Experiments in migration processing of SS precursor data to image upper mantle discontinuity structure, *J. Geophys. Res.*, **104**, 7229-7242, 1999.
- [44] Steinbach, V., Hansen, U., and Adolf Ebel, Compressible convection in the earth's mantle: a comparison of different approaches, *Geophys. Res. Lett.*, **16**, 633-636, 1989.
- [45] Steinbach, V. *Numerische experimente zur Konvektion in kompressiblen Medien*, PhD Thesis, Universität zu Köln, 1991.

- [46] Stixrude, L., and Bukowinski, M.S.T., Thermodynamic analysis of the system MgO – FeO – SiO₂ at high pressure and the structure of the lowermost mantle. In *Evolution of Earth and Planets*, eds. E. Takahashi, R. Jeanloz and D. Rubie, Geophysics Monograph Series 74, 131-142, AGU, Washington DC, 1993.
- [47] Stixrude, L., Structure and sharpness of phase transitions and mantle discontinuities, *J. Geophys. Res.*, **102**, 14,835-14,852, 1997.
- [48] Stixrude, L. and C. Lithgow-Bertelloni, Thermodynamics of mantle minerals-I Physical properties, *Geophys. J. Int.*, **162**, 610-632, 2005.
- [49] Turcotte, D.L. and G. Schubert, *Geodynamics*, Cambridge University Press, Cambridge, UK, 2002.
- [50] Vacher, P., Spakman, W. and M.J.R. Wortel, Numerical tests on the seismic visibility of metastable minerals in subduction zones, *Earth Planet. Sci. Lett.*, **170**, 335-349, 1999.
- [51] van den Berg, A.P., van Keken, P.E. and D.A. Yuen, The Effects of a composite non-Newtonian and Newtonian rheology on Mantle convection, *Geophysical Journal International*, **115**, 62-78, 1993.
- [52] van den Berg, A.P., Jacobs, M.H.G. and B.H.W.S. de Jong. Numerical models of mantle convection based on thermodynamic data for the MgO-SiO₂ olivine-pyroxene system, *EOS, Trans. AGU*, **83(47)**, F624, 2002.
- [53] van den Berg, A.P., Rainey, R.S.G. and D.A. Yuen, The combined influences of variable thermal conductivity, temperature- and pressure-dependent viscosity and core-mantle coupling on thermal evolution, *Phys. Earth Planet. Inter.*, **149**, 259-278, 2005.
- [54] van der Hilst, R.D., de Hoop, M.V., Wang, P., Shim S.-H., Ma, P. and L. Tenorio, Seismostratigraphy and thermal structure of Earth's core-mantle boundary region, *Science*, **315**, 1813-1817, 2007.
- [55] Vinet, P., Ferrante, J., Rose, J.H., Smith, J.R., Compressibility of solids, *J. Geophys. Res.*, **92**, 9319-9325, 1987.
- [56] Wang, P., de Hoop, M.V. and R.D. van der Hilst, Imaging of the lowermost mantle (D'') and the core-mantle boundary with SKKS coda waves, *Geophys. J. Int.*, **175**, 103-115, 2008.
- [57] Watanabe H., Thermochemical properties of synthetic high-pressure compounds relevant to the Earth's mantle, In: Manghnani M.H., Akimoto S (eds), *High-pressure research in geophysics*, Center for Academic Publications, Japan/Tokyo: Reidel/Dordrecht, pp 441-464, 1982.
- [58] Zhao, W., Yuen, D.A. and S. Honda, Multiple phase transitions and the style of mantle convection, *Phys. Earth Planet. Inter.*, **72**, 185-210, 1992.
- [59] Xu, W., Lithgow-Bertelloni, C., Stixrude, L. and J. Ritsema, The effect of bulk composition and temperature on mantle seismic structure, *Earth and Planetary Science Letters*, **275**, 70-79, 2008.

A Thermodynamic formalism and computation of thermophysical parameters and seismic wave velocities

A.1 Background of the thermodynamical model

In our modelling experiments thermophysical properties ρ, α, c_P and the seismic wave velocities were derived from a formalism based on thermodynamics and lattice dynamics. This phonon-thermodynamic formalism was developed in (Jacobs and de Jong, 2005, 2007). There are several important differences between this formalism and recently published alternative formulations and corresponding databases by Stixrude and Lithgow-Bertelloni (2005), Connolly (2005) and Piazzoni et al., (2007). The latter databases employ the Debye model to obtain thermal pressure, but polynomial expressions for heat capacity and thermal expansivity are used in the data base of Piazzoni et al. (2007).

An important difference is that the formalism used by Jacobs and de Jong (2005, 2007) incorporates more details of the phonon spectrum than the Debye formalism allows.

Such details are essential because the Debye model was shown to result in insufficient accuracy in the heat capacity from 0 K up to the melting point of forsterite (Jacobs and de Jong, 2005a). That compelled them to employ Kieffer's (1979) model to represent the experimental data associated with the vibrational density of state (VDoS), such as vibrational frequency-temperature-pressure measurements derived from Raman and infrared spectroscopy. Kieffer's model appeared also to be required for other minerals such as MgSiO_3 perovskite.

Secondly, the formalism of Jacobs and de Jong (2005, 2007) incorporates intrinsic mode anharmonicity. These data became available for a number of mantle materials, such as forsterite, perovskite and akimotoite through the work of Gillet et al. (1991), Gillet et al. (2000) and Reynard and Rubie (1996) respectively. Jacobs and de Jong (2005, 2007) demonstrated that intrinsic anharmonicity significantly affects slope and position of phase boundaries in the magnesium-olivine and magnesium-pyroxene system.

Thirdly the formalism is constructed in such a way that, at zero Kelvin, static lattice properties are calculated, which can be constrained by or compared with ab initio methods.

The fourth difference is that the phonon-thermodynamic formalism incorporates electronic and magnetic properties, which is shown by Jacobs and de Jong (2009) to be a requirement for accurate representation of Fe bearing mantle materials. They showed that the vibrational formalism is suitable to predicting mixing properties in olivine. Their incorporation of iron in the system $\text{MgO} - \text{SiO}_2$ is work in progress. Because the phonon-thermodynamic model incorporates more physical properties relative to polynomial models used by e.g. Fabrichnaya (1998), Holland and Powell (1998), Saxena (1996), Jacobs et al. (2005b), the thermodynamic description of substances is more unambiguously constrained.

A.2 Computation of thermophysical parameters with the phonon-thermodynamic model

The central equation of the phonon-thermodynamic model is the expression for the Helmholtz energy of a material. For insulating materials such as present in the system $\text{MgO} - \text{SiO}_2$, The Helmholtz

energy is written as:

$$A(T, V) = U^{ref}(V_0^{st}) + U^{st}(V) + A^{vib}(T, V) \quad (6)$$

All thermodynamic properties are derived from the Helmholtz energy by applying classical thermodynamic theory. The system $\text{MgO} - \text{SiO}_2$ contains insulator materials and for such case the Helmholtz energy is expressed as the sum of a static lattice and a vibrational contribution:

The first term on the right-hand side of eqn. (6) is a reference energy contribution at zero Kelvin and zero pressure for a substance in which no vibrational motion is present. The molar volume of this static lattice is denoted by V_0^{st} . The second term represents the change in energy resulting from the change in volume of the static lattice from V_0^{st} to V . This term is calculated by applying an equation of state, such as a third order Birch-Murnaghan equation of state or the universal equation of state developed by Vinet et al. (1987). The last term represents the vibrational contribution to the Helmholtz energy and is expressed in vibrational mode frequencies and their mode Grüneisen parameters. Taking the isothermal volume derivative of eqn. (6) results in the expression for the equation of state, which is expressed as the sum of the static lattice and vibrational (thermal) contribution:

$$P(T, V) = P^{st}(V) + P^{vib}(T, V) \quad (7)$$

At a selected condition of pressure and temperature the molar volume of the material is solved numerically from (7). Density, ρ , in (3) is then calculated from the molar volume using the molar masses of the endmembers. Once the volume is known, all other thermodynamic properties follow by taking the appropriate differential of eqn. (6).

The calculation of the heat capacity, c_P , in (5) is more involved and proceeds in several steps by making use of the well-known thermodynamic relation:

$$c_P = c_V + \alpha^2 K V T \quad (8)$$

The explicit expressions for the heat capacity at constant volume, c_V , isothermal bulk modulus, K , and thermal expansivity, α , are derived from eqn (6) by taking the appropriate differentials:

$$c_V = T \left(\frac{\partial S}{\partial T} \right)_V = -T \left(\frac{\partial^2 A}{\partial T^2} \right)_V \quad (9)$$

$$K = -V \left(\frac{\partial P}{\partial V} \right)_T = V \left(\frac{\partial^2 A}{\partial V^2} \right)_T \quad (10)$$

$$\alpha = \frac{1}{K} \left(\frac{\partial P}{\partial T} \right)_V = -\frac{1}{K} \left(\frac{\partial}{\partial T} \left[\frac{\partial A}{\partial V} \right]_T \right)_V \quad (11)$$

All differentiations in (9,10,11) can be derived analytically from the explicit expression for the Helmholtz energy (Jacobs and de Jong, 2007).

In Kieffers (1979) model the directionally averaged longitudinal and transverse sound velocities are related to the angular vibrational frequency $\omega_{i,max}$ by:

$$u_i = \frac{1}{4} \left(\frac{4\pi ZV}{3N_A} \right)^{1/3} \omega_{i,max} \quad (12)$$

In (12) u_1 and u_2 represent the sound velocities of the transverse (shear) waves and u_3 that of the Voigt-Reuss-Hill longitudinal wave. The shear wavespeed is calculated according to:

$$v_s^2 = \frac{G}{\rho} = \frac{3}{4} (u_3^2 - v_B^2) = \frac{3}{4} \left(u_3^2 - \frac{K_S}{\rho} \right) \quad (13)$$

The P - T phase diagram for a specific composition in the MgO – SiO₂ system is derived from the Gibbs energy of each mineral phase. The Gibbs energy determines the stability of the phase and is derived from the Helmholtz energy as:

$$G(P, T) = A(T, V) + P \cdot V \quad (14)$$

As stated above, the volume of the mineral phase at a specific pressure and temperature is derived from (7) resulting in a value for the $P \cdot V$ term in (14). This value combined with the value for the Helmholtz energy derived from (6) results in the value of the Gibbs energy of each phase. From the Gibbs energy values of each phase and applying general rules of classical thermodynamics it is straightforward to derive which phase assemblage is stable at given (P, T) , (Connolly, 2005).

A.3 Computational aspects

The model equations (1),(2) and (5) are solved on a two dimensional quarter circle cylindrical domain covering the full depth of the mantle, using finite element methods based on the SEPRAN finite element package (Segal and Praagman, 2000).

Free slip impermeable boundaries are prescribed for the Stokes equation (2). We apply cartesian coordinates and local transformations to implement the mechanical boundary conditions on the curved boundaries (Cuvelier et al., 1986). Temperature is prescribed at the bottom and top horizontal surfaces with a driving temperature contrast $\Delta T = 3500$ K. Zero heatflux symmetry conditions apply on the radial boundaries.

Approximately $2 \cdot 10^5$ nodal points are used spanning 64000 quadratic elements for the velocity field using 227 and 567 nodal points in the radial and horizontal direction respectively. A fourfold subdivision is applied on the vertices of the 7-point velocity elements to obtain linear temperature elements for the energy equation (van den Berg et al., 1993). Grid refinement is applied with nodal point spacing reduced to approximately 6 km in the depth interval containing the transition zone.

An integrated finite element method in terms of the velocity and pressure fields is combined with a direct solver for the solution of the Stokes equation where renumbering of the degrees of freedom is applied to reduce the profile of the stiffness matrix.

The energy equation is integrated in time with a predictor-corrector method, applying a Courant type time step control (van den Berg et al., 1993).

Necessary thermo-physical properties density, thermal expansivity and specific heat, are computed through a general interface procedure using equidistant tabulation and bi-linear interpolation of these properties in the P, T domain relevant for the mantle $300 \text{ K} < T < 4000 \text{ K}$, $\delta T = 2 \text{ K}$, $0 < P < 140 \text{ GPa}$, $\delta P = 0.066 \text{ GPa}$. This represents a flexible and efficient tool that facilitates the use of alternative thermodynamic data bases (van den Berg et al., 2002, Jacobs et al., 2006).

P, T domain tabulation is essential since pointwise calculation of thermo-physical properties is not feasible in view of the large number of evaluations of these properties. This becomes clear from the following assessment of the number of evaluations involved per integration time step, for a general model case with N_e Stokes elements.

For the Stokes equation only ρ needs to be computed. For the energy equation the values of ρ , c_P and α must be computed (factor 3 in (15)). A four times higher number of elements is used

840 for the energy equation (factor 4 in (15)). $n_g = 7$ gauss integration points are used per element.
 841 These evaluations are repeated two times per integration time step in a predictor/corrector scheme,
 842 resulting in the following number of thermo-table evaluation per time step.

$$843 \quad N_{eval} = (N_{stokes} + N_{energy}) \times 2 = (N_e \times n_g + 3 \times N_e \times 4 \times n_g) \times 2 = 26 \times N_e \times n_g \quad (15)$$

844 This amounts to $N_{eval} = 1.2 \cdot 10^7$ table evaluations per integration time step for the finite element
 845 mesh size used in our model calculations where $N_e = 64000$.

Figure 1: Pressure-depth dependence of the seismic, longitudinal (v_p), shear (v_s) and bulk (v_b) wavespeeds, after (Jacobs and de Jong, 2009). Solid red curves are model predictions for mantle adiabats (1420 K) and a similar mineral composition as in our present model. Grey fields represent wavespeeds for compositions ranging from Mg_2SiO_4 to MgSiO_3 . Green curves represent a model with $(\text{Mg}_{0.9}\text{Fe}_{0.1})_2\text{SiO}_4$. The dashed green curve is from a model based on calphad type parameterizations for the same olivine composition (Jacobs and de Jong, 2005b). Discrete symbols represent the PREM model.

Figure 2: Heat capacity of forsterite at 1 bar pressure from alternative mineral physics models compared with observational data. The curve labeled MGK-a corresponds to the model of Jacobs and de Jong (2007) that includes a Kieffer type VDoS and intrinsic anharmonicity. The curve labeled MGD is obtained from the Mie-Grüneisen-Debye model employed by Stixrude and Lithgow-Bertelloni (2005). Experimental data are from Orr (1953), triangles, Robie (1982), squares, Watanabe (1982), diamonds, Ashida et al. (1987), inverted triangles, Gillet et al. (1991) filled circles.

Figure 3: (a) Full domain temperature snapshot showing two colliding cold downwellings and a hot mantle upwelling. Black boxes represent zoom-in windows corresponding to frames (b,c). (b) zoom-in temperature of the cold downwellings flow. (c) zoom-in temperature of the shallow head of a hot upwelling. (d) Full domain distribution of the lateral variation of the shear wavespeed δv_s with respect to the smooth background profile. (e) zoom-in of δv_s of the cold downwelling. (f) zoom-in of δv_s of the hot mantle plume head.

849

Figure 4: Isobaric wavespeed-temperature data samples taken from the model output shown in Fig. 1. Samples correspond to finite element grid points in 20 km deep zones centered at 300, 636 and 797 km depth. Curve labels indicate the mineral associations that correspond to the stability fields in the phase diagram shown in Fig. 5. Sampling depths indicated in the plot labels correspond to model pressure values of 9.4, 21 and 27.5 GPa respectively.

Figure 5: Phase diagram of the olivine-pyroxene magnesium endmember model used in the convection model. Adiabats of potential temperature T_p between 1400 and 1800 K are plotted for reference. Solid curves represent radial temperature profiles, with angular position shown in the legend, and corresponding to the dashed lines in Fig. 6a. Horizontal dashed lines, labeled with a pressure value, correspond to the isobaric cross sections shown in Fig. 4.

Figure 6: (a) Zoom-in on the spatial mineral phase distribution near the cold downwelling shown in Fig. 3b,e. Dashed lines indicate the position of radial profiles. (b) Similar zoom-in as in (a) near the hot mantle region shown in 3c,f. Individual mineral domains in (a,b) have been labeled according to the phase diagram shown in Fig. 5. Frames (c,d) show the lateral variation of the shear wavespeed, δv_s for the same zoom-in windows as in (a,b) respectively.

Figure 7: Radial profiles in zoom-in windows of Fig. 6. Temperature (left) and shear wavespeed (right). The (global) laterally averaged temperature is plotted for reference as a black curve in frames (a,c). The dashed curve in the frames (b), (d) represents the smoothed background wavespeed profile. Sharp transitions in the velocity profiles are labeled by the corresponding phase transitions that can be identified in the profiles drawn in Fig. 6.

Figure 8: Spectral analysis of a single unfiltered reflectivity time trace plotted shown in frame (a), corresponding to the highlighted red trace in Fig. 9(a). (b) powerspectrum of the time trace in (a), showing that the peak power is below the frequency of 0.1 Hz, corresponding to wavelength's from 50 to 60 km in the transition zone.

Figure 9: Reflectivity-time traces derived from the s-wavespeed distribution corresponding to the zoom-in window shown in Fig. 6a,c. The corresponding s-wavespeed distribution is shown in Fig. 10. The red colored trace corresponds to Fig. 8. (a) shows the raw data derived from the convection model. (b) and (c) show corresponding time traces after application of low-pass Butterworth filters with corner-period settings of 20 sec. and 40 sec. respectively.

855

Figure 10: Shear wavespeed, in km/s, distribution for the zoom-in window of Fig. 6a,c. Labeled regions of contrasting shear wavespeed correspond to the contrasting mineral phase regions in Fig. 6. The wavespeed contrasts of the different mineral phase domains are closely correlated with the 'signal phases' of the reflectivity sections shown in Fig. 9.