Lipid profile of tenon's capsule in glaucoma patients

Alain M. Bron, Lionel Brétillon, Stéphane Grégoire, Niyazi Acar, Catherine Creuzot-Garcher

► To cite this version:

Alain M. Bron, Lionel Brétillon, Stéphane Grégoire, Niyazi Acar, Catherine Creuzot-Garcher. Lipid profile of tenon's capsule in glaucoma patients. Association for Research in Vision and Ophthalmology annual Congress, May 2009, Fort Lauderdale, Florida, United States. pp.E-Abstract 2677. hal-00758438

HAL Id: hal-00758438

<https://hal.science/hal-00758438>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Lipid Profile of Tenon's Capsule In Glaucoma Patients

A. M. Bron^{1,2}, L. Bretillon², S. Gregoire², N. Acar² and C. Creuzot-Garcher^{1,2}

¹Ophthalmology, University Hospital, Dijon, France

²INRA, University of Burgundy, Eye & Nutrition Research Group, Dijon, France

Commercial Relationships: A.M. Bron, None; L. Bretillon, None; S. Gregoire, None; N. Acar, None; C. Creuzot-Garcher, None.

Support: None.

Clinical Trial: www.clinicaltrials.gov NCT01021657

Abstract

Purpose: To provide data on the lipid profile of the Tenon's capsule in Human glaucomatous eyes and to establish possible relationships with the short term outcome of filtration surgery.

Methods: Fragments of Tenon's capsule were collected during glaucoma surgery in patients suffering from open angle glaucoma (10 trabeculectomies and 17 non penetrating deep sclerectomies). The polyunsaturated fatty acid (PUFA) composition of the human samples was determined by gas chromatography. A successful outcome after glaucoma surgery was defined by an intraocular pressure (IOP) ≤ 15 mmHg at 6 months without any IOP-lowering agent.

Results: We included 27 patients (12 males, 15 females) with a mean age of 65.5 ± 12.1 years. Twenty three surgeries were successful according to our criteria. The IOP before and 6 months after the surgery was 20.8 ± 7.1 mm Hg, range [13-37] and 11.7 ± 3.2 mmHg, range [6-18] respectively. The percentages of the main PUFAs were as follows: arachidonic acid $5.33 \pm 3.66\%$, docosaexaenoic acid $0.88 \pm 0.62\%$, total $\omega 3$ $1.86 \pm 1.05\%$, total $\omega 6$ $18.97 \pm 5.94\%$, $\omega 6/\omega 3$ ratio $11.86 \pm 3.95\%$. We did not find any statistically significant correlation between PUFAs content of Tenon's capsule, IOP at 6 months, delta IOP and surgical outcome.

Conclusions: Fatty acid composition of Tenon's capsule was highly variable among patients. In this small series we failed to identify a specific PUFA profile of Tenon's capsule as a risk factor for failure of glaucoma surgery.

Keywords: anterior segment • lipids • clinical (human) or epidemiologic studies: outcomes/complications