Fumonisin in Catalonia (Spain) and exposure assessment of special population groups.

German Cano Sancho, A.J. Ramos, S. Marin, V. Sanchis

► To cite this version:

German Cano Sancho, A.J. Ramos, S. Marin, V. Sanchis. Fumonisin in Catalonia (Spain) and exposure assessment of special population groups.. Food Additives and Contaminants, 2012, 29 (5), pp.1. 10.1080/19440049.2011.644813 . hal-00780447

HAL Id: hal-00780447

<https://hal.science/hal-00780447>

Submitted on 24 Jan 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Fumonisin in Catalonia (Spain) and exposure assessment of special population groups.

Journal:	<i>Food Additives and Contaminants</i>
Manuscript ID:	TFAC-2011-299.R1
Manuscript Type:	Original Research Paper
Date Submitted by the Author:	22-Nov-2011
Complete List of Authors:	Cano-Sancho, German; University of Lleida, Food Technology Ramos, Antonio; University of Lleida, Food Technology Marín, Sonia; University of Lleida, Food Technology Sanchis, Vicent; University of Lleida, Food Technology
Methods/Techniques:	Chromatography - HPLC, Risk assessment - modelling
Additives/Contaminants:	Fumonisin, Mycotoxins - fumonisin
Food Types:	Baby food, Beer
Abstract:	<p>Fumonisin B1 (FB1) and B2 (FB2) are mycotoxins produced by <i>F. verticillioides</i> and <i>F. proliferatum</i> that commonly contaminate corn. The objectives of this study were, in one hand, to study the occurrence of fumonisin in Catalonia (north-eastern region of Spain), and, in the other hand, to assess the exposure of Catalan population to these mycotoxins. Contamination data was provided by a wide survey, where 928 individual samples were pooled to analyze 370 composite samples. Fumonisin were extracted and purified using immunoaffinity columns, and determined by means of HPLC with fluorescence detector. The raw consumption data came from a nutritional study specifically designed to assess the dietary intake of the main foodstuffs related to fumonisin contamination for all population age groups. In addition, two special groups were selected to be specially related with maize consumption: immigrants and celiac sufferers. Contamination and consumption data were combined by simulation using an essentially parametric (P-P) method. The P-P method draws sampling values from distribution functions fitted to consumption and contamination datasets. Moreover, to quantify the accuracy and reliability of the statistics estimates, we built the related confidence intervals using a Pseudo-Parametric bootstrap method. Considering the results from this study, fumonisin can be commonly found in some commodities from Catalan market, such as beer, corn snacks or ethnic foods, however, in any case these values exceed the maximum EU</p>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

	levels. The most exposed group should be expected to be infants followed by immigrants, but in all cases, they were far from the TDI of 2 µg/ kg bw/day.

SCHOLARONE™
Manuscripts

For Peer Review Only

Fumonisin in Catalonia (Spain) and exposure assessment of special population groups

Cano-Sancho*, G., Ramos A.J., Marín S. and Sanchis V

Applied Mycology Unit, Food Technology Department, University of Lleida, XaRTA-UTPV, Rovira Roure 191, 25198, Lleida, Spain.

*Corresponding author: German Cano-Sancho. Telephone number: (+0034) 973702670.

Fax number: (+0034) 973702596. E-mail address: gcano@tecal.udl.cat

Abstract

Fumonisin B₁ (FB₁) and B₂ (FB₂) are mycotoxins produced by *F. verticillioides* and *F. proliferatum* that commonly contaminate corn. The objectives of this study were, on the one hand, to study the occurrence of fumonisins in Catalonia (north-eastern region of Spain), and, on the other hand, to assess the exposure of Catalanian population to these mycotoxins. Contamination data was provided by a wide survey, where 928 individual samples were pooled to analyse 370 composite samples. Fumonisin were extracted and purified using immunoaffinity columns, and determined by means of HPLC with fluorescence detection. The raw consumption data came from a nutritional study specifically designed to assess the dietary intake of the main foodstuffs related to fumonisins contamination for all population age groups. In addition, two special groups

were selected to be specially related with maize consumption: immigrants and celiac sufferers. Contamination and consumption data were combined by simulation using an essentially parametric (*P-P*) method. The *P-P* method draws sampling values from distribution functions fitted to consumption and contamination datasets. Moreover, to quantify the accuracy and reliability of the statistical estimates, we built the related confidence intervals using a Pseudo-Parametric bootstrap method. Considering the results from this study, fumonisins can be commonly found in some commodities from the Catalonian market, such as beer, corn snacks and ethnic foods, however, in any case these values were well below the maximum EU levels. The most exposed group should be expected to be infants followed by immigrants, but in all cases, they were far from the TDI of 2 µg/ kg bw/day.

Keywords:- Occurrence, exposure assessment, fumonisins, cereal food, gluten-free products, ethnic food

Introduction

Fumonisin B₁ (FB₁) and B₂ (FB₂) are mycotoxins mainly produced by *Fusarium verticillioides* and *F. proliferatum* that commonly contaminate corn (Nelson et al. 1992). Fumonisin (FBs) occur mainly in maize and maize-based foods, therefore populations with high maize consumption could be exposed to significant amounts of these mycotoxins through the ingestion of fumonisin contaminated maize (Marasas 1996; Shephard et al. 1996; Visconti et al. 1996; WHO 2001).

FBs have a remarkable structural similarity to sphingolipids (Merrill and Sweeley 1996; Riley et al. 2001). This group of mycotoxins, especially FB₁, potently inhibit the enzyme ceramide (CER) synthase which catalyzes the acylation of sphinganine (Sa) and reacylation of sphingosine (So). The inhibition of CER synthase by FBs increases the intracellular Sa concentration, main contributor to the toxicity and carcinogenicity of FB₁ (Wang et al. 1991; Merrill et al. 1993; Yoo et al. 1996; Riley et al. 2001). Based on this biological perturbation, elevation of Sa to So, or Sa 1-phosphate to So 1-phosphate, ratios in tissues, urine and blood, have been proposed as potential biomarkers of fumonisin exposure in various animal species (Wang et al. 1992; Riley et al. 1993; Wang et al. 1999; Van der Westhuizen et al. 2001; Kim et al. 2006; Tran et al. 2006; Cai et al. 2007; Cano-Sancho et al. 2011b).

Human exposure to fumonisin contaminated commodities has been linked to oesophageal and liver cancer in South Africa and China (Sydenham et al. 1990; Yoshizawa et al. 1994). Acute and chronic toxicity of FBs has been largely demonstrated in several animal species, including carcinogenicity and cardiovascular toxic effects (Gelderblom et al. 1988, 1991). FB₁ is a cancer promoter, but a poor cancer initiator. It is not genotoxic because FB₁ does not induce unscheduled DNA synthesis in primary rat hepatocytes (Norred et al. 1992).

Based on toxicological evidence, the International Agency for Research on Cancer (IARC) has classified FB₁ as possibly carcinogenic (group 2B) to humans (IARC 2006). The Joint FAO/WHO Expert Committee on Food Additives (JECFA) evaluated FBs and

allocated a provisional maximum tolerable daily intake (PMTDI) of 2 µg/kg body weight/day to FBs on the basis of the non observed effect level (NOEL) of 0.2 mg/kg body weight/day and a safety factor of 100 (WHO 2001). This PMTDI for FBs has recently been confirmed by the Scientific Committee on Food (SCF 2003).

The European Commission established a maximum FBs levels of 1000 µg/kg in maize and maize-based food intended for human consumption, 800 µg/kg in maize-based breakfast cereals and snacks and 200 µg/kg in maize-based infant food (European Commission 2006b, 2007).

In the Mediterranean countries, maize is more widely used for animal nutrition than for human consumption, but some countries traditionally use the maize in regional cooking as *broa* (maize-based bread from Portugal) and polenta (from Italy). As a consequence of the use of maize grits as adjunct in brewing process, aflatoxins and FBs can be also found in beer (Torres et al. 1998; Pietri et al. 2009). The natural occurrence of FBs in corn or corn-based foods has been studied largely (Sydenham et al. 1991; Shephard et al. 1996; SCOOP 2003; Castelo et al. 1998; Hlywka et al. 1999), and surveys have been conducted in several cases in corn-based products marketed in Spain (Sanchis et al. 1994, 1995; Torres et al. 1998; Castellá et al. 1999; Velluti et al. 2001). Dietary intake of FBs has been previously assessed in population from France, USA, Netherlands, Canada or South Africa (Kuiper-Goodman et al. 1996; De Nijs et al. 1998; Humphreys et al. 2001; Leblanc et al. 2005; Shephard et al. 2007).

In the framework of the Project to Assess the Exposure of Catalanian Population to the Mycotoxins, and following the line of our previous studies (Cano-Sancho et al. 2009, 2010, 2011c), the objectives of this study were, on the one hand, to study the occurrence of FBs in Catalonia (north-eastern region of Spain), and, on the other hand, to assess the exposure of Catalanian population to these mycotoxins.

Materials and methods

Fumonisin analysis

Samples

During 2008 and 2009, 185 samples of sweet-corn, 216 samples of corn-based snacks, 216 beer samples, 168 corn flakes, 90 baby food samples were obtained in six hypermarkets and supermarkets from twelve main cities (Tortosa, Tarragona, Reus, Vilanova i la Geltrú, l'Hospitalet de Llobregat, Barcelona, Terrassa, Sabadell, Mataró, Girona, Manresa and Lleida) of Catalonia, representative of 72 % of the population. From each supermarket or hypermarket, 3 samples (if present) of each product were randomly taken. The level of FB₁ and FB₂ was determined in a total of 72 composite samples of each category by equally pooling the 2 or 3 items taken from each store (depending of availability) to obtain an analytical sample or “composite”: 12 cities x 6 stores/city = 72 samples/category. However, in some cases, 3 items or any items were not available in the store, thus, less than 72 composites could be obtained. Additionally, 18 gluten-free food samples (mainly maize-based bread and pasta) and 35 ethnic food samples, such as Mexican tortillas, corn flour or cuscus, were purchased in specialised stores to be analysed individually. Regarding brands, thirty-one sweet corn, seventy-nine corn-based

snacks, sixty-four beers, thirty-two corn flakes different brands were purchased, which can be considered the majority of market share in Catalonia of these products, as well as in the rest of Spanish market. The samples were transported and stored under suitable conditions until analysis.

Analytical procedure

Fumoniprep[®] immunoaffinity cleanup columns (IAC) (R-Biopharm, Rhône Ltd Glasgow, UK) were used to extract FB₁ and FB₂ from beer samples. A volume of 5 mL of beer previously degassed in ultrasonic bath during 40 minutes was mixed with 15 mL of phosphate buffer solution (PBS; 0.8 % NaCl, 0.12% Na₂HPO₄, 0.02% KH₂PO₄, 0.02% KCl) and drained through the IAC. The column was washed with 20 mL of PBS solution and FBs were eluted with 1.5 mL of methanol grade HPLC and 1.5 mL of milli-Q water. Regarding solid maize-based samples, 10 g of ground sample composite was mixed with 1 g NaCl, and 50 mL of extract solution (50% water, 25% methanol, 25% acetonitrile) for 20 minutes and filtered. 10 mL of filtrate was diluted with 40 mL of PBS and drained through the IAC and follows as described previously.

Fluorescent derivatives of FB₁ and FB₂ were obtained using pre-column derivatisation with an *o*-phthaldialdehyde (OPA) solution prepared diluting 40 mg of *o*-phthaldialdehyde with 1 mL of methanol HPLC grade and mixed with 5 mL of Na₂B₄O₇·10H₂O (0.1M) and 50 µL de 2-mercaptoethanol. Derivatisation was conducted mixing 200 µL of eluate with 200 µL of OPA solution for 30 seconds in vortex. Chromatography equipment: Separations Module Alliance 2695 Waters®, analytical

column Waters Spherisorb® 5µm ODS2, 4.6 x 150 mm, Multi λ Fluorescence Detector Waters 2475®, kept at 35 °C and a flow-rate maintained at 1 mL/min. Injection volume was 100 µl. Mobile phase was based on a methanol and 0.1M sodium dihydrogen phosphate (77:23, v/v) solution. Excitation and emission wavelength were 335 nm and 440 nm, respectively.

The analytical method used for FBs was assessed for selectivity, linearity, and precision. Selectivity was checked by injecting 100 µl of mycotoxin standard solution three times before injecting extracted samples and comparing the peak retention times and the fluorescence spectra of the substances that produced these peaks. Standard curves were generated by linear regression of peak areas against concentrations.

Precision and recovery were established by determination of FB₁ and FB₂ levels, spiked in each food categories by triplicate. The concentrations of fumonisins fortified in tested samples were ranged from 25 µg/kg to 660 µg/kg. Recovery was determined by comparing the absolute responses of FBs, with the absolute responses of calibration standards. The limit of detection (LOD) was considered to be three fold the signal of blank noise, and the limit of quantification (LOQ) was considered equal to 3×LOD.

Recovery ranges were 76.5 ± 11.5 - 109.7±12.3 % in corn snacks; 101.5±11.8 - 104.0±7.3 % in sweet corn; 91.6±8.4 - 107.5±5.0 in corn flakes, 93.0±14.0 - 108.0±10.0 % in beer and 90.7±15.1 - 115.5±27.9 % in baby food. Repeatability was expressed as relative standard deviation (RSDr), it ranged between 11.3 and 15.0 % in corn snacks; between 7.0 and 11.6 % in sweet corn; between 4.6 and 9.1 % in corn flakes; between

10.0 and 15.0 % in beer; between 16.6 and 24.1 % in baby food. The mean limit of detection (LOD) was 3.9 µg/kg in beer and 5.2 in solid based foods. These values were in accordance to performance criteria established by Commission Regulation (EC) N° 401/2006 (European Commission 2006a).

Fumonisin dietary intake assessment

The main problem in assessing the intake of mycotoxins is related to few representative available dietary data regarding foods contaminated with mycotoxins. There are some methods developed to assess dietary intake overall known as market basket, 24-hour dietary recall and food record methods, food-frequency methods or dietary history. Given that corn-based products consumption is considered as sporadic or casual among Catalonian population, food-frequency methods should be chosen because it may be advantageous to sacrifice precise intake measurements in exchange for more crude information related to an extended period of time (Walter 1998). In this study, food dietary intake was assessed through a specific Food Frequency Questionnaire (FFQ) developed for Catalonian population including those foods typically consumed in the region which may be potentially contaminated with these mycotoxins. According to World Health Organization advices, studies to assess dietary intake of chemical contaminants, should show the significant intake among standard population, with all population groups that could have different dietary patterns. Therefore, five different population groups were considered: infants (0-3 years), children (4-9), adolescents (10-19 years), adults (20-65 years) and elders (> 65 years). Moreover, adult celiac sufferers and adults with ethnic dietary pattern were included in the nutritional study, and studied

separately. FFQ consisted of 38 items of specific foods worldwide known to be the most important food contaminated by mycotoxins studied, excluding those foods not consumed in the region. Concerning frequency of consumption, 5 response options ranging from never to annually, were considered. Quantities were assessed by portion size with the aid of a series of colour photograph models. Finally, 76 seniors, 720 adults, 235 adolescents, 69 parents of children, 164 parents of infants, 70 adult celiac sufferers and 56 adults with ethnic dietary pattern were interviewed in 2008-2009 by trained interviewers. In the present study, the adults were treated separately according to their sex, with a major population group of 336 adult males and another of 384 adult females. Individuals were from 89 cities and towns from Catalonia. Study level, dietary habits, sex, age, ethnics and anthropometric data were considered in the questionnaire.

Exposure Assessment through Direct Approach

If we assume independency between consumption ($C_{\pi j}$) and contamination (T_j), as well as between their products, we can estimate the mean exposure of the population π with the population sample π_0 as follows:

$$\hat{E}_{\pi_0} = \sum_{j=1}^p \bar{C}_{\pi_0, j} \bar{T}_j$$

Where $\bar{C}_{\pi_0, j}$ is the arithmetical mean of the normalised consumption of the foodstuff j in the population group π_0 , and $\bar{T}_j$ is the arithmetical mean of the available contamination data of the related food. We could consider this approach to be a theoretical approach, but it would have to be a (very) simplified theoretical approach. For example, this simplified approach does not enable us to calculate complicated statistics such as high quantiles. Simulation methods would therefore be required to obtain estimates of these statistics.

Finally, we will consider this approach to be a direct approach. Left-censored data (those samples below LOQ) was substituted by LOD/2 in order to calculate the mean toxin level used in the exposure assessment model.

Exposure Assessment through Stochastic Approach

Stochastic procedure used is based on the Parametric-Parametric (P-P) method reported extensively in Gauchi and Leblanc (2002) and Cano-Sancho et al. (2011b). In brief, for each consumption and contamination dataset, the gamma *pdf* was fitted with the method of maximum likelihood using the CAPABILITY procedure of SAS software (SAS 2010). The appearance of the histograms was continuous and only one distribution was considered pooling all age groups data, fitting all data to gamma *pdf*. In case of gluten-free foods and sweet-corn, the mean contamination was used in the simulations because not enough data were available for computations. Left-censored was treated as the previous section, replacing non quantified samples by 0.5xLOD.

The *P-P* method used to estimate the normalised exposure *k* of the *S* simulation set was built as follows:

$$\hat{E}_k^{[P-P]} = \sum_{j=1}^p \tilde{c}_{i(j)} \tilde{t}_j$$

where $\tilde{c}_{i(j)}$ is a random normalised consumption *i* for the foodstuff *j*, drawn from $\hat{C}_j$, the corresponding adjusted gamma *pdf*, and $\tilde{t}_j$ is a random contamination for the foodstuff *j*, drawn from $\hat{T}_j$, the corresponding adjusted gamma *pdf*.

The mean of normalised exposures over the simulation set *S* was then estimated using the following equation, where *n* was the number of random deviates drawn (10,000 in the present study):

$$\hat{E}_S^{[P-P]} = \frac{1}{n} \sum_{k=1}^n \hat{E}_k^{[P-P]}$$

Other statistics were directly computed on the histogram built with the simulations of the S set, and statistics were also estimated from the lognormal and gamma fitted *pdfs* (results not shown).

Pseudo-parametric bootstrap confidence intervals were built as reported by Gauchi and Leblanc (2002), by randomly drawing B samples of size n_{π_0} in the exposure simulation set S (being B equal to 10,000). The boundaries of the 95% confidence interval are calculated taking the 0.025th and 0.975th empirical quantiles of the final bootstrap distribution.

Results

Occurrence of total fumonisins in food

FB₁ was the major fumonisin found in all food categories. In beer and corn snacks both mycotoxins were present simultaneously, and the mean ratios FB₁/FB₂ were 2.3 and 1.9, respectively. According to Table 1 and Figure 1, highest occurrence of total fumonisins (FB₁+FB₂) was found in beer samples with 90 % of positive samples, followed by corn snacks (61 %), ethnic foods (51 %) and corn flakes (39 %). Baby foods and gluten-free foods showed low levels of positive samples (23.3 and 5.6 %, respectively) and they were not detected in sweet corn. Positive samples of corn snacks and corn flakes presented moderated mean values (119.1±83.1 µg/kg and 78.9±27.9 µg/kg, respectively). The highest values were found in ethnic food samples, with a mean of 196.7 ± 191.8 and a maximum level of 682.2 µg/kg. Despite the high maximum value, no sample exceeded

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

the limit established by EU (European Commission 2007). The gluten-free foods and baby food showed the lowest mean levels of positive samples.

Place Table 1 about here and Figure 1 about here.

Fumonisin dietary intake assessment

Through the specific nutritional study performed with a sample of Catalanian inhabitants during 2008 and 2009 we could better estimate the dietary patterns of those commodities related with fumonisin contamination. As shown in Table 2, the corn flakes and corn snacks were mainly consumed by children and adolescents, while beer was most widely consumed by immigrants and adult males. Celiac sufferers commonly consumed gluten-free products as bread and pasta, and other maize based food as breakfast cereals and corn snacks. Most of immigrants consumed food products imported from their origin countries, becoming a staple food in their current diet.

Place Table 2 about here.

Fumonisin dietary intake of the Catalanian population sample was assessed through a direct method and by means of the simulation method described above. The mean and variance of fumonisin dietary intake estimated through the direct approach for all population groups are presented in table 3.

Place Table 3 about here

1
2
3 Considering these estimates, the most exposed population group was the infants, followed
4
5 by immigrants, children and adult males. Differences were found between adult males
6
7 and females.
8
9

10
11
12 Concerning the dietary intake estimated through the simulation procedure, mean values
13
14 were commonly higher than those calculated using the direct method, with the exception
15
16 of celiac sufferers (See Table 4 and Figure 2). Stochastic method was not applied to
17
18 elders because few consumers of each item was found in this group and commonly those
19
20 consumers reported a low level of consumption in each category.
21
22
23

24
25
26
27 *Place Table 4 and Figure 2 about here about here.*
28
29
30

31
32 The highest means were estimated for infants, followed by the immigrants and adults.
33
34 Celiac sufferers showed the lowest exposure estimates. Pseudo-parametric bootstrap
35
36 confidence intervals permitted to assess the reliability and accuracy of the computations.
37
38 We observed that these intervals decreased when we estimated the highest quantiles.
39
40 Extremely wide confidence intervals were built for the 95th and 99th quantiles, especially
41
42 for immigrants and children. It is an accepted fact that it is very difficult to obtain good
43
44 accuracy for the 99th-quantile, in particular (Breiman et al. 1990).
45
46
47
48
49

50
51 All estimates were far from the TDI of 2000 ng/kg body weight/day, including the
52
53 highest quantiles of immigrants and infants (the most exposed groups).
54
55
56
57
58
59
60

Discussion

Occurrence of total fumonisins in corn-based food

This is the first study conducted in Spain with a highly representative sampling method, using a high number of individual samples of the main corn-based food categories collected around the Catalonian geography to assess a wide pool of analytical samples, including baby foods, ethnic foods or gluten-free foods not reported until now. Although our study was conducted in Catalonia, the main brands marketed in Spain have been purchased and analysed, therefore, results from our study could be extrapolated to the Spanish market.

A previous study conducted in Spain showed smaller concentrations than our study for corn snacks and corn flakes but the authors found contaminated samples of sweet corn with respective mean values of 46.27, 17.03 and 55.96 µg/kg, all of them determined by HPLC (Velluti et al. 2001). D'Arco et al. (2007) compared organic and conventional maize-based food and they found higher percentage of contaminated samples in organic commodities (78 %) than in conventional food (16 %), moreover, the maximum and highest mean was found in the organic category (235 vs 51 µg/kg). In the present study, we analyzed mainly conventional food because the consumption of organic maize foods is not widespread in Spain, and these products are supplied by a little number of retail stores. Regarding beer samples from Spain, a previous study performed by ELISA kit, showed similar values to our study in the range 85.53±0.34 – 4.76±1.21 µg/kg (Torres et al. 1998).

Previous studies carried out with corn-flakes, reported mean levels of contamination closest to our mean value of 78.9 ± 27.9 $\mu\text{g/kg}$, being those means 76.0 $\mu\text{g/kg}$ in France, 79.3 and 129.0 $\mu\text{g/kg}$ in Canada (Kim et al. 2003; Molinié et al. 2005; Roscoe et al. 2008); however Martins et al. (2008) did not found any sample from Portugal exceeding the detection level. Lombaert et al. (2003) showed mean levels of 33 and 14 $\mu\text{g/kg}$ for soy-based and multi-grain cereals for babies, respectively, in the line of our results. Moreover, FBs were previously analyzed in ethnic foods from United Kingdom, and only trace levels were found in cereal-based foods as noodles, rice, corn flour and pitta bread (Patel et al. 1996), quite different from our study where fumonisins were quantified in half of samples. In Spain, the ethnic food market started in the 90's, whilst the highest growth rates have occurred since 2000. Sales passed from 7.2 million dollars in 1999 to 40.6 million dollars in 2004 (Camarena and Sanjuán 2008), therefore this commodities should be taken into account as potential mycotoxin carriers.

In the present study only FB_1 and FB_2 were determined in food, however, it is a well known that FBs can be conjugated when food is thermally treated. These bound forms are not detectable using the conventional analytical techniques, but can be hydrolysed in the gastrointestinal tract and produce the toxic effect (Dall'Asta et al. 2009). Therefore, in future exposure assessment works, powerful techniques focused on direct or indirect analysis of conjugated forms should be used to improve the accuracy of the estimates.

Fumonisin dietary intake assessment

The lack of accurate and representative dietary data of corn-based food intake from Spanish population was the main problem to assess the exposure of the population to FBs. Through a specific FFQ administered to a population sample from the region, it has been possible to obtain a more accurate approximation to dietary intake habits from each population group.

Despite experts have recommended the use of stochastic approaches to provide a more realistic exposure assessment, to date, few studies have been published using this methodology to assess the exposure of populations to mycotoxins (Kroes et al. 2002; WHO 2005; EFSA 2006; Verger and Fabiansson 2008), the direct or deterministic methods being the most commonly used.

The results of exposure assessment obtained through the direct method showed infants as the main risk group, followed by immigrants, male adults and children with respective daily dietary intake of FBs of 0.156 ± 0.154 , 0.065 ± 0.069 and 0.063 ± 0.090 and 0.063 ± 0.064 $\mu\text{g/kg}$ body weight/day. A previous study conducted in Spain on dietary intake of FBs was developed using corn dietary intake data of Spanish population supplied by Food and Agriculture Organization (FAO) combined with FBs levels in non processed corn obtained from their study (Ariño et al. 2007). Despite this fact could result in overestimation of FBs intake, that study showed lower levels of exposure than our results with mean intake for adults of 0.0038 $\mu\text{g/kg}$ body weight/day. Leblanc et al. (2005) reported exposure of French population to FBs using dietary data from a strict epidemiological work ($n=3003$), showing results in the line of our study with values of

0.05 and 0.18 $\mu\text{g/kg}$ body weight/day for children mean and high consumers (percentile 95) and 0.01 and 0.06 $\mu\text{g/kg}$ body weight/day for adults mean and high consumers, respectively, our combined values (males and females) being: 0.048 and 0.16 $\mu\text{g/kg}$ body weight/day for adult and baby means. Results from SCOOP (2003) showed variable levels, some of them slightly higher than our results, as for Italian consumers or Norwegian babies values of 0.52 and 0.86 $\mu\text{g/kg}$ body weight/day were reported, respectively, while other results were in the line with ours, as those for population from Austria, Germany, Belgium or Netherlands. In all cases, these values are far from those estimated in other regions highly exposed to FBs as South-Africa, where mean daily intake of FBs ranged from 3.43 ± 0.15 to 8.67 ± 0.18 $\mu\text{g/kg}$ body weight/day (Shephard et al. 2007).

When we compared the direct method with stochastic method, we commonly found higher estimations through simulation. In the present study this was not crucial because in all cases the estimates were far from the safe level. However, this method can become mandatory in those cases when risk assessors need to accurately study the dietary intake of highly exposed populations.

To our knowledge, this is the first study to assess the exposure of special population groups to FBs. The special groups selected in the present study had dietary habits markedly different from the general population. In one hand, celiac sufferers substitute the wheat-based food for other gluten free cereals, and in the other hand, some ethnic groups partially maintain the dietary patterns from their country of origin. In Catalonia

during 2009, the immigrant population represented 15.9 % of the total population (Migracat 2010); therefore, their specificities should be taken into account to assess the exposure of this collective in other close regions.

Considering our results, despite FBs being found regularly in Catalanian market, especially in ethnic food, beer and corn snacks, all levels remain far from EU maximum levels. Babies should be expected to be the main risk group to fumonisin intake; however these results suggest that no human risk derived from FBs should be expected in any Catalanian population group. The stochastic method provides more accurate estimations than the direct method, including the variability and uncertainties in the model, as well as the possibility to compute confidence intervals to validate the estimations.

Acknowledgements

The authors would like to acknowledge Exposure Assessment of Spanish Population to *Fusarium* Toxins Project, National Plan of Spanish Government (AGL2008-05030-C02-01), Catalanian Food Safety Agency of ‘Generalitat de Catalunya’ Health Department and University of Lleida for their financial support, and the Catalanian Celiac Association for their collaboration.

Bibliography

Ariño A, Estopañan G, Juan T, Herrera A. 2007. Estimation of dietary intakes of fumonisins B₁ and B₂ from conventional and organic corn. Food Control 18:1058-1062.

Breiman L, Stone CJ, Kooperberg C. 1990. Robust confidence bounds for extreme upper quantiles. *J Stat Comp Simul.* 37:127-149.

Cai Q, Tang L, Wang J-S. 2007. Validation of fumonisin biomarkers in F344 rats. *Toxicol Appl Pharmacol.* 225:28-39.

Camarena DM, Sanjuán AI. 2008. El mercado de comida étnica en España: el caso de la comida mexicana. *Estudios Sociales: Revista de investigación científica* 16:7-37.

Cano-Sancho G, Marín S, Ramos AJ, Sanchis V. 2009. Survey of patulin occurrence in apple juice and apple products in Catalonia, Spain, and an estimate of dietary intake. *Food Addit Contam B* 2:59-65.

Cano-Sancho G, Marín S, Ramos AJ, Peris-Vicente J, Sanchis V. 2010. Occurrence of aflatoxin M₁ and exposure assessment in Catalonia (Spain). *Rev Iberoam Micol.* 27:130-135.

Cano-Sancho G, Gauchi J-P, Sanchis V, Marín, S, Ramos AJ. 2011a. Quantitative Dietary Exposure Assessment of the Catalanian Population (Spain) to the Mycotoxin Deoxynivalenol. *Food Addit Contam A* 1:1-12.

Cano-Sancho G, Marín S, Sanchis V, Colom C, Coronel MB, Ramos AJ. 2011b. Sphinganine and sphingosine levels and ratio in urine and blood samples from a Catalan population, Spain. Food Addit Contam A [In Press, doi: 10.1080/19440049.2011.576437]

Cano-Sancho G, Valle-Algarra FM, Jiménez M, Burdaspal P, Legarda TM, Ramos AJ, Sanchis V, Marín S. 2011c. Presence of trichothecenes and co-occurrence in cereal-based food from Catalonia (Spain). Food Control 22:490-495.

Castellá G, Bragulat MR, Cabañes FJ. 1999. Surveillance of fumonisins in maize-based feed and cereals from Spain. J Agric Food Chem 47:707-710.

Castelo MM, Summer SS, Bullerman LB. 1998. Occurrence of fumonisins in corn-based food products. J Food Protect. 61:704-707.

Dall'Asta C, Mangia M, Berthiller F, Molinelli A, Sulyok M, Schuhmacher R, Krska R, Galaverna G, Dossena A, Marchelli R. 2009. Difficulties in fumonisin determination: The issue of hidden fumonisins. Anal Bioanal Chem. 395:1335-1345.

D'Arco G, Fernández-Franzón M, Font G, Damiani P, Mañes J. 2008. Analysis of fumonisins B1, B2 and B3 in corn-based baby food by pressurized liquid extraction and liquid chromatography/tandem mass spectrometry. J Chromatogr A 31:188-194.

De Nijs M, Van Egmond HP, Nauta M, Rombouts FM, Notermans SHW. 1998. Assessment of human exposure to fumonisin B₁. J. Food Protect. 61:879-884.

EFSA (European Food Safety Authority). 2006. Guidance of the scientific committee on a request from EFSA related to uncertainties in dietary exposure assessment. EFSA J. 438:1-54.

European Commission (EC). 2006a. Regulation (EC) N° 401/2006 of 23/2/2006. OJEU. L70:12-34.

European Commission (EC). 2006b. Regulation (EC) N° 1881/2006 of 19/12/2006. OJEU. L364:5-24.

European Commission (EC). 2007. Regulation (EC) N° 1126/2007 of 28/9/2007. OJEU. L255:14-17.

FAO/WHO. 1997. Food consumption and exposure assessment of chemicals. Report of a FAO/WHO Consultation. Geneva, Switzerland, 10–14 February 1997. WHO/FSF/FOS/97.5. Geneva: WHO.

Gelderblom WCA, Jaskiewicz K, Marasas WFO, Thiel PG, Hora RM, Vleggar R, Kriek NP. 1988. Fumonisin—Novel mycotoxins with cancer-promoting activity produced by *Fusarium moniliforme*. Appl Environ Microbiol. 54:1806–1811.

Gelderblom WCA, Kriek NPJ, Marasas WFO, Thiel PG. 1991. Toxicity and carcinogenicity of the *Fusarium moniliforme* metabolite, fumonisin B₁ in rats. Carcinogenesis 12:1247–1251.

Hlywka JJ, Bullerman LI. 1999. Occurrence of fumonisin B₁ and B₂ in beer. Food Addit Contam. 16:319-324.

Humphreys SH, Carrington C, Bolger M. 2001. A quantitative risk assessment for fumonisins B₁ and B₂ in US corn. Food Addit Contam. 18:211-220.

IARC (International Agency for Research on Cancer). 2006. International Agency for Research on Cancer (IARC) Monographs (2006). 2 April 2006. Vols. 1–95.

Kim EK, Scott PM, Lau BP-Y. 2003. Hidden fumonisin in corn flakes. Food Addit Contam. 20:161-169.

Kim D-H, Yoo H-S, Lee Y-M, Kie J-H, Jang S, Oh S. 2006. Elevation of sphinganine 1-phosphate as a predictive biomarker for fumonisin exposure and toxicity in mice. J Toxicol Environ Health A 69:2071-2082.

Kroes R, Müller D, Lambe J, Löwik MRH, Van Klaveren J, Kleiner J, Massey R, Mayer S, Urieta I, Verger P, Visconti A. 2002. Assessment of intake from the diet. Food Chem

Toxicol. 40:27-85.

Kuiper-Goodman T, Scott PM, McEwen NP, Lombaert GA. 1996. Approaches to the risk assessment of fumonisins in corn-based foods in Canada. *Adv Exp Med Biol.* 392:369-393.

Leblanc J-C, Tard A, Volatier J-L, Verger P. 2005. Estimated dietary exposure to principal food mycotoxins from The First French Total Diet Study. *Food Addit Contam.* 22:652-672.

Lombaert GA, Pellaers P, Roscoe V, Mankotia M, Neil R, Scott PM. 2003. Mycotoxins in infant cereal foods from the Canadian retail market. *Food Addit Contam.* 20:494-504.

Marasas WFO. 1996. Fumonisins: History, world-wide occurrence and impact. *Adv Exp Med Biol.* 392:1-17.

Martins HM, Almeida I, Marques MF, Guerra MM. 2008. Fumonisins and deoxynivalenol in corn-based food products in Portugal. *Food Chem Toxicol.* 46:2585-2587.

Migracat (Observatori de la Imigració a Catalunya) [Internet] [Cited 2010 Aug 5]
Available from: <http://www.gencat.cat/dasc/publica/butlletiIMMI/xifres6/index.htm>.

Merrill AHJ, Wang E, Gilchrist DG, Riley RT. 1993. Advances in Lipid Research: Sphingolipids and Their Metabolites. Orlando: Academic Press, p. 215–234.

Merrill AH, Sweeley CC. 1996. Biochemistry of Lipids, Lipoproteins and Membranes. New York: Elsevier, p. 43–73.

Molinié A, Faucet V, Castegnaro M, Pfohl-Leszkowicz A. 2005. Analysis of some breakfast cereals on the French market for their contents of ochratoxin A, citrinin and fumonisin B₁: Development of a method for simultaneous extraction of ochratoxin A and citrinin. Food Chem. 92:391-400.

Nelson PE, Plattner RD, Shackelford DD, Desjardins AE. 1992. Fumonisin B₁ production by *Fusarium* species other than *F. moniliforme* in section *Liseola* and by some related species. Appl Environ Microbiol. 58:984-989.

Norred WP, Plattner RD, Vesonder RF, Bacon CW, Voss KA. 1992. Effects of selected secondary metabolites of *Fusarium moniliforme* on unscheduled synthesis of DNA by rat primary hepatocytes. Food Chem Toxicol. 30:233–237.

Patel S, Hazel CM, Winterton AGM, Mortby E. 1996. Survey of ethnic foods for mycotoxins. Food Addit Contam. 13:833-841.

Pietri A, Zanetti M, Bertuzzi T. 2009. Distribution of aflatoxins and fumonisins in dry-milled maize fractions. *Food Addit Contam.* 26:372–380.

Riley RT, An NH, Showker JL, Yoo HS, Norred WP, Chamberlain WJ, Wang E. 1993. Alteration of tissue and serum sphinganine to sphingosine ratio: an early biomarker of exposure to fumonisin-containing feeds in pigs. *Toxicol Appl Pharmacol.* 118:105–112.

Riley RT, Enongen E, Voss KA, Norred WP, Meredith FI, Sharma RP, Spitsbergen J, Williams DE, Carlson DB, Merrill AHJ. 2001. Sphingolipid perturbations as mechanisms for fumonisin carcinogenesis. *Environ Health Perspect.* 109:301–308.

Roscoe V, Lombaert G, Huzel V, Neumann G, Melietio J, Kitchen D, Kotello S, Krakalovich T, Trelka R, Scott PM. 2008. Mycotoxins in breakfast cereals from the Canadian retail market: a 3-year survey. *Food Addit Contam.* 25:347–355.

Sanchis V, Abadias M, Oncins L, Sala N, Viñas I, Canela R. 1994. Occurrence of fumonisins B₁ and B₂ in corn-based products from the Spanish market. *Appl Environ Microbiol.* 60:2147–2148.

Sanchis V, Abadias M, Oncins L, Sala N, Viñas I, Canela R. 1995. Fumonisins B₁ and B₂ and toxigenic *Fusarium* strains in feeds from the Spanish market. *Int J Food Microbiol.* 27:37–44.

SCF (Scientific Committee on Food). 2003. Updated opinion of the Scientific Committee on Food on Fumonisin B₁, B₂ and B₃. Brussels, Belgium, April.

SCOOP (Scientific Cooperation Task 3.2.10 of the European Commission). 2003. Collection of occurrence data of *Fusarium* toxins in food and assessment of dietary intake by the population of EU Member States. Final report. Directorate – General Health and Consumer Protection. Brussels, Belgium: European Commission.

Shephard GS, Thiel PG, Storckenstrom S, Sydenham EW. 1996. Worldwide survey of fumonisin contamination of corn and corn-based products. *J AOAC Int.* 79:671–687.

Shephard GS, Marasas WFO, Burger H-M, Somdyala NIM, Rheeder JP, Van der Westhuizen L, Gatyeni P, Van Schalkwyk DJ. 2007. Exposure assessment for fumonisins in the former Transkei region of South Africa. *Food Addit Contam.* 24:621-629.

Sydenham EW, Thiel PG, Marasas WFO, Shephard GS, Schalkwyk DJ, Koch KR. 1990. Natural occurrence of some *Fusarium* mycotoxins in corn from low and high esophageal cancer prevalence areas of Transkei, Southern Africa. *J Agric Food Chem.* 38:1900–1903.

Sydenham EW, Shephard GS, Thiel PG, Marasas WFO, Stockenström S. 1991. Fumonisin contamination of commercial corn-based human foodstuffs. *J Agric Food Chem.* 39:2014–2018.

Torres MR, Sanchis V, Ramos AJ. 1998. Occurrence of fumonisins in Spanish beers analysed by an enzyme-linked immunosorbent assay method. *Int J Food Microbiol.* 39:139-143.

Tran ST, Tardieu D, Auvergne A, Bailly JD, Babile R, Durand S, Benard G, Guerre P. 2006. Serum sphinganine and the sphinganine to sphingosine ratio as a biomarker of dietary fumonisins during chronic exposure in ducks. *Chem Biol Interact.* 160:41–50.

Van der Westhuizen L, Shephard GS, Van Schalkwyk DJ. 2001. The effect of a single gavage dose of fumonisin B₁ on the sphinganine and sphingosine levels in vervet monkeys. *Toxicon* 39:273–281.

Velluti A, Marín S, Sanchis V, Ramos AJ. 2001. Note. Occurrence of fumonisin B₁ in Spanish corn-based foods for animal and human consumption. *Food Sci Technol Int.* 7:433-437.

Verger P, Fabiansson S. 2008. Recent progress in exposure assessment and its interaction with the risk analysis process. *Trends Food Sci Technol.* 19:96-102.

Visconti A, Doko MB, Solfrizzo M, Pascale M, Boenke A. 1996. European Intercomparison Study for the Determination of Fumonisins in Maize. *Mikrochim Acta* 123:55-61.

Walter MD. 1998. Food Frequency Methods. In: J.L. Kelsey, M.G. Marmot, P.D. Stolley, & M.P. Vessey (Eds.), Nutritional Epidemiology, p. 74-94. Oxford: Oxford University Press.

Wang E, Norred WP, Bacon CW, Riley RT, Merrill AHJ. 1991. Inhibition of sphingolipid biosynthesis by fumonisins: implications for diseases associated with *Fusarium moniliforme*. J Biol Chem, 266:14486–14490.

Wang E, Ross PF, Wilson TM, Riley RT, Merrill JAH. 1992. Increases in serum sphingosine and sphinganine and decreases in complex sphingolipids in ponies given feed containing fumonisins, mycotoxins produced by *Fusarium moniliforme*. J Nutr. 122:1706–1716.

Wang E, Riley RT, Meredith FI, Merrill JAH. 1999. Fumonisin B₁ consumption by rats causes reversible, dose-dependent increases in urinary sphinganine and sphingosine. J Nutr. 129:214–220.

WHO (World Health Organization). 2001. Fumonisin. Safety Evaluation of Certain Mycotoxins in Food. WHO Food Additives Series 47, FAO Food and Nutrition Paper 74. Geneva, WHO.

WHO (World Health Organization). 2005. Dietary exposure assessment of chemicals in food. Report of a joint FAO/WHO consultation, Annapolis, Maryland, USA.

Yoo H-S, Norred WP, Showker JL, Riley RT. 1996. Elevated sphingoid bases and complex sphingolipid depletion as contributing factors in fumonisin-induced cytotoxicity. *Toxicol Appl Pharmacol.* 138:211–218.

Yoshizawa T, Yamashita A, Luo Y. 1994. Fumonisin occurrence in corn from high- and low-risk areas for human esophageal cancer in China. *Appl Environ Microbiol.* 60:1626–1629.

Table 1. Occurrence of total fumonisins (FB₁+FB₂) in corn-based food marketed in Catalonia

Food category	n	n	Positives	Mean± SD	Max
	Individual samples	“Composite”		µg/kg	µg/kg
Corn snacks	216	72	44	119.1 ± 83.1	475.5
Sweet corn	185	72	0	—	—
Corn Flakes	168	72	28	78.9 ± 27.9	139.5
Beer	216	72	64	36.9 ± 20.1	157.2
Baby foods	90	30	7	36.4 ± 8.5	50.7
Free-gluten food	18	18	1	22.5	22.5
Ethnic food	35	35	18	196.7 ± 191.8	682.2
Total	928	370	162		

Mean ± SD = Mean ± Standard Deviation

Table 2. Normalised consumption of foodstuffs by Catalanian population (expressed in g / kg body weight/day) and percentage of consumers in each category (between parentheses, %)

	Corn flakes	Corn snacks	Beer	Sweet corn	Ethnic food	Gluten-free bread	Gluten-free pasta
<i>Elders</i>	0.01 (6.6)	0.00 (25)	0.53 (29)	0.00 (0)			
<i>Adult females</i>	0.19 (40)	0.02 (33)	0.76 (53)	0.19 (37)	-	-	-
<i>Adult males</i>	0.13 (33)	0.02 (32)	1.69 (73)	0.14 (39)	-	-	-
<i>Adolescents</i>	0.50 (65)	0.13 (82)	0.29 (25)	0.15 (45)	-	-	-
<i>Children</i>	1.41 (72)	0.18 (84)	0.00 (0)	0.33 (38)	-	-	-
<i>Immigrants</i>	0.23 (38)	0.11 (45)	2.27 (61)	0.14 (54)	0.39 (80)	-	-
<i>Celiac sufferers</i>	0.34 (64)	0.07 (72)	0.32 (40)	0.14 (52)	-	1.19 (94)	1.31 (96)

Table 3. Exposure of each population group to fumonisins (units in ng kg⁻¹ bw day⁻¹) using the direct approach. Where $\hat{E}_{\pi_0}$ is the exposure mean of the population π with the population sample π_0 and $\hat{V}(E_{\pi_0})$ the variance estimate.

	$\hat{E}_{\pi_0}$	$\hat{V}(E_{\pi_0})$
Celiac sufferers	40.9	(35.0) ²
Immigrants	65.1	(69.0) ²
Adults females	34.5	(55.5) ²
Adults males	63.2	(90.3) ²
Adolescents	36.7	(44.3) ²
Children	63.0	(64.0) ²
Infants	156.0	(152.4) ²
(ng kg ⁻¹ bw day ⁻¹)		

Table 4. Results of exposure assessment ($\text{ng kg}^{-1} \text{bw day}^{-1}$) of the population groups from *P-P* simulation method. These results were obtained directly from the 10000 simulation outputs without *pdf* fittings. Bootstrap confidence intervals are presented between brackets (in $\text{ng kg}^{-1} \text{bw day}^{-1}$).

	<i>Adult females</i>	<i>Adult males</i>	<i>Immigrants</i>	<i>Celiac sufferers</i>
Mean	103.02 [91.84 - 115.27]	100.41 [88.62 – 113.33]	130.99 [71.64 – 221.57]	33.16 [20.13 - 52.18]
SD	116.47 [91.82 - 147.31]	114.21 [89.78 – 143.31]	291.26 [106.22 – 548.57]	61.62 [24.78 - 125.43]
Skewness	3.55	3.28	7.97	6.16
Kurtosis	22.52	17.59	135.88	68.44
Median	66.97 [58.92 - 76.33]	64.17 [55.99 – 73.87]	37.07 [18.35 – 66.20]	12.93 [8.19 - 19.66]
0.90 th quantile	226.73 [194.24 - 268.01]	225.48 [187.84 -264.92]	337.03 [171.85 – 631.48]	82.20 [43.72 - 153.27]
0.95 th quantile	320.76 [263.42 - 394.07]	313.33 [258.01 – 394.49]	560.64 [278.11 – 1264.82]	132.39 [68.71 - 276.57]
0.99 th quantile	563.08 [434.16 - 792.45]	577.19 [413.73 – 770.59]	1313.14 [479.93 – 3727.78]	294.88 [112.74 - 836.29]
	<i>Adolescents</i>	<i>Children</i>	<i>Infants</i>	
Mean	79.25 [69.01 – 90.36]	85.30 [59.30 - 119.96]	195.19 [168.91 – 223.68]	
SD	83.17 [64.81 – 103.93]	125.13 [62.71 - 250.98]	161.90 [130.01 – 201.47]	
Skewness	2.78	4.99	1.73	
Kurtosis	12.10	44.69	5.06	
Median	53.09 [44.45 – 61.61]	44.57 [30.66 – 63.69]	152.33 [125.27 - 185.52]	
0.90 th quantile	175.84 [143.52 – 211.80]	201.01 [128.33 – 314.25]	402.98 [335.03 – 484.66]	
0.95 th quantile	241.05 [191.29 – 295.83]	293.66 [172.74 - 489.71]	508.20 [410.86 - 650.41]	
0.99 th quantile	414.62 [282.63 – 559.11]	601.14 [280.30 – 1943.99]	748.56 [543.63 – 966.64]

Figure 1. Contamination histograms (relative frequencies), in $\mu\text{g kg}^{-1}$ for each food
952x952mm (96 x 96 DPI)

Figure 2. Fitted gamma (solid line) and lognormal (broken line) pdfs for the population groups with exception of celiac sufferers (relative frequencies). Exposure in ng kg⁻¹ bw day⁻¹.
533x682mm (96 x 96 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure Captions

Figure 1. Contamination histograms (relative frequencies), in $\mu\text{g kg}^{-1}$ for each food

Figure 2. Fitted gamma (solid line) and lognormal (broken line) *pdfs* for the population groups with exception of celiac sufferers (relative frequencies). Exposure in $\text{ng kg}^{-1} \text{ bw day}^{-1}$.