Immunological detection of osteocalcin in meat and bone meal: a novel heat stable marker for the investigation of illegal feed adulteration

Journal:	<i>Food Additives and Contaminants</i>
Manuscript ID:	TFAC-2011-420.R1
Manuscript Type:	Original Research Paper
Date Submitted by the Author:	22-Nov-2011
Complete List of Authors:	Kreuz, Grit; Federal Institute for Risk Assessment, Food Safety Zagon, Jutta; Federal Institute for Risk Assessment (BfR), Food Safety Broll, Hermann; Federal Institute for Risk Assessment, Food Safety Bernhardt, Christian; Federal Institute for Risk Assessment, Food Safety Linke, Bettina; Federal Institute for Risk Assessment, Food Safety Lampen, Alfonso; Federal Institute for Risk Assessment, Food Safety
Methods/Techniques:	Immunoassays, Traceability, Authenticity
Additives/Contaminants:	Animal products, Feeding
Food Types:	Animal products – meat, Animal feedingstuffs
Abstract:	A sandwich ELISA was developed for the detection of bovine meat and bone meal (BMBM) in feed, based on polyclonal rabbit antibodies raised against the synthetic N-terminal amino acid sequence 1–9 (YLDHWLGAP) of bovine osteocalcin. To set up a sandwich ELISA pair, a commercial mouse monoclonal capture antibody binding to a highly conserved epitope in the mid-fragment of the peptide was employed. It is shown that the bone marker osteocalcin is immunologically well detectable in BMBM extracts obtained by a simple EDTA based procedure even in a sample heated up to 145 °C. Furthermore a genus-specific restriction of the major specificity to cattle and horse was possible. The observed bi-specificity is consistent with theoretical predictions. The assay sensitivity with bovine osteocalcin of 1 ng was sufficient to enable the detection of 0.1% BMBM in compound plant feed or fish meal, for which no cross reaction was observed. In general the quantification of osteocalcin in extracts is possible using a standard curve procedure with pure bovine osteocalcin.

SCHOLARONE™
Manuscripts

For Peer Review Only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Immunological detection of osteocalcin in meat and bone meal: a novel heat stable marker for the investigation of illegal feed adulteration

G. KREUZ, J. ZAGON*, H. BROLL, C. BERNHARDT, B. LINKE, A. LAMPEN

Federal Institute for Risk Assessment, D-14195 Berlin, Germany

* Corresponding author. E-mail: jutta.zagon@bfr.bund.de

Abstract

A sandwich ELISA was developed for the detection of bovine meat and bone meal (BMBM) in feed, based on polyclonal rabbit antibodies raised against the synthetic N-terminal amino acid sequence 1–9 (YLDHWLGAP) of bovine osteocalcin. To set up a sandwich ELISA pair, a commercial mouse monoclonal capture antibody binding to a highly conserved epitope in the mid-fragment of the peptide was employed. It is shown that the bone marker osteocalcin is immunologically well detectable in BMBM extracts obtained by a simple EDTA based procedure even in a sample heated up to 145 °C. Furthermore a genus-specific restriction of the major specificity to cattle and horse was possible. The observed bi-specificity is consistent with theoretical predictions. The assay sensitivity with bovine osteocalcin of 1 ng was sufficient to enable the detection of 0.1% BMBM in compound plant feed or fish meal, for which no cross reaction was observed. In general the quantification of osteocalcin in extracts is possible using a standard curve procedure with pure bovine osteocalcin.

KEYWORDS: Osteocalcin, Meat-and-Bone-Meal (MBM), Bovine spongiform encephalopathy (BSE), Animal feed, Processed animal proteins (PAPs), ELISA, Species identification

For Peer Review Only

30 Introduction

31 Feeding meat and bone meal (MBM) to farmed animals is prohibited in Europe by Regulation
32 (EC) No 999/2001. In addition, an overarching species-to-species feed ban has been laid
33 down by Regulation (EC) No 1069/2009. The only exemption from the strict MBM ban is fish
34 meal, which has been allowed for pig and fowl feeding and recently also for calves in milk
35 replacing feed mixtures (Regulation (EC) No 999/2001). According to the recitals of the
36 above mentioned Regulation it might be assumed that the feed ban for ruminants will not be
37 lifted in the near future whereas the prohibition of processed animal proteins (PAP) feeding
38 to other animals than ruminants might be relaxed as far as analytical tools for species
39 identification are available.

40
41 The official method for the detection of MBM in feeding stuff is based on classical light
42 microscopy (Regulation (EC) No 152/2009). A second test may also be carried out using
43 variant or alternative methods, in order to improve the detection of certain types of animal
44 constituents or to further specify the origin of the animal constituents. The microscopic-based
45 method reliably detects heat- and pressure-stable particles of animal origin like bone
46 fragments, teeth, hair, gills or scales which are resisting the legally required MBM treatment
47 of 133 °C for 20 min at 300 kPa (Liu et al. 2011). However, by applying the microscope
48 approach it is not possible to distinguish between species or genus in samples containing
49 PAPs. Thus, up to now the implemented species-to-species ban cannot be controlled.
50 Furthermore, the microscopic investigation depends solely on individual skills and allows an
51 estimation but not quantification of the MBM content in a feed sample. Therefore - and
52 against the background of recently discussed thresholds and a further lifting of the feed ban
53 in the European Union (COMMISSION STAFF WORKING DOCUMENT 2010) - there is an
54 urgent need for alternative methods which might overcome present drawbacks to enable
55 legally consistent feed survey and law enforcement.

56

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

57 The immunological detection of PAPs is for long used as an alternative to microscopy.
58 Immunoassays can be performed in any laboratory independently from sophisticated
59 equipment or particularly trained staff. Antibody based assays are convertible into rapid
60 formats like lateral flow assays (dip stick tests) representing most convenient field methods
61 for rapid screening purposes. Ruminant or pork specific antibodies binding to PAPs, so far
62 have been raised either directly against bovine meat and bone meal (MBM) (Muldoon et al.
63 2004; Kim et al. 2004), mixtures of muscle proteins (Kim et al. 2005; Chen et al. 1998;
64 Ansfield 1994), purified troponin I (Chen et al. 2002) or *h*-caldesmon (Kim et al. 2004),
65 isolated from skeletal or smooth muscle of the bovine intestine, respectively. Four
66 commercial PAP detecting test kits (two dip stick tests and two ELISA) based on antibodies
67 binding to the described targets have been repeatedly evaluated in studies conducted by
68 official institutions in the USA or Europe. All of them revealed different shortcomings in terms
69 of specificity, selectivity or sensitivity (Fumière et al. 2009). Moreover, from the viewpoint of
70 validation, it is not satisfying that neither the kit manuals nor accompanying scientific
71 publications disclose the precise nature of the targeted epitopes, neither amino acid
72 sequences nor conformational molecular reaction sites, involved in the antigen-antibody
73 binding. As a consequence, potential cross reactivity cannot be deduced *in silico* and the
74 likelihood to reproduce immunization results obtained with changing or undefined antigen
75 mixtures is low.
76
77 The polymerase chain reaction (PCR) is another alternative approach to identify animal
78 species-specific components in feedstuffs. PCR is highly-specific and is amplifying target
79 DNA sequences in an exponential way. However PCR is not tissue-specific due to the fact
80 that the DNA of an individual organism does not differ between different tissues. Therefore it
81 is not possible to distinguish if the detected DNA is derived from the use of MBM or from
82 allowed animal products such as milk or egg protein.
83

1
2
3 84 In the present study we employed defined target sequences of bovine osteocalcin, consisting
4
5 85 of the amino acid sequence 1-9 and the epitope 17 of the peptide, to establish an ELISA for
6
7 86 the purpose of MBM detection in feed. Osteocalcin is a small extracellular bone matrix
8
9 87 protein produced by osteoblasts and a common constituent in calcified bone tissue all over
10
11 88 the vertebrate group including bone fish (Hauschka et al. 1989; Nishimoto et al. 2003; Laizé
12
13 89 et al. 2005). The mature peptide of approx. 5800-5900 Da size is composed of 45 – 49
14
15 90 amino acids. Up to 90 % of osteocalcin is secreted inside the bone tissue where it is, apart
16
17 91 from collagen, the most abundant bone protein accounting up to 2 mg/g bone matrix or 1 – 5
18
19 92 % (w/w) of the total bone protein, respectively 10 – 20 % (w/w) of the non-collagenous bone
20
21 93 proteins (Hauschka et al. 1989). A minor share of osteocalcin (10 - 40 %) is circulating in the
22
23 94 blood where it is - once detached from the bone mineral - quickly degraded in kidney and
24
25 95 liver (Lee et al. 2000). The function of osteocalcin is not completely understood but the
26
27 96 molecule seems to play a role in the molecular organisation of the hydroxyapatite skeleton as
28
29 97 well as in bone turnover (Dowd et al. 2003; Ducy et al. 1996). Three glutamic acid residues
30
31 98 are located at amino acid position 17, 21 and 24, constituting a highly conserved motif of the
32
33 99 peptide. These glutamic acids are transformed into γ -carboxyglutamic acids in a post-
34
35 100 translational carboxylation step and function as precipitation sites for Ca^{2+} ions. In the
36
37 101 presence of calcium the molecule forms a compact structure comprising of two α -helical
38
39 102 regions and a hydrophobic core near the C-terminus (Dowd et al. 2003). Both, N- and C-
40
41 103 terminal sequences of osteocalcin reveal species specific differences (Figure 1).
42
43 104 Osteocalcin is known to be heat stable. Human or bovine bone osteocalcin which was
44
45 105 extracted at 120°C overnight, lyophilized and decarboxylated at 110 °C for five hours, still
46
47 106 retained its substrate specificity towards enzymatic vitamin K-dependent carboxylase
48
49 107 (Vermeer et al. 1984). Osteocalcin could be identified in 75.000-year-old Neanderthal bone
50
51 108 by MALDI-TOF/TOF-MS (Nielsen-Marsh et al. 2005) and in fossil bones and teeth by
52
53 109 immunoassay-based investigation (Ulrich et al. 1987; Collins et al. 2000; Muyzer et al. 1992).
54
55 110 We recently demonstrated that osteocalcin can be identified and differentiated in MBM of
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

111 bovine and pig origin using high-resolution (MALDI/TOF-MS and HR-ESI Q/TOF) mass
112 spectrometry (Balizs et al. 2011).
113
114 The aim of the present study was to investigate i) if osteocalcin is still immunologically
115 detectable in MBM treated according to the legal requirements, ii) how far a discrimination of
116 farmed animals, especially cattle in feed, is possible and iii) to evaluate if a selectivity in an
117 ELISA down to 0.1 % MBM either in compound feed or fish meal would be achievable.

118
119 **Materials and Methods**

120 *Materials*

121 Pure bovine osteocalcin (OC) was purchased from Merck Calbiochem (Merck, Darmstadt,
122 Germany). Bone from farmed animals and pasteurized egg- and milk powder were
123 purchased from local retailer in Berlin. 100 % pure bovine meat and bone meal (BMBM),
124 treated 145 °C, with a bone content of approx. 80 %, as well as 100 % sheep meat and bone
125 meal (OMBM) were kindly submitted by the European Reference Laboratory for Animal
126 Proteins EU-RL AP, Gembloux, Belgium. From the same source a spiked 0.1 % BMBM
127 sample in plant feed (Plant Feed-0.1 % BMBM), serving as reference and training material
128 for microscopists, was obtained. This material was purity checked by classical microscopy
129 and PCR. Based on the results the EU-RL AP specified the material as a mix of four pure
130 bovine meat and bone meals from different heat treatments with a bone content of ~ 54 %.
131 On request, the EU-RL AP also kindly made available a sample of the non-contaminated
132 plant feed background (Plant Feed-0 % BMBM), used for spiking consisting of a classical
133 feed mix containing wheat, tapioca, soybean, rapeseed, palm kernel and barley meals, beet
134 pulp, molasses, bakery by-products, vegetable fat, limestone, salts and vitamins. A
135 commercial dog feed BMBM (100 % cattle according to the label) was purchased from
136 PerNaturam V.O.F. (Aalten, The Netherlands). Fish meal (FM) and poultry meal (PM)
137 produced under unknown processing parameters were obtained from diverse rendering
138 plants in Germany (SARIA Bio-Industries GmbH & Co., Selm, Germany). An anonymous

sample of BMBM treated in accordance with legal requirements (133 °C) was obtained from the Department of Feed and Feed Additives of the Federal Institute for Risk Assessment, Berlin. Pure pig meal from an anonymous source was kindly supplied by LUFA Nord-West (Oldenburg, Germany). Samples containing 0.1, 1 and 10 % (v/v) BMBM (dog feed) extract were prepared by mixing respective volumes of MBM extracts containing a comparable total protein concentration.

Sample preparation and protein extraction

For the production of reference proteins, native bones from different species were cleaned from any fat, meat residues and bone marrow and roughly trenched. Bone fragments were degreased by Soxhlet extraction in a Soxtec™ 1045 apparatus (FOSS GmbH, Rellingen, Germany) with boiling petroleum ether (Merck, Darmstadt, Germany). The defatted material was transferred into steel cartridges, cooled down for 2-3 minutes with liquid nitrogen and ground to yield a fine powder using a steel ball mill (TissueLyser Qiagen, Cat.No.85220, Retsch GmbH, Haan, Germany) for 3 min and 30 Hz.

MBM, plant feed or bone powder was extracted by EDTA-extraction. All centrifugation steps were performed at 15000 g, 10-20 min, 4 °C in a Heraeus Multifuge® 3S+/3SR+ (Thermo Scientific, Schwerte, Germany). In brief, 2 g sample were extracted with 8 ml 0.5 M Ammonium EDTA (pH 6.1) by stirring at 4 °C over night. The slurry was transferred into a conical Falcon™ tube and centrifuged. The supernatant was transferred into ultrafiltration tubes with an exclusion size of 3000 Da (Amicon® Ultra Centrifugal Filters Ultracel-3K, Millipore, Ireland) and the retained protein fraction was washed with 2 ml phosphate buffered saline (PBS, 10 mM NaCl, 10 mM KCl, 10 mM Na₂HPO₄, -NaH₂PO₄, pH 7.4) to exchange the buffer.

The protein concentration was measured for all extracts in triplicates using a Bicinchononic Acid Protein Assay Kit (Sigma-Aldrich, Saint Louis, Missouri, USA) according to the manufacturer's instruction. All extracts (Table 1) were aliquoted and stored at - 80 °C until use.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

167

168 *Antibodies*

169 A commercial mouse monoclonal antibody binding to the epitope aa17 of bovine osteocalcin

170 (OC4-30, Takara Bio Inc., Otsu/Shiga, Japan), specified to cross react with human, bovine,

171 rabbit, sheep, dog, monkey and goat but not rat or mouse, was used as animal specific

172 capture antibody in a two-side sandwich ELISA. Polyclonal antibodies (PAB) were raised in

173 two rabbits against the synthetic and LPH- (*Limulus polyphenus hemocyanin*) coupled

174 osteocalcin amino acid sequence OC 1-9 (YLDHWLGAP) by a service laboratory (Biogenes,

175 Berlin). The best performing serum in a pre-test against BSA-coupled amino acid sequence

176 (PAB OC1-9) was purified by affinity chromatography on CNBr-activated sepharose by the

177 service laboratory to isolate a monospecific IgG-fraction which was used in the sandwich

178 ELISA as secondary antibody. For this purpose PAB OC1-9 was conjugated with streptavidin

179 (PAB OC1-9-Strep) using a commercial kit according to the manufacturer's instruction

180 (Lightning-Link™ Streptavidin Conjugation Kit, Innova Biosciences, Cambridge, UK,

181 Technical Bulletin 1086, Release 02, June 2010).

182

183 *Sandwich ELISA*

184 Capture, secondary and detecting agent concentrations as well as the incubation time and

185 temperature were optimized in grid experiments. The final protocol was performed as follows.

186 Washing steps were repeated in general four times with 380 µl PBS-Tween 20 (154 mM

187 NaCl, 7.7 mM Na₂HPO₄, 2 mM NaH₂PO₄, 0.05% Tween 20, pH 7.4) using an automatic

188 ELISA Washer (Wellwash, Thermo Fisher Scientific). All incubation steps were carried out in

189 a volume of 100 µl/well. 1 µg/ml of the capture antibody (OC4-30) was bound to a microtiter

190 plate (Maxisorp, Nunc™, Denmark) for 48h at 4°C. The plate was washed and then blocked

191 by adding 200 µl/well Blocking Buffer made of 1.5% (w/v) bovine albumine fraction V (BSA;

192 Merck, Darmstadt, Germany) in 10 mM PBS, pH 7.4, overnight at 4°C. The plate was

193 washed and incubated with standard dilutions in PBS (1, 10, 100 and 1000 ng/ml

194 osteocalcin) or undiluted sample extract (Table 1) for 2 h at room temperature (RT). The

plate was washed and incubated with detection antibody (PAB OC1-9-Strep; 1:500 in PBS) for 2 h/RT, washed and incubated 1 h/RT with Biotinylated HRP (Invitrogen Corporation, Camarillo, CA; 1:10.000 in PBS). The plate was washed and the assay developed with chromogenic solution prepared as follows: 15 ml of citrate buffer (210 mM citric acid monohydrate, 300 mM KOH, pH 3.95) and 13.6 mM freshly added H₂O₂ were mixed 20:1 (v/v) with 21.6 mM TMB (3,3',5,5'-Tetramethylbenzidine, BioChemica, AppliChem, Darmstadt, Germany) diluted in 2.5 ml acetone and 22.5 ml methanol (p.A.). After incubation for 20 min/RT in the dark the reaction was stopped by adding 1N H₂SO₄ to each well. The optical density was determined at 450 nm against air in a multimode plate reader (Mithras LB 940, Berthold Tech. GMBH & Co. KG, Bad Wildbad, Germany). All samples and standards were measured in triplicates, statistical relevance between two data sets was calculated by two-side unpaired Student's t-test.

SDS-PAGE and Western blot

Discontinuous sodium dodecylsulfate polyacrylamide gel electrophoresis (SDS-PAGE) was performed with Tris-Glycin buffer according to Laemmli (1970) using a Mini Protean® Tetra System electrophoresis cell (Bio-Rad, Laboratories Inc., Germany). For the separation of low molecular weight proteins the separating gel contained 16.5% and the stacking gel 7.5% acrylamide (Rotiphorese®Gel 30 [37.5:1]), Carl Roth GmbH & Co KG, Karlsruhe, Germany). Separated proteins in the gel were electro-transferred onto a nitrocellulose membrane (Amersham™ Hybond™-ECL, GE Healthcare) with a blotter unit (Bio-Rad, Laboratories Inc.) at 4 °C and 350mA/ 400V for 1 h using transfer buffer as described by the blotter unit's manual. The blot was blocked in PBS/3 % BSA for 45 min at RT followed by an overnight incubation at 4 °C with PAB OC1-9-Strep diluted 1:500 in PBS/1.5 % BSA. The membrane was washed three times with TBST and PAB OC1-9-Strep binding onto the blot was detected by biotinylated HRP (1:5.000 in PBS containing 1.5 % BSA) after 1 h/RT incubation. After washing the blot was developed by adding 750 µl of a ready-mixed developing solution (Super Signal West Femto Maximum Sensitivity Substrate, Thermo Scientific) and chemo

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

luminescence was measured with the Imaging System VersaDoc MP 4000 (Bio-Rad Laboratories, Munich, Germany).

Results

In order to develop a sandwich ELISA for the detection of processed bovine osteocalcin (OC), a rabbit polyclonal antibody was raised against the amino acid sequence 1-9 of pure bovine osteocalcin. The best performing pre-tested serum (PAB OC1-9) was mono-specifically purified. To confirm the reactivity of the isolated mono-specific IgG fraction, an immunoblot was carried out with an EDTA protein extract produced from native bovine bone. Solely one major low-molecular weight protein band (~5.8 kDa) was detectable corresponding to the size of intact mature reference protein (Figure 2). No significant unspecific binding to other bone proteins occurred in bovine bone extract so that the antibody PAB OC1-9 could be used to establish a sandwich ELISA pair with a monoclonal mouse capture antibody (OC4-30).

The intra- and inter- assay variability and the sensitivity of the final ELISA set up was validated with standard dilutions of bovine osteocalcin in triplicates on three different days by the same operator (Figure 3). According to this validation experiment, the two-side assay is characterized by an average background signal ranging between $A_{450\text{ nm}}$ 0.170 – 0.260 for the non template control (NTC; PBS instead of sample) and blank (PBS instead of capture MAB). A significant signal above NTC and blank was obtained - with a probability of $p < 0.05$ in two-side Student's t-test - starting from an OC amount of 1 ng per well. Above this value the standard deviation is rising along with increasing signal strength, but the calculated inter assay coefficient of variation (2.8 – 23.3 %) (Figure 3) was considered satisfactory for all concentration levels and the assay accepted for the scope of investigation.

EDTA extracts of native bone derived from most significant farmed animals as well as egg and milk protein and a compound plant feed used for BMBM spiking (Plant Feed-0 % BMBM)

251 were measured to evaluate the specificity of the test. All EDTA-extracts were applied
252 undiluted in ELISA, to ensure a maximum achievable response with highly processed
253 materials or matrices of low MBM content. Protein yields typically varied between 0.6 – 2.8
254 mg/ml for different bone and MBM extracts (Table 1). The EDTA extract with the highest
255 protein amount was gained from egg powder with approximately 4 mg/ml.
256 The results of the specificity testing are summarized in Figure 4. The Limit of detection (LOD)
257 was calculated as the mean absorption A_{450} value plus 3-fold standard deviation of 13
258 different negative samples (measured in triplicate) which constitute typical feed compounds
259 (plant feed, whole milk powder, hen`s egg) or potential contamination with other species than
260 cattle (fish meal, poultry meal, sheep MBM, sheep bone, pork MBM, pork bone, chicken
261 bone, goat bone, rabbit bone) as well as the NTC. The horse sample has been excluded
262 from the calculation due to the 100 % amino acid identity with bovine osteocalcin and could
263 therefore not been considered as a negative sample. In all following experiments the LOD
264 indicates the baseline to discriminate true positive from negative results. The strongest
265 ELISA signals ($A_{450} \text{ nm} > 3$) were detected for cattle and horse bone extracts. For these
266 samples absolute protein amounts of 55 to 75 $\mu\text{g}/\text{reaction}$ were sufficient to elicit a strong
267 ELISA response with signals rapidly out-of-range if using further increased input amounts.
268 Regarding bone extracts from other animals, a weaker but significant ELISA signal pointing
269 to an evident cross reaction was measured with 100 % native pork bone extract. In a western
270 blot investigation with pure pig bone extract and PAB OC1-9 it was noticed, that a cross
271 reaction is directed against an unidentified protein of approximately 31 kDa (data not shown).
272 In addition, a weak cross reaction was also monitored for 100 % native sheep bone extract
273 (Figure 4). Bone extracts from chicken, goat or rabbit so as EDTA-extractable proteins from
274 hen`s egg, whole milk powder or compound plant feed all yielded negative results below
275 LOD-Baseline. EDTA-extracts derived from goose- or salmon bone, feather meal, as well as
276 rye, maize and soybean were tested negative (data not shown).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

In the next step, the capacity of the immunoassay to detect OC in heat treated meat and bone meal extracts was investigated, to test the applicability of the ELISA on real life samples. Since no horse MBM was available, horse serum was used instead as an exemplary matrix containing only trace amounts of osteocalcin. The total protein amount of all employed undiluted MBM-extracts per ELISA well ranged in a comparable order of magnitude between approx. 100 (poultry meal) – 280 µg (BMBM). The results are depicted in Figure 5. All BMBM samples showed a pronounced immunological response, even if the material was heated up to 145°C, whereas fish, poultry and sheep meal were tested negative. Therefore the cross reaction with native sheep bone is judged negligible since it is not interfering with the scope of the assay. A positive reaction was only detectable for undiluted horse serum and – in contrast to sheep meal - for pork MBM extract. Though the immune response against porcine MBM was rather weak compared with BMBM (c.f. dog feed), the obtained pork MBM signal exceeded the LOD-baseline (Figure 5).

The selectivity of the test was evaluated with extracts yielded from a standard plant feed spiked with 0.1 % (w/v) BMBM in comparison with the non-spiked matrix. In addition, since no homogeneity approved material was available, dilutions of 0.1 % BMBM (dog feed) in fish meal extracts were measured (Figure 6). The test discriminated 0.1 % (w/v; v/v) BMBM-spiked plant feed or fish meal, respectively, from the non-spiked matrix. The quantitative content of osteocalcin in the plant feed spiked with 0.1 % (w/v) BMBM corresponds to 6.7 ng calculated in relation to the standard curve. Due to the observed cross reaction of the ELISA with pig meal, a separate selectivity experiment was performed. 0.1 %, 1 % and 10 % (v/v) BMBM (dog feed) extract in pig meal extract was measured on three days (Table 2). A significant positive response was measurable for the 10 % (v/v) BMBM spiking level but not for the 1 % (v/v) spiked sample. The absorption measured with 100 % pork meal extract did not exceed the values corresponding to 10 ng bovine OC per well.

Discussion

306 A pre-requisite for the development of an immunological assay to detect fraudulent
307 application to or unintended traces of MBM in feed is the thorough choice of a protein which
308 exhibits exceptional resistance against mechanical forces, pressure and peak temperatures
309 up to 140 – 150°C evoked by the rendering process (Pérez-Calvo et al. 2010; Garcia et al.
310 2006). In this study the heat stable bone protein osteocalcin has proved to be a well suited
311 MBM marker since a strong immunological signal was still detectable in BMBM even if
312 processed at 145°C. In addition, osteocalcin represents a genuine MBM- as well as animal
313 marker protein since it is exclusively expressed in bone tissue or dentin in all vertebrates
314 including bone fish (Hauschka et al. 1989).

315

316 Though fairly stable if bound to the hydroxyapatite skeleton, osteocalcin under natural
317 conditions undergoes rapid degradation in a turnover process in the blood plasma. Two
318 arginine-arginine cleavage sites are situated at the amino acid positions 19(Arg)-20(Arg) and
319 43(Arg)-44(Arg) (Figure 1) resulting in two to three fragments (aa1-19, aa20-43, aa44-49) on
320 tryptic hydrolysis. The cleavage site near the C-terminal sequence is quickly hydrolysed
321 cutting off the latter six amino acid residues whereas the resting aa1-43 fragment (N-MID
322 fragment) is far more stable (Lee et al. 2000). Respecting the binding sites of the antibodies
323 in our assay (aa1-9 and the epitope aa17) the results obtained with three different bovine
324 MBM samples underline that a significant portion of at least the N-MID (aa1-43) is still
325 immunologically accessible in MBM.

326

327 To restrict the assay specificity to cattle, the synthetic N-terminal amino acid sequence aa1-9
328 of bovine osteocalcin was used for the immunization of rabbits to exclude other most relevant
329 farmed animals. The N-terminal sequences of osteocalcin in the genus *Bos*, *Sus*, *Ovis* and
330 *Capra* differ only in one to three amino acids whereas this sequence is completely identical
331 among *Bos* and *Equus* (Figure 1). The specificity testing confirmed that minor sequence
332 differences were sufficient to limit the major polyclonal antibody reactivity to *Bos* with the
333 predictable exception of *Equus*. However, an increasing tendency towards cross reaction is

observed in accordance with the number of amino acid exchanges in relation to *Bos* OC aa1-9 ranking from *Capra* (3 exchanges): no cross reaction – *Ovis* (2 exchanges): very weak cross reaction – *Sus* (1 exchange): weak cross reaction - *Equus* (100 % sequence identity): strong cross reaction. The observed binding of the polyclonal rabbit IgG fraction to a 31 kDa-protein in western blot studies with PAB OC1-9 and porcine bone extract (data not shown) points rather to an unspecific side reaction to another protein than osteocalcin. To unravel the nature of this cross reaction a mass spectroscopic investigation would be required which was not subject of this study. However, the exchange of the voluminous aromatic amino acid tryptophan at position aa5 in bovine OC against glycine in *Sus* could be targeted by the production of a high affinity monoclonal antibody.

In the case of horse and cattle a sequence difference between both is given only at position aa19 where lysine in bovine osteocalcin is exchanged by arginine in the genus *Equus* (Figure 1). Therefore elimination of cross reaction in theory could be possible by an exchange of the capture antibody. However, an attempt to boost a polyclonal rabbit capture antibody against the synthetic mid-fragment spanning aa16 – 22 of bovine osteocalcin was not successful. This might be due to the lacking carboxylation of the glutamic acid residues at position 17, 21 and 24 of the synthetic peptide used for immunization, preventing appropriate molecule folding. In addition the hydrophobic core of the molecule (Dowd et al. 2003) most probably requires a conformational epitope rather than a sequence based recognition site for antibody binding. The latter consideration is supported by a study using monoclonal and polyclonal OC- directed antibodies which were tested with differently sized fragments of the molecule. The authors demonstrated a generally poor affinity of antibodies against OC fragments comprising solely of aa11-29 or aa21-43 (Gundberg et al. 1998). Accordingly, the exclusion of horse- but also a reduction of pig cross reactivity could be tackled by a compelling anti-protein production mode, e.g. as given by stringent phage display or monoclonal selection against native bovine osteocalcin fragments aa1-43, aa1-19, aa 4-19 or intact osteocalcin. The latter is planned for further investigations.

362

363 Apart from osteocalcin, γ -carboxylated glutamic acids are occurring e.g. in proteins involved
364 in the blood coagulation cascade like prothrombin, but could not be detected so far in
365 hemolymph of annelids, crustaceous organisms, insects, soybean leafs, *E. coli* extract or
366 bovine milk (Zytkowicz and Nelsestuen 1976). Accordingly, the ELISA developed in this study
367 verified a remarkable selectivity for osteocalcin in plant background. A protein extract
368 prepared from a classical feed mix used for the preparation of 0.1 % BMBM spiked feed
369 samples, yielded a background signal comparable with blank and NTC. Similar results were
370 obtained with 100 % soybean-, wheat- or corn protein extracts and a further plant based milk
371 replacer containing 12 % citrus and 5 % cocoa shell (data not shown). The presence of beet
372 or citrus pulp obviously does not pose a problem in contrast to test interference observed for
373 these ingredients in a previous validation study on commercial test kits (Boix et al. 2004).
374 The selectivity of the assay was also excellent in fish meal background which is explainable
375 by the pronounced sequence differences between osteocalcin from bone fish compared with
376 terrestrial animals (Figure 1). In contrast, the observed weak but significant cross reactivity
377 with pure pig bone or pig meal decreased the selectivity of the assay. However, only one
378 sample of pure pork meal was available for testing so that further pork MBM materials as well
379 as defined mixtures have to be investigated to verify this result. Due to the nature of the
380 material a cross-contamination with bovine material could not be excluded. The identification
381 of MBM traces derived from other species – e.g. BMBM in pork or fish meal – seems to be a
382 most challenging issue associated with MBM targeted immunological methods which was
383 also observed for commercial cooked meat kits. In studies on the performance on two
384 broadly distributed dip stick tests increasingly false positive results were monitored in the
385 presence of 5 % pork meal or animal fat or with fish meal at levels as low as 1.5 % (Fumière
386 et al. 2009; Boix et al. 2004).

387

388 The percentage of bone fragments in two defined BMBM samples which we used in our
389 experiments was about 54 and 80 % (w/w), which is the common range reported for

1
2
3 390 materials from rendering plants (Liu et al. 2010). The good results yielded with EDTA as a
4
5 391 bone solvent is in line with recently published results on an improved extraction method for
6
7 392 the detection of ruminant material. Due to an EDTA-based extraction mode the performance
8
9 393 of a commercial dip stick test could be improved from 2 % down to a detection level of 0.1 %
10
11 394 BMBM in feed (Karanam 2011). In contrast to our approach the method includes a
12
13 395 concentration step of the bone fraction by sedimentation with 100 ml tetra chloroethylene
14
15 396 (TCE) as used in sample preparation for microscopy (Regulation (EC) No 152/2009). An
16
17 397 additional density based concentration of bone fragments is not necessary for the ELISA
18
19 398 method described here avoiding an agent supposed to be carcinogenic (IPCS INCHEM;
20
21 399 <http://www.inchem.org/>). However, an over night bone digestion step which does not allow
22
23 400 rapid sample analysis within a few hours, was also applied in our study. To speed up the
24
25 401 extraction method used here the procedure could possibly be optimized by hot and/or acid
26
27 402 extraction or augmentation of the sample amount or further homogenization of coarse
28
29 403 fractions.
30
31 404
32
33 405 The bone content might play an essential role regarding the sensitivity of the method. The
34
35 406 question if an osteocalcin based method would also perform with samples containing < 54 %
36
37 407 (w/w) bone or even with 100 % muscle meal remains to be clarified with appropriate
38
39 408 materials which were not available in this study. With respect to the expectable great
40
41 409 variance of MBM from different rendering plants it would be anyhow advisable not only to rely
42
43 410 on one single target molecule but to apply a panel of well defined marker-proteins to capture
44
45 411 the great diversity of animal by-products produced by the rendering industry.
46
47 412
48
49 413 Although the sensitivity of our assay 1 ng bovine OC is about one order of magnitude below
50
51 414 commercial osteocalcin ELISA validated for pharmaceutical applications, the sensitivity with
52
53 415 MBM looks encouraging compared with commercial ELISA test kits developed for the scope
54
55 416 of species detection in processed meat. For these, sensitivities of 0.1, 0.5 or ≥ 2 % MBM in
56
57 417 feed or even a complete failure to detect legally treated MBM have been reported from
58
59
60

diverse validation studies (Boix et al. 2004; Karanam 2011; Meyers et al. 2007; von Holst et al. 2006; Rao and Hsieh 2007; Love and Carey-Smith 2004; Gizzi and von Holst 2004). The sensitivity of the ELISA procedure used in our study primarily might have been decreased during the mono-specific purification step retaining most potent binding IgG from the polyclonal sera. The linkage to the voluminous streptavidine molecule had no negative impact on the sensitivity compared with horseradish peroxidase labelling (data not shown) but an enhancing effect due to multiple biotinylation sites.

Conclusions

In conclusion our results demonstrate that osteocalcin is a promising tissue specific target protein for the sensitive immunological detection of meat and bone meal in compound feed or fish meal with an excellent selectivity for BMBM in these matrices. Furthermore environmental bone fragment contaminants in sugar beet epidermis, e.g. derived from rabbit, rat or mouse can be easily discriminated because of marked sequence divergences to e.g. bovine osteocalcin. The here described antibody-based osteocalcin approach might be the basis for a validated detection tool for official law enforcement of the feed ban. Moreover the general approach using epitop-specific antibodies is target-directed and might be an improvement in terms of the acceptance as national or international standard in the future. Moreover quantification is possible based on a standard curve established by commercially available pure bovine osteocalcin.

References

- Ansfield M. 1994. Production of a Sensitive Immunoassay for Detection of Ruminant Proteins in Rendered Animal Material Heated to > 130 °C. Food & Agricultural Immunology. 6: 419-433
- Balitz G, Weise C, Rozycki C, Opialla T, Sawada S, Zagon J, Lampen A. 2011. Determination of osteocalcin in meat and bone meal of bovine and porcine origin using

- 447 matrix-assisted laser desorption ionization/time-of-flight mass spectrometry and high-
448 resolution hybrid mass spectrometry. *Analytica Chimica Acta*. 693: 89-99
449
- 450 Boix A, Serano F, Bellorini S, von Holst C. 2004. RUGGEDNESS STUDY OF
451 IMMUNOASSAYS FOR PROCESSED ANIMAL PROTEINS DETECTION IN FEED: Neogen
452 Reveal for Ruminant Feed Test System. European Commission, DG JRC-IRMM, Geel,
453 Belgium. GE/R/FSQ/03/2006/09/04
454
- 455 Chen F-C, Hsieh Y-HP, Bridgman RC. 1998. Monoclonal Antibodies to Porcine Thermal-
456 Stable Muscle Protein for Detection of Pork in Raw and Cooked Meats. *Journal of Food*
457 *Science*. 63(2): 201-205
458
- 459 Chen F-C, Hsieh Y-HP, Bridgman RC. 2002. Monoclonal antibodies against troponin I for the
460 detection of rendered muscle tissues in animal feedstuffs. *Meat Science*. 62: 405-412
461
- 462 Collins MJ, Gernaey AM, Nielsen-Marsh CM, Vermeer C, Westbroek P. 2000. Slow rates of
463 degradation of osteocalcin: Green light for fossil bone protein? *Geology*. 28(12): 1139–1142
464
- 465 COMMISSION STAFF WORKING DOCUMENT, accompanying the
466 Communication from the Commission to the European Parliament and the Council
467 on the TSE Roadmap 2. COM(2010)384. Brussels. SEC(2010) 899 final
468
- 469 Dowd TL, Rosen JF, Li L, Gundberg C M. 2003. The Three-Dimensional Structure of Bovine
470 Calcium Ion-Bound Osteocalcin Using ^1H NMR Spectroscopy. *Biochemistry*. 42: 7769-7779
471
- 472 Ducy P, Desbois C, Boyce B, Pinero G, Story B, Dunstan C, Smith E, Bonadio J, Goldstein
473 S, Gundberg C, Bradley A, Karsenty G. 1996. Increased bone formation in osteocalcin-
474 deficient mice. *Nature*. 382: 448-452
475
- 476 Fumière O, Veys P, Boix A, von Holst C, Baeten V, Berben G. 2009. Methods of detection,
477 species identification and quantification of processed animal proteins in feedingstuffs.
478 *Biotechnol. Agron. Soc. Environ.* 13(S): 59-70
479
- 480 Garcia RA, Rosentrater KA, Flores RA. 2006. Characteristics of North American Meat and
481 Bone Meal Relevant to the Development of Non-Feed Applications. *American Society of*
482 *Agricultural and Biological Engineers*. 22(5): 729-736
483

- 484 Gizzi G, von Holst C. 2004. Determination of Processed Animal Proteins, Including Meat and
485 Bone Meal, in Animal Feed. *Journal of AOAC International*. 87(6): 1334-1340
486
- 487 Gundberg CM, Nieman SD, Abrams S, Rosen H. 1998. Vitamin K Status and Bone Health:
488 An Analysis of Methods for Determination of Undercarboxylated Osteocalcin. *Journal of*
489 *Clinical Endocrinology and Metabolism*. 83(9): 3258-3266
490
- 491 Hauschka PV, Lian JB, Cole DEC, Gundberg CM. 1989. Osteocalcin and Matrix Gla Protein:
492 Vitamin K-Dependent Proteins in Bone. *Physiological Reviews*. 69(3): 990-1047
493
- 494 Karanam M. 2011. An improved protein extraction method for detecting ruminant material in
495 feed using lateral flow device dipsticks. *Biotechnologie, Agronomie, Société et*
496 *Environnement*. 15(S1): 25-29
497
- 498 Kim S-H, Huang T-S, Seymour TA, Wei C-I, Kempf SC, Bridgman CR, Clemens RA, An H.
499 2004. Production of Monoclonal Antibody for the Detection of Meat and Bone Meal in Animal
500 Feed. *Journal of Agricultural and Food Chemistry*. 52: 7580-7585
501
- 502 Kim S-H, Huang T-S, Seymour TA, Wei C-I, Kempf SC, Bridgman CR, Momcilovic D,
503 Clemens RA, An H. 2005. Development of Immunoassay for Detection of Meat and Bone
504 Meal in Animal Feed. *Journal of Food Protection*. 68(9): 1860-1865
505
- 506 Laemmli UK. 1970. Cleavage of Structural Proteins during the Assembly of the Head of
507 Bacteriophage T4. *Nature*. 227: 680-685
508
- 509 Laizé V, Martel P, Viegas CSB, Price PA, Cancela ML. 2005. Evolution of Matrix and Bone γ -
510 Carboxyglutamic Acid Proteins in Vertebrates. *Journal of Biological Chemistry*. 280(29):
511 26659-26668
512
- 513 Lee AJ, Hodges S, Eastell R. 2000. Measurement of osteocalcin. *Ann Clin Biochem*. 37: 432-
514 446
515
- 516 Liu X, Han L, Veys P, Baeten V, Jiang X, Dardenne P. 2011. An Overview of the Legislation
517 and Light Microscopy for Detection of Processed Animal Proteins in Feeds. *Microscopy*
518 *Research and Technique*. 74(8):735-743
519

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

520 Love JL, Carey-Smith GV. 2004. Immunoassay Kit Used to Detect the Presence of Bovine
521 Material in Processed Foods. *Journal of AOAC International*. 87(5): 1143-1147
522
523 Meyers MJ, Yancy HF, Farrell DE, Washington JD, Deaver CM, Frobish RA. 2007.
524 Assessment of Two Enzyme-Linked Immunosorbent Assay Tests Marketed for Detection of
525 Ruminant Proteins in Finished Feed. *Journal of Food Protection*. 70(3): 692-699
526
527 Muldoon MT, Onisk DV, Brown MC, Stave JW. 2004. Targets and methods for the detection
528 of processed animal proteins in animal feedstuffs. *International Journal of Food Science and*
529 *Technology*. 39: 851-861
530
531 Muyzer G, Sandberg P, Knapen MHJ, Vermeer C, Collins M, Westbroek P. 1992.
532 Preservation of the bone protein osteocalcin in dinosaurs. *Geology*. 20(10): 871-874
533
534 Nielsen-Marsh CM, Richards MP, Hauschka PV, Thomas-Oates JE, Trinkaus E, Pettitt PB,
535 Karavanic I, Poinar H, Collins MJ. 2005. Osteocalcin protein sequences of Neanderthals and
536 modern primates. *PNAS*. 102(12): 4409-4413
537
538 Nishimoto SK, Waite JH, Nishimoto M, Kriwacki RW. 2003. Structure, Activity, and
539 Distribution of Fish Osteocalcin. *Journal of Biological Chemistry*. 278(14): 11843-11848
540
541 Pérez-Calvo E, Castrillo C, Baucells MD, Guada JA. 2010. Effect of rendering on protein and
542 fat quality of animal by-products. *Journal of Animal Physiology and Animal Nutrition*. 94(5):
543 e154-e163
544
545 Rao Q, Hsieh Y-HP. 2007. Evaluation of a commercial lateral flow feed test for rapid
546 detection of beef and sheep content in raw and cooked meats. *Meat Science*. 76: 489-494
547
548 Regulation (EC) No 999/2001 OF THE EUROPEAN PARLIAMENT AND OF THE COUNCIL
549 of 22 May 2001 laying down rules for the prevention, control and eradication of certain
550 transmissible spongiform encephalopathies. *Official Journal of the European Communities*.
551 2001R0999— EN— 20.04.2009 — 033.001— 5.
552
553 Regulation (EC) No 1069/2009 OF THE EUROPEAN PARLIAMENT AND OF THE
554 COUNCIL of 21 October 2009 laying down health rules as regards animal by-products and
555 derived products not intended for human consumption and repealing Regulation (EC) No

- 1
2
3 556 1774/2002 (Animal by-products Regulation). Official Journal of the European Union. L 300:
4 557 1-33
5
6 558
7
8 559 Regulation (EC) No 152/2009 of 27 January 2009 laying down the methods of sampling and
9 560 analysis for the official control of feed. Official Journal of the European Union. L 54: 1-130
10 561
11 562 Ulrich MMW, Perizonius WRK, Spoor CF, Sandberg P, Vermeer C. 1987. Extraction Of
12 563 Osteocalcin From Fossil Bones And Teeth. Biochemical and Biophysical Research
13 564 Communications. 149(2): 712-719
14
15 565
16
17 566 Vermeer C, Soute BAM, Hendrix H, de Boer-van den Berg MAG. 1984. Decarboxylated bone
18 567 Gla-protein as a substrate for hepatic vitamin K-dependent carboxylase. Federation of
19 568 European Biochemical Societies. 165(1): 16-20
20
21 569
22
23 570 von Holst C, Boix A, Baeten V, Vancutsem J, Berben G. 2006. Determination of processed
24 571 animal proteins in feed: The performance characteristics of classical microscopy and
25 572 immunoassay methods. Food Additives and Contaminants. 23(3): 252-264
26
27 573
28
29 574 Zytковicz TH, Nelsestuen GL. 1976. γ -Carboxyglutamic Acid Distribution. Biochimica et
30 575 Biophysica Acta. 444: 344-348
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1. Comparison of osteocalcin sequences published for farmed animals, pets, rodents, amphibian and bone fish species. Conserved carboxylated glutamic acid residues at position 17, 21 and 24 are asterisked. Black arrows point to tryptic cleavage sites. Amino acid exchanges between *Bos* and other farmed animals are highlighted in bold types. The position of the sequence targeted by the detection antibody PAB OC1-9 and the N-MID-fragment (1-43) are marked by black lines. The capture antibody OC4-30 (Takara) used in the sandwich assay in this study binds to the epitope around position 17.

Figure 2. Specificity of PAB OC1-9 (polyclonal anti-osteocalcin IgG-fraction) towards native bovine bone protein in western blot analysis. M, Lane 2: Low-Range Molecular Weight Marker (3.5-40 kDa), Amersham, GE Healthcare; Lane 1: NTC (water instead protein); Lane 3: OC=Positive Control (bovine osteocalcin, 1.3 µg/slot); Lane 4: BB=bovine bone EDTA extract (7.2 µg/slot)

Figure 3. Osteocalcin specific sandwich ELISA with MAB OC4-30 (Takara) as capture and PAB OC1-9 as detection antibody. Sensitivity with purified bovine osteocalcin. Error bars indicate a one-fold standard deviation. Measurements were done in three replicates on three different days. Blank=PBS instead of capture; NTC= PBS instead of sample; *significant difference ($p<0.05$) against NTC and Blank.

Figure 4. Osteocalcin specific sandwich ELISA with MAB OC4-30 (Takara) as capture and PAB OC1-9 as detection antibody. Specificity with EDTA-extracts from native bone and feed components. Error bars indicate a one-fold standard deviation. Undiluted EDTA extracts were measured in triplicate. Horizontal line = LOD = Mean + 3x SD from 13 negative samples as indicated in the results section.

Figure 5. Osteocalcin specific sandwich ELISA with MAB OC4-30 (Takara) as capture and PAB OC1-9 as detection antibody. Method performance with meat and bone meal extracts. Error bars indicate a one-fold standard deviation. Undiluted EDTA extracts were measured in triplicate. Horizontal line = LOD = Mean + 3x SD from 13 negative samples as indicated in the results section.

Figure 6. Osteocalcin specific sandwich ELISA with MAB OC4-30 (Takara) as capture and PAB OC1-9 as detection antibody. Selectivity in mixtures containing 0.1 % bovine MBM. Error bars indicate a one-fold standard deviation. Undiluted EDTA extracts were measured in triplicate. Horizontal line = LOD = Mean + 3x SD from 13 negative samples as indicated in the results section.

Species:	N-MID-fragment		NCBI Accession:
	1	43	
	1	9	
		* ▼ *	
<i>Bos taurus</i>	YLDH--WLGAPAPYPDPLEPKREVCELNPDCELDADHIGFQEAYRRFYGPV--		ABU88822.1
<i>Ovis aries</i>	YLDP--GLGAPAPYPDPLEPKREVCELNPDCELDADHIGFQEAYRRFYGPV--		ABD83814.1
<i>Capra hircus</i>	YLDP--GLGAPAPYPDPLEPKREVCELNPDCELDADHIGFQEAYRRFYGLA--		1005180A
<i>Sus scrofa</i>	YLDH--GLGAPAPYPDPLEPKREVCELNPDCELDADHIGFQEAYRRFYGLA--		Q8HY9.2
<i>Equus caballus</i>	YLDH--WLGAPAPYPDPLEPKREVCELNPDCELDADHIGFQEAYRRFYGPV--		P83005.1
<i>Gallus gallus</i>	HYAQDSGVAGA--PPNPLEAQREVCELNPDCELDADQIGFQEAYRRFYGPV--		AAA78809.1
<i>Dromaius novaehollandiae</i>	-SFAVGSSYGA--APDPLEAQREVCELNPDCELDADHIGFQEAYRRFYGPV--		P15504.1
<i>Rattus norvegicus</i>	YLMN--GLGAPAPYPDPLEPHREVCELNPDCELDADHIGFQDAYKRIYGTTV-		AAA53280.1
<i>Mus musculus</i>	-----YLGASVSPDPLEPTREQCELNPADELSDQYGLKTAYKRIYGITI-		AAA39856.1
<i>Canis familiaris</i>	YLDS--GLGAPVPYPDPLEPKREVCELNPDCELDADHIGFQEAYRRFYGPV--		AAB27444.1
<i>Felis sylv. catus</i>	YLAP--GLGAPAPYPDPLEPKREVCELNPDCELDADHIGFQDAYRRFYGTV--		P02821.1
<i>Oryctolagus cuniculus</i>	QLIN--GQGAPAPYPDPLEPKREVCELNPDCELDADQVGLQDAYRRFYGPV--		P39056.1
<i>Xenopus laevis</i>	--SYGNVVGQGAAGVSPLESQREVCELNPDCELDADHIGFQEAYRRFYGPV--		AAB36024.1
<i>Salmo salar</i>	AGTALADLSL-----TQLESREVCELNWCENMMDTAGIIAAYTEFYGPPIY		NP_001130023.1
<i>Oncorhynchus mykiss</i>	ASTALADLYL-----TQLESREACELNWCENMMDTAGIIAAYTEFYGPPIY		AAPO4487.1
<i>Scophthalmus maximus</i>	---AAGELSL-----TQLESREVCELNACEEMMDTGIIAAYNAFYGPPIQ		AAV80230.1
<i>Diplodus puntazzo</i>	---AAGQLSL-----TQLESREVCELNACEHMDTEGIIAAYTAYYGPPIY		AAV17284.1
<i>Sparus aurata</i>	---AAGQLSL-----TQLESREVCELNACEHMDTEGIIAAYTAYYGPPIY		AAK66568.1
<i>Pagrus auriga</i>	---AAGQLSL-----TQLESREVCEANLACEHMDTEGIIAAYTAYYGPPIY		AAV80232.1
<i>Halobatrachus didactylus</i>	---AAELSL-----VQLESREVCETNVACDEMMDTGIIAAYTAYYGPPIF		AAQ13575.1
<i>Argyrosomus regulus</i>	---AAKELTL-----AQTESREVCETNMACDEMADAQGIVAAYQAFYGPPIF		AAO48725.1
<i>Macropus eugenii</i>	--YLYQTLGFAPYPDPQENKREVCELNPDCELDADHIGFQEAYRRFYGTA--		P0C226.1
<i>Lepomis macrochirus</i>	---AAGELTL-----TQLESREVCEANLACEHMDAQQIIAAYTAYYGPPIY		AAB22263.1

Comparison of osteocalcin sequences published for farmed animals, pets, rodents, amphibian and bone fish species. Conserved carboxylated glutamic acid residues at position 17, 21 and 24 are asterisked. Black arrows point to tryptic cleavage sites. Amino acid exchanges between *Bos* and other farmed animals are highlighted in bold types. The position of the sequence targeted by the detection antibody PAB OC1-9 and the N-MID-fragment (1-43) are marked by black lines. The capture antibody OC4-30 (Takara) used in the sandwich assay in this study binds to the epitope around position 17.

248x164mm (150 x 150 DPI)

Specificity of PAB OC1-9 (polyclonal anti-osteocalcin IgG-fraction) towards native bovine bone protein in western blot analysis. M, Lane 2: Low-Range Molecular Weight Marker (3.5-40 kDa), Amersham, GE Healthcare; Lane 1: NTC (water instead protein); Lane 3: OC=Positive Control (bovine osteocalcin, 1.3 µg/slot); Lane 4: BB=bovine bone EDTA extract (7.2 µg/slot)
100x106mm (150 x 150 DPI)

Osteocalcin specific sandwich ELISA with MAB OC4-30 (Takara) as capture and PAB OC1-9 as detection antibody. Sensitivity with purified bovine osteocalcin. Error bars indicate a one-fold standard deviation. Measurements were done in three replicates on three different days. Blank=PBS instead of capture; NTC= PBS instead of sample; *significant difference ($p < 0.05$) against NTC and Blank.

Osteocalcin specific sandwich ELISA with MAB OC4-30 (Takara) as capture and PAB OC1-9 as detection antibody. Specificity with EDTA-extracts from native bone and feed components. Error bars indicate a one-fold standard deviation. Undiluted EDTA extracts were measured in triplicate. Horizontal line = LOD = Mean + 3x SD from 13 negative samples as indicated in the results section.

165x92mm (150 x 150 DPI)

Osteocalcin specific sandwich ELISA with MAB OC4-30 (Takara) as capture and PAB OC1-9 as detection antibody. Method performance with meat and bone meal extracts. Error bars indicate a one-fold standard deviation. Undiluted EDTA extracts were measured in triplicate. Horizontal line = LOD = Mean + 3x SD from 13 negative samples as indicated in the results section.

166x90mm (150 x 150 DPI)

Osteocalcin specific sandwich ELISA with MAB OC4-30 (Takara) as capture and PAB OC1-9 as detection antibody. Selectivity in mixtures containing 0.1 % bovine MBM. Error bars indicate a one-fold standard deviation. Undiluted EDTA extracts were measured in triplicate. Horizontal line = LOD = Mean + 3x SD from 13 negative samples as indicated in the results section.

166x90mm (150 x 150 DPI)

Table 1. Protein concentration of EDTA extracts used in ELISA

Sample	Protein concentration of the undiluted extract used per well [mg]
BMBM. 145°C	0.278
BMBM. 133°C	0.283
BMBM dog feed	0.124
Plant Feed 0.1% (w/w) BMBM	0.094
Plant Feed 0% BMBM	0.068
fish meal	0.193
pork MBM	0.124
sheep MBM	0.182
poultry meal	0.098
bovine bone	0.055
sheep bone	0.163
goat bone	0.267
pork bone	0.242
chicken bone	0.182
rabbit bone	0.117
horse bone	0.075
horse serum	0.878
hen`s egg	0.406
whole milk powder	0.231

Table 2. Selectivity in osteocalcin specific sandwich ELISA. using MAB OC4-30 (Takara) as capture and PAB OC1-9 as detection antibody. for hidden BMBM in pork meal in three independent measurements

Extract	Day 1	Day 2	Day 3	Mean	SD
BMBM dog feed	3.469	3.352	3.501	3.441	0.078
Pork MBM	0.677	0.753	0.586	0.672	0.084
10% BMBM in pork MBM	0.887	1.116	1.232	1.078	0.176
1% BMBM in pork MBM	0.525	0.561	0.595	0.560	0.035
0.1% BMBM in pork MBM	0.476	0.524	0.514	0.505	0.025