Effect of household and industrial processing on levels of pesticide residues and degradation products in melons

Aurore Bonnechère, Vincent Hanot, Claude Bragard, Thomas Bedoret, Joris van Loco

► To cite this version:

Aurore Bonnechère, Vincent Hanot, Claude Bragard, Thomas Bedoret, Joris van Loco. Effect of household and industrial processing on levels of pesticide residues and degradation products in melons. Food Additives and Contaminants, 2012, 29 (7), pp.1058-1066. 10.1080/19440049.2012.672339 . hal-00811703

HAL Id: hal-00811703

<https://hal.science/hal-00811703>

Submitted on 11 Apr 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Effect of household and industrial processing on levels of pesticide residues and degradation products in melons

Journal:	<i>Food Additives and Contaminants</i>
Manuscript ID:	TFAC-2011-530.R1
Manuscript Type:	Original Research Paper
Date Submitted by the Author:	20-Feb-2012
Complete List of Authors:	Bonnechère, Aurore; Scientific Institute of Public Health, Pesticides Hanot, Vincent; Scientific Institute of Public Health, Bragard, Claude; Faculté d'ingénierie biologique, agronomique et environnementale UCL, Bedoret, Thomas; Redebel, Van Loco, Joris; Scientific Institute of Public Health,
Methods/Techniques:	LC/MS, GC
Additives/Contaminants:	Pesticide residues
Food Types:	Fruit, Vegetables
Abstract:	Two varieties of melons (<i>Cucumis melo</i>) were treated by two fungicides (carbendazim and maneb) and four insecticides (acetamiprid, cyromazin, imazalil and thiamethoxam) to quantify the effect of household processing on the pesticide residues. To ensure sufficiently high levels of residues in flesh and peels, the most concentrated formulations were applied pursuant to Good Agricultural Practices (GAPs). The peeling step decreased the concentration of pesticide residues for maneb, imazalil and acetamiprid by more than 90%. Cyromazin, carbendazim and thiamethoxam were reduced by ~50%. The reduction of the pesticides could not be fully explained by the systemic character of the pesticides. However, the agricultural practices (time of application), the solubility and mode of action (systemic vs. contact pesticide) of the pesticides allowed to make assumptions to explain the difference in processing factors for the studied pesticides. Degradation products (melamine and ethylenethiourea) were also investigated in this study but were not detected.

SCHOLARONE™
Manuscripts

For Peer Review Only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Effect of household and industrial processing on levels of pesticide residues and degradation products in melons

Aurore Bonnechère ^a, Vincent Hanot ^a, Claude Bragard ^b, Thomas Bedoret ^c and Joris Van Loco ^a

^a Food, Medicines and Consumer Safety Department, Scientific Institute of Public Health, Rue Juliette Wytsman 14, BE-1050 Ixelles, Belgium

^b Earth & Life Institute, Applied Microbiology-Phytopathology, Université Catholique de Louvain, Croix du Sud 2, L7.05.03, BE-1348 Louvain-la-Neuve, Belgium

^c Redebel sa, Rue de Chassart 4, 6221 Saint-Amand

Corresponding author: Vincent Hanot

Tel.: 00322/642.51.89

Fax: 00322/642.56.91

Email address: vincent.hanot@wiv-isp.be

Postal address: Rue Juliette Wytsman 14, BE-1050 Ixelles, Belgium

Abstract

Two varieties of melons (*Cucumis melo*) were treated by two fungicides (carbendazim and maneb) and four insecticides (acetamiprid, cyromazin, imazalil and thiamethoxam) to quantify the effect of household processing on the pesticide residues. To ensure sufficiently high levels of residues in flesh and peel, the most concentrated formulations were applied observing Good Agricultural Practice (GAP). The peeling step decreased the concentration of pesticide residues for maneb, imazalil and acetamiprid by more than 90%. Cyromazin, carbendazim and thiamethoxam were reduced by ~50%. The reduction of the pesticides could not be fully explained by the systemic character of the pesticides. However, the agricultural practices (time of application), solubility and mode of action (systemic vs. contact pesticide) of the pesticides could be used to explain the difference in processing factors for the studied

pesticides. Degradation products (melamine and ethylenethiourea) were also investigated in this study but were not detected.

Keywords: Pesticide residues, Processing factor, Fungicides, Insecticides, Degradation products

Introduction

Melon (*Cucumis melo*) belongs to the same family as the cucumber, squash, pumpkin and gourd and grows on the ground on a trailing vine. It is a sweet, juicy and very nutritious fruit. The major producing countries are China (which produces more than 50 % of the world production), Turkey, Iran, Spain, the United States, Rumania, Egypt and India (FAO, 2011). This crop is susceptible to *fusarium* wilt disease (destructive vascular disease), downy mildew (killing plants prematurely and reducing sugar content of melons) and gummy stem blight (lesion on stems and leaves). To protect plants particularly during autumn growing seasons and avoid these type of diseases, fungicide applications are necessary (Keinath et al. 2007, Zhao et al. 2011). Furthermore, insecticides on melons are necessary to suppress high whitefly populations and aphids (Tong-Xian 2004).

Pesticides have been linked to a wide spectrum of human health hazards, ranging from short-term impacts such as headaches and nausea to chronic impacts like cancer, reproductive harm, and endocrine disruption. Chronic health effects may occur years after even minimal exposure to them in the environment, or result from their residues ingested through food and water (Berrada et al. 2010, Claeys et al. 2011, Keikotlhaile et al. 2010). Research focuses on the proper use of pesticides in terms of authorization and registration and on compliance with Maximum Residue Limits (MRLs). Given the potential risk of pesticides for public health, the use of pesticides in agriculture is subjected to constant monitoring. (Chen et al. 2011, Claeys

et al., 2011). It is important for the consumer to know the intake of pesticide residues. The processing factors (PF: ratio between residues concentration in the processed commodity and the same in the raw commodity) are useful to estimate the level of pesticide exposure at the point of consumption after processing (Ling et al. 2011). Until now, no processing factors are available for melons. However, melon consumption is increasing in the EU (AND International 2010). Indeed, among the EU 27, the volumes of import of melons increased twofold between 1999 and 2008.

It is well known that peeling plays a role in the reduction of residues (Holland et al. 1994, Kaushik et al. 2009, Timme et al. 2004). Several studies of peeling have been done on potatoes to measure some pesticides and minimum 90% reduction of pesticide residues were observed (Kaushik et al. 2009, Lentza-Rizos et al. 2001, Lewis et al. 1996, Randhawa et al. 2007, Schattenberg et al. 1996). The peeling of apples, tomatoes, peppers, carrots, bananas, oranges and lemons were also investigated in several studies (Balnova et al. 2006, Burchat et al. 1998, Chavarri et al. 2005, Rasmussen et al. 2003, Timme & Walz-Tylla 2004). In these experiments, peeling decreased residues (70 to 100%). However, some pesticides were not influenced by this process because of their systemic activity (Krol et al. 2000).

Melons, frequently consumed in Belgium, were chosen in cooperation with the Belgian Federal Agency for Safety of the Food Chain (FASFC). Pesticides were chosen according to their frequency of MRL exceedance, frequency of detection and their toxicity (FASFC 2007, FASFC 2008). Selected pesticides included acetamiprid, carbendazim, cyromazin, imazalil, maneb and thiamethoxam.

To establish of processing factors for these pesticide residues, melons were grown, sprayed, harvested, peeled, homogenized and analyzed. The selected pesticides and the degradation products (ETU from maneb and melamine from cyromazin) were analysed before and after peeling with several analytical methods (GC-MS/MS, GC-ECD, UHPLC-MS/MS).

1
2
3 1
4
5 2
6
7 3 **Materials and Methods**

8
9 4 *Materials*

10
11 5 *Pesticides*

12
13
14 6 Deuterated ethylenethiourea (d₄-ETU), ethylenethiourea, carbendazim and acetamiprid
15
16 7 were purchased from Dr. Ehrenstorfer GmbH (Augsburg, Germany) with certified purity of
17
18 8 99.0%. Thiram, cyromazin, melamine, imazalil and thiamethoxam were purchased from
19
20 9 Sigma-Aldrich (Seelze, Germany) with certified purity of 99.0%, 99.9%, 99.0%, 99.8% and
21
22 10 99.7%, respectively. Carbon disulfide (99.9%) was obtained from Merck (Darmstadt,
23
24 11 Germany).

25
26 12
27
28
29 13 *Chemicals for analysis*

30
31 14 Ultrapure water (< 8 MΩ·cm resistance) was supplied by a Millipore purification system
32
33 15 (Millipore Milli-Q Water System, Bedford, USA). Methanol, dichloromethane and
34
35 16 acetonitrile were of HPLC grade from Biosolve (Valkenswaard, The Netherlands). Iso-octane,
36
37 17 sodium hydroxide and ammoniac (25%) came from Merck (Darmstadt, Germany).
38
39 18 Trifluoroacetic acid (TFA) and acid acetic (with certified purity of at least 99% for both) as
40
41 19 well as ammonium acetate were purchased from Aldrich (Steinheim, Germany). Hydrochloric
42
43 20 acid (37%) and tin(II)-chloride were obtained from VWR (Fontenay sous Bois, France).
44
45 21 Ethanol (96%), ammonium chloride (99.5%), celite® 545, sodium sulphate (99%) and sodium
46
47 22 chloride (99%) were purchased from VWR (Leuven, Belgium).

48
49
50 23 The stock solution of carbon disulfide (CS₂), for analysis of maneb, was prepared at 4 mg
51
52 24 mL⁻¹ in iso-octane. Dilute standards at 40 µg mL⁻¹ were prepared by dilution of the stock

1 solution with iso-octane. A solution of thiram was prepared with ethanol to obtain an
2 equivalent concentration in CS₂ of 63.23 mg L⁻¹.

3 The stock solutions of acetamiprid, carbendazim, cyromazin, imazalil and thiamethoxam
4 for LC-MS/MS were prepared at 1 mg mL⁻¹ in acetonitrile containing 0.1% of acetic acid.
5 Dilute standards at 10 µg mL⁻¹, 1 µg mL⁻¹ and 100 ng mL⁻¹ were prepared by dilution of the
6 stock solution with methanol.

7 The stock solution of internal standard, ETU D4, was prepared at 100 µg mL⁻¹ in
8 methanol. The stock solution of ETU was prepared at 1 mg mL⁻¹ in methanol. Dilute
9 standards were prepared by dilution of the stock solution with water.

10 The stock solution of melamine was prepared at 1 mg mL⁻¹ in NaOH 0.1%. Dilute
11 standards at 10 µg mL⁻¹ and 1 µg mL⁻¹ were prepared by dilution of the stock solution with
12 water.

13 Solutions for calibration were prepared with spiking of diluted standard solution on blank
14 extract of melon. The blank extract came from melons previously tested in which no
15 pesticides were detected above the limit of detection. All solutions were stored at -18 °C.

16 17 **Produce**

18 *Cultivation*

19 The purpose of the field experiment was to produce in parallel two different melons
20 varieties, Mohican and Pancha, exposed to six selected pesticides.

21 Field melon was grown at Villers-Perwin, Belgium by Redebel sa. under greenhouse
22 conditions. Four applications of pesticides were performed on different crop growth stages as
23 shown in Table 1. The concentrations of the six active substances and the number of
24 applications were as high as possible, but following the GAPs (pre-harvest time, time between
25 two applications...), to ensure sufficiently high pesticide levels for the effects of processing

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

practices to be studied. It should be noted that some pesticides are not approved for the melon crops in Belgium but are in other countries. The formulations were applied in two passing using vertical spraying boom equipment at the proposed normal settings and timing.

Harvest

Fruit specimens were taken manually. Sterile disposable (nitrile) gloves were worn for the sampling. The first and the last plants of plot, diseased and undersized fruits were excluded from sampling. To obtain more than 70 kg of melons, 67 melons Pancha and 103 melons Mohican were randomly collected from the whole plot. A number identified each variety. Melons were stored in a refrigerator room at 4 °C directly after the sampling. They were transported in refrigerated condition to the laboratory on the day of harvest.

Sample preparation and processing

Overview of processing and sampling steps

Figure 1 shows the sampling flow diagram. To minimize the factor of variability, the two melon varieties were each divided into two batches of 18 kg of raw melons, giving four batches in total (Figure 1). As shown in Figure 1, part of melons did not undergo any processing and was used as “positive control sample”. Peeling step for all batches was executed on the same day.

At each step, melons were homogenized and 500 g were collected and stored at -20 °C until pesticide residue analysis. Frozen samples were analyzed within one month.

Unprocessed melon

These melons, which did not undergo any processing are the “positive control sample”. It is the raw melon used to calculate the PFs. The harvested melons (18 kg of melons taken

randomly) were cut into pieces and comminuted with the Robot Coupe® R23 (Mont-Saint-Genevieve, Belgium). A sample was taken after a rough cut (~ 1 cm) and frozen for the analysis of mancozeb. The rest was comminuted for a longer time and frozen until analysis.

Peeling

The peeling was realised with a knife with a blade of 10 cm and corresponded to the peeling done by the consumer at home. We left the green part on the peels (< 0.5 cm). Melons Mohican were smaller than melons Pancha but both varieties were highly variegated.

After removal of the peels, flesh (about 10 kg) and peels (about 7 kg) were comminuted separately with the Robot Coupe® R23 (Mont-Saint-Genevieve, Belgium) and frozen until analysis.

Extraction and analysis of pesticides

Analysis of maneb by GC-ECD

The analysis of maneb as CS₂ (compound obtained when dithiocarbamates are heated in presence of acid and tin chloride) is based on methods normalized by the European Committee for Standardization and modified by de Kok to replace the existing headspace chromatography method by a liquid injection in GC-ECD (de Kok 2001, European Committee for Standardization 1998).

The extraction of 25 g of matrix was done in presence of hydrochloric acid, iso-octane and tin chloride. A closed vessel was stirred during 2 h in a water bath at 80 °C. After cooling, 1 µl of the supernatant containing 0.28 g matrix per mL was injected on a Varian 3400 gas chromatographic system coupled with a ⁶³Ni Electron Capture Detector and a Varian 8200C autosampler. The GC separation of CS₂ was achieved on a DB-5 column (60 m, 0.53 mm and 1.5 µm) from Grace (Deerfield, USA). The helium gas flow was at 2.6 mL min⁻¹ at 5 psi. The

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 injector and detector temperatures were 250 °C and 300 °C, respectively. The column
2 temperature was at 50 °C for 5 min, after that it increased by 50 °C per min until 250 °C. This
3 temperature was maintained for 5 min to obtain a total run time of 19 min. The retention time
4 of CS₂ was 3.3 min.

6 *Analysis of acetamiprid, carbendazim, cyromazin, imazalil and thiamethoxam by LC-MS/MS*

7 This method followed the Granby principle (Granby et al. 2004) with some adaptations. A
8 10 g sample of melons was extracted with a mixture of an extraction solvent (5mM
9 ammonium acetate, 90% MeOH-10% water), blended during 1 min with an ultra-turrax
10 homogenizer (Ultra-turrax IKA) and filtrated on büchner. Oxfendazole, the internal standard
11 was added to the filtrate. Mixture of extraction was used to give an extract volume of 60 mL.
12 The raw extract was diluted with mobile phase A (10% MeOH, 1% ammonium acetate 0.5 M
13 and 89% water) to a final concentration equivalent to 0.1 g matrix per mL and injected after
14 filtration on 0.2 µm. The LC separation was achieved on an Acquity UPLC BEH C₁₈ column
15 (1.7 µm, 2.1 × 100 mm) from Waters (Ireland). A gradient of mobile phase A and mobile
16 phase B (containing 10% water, 1% ammonium acetate 0.5M and 89% of methanol) supplied
17 at 0.45 mL min⁻¹ was applied. Starting from 99.9% of mobile phase A, the mobile phase was
18 linearly increased to 99.9% of mobile phase B in 10 min and maintained during two min. The
19 gradient was then returned to initial conditions in 0.1 min and held there for 2.9 min for
20 stabilization. The mass spectrometer operated in a positive electrospray ionization (ESI) mode
21 and acquired two transitions for each pesticide. Typical optimized ESI voltage settings are
22 presented in Table 2. Argon was used as the collision gas at a setting of 0.35 mL min⁻¹. The
23 dwell time was 10 ms per channel for quantifier ions and 5 ms for second transition.

25 *Analysis of ethylenethiourea by LC-MS/MS*

The method of analysis is described in a paper of Bonnechère *et al.* (Bonnechère *et al.* 2011). Briefly, ETU was extracted from the homogenized melon (10 g), after thawing of this sample, by a 3:1 mixture of methanol-water. After 0.2 µm filtration, the extract containing 0.333 g matrix per mL was injected in LC-MS/MS (conditions in Table 2) on a Uptisphere 5MM1 mixed-mode chromatographic column (100 × 2.1mm, 5µm) with two different silanes (C8/SCX) bonded on silica (Interchim, Montluçon, France), using an isocratic mobile phase of 0.1% trifluoroacetic acid / methanol (95/5). The injection volume was 5 µL in full loop and the run time was 2 min.

Analysis of melamine by LC-MS/MS

The method of analysis is described in a paper of Goscinny *et al.* (Goscinny *et al.* 2011). The extraction of melamine is carried out using an acetonitrile-water-dichloromethane in an acidic environment. After SPE and 0.2 µm filtration, the extract was injected in LC-MS/MS (conditions in Table 2) on a Acquity HILIC column (100 × 2.1mm, 1.7µm) (Waters, Ireland), using an isocratic mobile phase containing 2% ammonium acetate 0.5M, 8% water and 90% of acetonitrile. The injection volume was 1 µL and the run time was 2 min.

Quantification and quality control

Limits of Quantification (LOQ) were estimated from the lowest calibration level. Based on the signal-to-noise ratio upper or equal to 6, LOQs were at 40 µg kg⁻¹ for maneb, 2 µg kg⁻¹ for imazalil, 5 µg kg⁻¹ for cyromazin, 2 µg kg⁻¹ for thiamethoxam, and 1 µg kg⁻¹ for acetamiprid and carbendazim, 50 µg kg⁻¹ for ETU and 25 µg kg⁻¹ for melamine. The linearity of the instrument was evaluated with 5 or 6 points in the range of 50-1000 µg kg⁻¹ (wet weight basis) for ETU, 25-600 µg kg⁻¹ for melamine, 50-5000 µg kg⁻¹ for maneb and 10-1000

1 $\mu\text{g kg}^{-1}$ for the pesticide residues analyzed by LC-MS/MS. In all cases, good linearity was
2
3
4
5 achieved with correlation coefficients > 0.995 .
6

7 For each batch of analysis, a control (a blank of melon) and a sample were spiked prior to
8
9 extraction and run. In the control, no pesticide must be found with a concentration greater
10
11 than the LOQ. If it is not the case that means a contamination and the analysis of the samples
12
13 must be redone. The spiked sample must have recovery of extraction between 70% and 120%
14
15 (SANCO/10684/2009) and the concentrations found in this sample were reported on the
16
17 control chart, which allows following the trend. Recoveries during the analysis of melon were
18
19 of 95.7% for ETU, 73.2% for maneb, 83.4% for melamine, 87.3% for cyromazin, 87.3% for
20
21 acetamiprid, 84.0% for carbendazim and 83.4% for imazalil. The calibration was checked
22
23 using a standard injected at the end of the sequence and could not deviate by more than 20%
24
25 from the injection of the same standard at the beginning of the sequence.
26
27
28
29
30
31

32 ***Calculation of processing factors***

33
34 Processing factors (PFs) were calculated for all transformation steps by a ratio between
35
36 the pesticide residue concentration (mg kg^{-1}) in the processed commodity and the pesticide
37
38 residue concentration (mg kg^{-1}) in the raw commodity.
39
40
41
42

43 **Results and discussion**

44
45 Sprayed pesticides were detected at various concentrations. In spite of the high
46
47 concentration of active substances applied on the raw melon, the concentrations for
48
49 thiamethoxam, carbendazim and imazalil were rather low. The lowest concentrations in the
50
51 raw product were observed for thiamethoxam and carbendazim (Table 3). Thiamethoxam was
52
53 applied at the lowest concentration in this study and carbendazim is the only which was
54
55 applied once, as early as the opening of the main stem (Table 1).
56
57
58
59
60

In this study, the peeling was done with a knife. Table 4 represents all processing factors for the peeling step for the six detected pesticides for both varieties. The peeling removed an important part of the pesticide residues between 62% and 95% for melon Mohican and between 52% and 95% for melon Pancha.

Maneb and cyromazin are contact pesticides and form deposit on the surfaces of leaves and fruits (Table 4). After the peeling step, in spite of an important reduction for the maneb (91% for Mohican and 93% for Pancha), the decrease of cyromazin did not exceed 62% and 52% for Mohican and Pancha respectively. However this smaller reduction could be explained by the higher solubility of cyromazin compared with the other pesticides (Table 4). Compounds applied to the plants (leaves) do not have to cross the symplast to arrive in the xylem contrary to compounds passing by roots. On the other hand, they have to cross the more or less waxy cuticle according to the type of plant and the age. With the age, fractures in the waxy coat form hydrophilic pores allowing the passage of the most hydrophilic molecules as cyromazin (Al-Sayed 2007). With this particularity, cyromazin could enter in the fruit by the flow of xylem and be found in the flesh. As Krol *et al.* has also confirmed it is possible that pesticide residues are incorporated into plant tissue proportional to the time they remain on biologically active crops in the field. This may even be true of pesticides that are not specifically labelled as systemic (Krol et al. 2000). The other four pesticides (acetamiprid, carbendazim, imazalil and thiamethoxam) are systemic. Nevertheless, acetamiprid and imazalil were strongly reduced by 91% for Mohican, 89% for Pancha and 91% for Mohican and Pancha respectively. The reduction was somewhat less important for the thiamethoxam (67% & 58%) and carbendazim (63% & 47%) (Table 4). The difference of behaviour could be explained by the time of application of these 4 active substances (Table 1). Indeed, carbendazim was the first pesticide applied when the plant of melons had it's 5th flower open on main stem. With this early application, carbendazim could enter in the plant and was

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 present in it before the apparition of fruit. The others were applied in the latest stage of
2 development when 10% of fruits showed typical fully ripe color (Table 1). Otherwise,
3 thiamethoxam has a good solubility and an octanol-water partition coefficient which allow
4 him at the same time to circulate in the phloem and in the xylem and consequently to be
5 present in the fruit (Table 4). On one hand, more than 90 % of acetamiprid, imazalil and
6 maneb, contact as well systemic pesticides, were removed with this process. On the other
7 hand, ~50% of carbendazim and thiamethoxam (systemic pesticide) and cyromazin (contact
8 pesticide) were eliminated. These 2 types of reduction could not be explained by the systemic
9 character of the pesticides. However, the agricultural practices (time of application) and the
10 water solubility could explain statistically these differences between the pesticides. The
11 octanol-water partition coefficient is also necessary to explain the behaviour of cyromazin. It
12 is a particular case, not shown in the statistical test which is a global approach. Indeed a
13 systemic pesticide applied late on the plants is found with a low concentration in the flesh.
14 Until now, several studies have explained processing factor with the physico-chemical
15 properties but they did not reflect the moments of application on the harvest (Rasmussen et
16 al. 2003, Burchat et al. 1998, Watts et al. 1974, Lee et al. 2009). These results show that it
17 really important to take into account the time of application of active substances for the risk
18 assessment. Indeed as Krol (2000) said, a systemic pesticide applied late could not act as a
19 systemic. For risk assessment, it is also necessary to be aware that the levels of residues in
20 melons depend on the time of application quite as the statistical test reflects it.

21 Reductions of 90% are consistent with previous studies on the peeling of various fruit and
22 vegetables (Balinova et al. 2006, Boulaid et al. 2005, Chavarri et al. 2005, Kaushik et al.
23 2009, Lentza-Rizos & Balokas 2001, Rasmussen et al. 2003, Schattenberg et al. 1996,
24 Timme & Walz-Tylla 2004, Fernandez-Cruz et al. 2006). It was indeed demonstrated that a
25 majority of the insecticides or fungicides applied directly to crops undergo very limited

1 movement or penetration of the cuticle. However, residues of these materials are confined to
2
3 the outer surfaces where they are amenable to removal by peeling (Holland et al. 1994,
4
5 Timme & Walz-Tylla 2004, Burchat et al. 1998).
6
7

8
9 In this study, we also investigated the degradation products of cyromazin and maneb,
10
11 being melamine and ethylenethiourea (ETU), respectively. In the unprocessed melon, these
12
13 two by-products were not detected (Table 3). In general, ETU is formed in an acidic, heated
14
15 environment and it is not the case with the peeling (Dubey et al. 1997, Hwang et al. 2003,
16
17 Kontou et al. 2001, Sottani et al. 2003).
18
19

20
21 If we compare the concentrations applied on the plants and the concentrations found in
22
23 melon, there is high variability between the active substances under investigation (Table 1).
24
25 For example, the concentration in maneb applied on plants was more than 50 times higher
26
27 than the concentration in cyromazin. The concentration retrieved in the melons (Table 3)
28
29 showed that the concentration in maneb is 1.5 times lower than for cyromazin. However, the
30
31 correlation of the results of both varieties was good.
32
33

34
35 Mass balance recoveries were calculated to compare the total mg of each pesticide in the
36
37 whole commodity to the residue in the flesh and peels after peeling. Measured concentrations
38
39 available in Table 4 were corrected with the mass of the whole melons, the flesh and peels
40
41 comminuted to obtain reconstituted samples. Taking account the measured uncertainties, the
42
43 measured concentration and calculated concentration were close, as shown in Table 5, The
44
45 Table 6 showed all corrected PFs. These PFs shows some difference between varieties. An
46
47 explanation could be the difference of size between both varieties. The proportion peels/flesh
48
49 is more important and engenders variations between the PFs of both varieties for certain
50
51 pesticides.
52
53
54
55
56
57
58
59
60

Conclusion

During this study, processing factors for peeling of melons were determined for acetamiprid, carbendazim, cyromazin, imazalil, maneb and thiamethoxam. Mechanical peeling, a typical household process, will remove 52 to 91% of the pesticides on the melon. The reduction of the pesticides could not be fully explained by the systemic character of the pesticides. However, the agricultural practices (time of application), the solubility and mode of action (systemic vs. contact pesticide) of the pesticides allowed to make assumptions to explain the difference in processing factors for the studied pesticides.

Two degradation products (melamine and ETU) with a higher toxicity than the parent compounds were also studied and were not detected before and after peeling.

Acknowledgement

This study was funded by the Belgian Federal Public Service of Health, Food Chain Safety and Environment (contract RT 08/3 PESTRANS). We extend our grateful thanks to Redebel for setting up field trials to obtain commodities, to Martine Deridder and Marie-Anne Van den Bergh for their help during the analysis of the samples.

References

- Al-Sayeda H. (2007). Transfert d'un insecticide systémique, l'imidaclopride, chez la tomate: implication du transport phloémien.
- AND International. 2010. Normes de commercialisation dans le secteur des fruits et légumes.
- Balinova AM, Mladenova RI, & Shtereva DD. 2006. Effects of processing on pesticide residues in peaches intended for baby food. *Food Addit. Contam.* 23(9): 895-901.
- Berrada H, Fernandez M, Ruiz MJ, Molto JC, & Font G. 2010. Surveillance of pesticide residues in fruits from Valencia during twenty months (2004/05). *Food Control.* 21(1): 36-44.
- Bonnechere, A., Hanot, V., & Van Loco, J. A rapid and environmental friendly determination of the dithiocarbamates metabolites ethylenethiourea and propylenethiourea in fruit and vegetables by ultra high performance liquid chromatography tandem mass spectrometry. *J. Chromatogr. A.* 1218(29):4627-4631.
- Boulaid M, Aguilera A, Camacho F, Soussi M, & Valverde A. 2005. Effect of household processing and unit-to-unit variability of pyrifenoxy, pyridaben, and tralomethrin residues in tomatoes. *J. Agric. Food Chem.* 53(10): 4054-4058.
- Burchat CS, Ripley BD, Leishman PD, Ritcey GM, Kakuda Y, & Stephenson GR. 1998. The distribution of nine pesticides between the juice and pulp of carrots and tomatoes after home processing. *Food Addit. and Contam.* 15(1): 61-71.
- Chavarri MJ, Herrera A, & Arino A. 2005. The decrease in pesticides in fruit and vegetables during commercial processing. *Int. J. Food Sci. Tech.* 40(2): 205-211.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Chen C, Qian Y, Chen Q, Tao C, Li C, & Li Y. 2011. Evaluation of pesticide residues in
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Chen C, Qian Y, Chen Q, Tao C, Li C, & Li Y. 2011. Evaluation of pesticide residues in
2 fruits and vegetables from Xiamen, China. Food Control. 22(1114-1120.
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Claeyls WL, Schmit JF, Bragard C, Maghuin-Rogister G, Pussemier L, & Schiffers B. 2011.
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Exposure of several Belgian consumer groups to pesticide residues through fresh fruit
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 and vegetable consumption. Food Control. 22(3-4): 508-516.
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 de Kok. Developments in GC-MS(/MS) and LC-MS(/MS), Methodology for pesticide residue
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 analysis [Internet]. Rome. Available from:
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 <http://workshop.iss.it/DiMuccioV/Testi/Relazioni/adekok.ppt>
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Dubey JK, Heberer T, & Stan HJ. 1997. Determination of ethylenethiourea in food
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 commodities by a two-step derivatization method and gas chromatography with
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 electron-capture and nitrogen-phosphorus detection. J. Chromatogr. A. 765(1): 31-38.
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 European Committee for Standardization. 1998. Final Draft prEN 12396-1, prEN 12396-2,
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 prEN 12396-3; Non-fatty foods - Determination of dithiocarbamate and thiuram
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 disulfide residues.
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 FASFC. 2007. Controls of pesticide residues in food Belgium 2006. Brussels (Belgium)
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 FASFC. 2008. Controls of pesticide residues in food Belgium 2007. Brussels (Belgium)
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Fernandez-Cruz ML, Grimalt S, Villarroya M, Lopez FJ, Llanos S, & Garcia-Baudin JM.
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 2006. Residue levels of captan and trichlorfon in field-treated kaki fruits, individual
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 versus composite samples, and after household processing. Food Addit. and Contam.
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 23(6): 591-600.
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1 Food and agriculture organization of the United Nations (FAO). FAOSTAT [Internet].
2 <http://faostat.fao.org/site/567/DesktopDefault.aspx?PageID=567>
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- 1 Goscinnny S, Hanot V, Halbardier J-F, Michelet J-Y, & Van Loco J. 2011. Rapid analysis of
2 melamine residue in milk, milk products, bakery goods and flour by ultra-performance
3 liquid chromatography/tandem mass spectrometry: From food crisis to accreditation.
4 Food Control. 22(226-230).
5
6
7 Granby K, Andersen JH, & Christensen HB. 2004. Analysis of pesticides in fruit, vegetables
8 and cereals using methanolic extraction and detection by liquid chromatography-
9 tandem mass spectrometry. Analytica Chimica Acta. 520(1-2): 165-176.
10
11
12 Holland PT, Hamilton D, Ohlin B, & Skidmore MW. 1994. Effects of storage and processing
13 on pesticide residues in plant products. Pure Appl. Chem. 66(2): 335-356.
14
15
16 Hwang ES, Cash JN, & Zabik MJ. 2003. Determination of degradation products and
17 pathways of mancozeb and ethylenethiourea (ETU) in solutions due to ozone and
18 chlorine dioxide treatments. J.Agric.Food Chem. 51(5): 1341-1346.
19
20
21 Kaushik G, Satya S, & Naik SN. 2009. Food processing a tool to pesticide residue dissipation
22 - A review. Food Res.Int. 42(26-40).
23
24
25 Keikotlhaile BM, Spanoghe P, & Steurbaut W. 2010. Effects of food processing on pesticide
26 residues in fruits and vegetables: A meta-analysis approach. Food Chem.Toxicol.
27 48(1): 1-6.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- 17

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 control of downy mildew and gummy stem blight on melon. Crop Protection. 26(2):
2
3 83-88.
4
5
6
7
8
9 3 Kontou S, Tsipi D, Oreopoulou V, & Tzia C. 2001. Determination of ETU in tomatoes and
10
11 4 tomato products by HPLC-PDA. Evaluation of cleanup procedures. J.Agric.Food
12
13 5 Chem. 49(3): 1090-1097.
14
15
16 6 Krol WJ, Arsenault TL, Harry M, & Incorvia Mattina MJ. 2000a. Reduction of Pesticide
17
18 7 Residues on Produce by Rinsing. J.Agric.Food Chem. 48(10): 4666-4670.
19
20
21 8 Lee MG & Jung D. 2009. Processing factors and removal ratios of select pesticides in hot
22
23 9 pepper leaves by a successive process of washing, blanching, and drying. Food Sci.
24
25 10 Biotechnol. 18(5): 1078-1082.
26
27
28
29 11 Lentza-Rizos C & Balokas A. 2001a. Residue levels of chlorpropham in individual tubers and
30
31 12 composite samples of post-harvest treated potatoes. J.Agric.Food Chem. 49: 710-714.
32
33
34 13 Lewis DJ, Thorpe SA, & Reynolds SL. 1996. The carry-through of residues of thiabendazole,
35
36 14 tecnazene and chlorpropham from potatoes following manufacture into potato crisps
37
38 15 and jacket potato crisps. Food Addit. Contam. 13(2): 221-229.
39
40
41
42 16 Ling Y, Wang H, Yong W, Zhang F, Sun L, Yang ML, Wu YN, & Chu XG. 2011. The
43
44 17 effects of washing and cooking on chlorpyrifos and its toxic metabolites in vegetables.
45
46 18 Food Control. 22(1): 54-58.
47
48
49
50 19 Randhawa MA, Anjum FM, Asi MR, Butt MS, Ahmed A, & Randhawa MS. 2007. Removal
51
52 20 of endosulfan residues from vegetables by household processing. J. Sci. Ind. Res.
53
54 21 66(10): 849-852.
55
56
57
58
59
60

- 1 Rasmussen RR, Poulsen ME, & Hansen HC. 2003. Distribution of multiple pesticide
residues in apple segments after home processing. Food Addit.Contam. 20(11): 1044-
1063.
- SANCO/10684/2009. Method validation and quality control procedures for pesticide residues
analysis in food and feed [Internet]. 2009.
http://ec.europa.eu/food/plant/protection/resources/qualcontrol_en.pdf
- Schattenberg HJ, Geno PW, Hsu JP, Fry WG, & Parker RP. 1996. Effect of household
preparation on levels of pesticide residues in produce. J. AOAC Int. 79(6): 1447-1453.
- Sottani C, Bettinelli M, Fiorentino ML, & Minoia C. 2003. Analytical method for the
quantitative determination of urinary ethylenethiourea by liquid
chromatography/electrospray ionization tandem mass spectrometry. Rapid
Commun.Mass Spectrom. 17(20): 2253-2259.
- Timme G & Walz-Tylla B (2004). Effects of food preparation and processing on pesticide
residues in commodities of plant origin. In Denis Hamilton & Stephen Crossley (Ed.),
Pesticides residues in food and drinking water: human exposure and risks (pp. 121-
148). John Wiley & Sons.
- Tong-Xian L. 2004. Toxicity and efficacy of spiromesifen, a tetrone acid insecticide, against
sweetpotato whitefly (homoptera: aleyrodidae) on melons and collards. Crop
Protection. 23(6): 505-513.
- Watts RR, Storherr RW, & Onley JH. 1974. Effects of cooking on ethylenebisdithiocarbamate
degradation to ethylenethiourea. B. Environ. Contam. Tox. 12(2): 224-226.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Zhao Q, Dong C, Yang X, Mei X, Ran W, Shen Q, & Xu Y. 2011. Biocontrol of Fusarium
2 wilt disease for Cucumis melo melon using bio-organic fertilizer. Appl. Soil Ecol.
3 47(1): 67-75.
4
5
6

For Peer Review Only

Figure Caption

Figure 1: Flow diagram of processing steps and sampling for one variety of melon.

For Peer Review Only

Figure 1: Flow diagram of processing steps and sampling for one variety of melon.
90x60mm (300 x 300 DPI)

Table 1: Pesticides with their product name, rate of application and the crop stage at the moment of application

Pesticides	Product Name	Active ingredient (g ha ⁻¹)	Application date	Crop Stage (BBCH ^a)	Number of days before harvest	MRL (mg/kg)
Acetamiprid	Gazelle SP	105	9/08/2010	BBCH 83 = 30% of fruits show typical fully ripe colour (FTRC)	10	<i>0.01</i> *
		98	16/08/2010	BBCH 87 = 70% FTRC	3	
Carbendazim	Punch SE	319	5/07/2010	BBCH 65 = 5th flower open on main stem	45	<i>0.1</i> *
Cyromazine	Trigard 100 SL	58	2/08/2010	BBCH 81 = 10% FTRC	17	<i>0.3</i>
		63	9/08/2010	BBCH 83 = 30% FTRC	10	
		59	16/08/2010	BBCH 87 = 70% FTRC	3	
Imazalil	Fungaflor EC	97	2/08/2010	BBCH 81 = 10% FTRC	17	<i>2</i>
		105	9/08/2010	BBCH 83 = 30% FTRC	10	
		98	16/08/2010	BBCH 87 = 70% FTRC	3	
Maneb	Trimangol 75 WG	1950	2/08/2010	BBCH 81 = 10% FTRC	17	<i>1</i>
		2108	9/08/2010	BBCH 83 = 30% FTRC	10	
		1971	16/08/2010	BBCH 87 = 70% FTRC	3	
Thiamethoxam	Actara 25 WG	5	2/08/2010	BBCH 81 = 10% FTRC	17	<i>0.2</i>
		5	9/08/2010	BBCH 83 = 30% FTRC	10	
		5	16/08/2010	BBCH 87 = 70% FTRC	3	
			19/08/2010	BBCH 89 = fully ripe: fruit have typical fully ripe colour	HARVEST	

^a The BBCH-scale is a system for a uniform coding of phenologically similar growth stages of all mono- and dicotyledonous plant species.

* Indicates lower limit of analytical determination (Reg. (EU) No 600/2010)

Italics characters indicate that these substances are not authorized on melon in Belgium

Table 2
MS detection and selected ion for multi-residues analysis by LC-MS/MS. Quantifier daughter ions are in bold.

MS detection								
Extractor (V)	4							
RF Lens (V)	0.2							
Cone Gas Flow (L/Hr)	50							
Desolvation Gas Flow (L/Hr)	800							
Selected Ion	Parents (m/z)	Daughter (m/z)	Capillary (kV)	Source Temperature (°C)	Desolvation Temperature (°C)	Cone (V)	Collision (V)	Retention time (min)
Cyromazin	167.01	84.9	0.6	130	400	28	16	1.53
		60.3					18	
	292.3	211.0					19	
Thiamethoxam		180.9	0.6	130	400	19	24	2.49
	294.3	211.0					12	
Acetamiprid		125.8	0.6	130	400	26	21	3.65
	223.0	89.9					35	
Carbendazim		160.1	0.6	130	400	25	18	3.92
	192.1	132.1					30	
Oxfendazole	316.0	159.0	0.6	130	400	34	28	5.35
Imazalil		159.0	0.6	130	400	30	20	8.67
	297.1	69.1					20	
ETU	102.9	44.3	0.3	130	450	31	15	0.91
		85.9					15	
ETU D4		45.1	0.3	130	450	31	15	0.89
	106.9	48.2					15	
Melamine		84.8	0.8	150	400	42	22	0.91
	126.9	67.8					17	

Table 3

Concentrations (mg kg^{-1}) of pesticides and degradation products for the two melon varieties after peeling step. Mean values ($\pm$ SD, $n = 2$) of two melon batches are depicted.

Variety 1: Mohican				
Pesticides	LOQ (mg/kg)	Raw Mean values ($\pm$ SD)	Flesh Mean values ($\pm$ SD)	Peels Mean values ($\pm$ SD)
Acetamiprid	0.001	0.042 ($\pm$ 0.00057)	0.0039 ($\pm$ 0.00007)	0.081 ($\pm$ 0.0087)
Carbendazim	0.001	0.0014 ($\pm$ 0.00049)	< LOQ	0.0042 ($\pm$ 0.0018)
Cyromazin	0.005	0.078 ($\pm$ 0.0022)	0.029 ($\pm$ 0.0031)	0.110 ($\pm$ 0.0024)
Imazalil	0.002	0.0092 ($\pm$ 0.00085)	< LOQ	0.017 ($\pm$ 0.0039)
Maneb	0.040	0.48 ($\pm$ 0.075)	0.043 ($\pm$ 0.032)	0.82 ($\pm$ 0.086)
Thiamethoxam	0.002	0.0045 ($\pm$ 0.0000)	< LOQ	0.0074 ($\pm$ 0.0003)
ETU	0.050	< LOQ	< LOQ	< LOQ
Melamine	0.025	< LOQ	< LOQ	< LOQ
Variety 2: Pancha				
Pesticides	LOQ (mg/kg)	Raw Mean values ($\pm$ SD)	Flesh Mean values ($\pm$ SD)	Peels Mean values ($\pm$ SD)
Acetamiprid	0.001	0.046 ($\pm$ 0.0029)	0.0051 ($\pm$ 0.0018)	0.083 ($\pm$ 0.00092)
Carbendazim	0.001	0.0012 ($\pm$ 0.00021)	< LOQ	0.0034 ($\pm$ 0.00014)
Cyromazin	0.005	0.081 ($\pm$ 0.013)	0.039 ($\pm$ 0.0026)	0.084 ($\pm$ 0.00064)
Imazalil	0.002	0.011 ($\pm$ 0.00014)	< LOQ	0.017 ($\pm$ 0.0032)
Maneb	0.040	0.55 ($\pm$ 0.050)	< LOQ	0.91 ($\pm$ 0.062)
Thiamethoxam	0.002	0.0036 ($\pm$ 0.0003)	< LOQ	0.0074 ($\pm$ 0.0001)
ETU	0.050	< LOQ	< LOQ	< LOQ
Melamine	0.025	< LOQ	< LOQ	< LOQ

Table 4: Mean values ($\pm$ SD, n = 2) of processing factors (PF) for peeling step for six pesticides in two varieties of melons.

Pesticides	PF peeling ($\pm$ SD)		Action mode	log-octanol-water-partitioning coefficients	Water solubility at 20 °C (mg/L)
	Mohican	Pancha			
Acetamiprid	0.091 ($\pm$ 0.002)	0.11 ($\pm$ 0.040)	Systemic	0.8	2950
Carbendazim	0.37 ($\pm$ 0.14)	0.43 ($\pm$ 0.08)	Systemic	1.48	8
Cyromazin	0.38 ($\pm$ 0.04)	0.48 ($\pm$ 0.08)	Non systemic	0.069	13000
Imazalil	0.11 ($\pm$ 0.010)	0.093 ($\pm$ 0.001)	Systemic	3.82	22.4
Maneb	0.088 ($\pm$ 0.068)	0.073 ($\pm$ 0.007)	Non systemic	-0.45	178
Thiamethoxam	0.33 ($\pm$ 0.16)	0.42 ($\pm$ 0.20)	Systemic	-0.13	4100

Table 5: Active substances (mg) accounting for mass balance of pesticides and degradation products for the two melon varieties after peeling step. Mean values ($\pm$ SD, $n = 2$) of two melon batches are depicted.

Variety 1: Mohican				
Pesticides	Raw (measured)	Flesh	Peels	Raw (calculated)
	Mean values ($\pm$ SD)	Mean values ($\pm$ SD)	Mean values ($\pm$ SD)	Mean values ($\pm$ SD)
Acetamiprid	0.75 ($\pm$ 0.058)	0.037 ($\pm$ 0.0001)	0.57 ($\pm$ 0.084)	0.60 ($\pm$ 0.084)
Carbendazim	0.024 ($\pm$ 0.0073)	0.048 ($\pm$ 0.0001)	0.029 ($\pm$ 0.014)	0.034 ($\pm$ 0.014)
Cyromazine	1.38 ($\pm$ 0.049)	0.281 ($\pm$ 0.026)	0.77 ($\pm$ 0.014)	1.05 ($\pm$ 0.04)
Imazalil	0.16 ($\pm$ 0.026)	0.010 ($\pm$ 0.0001)	0.12 ($\pm$ 0.032)	0.13 ($\pm$ 0.032)
Maneb	8.60 ($\pm$ 1.88)	0.41 ($\pm$ 0.31)	5.78 ($\pm$ 0.84)	6.19 ($\pm$ 1.15)
Thiamethoxam	0.080 ($\pm$ 0.0051)	0.010 ($\pm$ 0.013)	0.052 ($\pm$ 0.0001)	0.061 ($\pm$ 0.013)
ETU	< LOQ	< LOQ	< LOQ	< LOQ
Mélamine	< LOQ	< LOQ	< LOQ	< LOQ
Variety 2: Pancha				
Pesticides	Raw	Flesh	Peels	Raw (calculated)
	Mean values ($\pm$ SD)	Mean values ($\pm$ SD)	Mean values ($\pm$ SD)	Mean values ($\pm$ SD)
Acetamiprid	0.86 ($\pm$ 0.0016)	0.065 ($\pm$ 0.025)	0.59 ($\pm$ 0.077)	0.653 ($\pm$ 0.10)
Carbendazim	0.022 ($\pm$ 0.0052)	0.0063 ($\pm$ 0.0002)	0.024 ($\pm$ 0.0019)	0.030 ($\pm$ 0.0021)
Cyromazine	1.50 ($\pm$ 0.15)	0.49 ($\pm$ 0.050)	0.59 ($\pm$ 0.067)	1.086 ($\pm$ 0.12)
Imazalil	0.20 ($\pm$ 0.015)	0.013 ($\pm$ 0.0004)	0.12 ($\pm$ 0.037)	0.13 ($\pm$ 0.037)
Maneb	10.17 ($\pm$ 0.31)	0.50 ($\pm$ 0.017)	6.46 ($\pm$ 1.21)	6.97 ($\pm$ 1.23)
Thiamethoxam	0.067 ($\pm$ 0.0012)	0.019 ($\pm$ 0.0095)	0.051 ($\pm$ 0.0066)	0.070 ($\pm$ 0.016)
ETU	< LOQ	< LOQ	< LOQ	< LOQ
Mélamine	< LOQ	< LOQ	< LOQ	< LOQ

Table 6: Mean values ($\pm$ SD, n = 2) with mass balance of processing factors (PF) for peeling step for six pesticides in two varieties of melons.

Pesticides	PF peeling ($\pm$ SD)	
	Mohican	Pancha
Acetamiprid	0.049 ($\pm$ 0.0038)	0.075 ($\pm$ 0.030)
Carbendazim	0.20 ($\pm$ 0.062)	0.29 ($\pm$ 0.072)
Cyromazine	0.20 ($\pm$ 0.020)	0.33 ($\pm$ 0.046)
Imazalil	0.058 ($\pm$ 0.0091)	0.063 ($\pm$ 0.0051)
Maneb	0.048 ($\pm$ 0.038)	0.050 ($\pm$ 0.0023)
Thiamethoxam	0.12 ($\pm$ 0.16)	0.29 ($\pm$ 0.14)