

# On the Complexity of two Problems on Orientations of Mixed Graphs

Guillaume Fertin, Hamed Mohamed-Babou, Irena Rusu

► **To cite this version:**

Guillaume Fertin, Hamed Mohamed-Babou, Irena Rusu. On the Complexity of two Problems on Orientations of Mixed Graphs. In Proc. 5èmes Journées Ouvertes Biologie Informatique Mathématiques (JOBIM 2012), Jul 2012, Rennes, France. pp.161-170, 2012. <hal-00826863>

**HAL Id: hal-00826863**

**<https://hal.archives-ouvertes.fr/hal-00826863>**

Submitted on 28 May 2013

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# On the Complexity of two Problems on Orientations of Mixed Graphs

Guillaume FERTIN, Hafedh MOHAMED-BABOU and Irena RUSU

LINA, UMR 6241 CNRS, 2 rue de la Houssinière BP 92208, Nantes, 44322, Cedex 03, France  
prenom.nom@univ-nantes.fr

**Abstract** *Interactions between biomolecules within the cell can be modeled by biological networks, i.e. graphs whose vertices are the biomolecules (proteins, genes, metabolites etc.) and whose edges represent their functional relationships. Depending on their nature, the interactions can be undirected (e.g. protein-protein interactions, PPIs) or directed (e.g. protein-DNA interactions, PDIs). A physical network is a network formed by both PPIs and PDIs, and is thus modeled by a mixed graph. External cellular events are transmitted into the nucleus via cascades of activation/deactivation of proteins, that correspond to paths (called signaling pathways) in the physical network from a source protein (cause) to a target protein (effect). There exists experimental methods to identify the cause-effect pairs, but such methods do not provide the signaling pathways. A key challenge is to infer such pathways based on the cause-effect informations. In terms of graph theory, this problem, called MAXIMUM GRAPH ORIENTATION (MGO), is defined as follows: given a mixed graph  $G$  and a set  $\mathcal{P}$  of source-target pairs, find an orientation of  $G$  that replaces each (undirected) edge by a single (directed) arc in such a way that there exists a directed path, from  $s$  to  $t$ , for a maximum number of pairs  $(s, t) \in \mathcal{P}$ . In this work, we consider a variant of MGO, called S-GO, in which we ask whether all the pairs in  $\mathcal{P}$  can be connected by a directed path. We also introduce a minimization problem, called MIN-DB-GO, in which all the pairs in  $\mathcal{P}$  must be connected by a directed path, while we allow some edges of  $G$  to be doubly oriented (i.e. replaced by two arcs in opposite directions). We investigate the complexity of S-GO and MIN-DB-GO by considering some restrictions on the input instances (such as the maximum degree of  $G$  or the cardinality of  $\mathcal{P}$ ). We provide several polynomial-time algorithms, hardness and inapproximability results that together give an extensive picture of tractable and intractable instances for both problems.*

**Keywords** Biological networks, computational biology, graph orientation, NP-completeness, APX-hardness.

## 1 Introduction

A *physical network* [16] is a biological network formed by protein-protein interactions (PPIs) and protein-DNA interactions (PDIs). While PPIs are undirected [5], PDIs are directed from the transcription factors to their target genes [8]. Thus, a physical network can be modeled by a mixed graph whose vertices are proteins, and edges (resp. arcs) are PPIs (resp. PDIs). Such a network is helpful to understand the processes that occur between a cell and its external environment, notably when an external stimulus is to be propagated into the nucleus. Indeed, this propagation is realized via cascades of activation/deactivation of proteins, and these cascades correspond to paths - in the physical network - from a source protein (cause) to a target protein (effect) [16]. The cause-effect pairs can be identified experimentally, for instance by the measure of transcription changes in response to a gene knock-out [16]. However, experimental methods do not provide the paths going from the source to the target proteins. A key problem in biology is to infer these paths by combining causal information on cellular events [10]. This question leads to the well-studied MAXIMUM GRAPH ORIENTATION problem (MGO) [3,4,6,10,14]: given a mixed graph  $G$  and a set  $\mathcal{P}$  of source-target (cause-effect) pairs of vertices, replace each (undirected) edge  $(u, v)$  by a single (directed) arc (either  $uv$  or  $vu$ ), so that in the new graph, there exists a directed path for a maximum number of source-target pairs. In this work, we focus on a variant of MGO, called S-GO, in which we ask whether *all* the pairs in  $\mathcal{P}$  can be connected by a directed path [1,7]. We also introduce a minimization problem, called MIN-DB-GO, in which we allow some edges of  $G$  to be *doubly*

*oriented* (i.e. replaced by two arcs in opposite directions), in such a way that all pairs of  $\mathcal{P}$  are satisfied (i.e., can be connected by a directed path). In the context of biology, a doubly oriented edge reflects the presence of a reversible reaction. Furthermore, in a dynamic biological system, most reactions tend to be irreversible [9]. For this reason, MIN-DB-GO asks that the number of doubly oriented edges be minimized.

## 2 Problem Formulation

All along this paper,  $G = (V, E, A)$  denotes a mixed graph without loops and with simple edges and arcs, where  $V(G)$  (resp.  $E(G)$ ,  $A(G)$ ) is the vertex set (resp. edge set, arc set) of  $G$ . The *underlying* graph of  $G$ , denoted  $G^*$ , is defined as follows:  $V(G^*) = V(G)$  and  $E(G^*) = E(G) \cup \{(u, v) : uv \in A(G)\}$ . Finally,  $\Delta(G^*)$  is the maximum degree over all vertices in  $G^*$ .

A *path*  $P$  in  $G = (V, E, A)$  from vertex  $v_1$  to vertex  $v_m$  is a sequence  $P = v_1, v_2, \dots, v_{m-1}, v_m$  of vertices  $v_i \in V$  such that for all  $1 \leq i \leq m-1$ ,  $(v_i, v_{i+1}) \in E$  or  $v_i v_{i+1} \in A$ . A *cycle*  $C$  in  $G$  is a path  $v_1, v_2, \dots, v_{m-1}, v_m$  such that  $v_1 = v_m$ . A *circuit* in  $G$  is a special case of cycle  $v_1, v_2, \dots, v_{m-1}, v_1$  where  $v_i v_{i+1} \in A$  for all  $1 \leq i \leq m-1$ . A *Mixed Acyclic Graph* [4] (or *MAG*) is a mixed graph that contains no cycle (and therefore no circuit).

An *orientation*  $G'$  of  $G$  is a directed graph  $G'$  obtained from  $G$  by replacing each edge  $(u, v) \in E$  by an arc  $uv$ , or an arc  $vu$ , or by  $uv$  and  $vu$  simultaneously. An edge  $(u, v)$  replaced by both arcs  $uv$  and  $vu$  is called a *doubly oriented* edge. Any orientation  $G'$  of  $G$  that contains no doubly oriented edge will be called a *simple orientation*. A pair of vertices  $(u, v) \in V \times V$  is said to be *satisfied* by the orientation  $G'$  of  $G$  if there is a (directed) path from  $u$  to  $v$  in  $G'$ . Let  $P = v_1, v_2, \dots, v_{m-1}, v_m$  be a path in  $G$ . In the following, we will often write *the orientation of  $P$  from  $v_1$  towards  $v_m$*  to refer to the orientation that replaces every edge of the form  $(v_i, v_{i+1})$ ,  $1 \leq i \leq m-1$ , by the arc  $v_i v_{i+1}$ .

DEFINITION 2.1. [1] Let  $G = (V, E, A)$  be a mixed graph and let  $\mathcal{P} \subseteq V \times V$  be a set of source-target pairs of vertices. The graph  $G$  is said to be  $\mathcal{P}$ -connected if for all  $(u, v) \in \mathcal{P}$ , there is a path in  $G$  from  $u$  to  $v$ .

DEFINITION 2.2. [1] Let  $G = (V, E, A)$  be a mixed graph and let  $\mathcal{P} \subseteq V \times V$  s.t.  $G$  is  $\mathcal{P}$ -connected. A  $\mathcal{P}$ -orientation  $G'$  of  $G$  is a simple orientation of  $G$  that satisfies all pairs in  $\mathcal{P}$ .

We call S-GO the problem of deciding whether a graph  $G$  admits a  $\mathcal{P}$ -orientation.

S-GO [1,7]

**Instance :** A mixed graph  $G = (V, E, A)$  and  $\mathcal{P} \subseteq V \times V$  s.t.  $G$  is  $\mathcal{P}$ -connected.

**Question:** Does  $G$  admit a  $\mathcal{P}$ -orientation ?

Analogously to a  $\mathcal{P}$ -orientation, we define a  $(\mathcal{P}, k)$ -DB-orientation as follows.

DEFINITION 2.3. Let  $G = (V, E, A)$  be a mixed graph and let  $\mathcal{P} \subseteq V \times V$  s.t.  $G$  is  $\mathcal{P}$ -connected. Let  $k \geq 0$  be an integer. A  $(\mathcal{P}, k)$ -DB-orientation  $G'$  of  $G$  satisfies the two following conditions: (i)  $G'$  is an orientation of  $G$  satisfying all the pairs in  $\mathcal{P}$  and (ii)  $G'$  contains exactly  $k$  doubly oriented edges.

We are now able to formulate the problem MIN-DB-GO (an illustration of such a problem is provided in Fig. 1).

MIN-DB-GO

**Instance :** A mixed graph  $G = (V, E, A)$ ,  $\mathcal{P} \subseteq V \times V$  s.t.  $G$  is  $\mathcal{P}$ -connected.

**Question:** Find a  $(\mathcal{P}, k)$ -DB-orientation of  $G$  that minimizes  $k$ .

In Section 3, we show that for both problems, we can always assume, without loss of generality, that  $G$  is a Mixed Acyclic Graph (MAG). In Section 4, we give some complexity results for S-GO. We study the complexity of MIN-DB-GO in Section 5. Section 6 is the conclusion, together with several open questions.


### 3 Reduction to Mixed Acyclic Graphs

It has been shown in [14] that starting with any instance  $(G_1, \mathcal{P}_1)$  of problem MGO (defined in Section 1), one can construct an equivalent instance  $(G_2, \mathcal{P}_2)$  s.t.  $G_2$  is a MAG.

PROPERTY 1. [14] Let  $G_1 = (V_1, E_1, A_1)$  be a mixed graph and let  $\mathcal{P}_1 \subseteq V_1 \times V_1$ . One can construct a MAG  $G_2 = (V_2, E_2, A_2)$  and a set  $\mathcal{P}_2 \subseteq V_2 \times V_2$  with  $|\mathcal{P}_2| = |\mathcal{P}_1|$  s.t. for every integer  $k \geq 0$ , there exists a simple orientation of  $G_1$  satisfying  $k$  pairs in  $\mathcal{P}_1$  if and only if there exists a simple orientation of  $G_2$  satisfying  $k$  pairs in  $\mathcal{P}_2$ .

Applying Property 1 with  $k = |\mathcal{P}_1|$ , we obtain that  $G_1$  admits a  $\mathcal{P}_1$ -orientation if and only if  $G_2$  admits a  $\mathcal{P}_2$ -orientation. Hence, in the S-GO problem, without loss of generality we can always consider the input mixed graph to be a MAG. We will show that this property is also valid for the MIN-DB-GO problem. For this, we first show in Property 2 that, for any mixed graph  $G$ , we can find a orientation  $G'$  such that the edges of any cycle  $C$  in  $G$  become arcs of a circuit  $C'$  in  $G'$ . We then show in Property 4 that the vertices of any circuit in  $G'$  can be contracted in a single vertex without changing the nature of the problem. This proof closely follows the one in [14].

PROPERTY 2 (ORIENTATION OF CYCLES). Let  $G = (V, E, A)$  be a mixed graph and let  $\mathcal{P} \subseteq V \times V$ . Let  $C$  be a cycle in  $G$ . There exists an optimal solution  $G''$  for MIN-DB-GO, with inputs  $G$  and  $\mathcal{P}$ , in which  $C$  becomes a circuit in  $G''$ .



**Figure 1.** (a)  $(G, \mathcal{P})$  is an instance of MIN-DB-GO with  $\mathcal{P} = \{(12, 17), (18, 19)\}$ ,  $C$  is a cycle in  $G$  with vertex set  $\{1, 2, \dots, 11\}$  (b)  $G'$  is an optimal  $(\mathcal{P}, 1)$ -DB-orientation of  $G$  in which the orientation  $C'$  of  $C$  is not a circuit (c)  $G''$  is an optimal  $(\mathcal{P}, 1)$ -DB-orientation of  $G$  in which the orientation  $C'$  of  $C$  is a circuit. Gray vertices in  $G'$  (resp.  $G''$ ) induce a path satisfying the pair  $(12, 17)$ .

*Proof.* First, since  $G$  is  $\mathcal{P}$ -connected, there must exist an optimal solution  $G'$  for MIN-DB-GO with inputs  $G$  and  $\mathcal{P}$ . Let  $G''$  be an orientation of  $G$  s.t. (i)  $C$  becomes a circuit  $C''$  and (ii) the edges in  $E(G) \setminus E(C)$  are oriented similarly as in  $G'$  (see Fig. 1 for an illustration). We now show that  $G''$  is also an optimal solution for MIN-DB-GO with inputs  $G$  and  $\mathcal{P}$ . Let  $(u, v) \in \mathcal{P}$ . If  $u, v \in V(C)$  then obviously the pair  $(u, v)$  is satisfied in  $G''$ . If  $u \notin V(C)$  or  $v \notin V(C)$ , then let us consider a path  $P' = a_1, a_2, \dots, a_m$  in  $G'$ , from  $u = a_1$  to  $v = a_m$ , that satisfies the pair  $(u, v)$ . Let  $x = \min \{i : a_i \in V(C)\}$  and let  $y = \max \{i : a_i \in V(C)\}$ . Then the pair  $(u, v)$  is satisfied in  $G''$  by the path formed by (1) the path in  $G''$  induced by the vertex set of the subpath of  $P'$  going from  $a_1$  to  $a_x$ , (2) the subpath of  $C''$  going from  $a_x$  to  $a_y$  and (3) the path of  $G''$  induced by the vertex set of the subpath of  $P'$  going from  $a_y$  to  $a_m$  (see for example Fig. 1, in which  $a_x = 3$  and  $a_y = 1$ ). Let  $E_{DB}$  denote the set of doubly oriented edges in  $G'$ . The set of doubly oriented edges in  $G''$  is  $E_{DB} \setminus E(C)$ . However, by minimality of  $|E_{DB}|$ , we know that  $E_{DB} \cap E(C) = \emptyset$ , and consequently  $G''$  is also an optimal solution for MIN-DB-GO.  $\square$

In the following, we say that two instances  $(G_1, \mathcal{P}_1)$  and  $(G_2, \mathcal{P}_2)$  of MIN-DB-GO are equivalent if and only if for every integer  $k \geq 0$ , there exists a  $(\mathcal{P}_1, k)$ -DB-orientation of  $G_1$  if and only if there exists a  $(\mathcal{P}_2, k)$ -DB-orientation of  $G_2$ .

Let  $G = (V, E, A)$  be a mixed graph and let  $\mathcal{P} \subseteq V \times V$ . Let  $G_1 = (V_1, E_1, A_1)$  be the mixed graph obtained from  $G$  by orienting, iteratively, each cycle into a circuit. According to Property 2,  $(G, \mathcal{P})$  and  $(G_1, \mathcal{P})$  are equivalent. Let also  $\mathcal{P}_1$  denote the set obtained from  $\mathcal{P}$  by removing each pair  $(u, v) \in \mathcal{P}$  s.t. there is a

directed path in  $G_1$  from  $u$  to  $v$ . In that case, the instance  $(G_1, \mathcal{P}_1)$  of MIN-DB-GO obtained from  $(G, \mathcal{P})$  will be called a *reduced instance*. Clearly,  $(G_1, \mathcal{P}_1)$  and  $(G_1, \mathcal{P})$  are equivalent, and thus the following property holds.

**PROPERTY 3 (REDUCED INSTANCES).** *Let  $(G_1, \mathcal{P}_1)$  be a reduced instance of MIN-DB-GO obtained from instance  $(G, \mathcal{P})$ . Then  $(G, \mathcal{P})$  and  $(G_1, \mathcal{P}_1)$  are equivalent.*

**PROPERTY 4 (CONTRACTION OF CIRCUITS).** *Let  $(G_1, \mathcal{P}_1)$  be a reduced instance of MIN-DB-GO, and let  $C'$  be a circuit in  $G_1$ . Let  $(G_2, \mathcal{P}_2)$  be the instance of MIN-DB-GO defined as follows: (i)  $\mathcal{P}_2 = \mathcal{P}_1$  and (ii)  $G_2$  is the graph obtained from  $G_1$  by contracting the vertices of  $C'$  in a single vertex. Then,  $(G_1, \mathcal{P}_1)$  and  $(G_2, \mathcal{P}_2)$  are equivalent.*

*Proof.* The graph  $G_2 = (V_2, E_2, A_2)$  is defined as follows:  $V_2 = (V_1 \setminus V(C')) \cup \{x_0\}$ ,  $E_2 = (E_1 \setminus \{(u, v) : u \in V(C') \text{ or } v \in V(C')\}) \cup \{(u, x_0) : u \notin V(C') \text{ and } \exists v \in V(C') \text{ s.t. } (u, v) \in E_1\}$ ,  $A_2 = (A_1 \setminus \{uv : u \in V(C') \text{ or } v \in V(C')\}) \cup \{ux_0 : u \notin V(C') \text{ and } \exists v \in V(C') \text{ s.t. } uv \in A_1\} \cup \{x_0v : v \notin V(C') \text{ and } \exists u \in V(C') \text{ s.t. } uv \in A_1\}$ . In other words,  $G_2$  is obtained from  $G_1$  by contracting the circuit  $C'$  in a single vertex  $x_0$ . Obviously,  $|E_1| = |E_2|$ .

Let  $\mathcal{P}_2 = \mathcal{P}_1$ . Now, let us show that the two instances  $(G_1, \mathcal{P}_1)$  and  $(G_2, \mathcal{P}_2)$  are equivalent. Let  $G'_1 = (V_1, A'_1)$  be a  $(\mathcal{P}_1, k)$ -DB-orientation of  $G_1$ . We construct an orientation  $G'_2$  of  $G_2$  as follows. Let  $(u, v) \in E_2$ . If  $u \neq x_0$  and  $v \neq x_0$ , then  $(u, v)$  is oriented in  $G'_2$  similarly as in  $G'_1$ . If  $u = x_0$ , then there is a vertex  $w \in V(C')$  s.t.  $(w, v) \in E_1$ . If  $uw \in A'_1$  (resp.  $vw \in A'_1$ ) we replace in  $G_2$  the edge  $(x_0, v)$  by the arc  $x_0v$  (resp.  $vx_0$ ). The case  $v = x_0$  is similar. Let  $(u, v) \in \mathcal{P}_1$  and let  $P'_1 = a_1, a_2, \dots, a_m$  be a directed path in  $G'_1$  from  $u = a_1$  to  $v = a_m$  satisfying the pair  $(u, v)$ . Let  $x = \min \{i : a_i \in V(C')\}$  and let  $y = \max \{i : a_i \in V(C')\}$ . Then the pair  $(u, v)$  is satisfied in  $G'_2$  by the path formed by (1) the path in  $G'_2$  induced by the vertex set of the subpath of  $P'_1$  going from  $a_1$  to  $a_{x-1}$ , (2) the vertex  $x_0$  (3) the path of  $G'_2$  induced by the vertex set of the subpath of  $P'_1$  going from  $a_{y+1}$  to  $a_m$ . Obviously,  $G'_2$  is a  $(\mathcal{P}_2, k)$ -DB-orientation of  $G_2$ . Reciprocally, starting with a  $(\mathcal{P}_2, k)$ -DB-orientation of  $G_2$ , by the same way, one can construct a  $(\mathcal{P}_1, k)$ -DB-orientation of  $G_1$ . Hence, the property follows.  $\square$

Now, using the previous properties, we are able to show that in the MIN-DB-GO we may, without loss of generality, assume that the input mixed graph is a MAG.

**PROPERTY 5 (REDUCTION TO MAG).** *Let  $(G, \mathcal{P})$  be an instance of the MIN-DB-GO problem. Then there exists an equivalent instance  $(G_M, \mathcal{P}_M)$  of MIN-DB-GO s.t.  $G_M$  is a MAG.*

*Proof.* We construct the graph  $G_M$  and the set  $\mathcal{P}_M$  by applying the following process:

1. Construct the reduced instance  $(G_1, \mathcal{P}_1)$  obtained from  $(G, \mathcal{P})$
2. Construct the graph  $G_2$ , obtained by contracting, in  $G_1$ , every circuit into a single vertex, and let  $\mathcal{P}_2 = \mathcal{P}_1$
3. If  $G_2$  is a MAG then set  $G_M = G_2$  and  $\mathcal{P}_M = \mathcal{P}_2$ . Otherwise, set  $G = G_2$  and  $\mathcal{P} = \mathcal{P}_2$ , and return to step 1.

Properties 3 and 4 ensure that  $(G_M, \mathcal{P}_M)$  is equivalent to  $(G, \mathcal{P})$ , which proves the property.  $\square$

Therefore, we will always assume, in the remaining of the paper, that for any instance  $(G, \mathcal{P})$  of MIN-DB-GO (resp. S-GO),  $G = (V, E, A)$  is MAG and  $G$  is  $\mathcal{P}$ -connected. Note that we can also assume that  $G^*$  to be connected. Otherwise, we can consider separately each graph  $G_1, G_2, \dots, G_r$  induced, in  $G$ , by the vertices of the connected components of  $G^*$ .

Let  $G = (V, E, A)$  be a MAG and  $\mathcal{P} = \{(s_i, t_i) \in V \times V : 1 \leq i \leq m\}$  be a set of pairs of vertices. For each  $i$ ,  $1 \leq i \leq m$ , we note by  $n_i$  the number of distinct paths in  $G$  from  $s_i$  to  $t_i$ . All along the paper, the integer  $B$  is the following value:  $B = \max\{n_i : 1 \leq i \leq m\}$ .

In the next sections, we study the complexity of S-GO (Section 4) and MIN-DB-GO (Section 5), by considering different constraints on the three following parameters:  $\Delta(G^*)$ ,  $B$  and  $|\mathcal{P}|$ .

Remark that when  $B = 1$  we can easily solve the S-GO and the MIN-DB-GO problems. Indeed,  $G$  is  $\mathcal{P}$ -connected and  $G^*$  is connected, thus if in addition we have  $B = 1$ , then for each pair  $(s, t) \in \mathcal{P}$  there is a unique path  $P_i$  in  $G$  from  $s_i$  to  $t_i$ . Consequently, in order to satisfy the pair  $(s_i, t_i)$  we must orient  $P_i$  from  $s_i$  towards  $t_i$ . Then, we orient each remaining edge  $(u, v)$ , in  $G$ , in a unique arbitrary direction. Obviously, in this orientation we create a minimum number of doubly oriented edges, and thus MIN-DB-GO is optimally solved. If there is no doubly oriented edge at the end of the process, then we obtain a  $\mathcal{P}$ -orientation of  $G$ . Otherwise,  $G$  has no  $\mathcal{P}$ -orientation. The case  $\Delta(G^*) = 1$  is obvious.

	$\Delta(G^*) = 2$	$\Delta(G^*) = 3$		
		$B = 2$	$B = 3$	$B$ is unbounded
S-GO	P [Cor. 1]	P [Th. 1]	NPC [Thm. 3]	NPC [Th. 3]
MIN-DB-GO	Open	APX-h [Th. 9]	NPC [Th. 6], Non-approx. [Th. 7], APX-h [Th. 9]	NPC [Th. 6], Non-approx. [Th. 7], APX-h and W[1]-h [Th. 8]

**Table 1.** Complexity of S-GO and MIN-DB-GO when  $G$  is a MAG and  $G^*$  is a bounded degree graph. Recall that  $B = \max\{n_i, 1 \leq i \leq |\mathcal{P}|\}$ , where  $n_i$  is the number of distinct paths in  $G$  from  $s_i$  to  $t_i$ . Note also that the result provided in Theorem 1 remains valid even when  $\Delta(G^*)$  is unbounded.

	$ \mathcal{P}  \leq 2$	$ \mathcal{P}  \geq 3$ (and $ \mathcal{P}  = \mathcal{O}(1)$ )	
		$B = \mathcal{O}(1)$	$B$ is unbounded
S-GO	P [Arkin et al. [1]]	P [Th. 2]	Open
MIN-DB-GO	P [Th. 4]	P [Th. 5]	Open

**Table 2.** Complexity of S-GO and MIN-DB-GO when  $G$  is a MAG and  $|\mathcal{P}|$  is a constant.

The complexity results, when  $B \geq 2$  and  $\Delta(G^*) \geq 2$ , are summarized in Table 1 and Table 2. Interestingly, Table 1 shows that parameter  $B$  defines the border between easy ( $B = 2$ ) and difficult ( $B = 3$ ) instances of S-GO, even when  $G^*$  is of small degree ( $\Delta(G^*) = 3$ ). Unlike S-GO, the problem MIN-DB-GO is difficult even when  $B = 2$ . In Table 2, we show the complexity of both problems when  $|\mathcal{P}|$  is a constant. Due to space constraints, some proofs are omitted in this paper.

## 4 Complexity of the S-GO problem

It has been shown that the S-GO problem is polynomial-time solvable on undirected graphs [7] and NP-complete on general MAGs [1]. In this section, we investigate the complexity of the S-GO problem for MAGs with bounded  $\Delta(G^*)$  and/or bounded  $B$  (see Table 1), and for bounded  $|\mathcal{P}|$  (see Table 2).

### 4.1 Easy cases

**THEOREM 1.** *The S-GO problem is polynomial-time solvable when  $G$  is a MAG and  $B = 2$ .*

*Proof.* For each pair  $(s_i, t_i) \in \mathcal{P}$  there are in  $G$  at most two paths from  $s_i$  to  $t_i$  and such paths can be computed in polynomial-time. If for a pair  $(s_i, t_i) \in \mathcal{P}$ , there is only one path from  $s_i$  to  $t_i$ , then we orient it from  $s_i$  towards  $t_i$  and we remove the pair  $(s_i, t_i)$  from the set  $\mathcal{P}$ . We continue this process until (1)  $G$  is no longer  $\mathcal{P}$ -connected or (2)  $\mathcal{P} = \emptyset$  or (3) for each pair  $(s_i, t_i) \in \mathcal{P}$  there are exactly two paths from  $s_i$  to  $t_i$ . The first case implies that  $G$  has no  $\mathcal{P}$ -orientation. In the second case we arbitrarily orient each edge, in the resulting graph, in a unique direction to obtain a  $\mathcal{P}$ -orientation. Finally, in the last case we reduce to an instance of the S-GO problem in which there are, in  $G$ , exactly two paths from  $s_i$  to  $t_i$  for all  $(s_i, t_i) \in \mathcal{P}$ .

We note by  $X_{i1}$  and  $X_{i2}$  the two paths, in  $G$ , from  $s_i$  to  $t_i$ . Given  $i, j \in \{1, 2, \dots, |\mathcal{P}|\}$ ,  $i \neq j$ , and  $a, b \in \{1, 2\}$ , we say that the two paths  $X_{ia}$  and  $X_{jb}$  are in *conflict* if orienting  $X_{ia}$  from  $s_i$  towards  $t_i$  and  $X_{jb}$  from  $s_j$  towards  $t_j$ , creates a doubly oriented edge. Now, we construct an instance  $(\mathcal{X}, \mathcal{C})$  of the problem 2-SAT as follows. Let  $\mathcal{X} = \{x_{i1}, x_{i2} : 1 \leq i \leq |\mathcal{P}|\}$  be the variable set. For all  $i \in \{1, 2, \dots, |\mathcal{P}|\}$ , we add the clause  $c_i = (x_{i1} \vee x_{i2})$ . For all  $i, j \in \{1, 2, \dots, |\mathcal{P}|\}$ ,  $i \neq j$ , and  $a, b \in \{1, 2\}$ , we add the clause

$(\overline{x_{ia}} \vee \overline{x_{jb}})$ , if paths  $X_{ia}$  and  $X_{jb}$  are in conflict. Let us show that there is an assignment of the variables in  $\mathcal{X}$  that satisfies all clauses in  $\mathcal{C}$  if and only if  $G$  has a  $\mathcal{P}$ -orientation. Indeed, consider a truth assignment of clauses in  $\mathcal{C}$  and let  $x_{ih_i}$ ,  $1 \leq h_i \leq 2$ , be a true literal of clause  $c_i$ ,  $1 \leq i \leq |\mathcal{P}|$ . We orient in  $G$  the path  $X_{ih_i}$ , from  $s_i$  towards  $t_i$ , for all  $i$ ,  $1 \leq i \leq |\mathcal{P}|$ . This orientation cannot create any doubly oriented edges. Otherwise, there are  $i, j \in \{1, 2, \dots, |\mathcal{P}|\}$ ,  $i \neq j$  such that the paths  $X_{ih_i}$  and  $X_{jh_j}$  are in conflict, implying that the clause  $(\overline{x_{ih_i}} \vee \overline{x_{jh_j}})$  is unsatisfied. To finish the orientation of  $G$ , we orient arbitrarily the remaining edges in  $G$  without creating any doubly oriented edge.

Now, let us show the reverse implication. We consider the set  $\{Y_{1h_1}, Y_{2h_2}, \dots, Y_{|\mathcal{P}|h_{|\mathcal{P}|}}\}$  s.t.  $Y_{ih_i}$  is a directed path from  $s_i$  to  $t_i$ , in a  $\mathcal{P}$ -orientation of  $G$ . Each path  $Y_{ih_i}$  is the orientation (from the source towards the target vertex) of a mixed path  $X_{ih_i} = G[V(Y_{ih_i})]$ ,  $h_i \in \{1, 2\}$ , for all  $1 \leq i \leq |\mathcal{P}|$ . We set to *true* the variable set  $\{x_{ih_i} : 1 \leq i \leq |\mathcal{P}|\}$  and we set to *false* the remaining variables. Obviously, this assignment satisfies the clause set  $\{c_i : 1 \leq i \leq |\mathcal{P}|\}$ . By contradiction, assume now that some clause  $(\overline{x_{ia}} \vee \overline{x_{jb}})$  is not satisfied. Then  $x_{ia} = \text{true}$  and  $x_{jb} = \text{true}$ . Consequently, in the resulting  $\mathcal{P}$ -orientation of  $G$ , the path  $X_{ia}$  (resp.  $X_{jb}$ ) is oriented from  $s_i$  towards  $t_i$  (resp. from  $s_j$  towards  $t_j$ ). A contradiction, because a  $\mathcal{P}$ -orientation cannot use, simultaneously, two paths that are in conflict. As the problem 2-SAT is polynomial-time solvable [2], we deduce that one can solve in polynomial-time the S-GO problem when graph  $G$  is a MAG s.t. there are in  $G$  at most two paths from  $s_i$  to  $t_i$ , for all  $(s_i, t_i) \in \mathcal{P}$ .  $\square$

**COROLLARY 1.** *The S-GO problem is polynomial-time solvable when  $G$  is a MAG and  $\Delta(G^*) = 2$ .*

*Proof.* The graph  $G^*$  is connected. Thus when  $\Delta(G^*) = 2$ , the graph  $G^*$  must be a path or a cycle, and consequently  $B \leq 2$ . If  $B = 1$  the S-GO problem is trivial. If  $B = 2$ , we deduce from the previous result (Theorem 1) that the S-GO problem is polynomial-time solvable.  $\square$

Now, we show that the S-GO problem is polynomial-time solvable when both parameters  $B$  and  $|\mathcal{P}|$  are bounded.

**THEOREM 2.** *The S-GO problem is polynomial-time solvable when  $G$  is a MAG,  $|\mathcal{P}| = \mathcal{O}(1)$  and  $B = \mathcal{O}(1)$ .*

## 4.2 Difficult cases


We showed in Theorem 1 that the S-GO problem is easy when  $B \leq 2$ . However, in the following theorem, we show that when  $B = 3$  the problem S-GO becomes difficult.

**THEOREM 3.** *The S-GO problem is NP-complete even when the graph  $G$  is a MAG,  $\Delta(G^*) = 3$  and  $B = 3$ .*

*Proof.* Arkin et al. [1] provided an NP-completeness proof of the S-GO problem on general MAGs. Their proof is based on a reduction for the Satisfiability problem (SAT). Here, we modify the mixed graph  $G$  constructed from their reduction to ensure that  $\Delta(G^*) = 3$ . For this, we reduce to our problem, the problem 3-SAT in which each variable appears at most in four clauses [15]. Let  $(\mathcal{C}_m, \mathcal{V}_n)$  be an instance of 3-SAT s.t.  $\mathcal{C}_m = \{c_1, \dots, c_m\}$  is a set of clauses and  $\mathcal{V}_n = \{x_1, \dots, x_n\}$  is a variable set, and for all  $j$ ,  $1 \leq j \leq n$ , the variable  $x_j$  satisfies the following condition: (1)  $x_j$  and  $\overline{x_j}$  appear at most in four clauses. In addition, one can require second condition: (2) for each variable  $x_j$ , there is at least one clause that contains  $x_j$  and at least one clause that contains  $\overline{x_j}$ . Otherwise, w.l.o.g the variable  $x_j$  can be fixed to *true* or *false*. Now, let us construct an instance  $(G, \mathcal{P})$  of the S-GO problem. For each clause  $c_i$ , we create two vertices  $s_i$  and  $t_i$ ,  $1 \leq i \leq m$ . For each variable  $x_j$ , we create these 15 vertices:  $\{u_j, v_j\} \cup \{a_{jk}, b_{jk}, a'_{jk}, b'_{jk}\}_{1 \leq k \leq 3}$ . Then, we add an edge  $(u_j, v_j)$  and the four following directed paths:  $a_{j1}a_{j2}a_{j3}u_j$ ,  $v_j b_{j3}b_{j2}b_{j1}$ ,  $a'_{j1}a'_{j2}a'_{j3}v_j$  and finally  $u_j b'_{j3}b'_{j2}b'_{j1}$ , for all  $1 \leq j \leq n$ . For each variable  $x_j$ , there are  $k_j$  clauses containing  $x_j$  and  $k'_j$  clauses containing  $\overline{x_j}$  s.t.  $1 \leq k_j \leq 3$ ,  $1 \leq k'_j \leq 3$  and  $k_j + k'_j \leq 4$ . Let  $\{c_{i_1}, c_{i_2}, \dots, c_{i_{k_j}}\}$  (resp.  $\{c'_{i'_1}, c'_{i'_2}, \dots, c'_{i'_{k'_j}}\}$ ) be the set of clauses that contain  $x_j$  (resp.  $\overline{x_j}$ ). We add an arc  $s_{i_\alpha} a_{j\alpha}$  and an arc  $b_{j\alpha} t_{i_\alpha}$ , for all  $\alpha \in \{1, 2, \dots, k_j\}$ . Also, we add an arc  $s_{i'_\beta} a'_{j\beta}$  and an arc  $b'_{j\beta} t_{i'_\beta}$ , for all  $\beta \in \{1, 2, \dots, k'_j\}$ . To finish our construction, we set  $\mathcal{P} = \{(s_i, t_i), 1 \leq i \leq m\}$ . An example of construction is illustrated in Fig. 2. According to conditions (1) and (2), one can easily show that  $\Delta(G^*) = 3$ . In addition, for each pair  $(s_i, t_i)$  there are exactly three paths in  $G$  from  $s_i$  to  $t_i$ , because each clause in  $\mathcal{C}_m$  contains exactly three literals. Thus  $B = 3$ .

We claim that there is an assignment satisfying all the clauses in  $\mathcal{C}_m$  if and only if there exists a  $\mathcal{P}$ -orientation of  $G$ . Indeed, consider an assignment satisfying all the clauses in  $\mathcal{C}_m$ , similarly to the proof presented in [1], if  $x_j = \text{true}$  (resp.  $x_j = \text{false}$ ) then we orient the edge  $(u_j, v_j)$  from  $u_j$  to  $v_j$  (resp. from  $v_j$  to  $u_j$ ). Let  $l_i$  be a true literal of clause  $c_i$ . Then, there is a variable  $x_j$  s.t.  $l_i = x_j$  or  $l_i = \overline{x_j}$ . If  $l_i = x_j$  (resp.  $l_i = \overline{x_j}$ ) then there is an integer  $k_i$ ,  $1 \leq k_i \leq 3$ , such that  $s_i a_{j k_i}, b_{j k_i} t_i \in A(G)$  (resp.  $s_i a'_{j k_i}, b'_{j k_i} t_i \in A(G)$ ). Thus, the pair  $(s_i, t_i)$  is satisfied by the path  $s_i a_{j k_i} a_{j(k_i+1)} \dots u_j v_j b_{j 3} \dots b_{j k_i} t_i$  (resp.  $s_i a'_{j k_i} a'_{j(k_i+1)} \dots v_j u_j b'_{j 3} \dots b_{j k_i} t_i$ ).

Now, let us prove the reverse implication. Given a  $\mathcal{P}$ -orientation  $G'$  of  $G$ , we set the variable  $x_j$  to *true* (resp. to *false*) if the edge  $u_j v_j \in A(G')$  (resp.  $v_j u_j \in A(G')$ ). Let  $c_i$  be a clause in  $\mathcal{C}_m$ . Then the pair  $(s_i, t_i)$  is satisfied in by a directed path  $P$  in  $G'$ , from  $s_i$  to  $t_i$ , going through an arc  $u_j v_j$  or an arc  $v_j u_j$ . If  $P$  contains the arc  $u_j v_j$  then the clause  $c_i$  must contain the literal  $x_j$  and thus  $c_i$  is satisfied. If  $P$  contains the arc  $v_j u_j$  (consequently  $x_j = \text{false}$ ) thus the clause  $c_i$  must contain the literal  $\overline{x_j}$  and thus  $c_i$  is also satisfied.  $\square$



**Figure 2.** Construction of an instance  $(G, \mathcal{P})$  of the S-GO problem, from an instance of 3-SAT in which each variable appears at most in four clauses. Here, the variable set is  $\mathcal{V} = \{x_j, 1 \leq j \leq 6\}$  and the clause set is  $\mathcal{C} = \{c_i, 1 \leq i \leq 4\}$  s.t.  $c_1 = (x_1 \vee \overline{x_2} \vee x_3)$ ,  $c_2 = (x_1 \vee x_4 \vee x_5)$ ,  $c_3 = (x_1 \vee \overline{x_4} \vee \overline{x_6})$  and  $c_4 = (\overline{x_1} \vee \overline{x_5} \vee x_6)$ . The set of pairs of vertices is  $\mathcal{P} = \{(s_i, t_i), 1 \leq i \leq 4\}$ . In this figure, we show only the subgraph corresponding to variable  $x_1$ .

## 5 Complexity of MIN-DB-GO

Let DB-GRAPHORIENTATION denote the decision version of the minimization problem MIN-DB-GO. The S-GO problem, investigated in the previous section (Section 4), is a particular case of the problem DB-GRAPHORIENTATION when no doubly oriented edge is allowed. Hence, each  $\mathcal{P}$ -orientation of  $G$  is a solution of MIN-DB-GO. However, if there is no  $\mathcal{P}$ -orientation of  $G$ , then we conclude just that at least one edge must be doubly oriented in a solution of MIN-DB-GO, but in general that gives no information about the number of edges to be doubly oriented to solve the MIN-DB-GO problem.

In this section, we study the complexity of MIN-DB-GO when the input graph is a MAG (see Table 1 and Table 2). As in the previous section, we suppose that  $G$  is a  $\mathcal{P}$ -connected MAG.

### 5.1 Easy cases

We first show that, similarly to the S-GO problem, the MIN-DB-GO problem is also polynomial-time solvable for general MAGs when  $|\mathcal{P}| \leq 2$ .

**THEOREM 4.** *The MIN-DB-GO problem is polynomial-time solvable when  $G$  is a MAG and  $|\mathcal{P}| \leq 2$ .*

*Proof.* The case  $|\mathcal{P}| = 1$  is obvious. Let  $G = (V, E, A)$  be a MAG and  $\mathcal{P} = \{(s_1, t_1), (s_2, t_2)\} \in V \times V$ .

A  $\mathcal{P}$ -essential edge is an edge  $e = (u, v) \in E$ , s.t. if we orient  $e$  from  $u$  to  $v$  or from  $v$  to  $u$ , the graph  $G$  is no longer  $\mathcal{P}$ -connected. One can compute the  $\mathcal{P}$ -essential edges in polynomial-time [1].

Let  $E_{ess}$  (resp.  $E_{min}$ ) be the set of  $\mathcal{P}$ -essential edges (resp. the set of doubly oriented edges in a solution of the MIN-DB-GO problem).

We show that  $E_{min} = E_{ess}$ . Let  $e = (u, v) \in E_{ess}$ . If we orient  $e$  in a unique direction (from  $u$  to  $v$  or from  $v$  to  $u$ ) then, by definition of  $\mathcal{P}$ -essential edges, there is an integer  $i$ ,  $1 \leq i \leq 2$ , s.t. there is no path in  $G$  from  $s_i$  to  $t_i$ . Thus, whatever the orientation of edges in  $E - \{e\}$ , if  $e$  is not doubly oriented, then the pair  $(s_i, t_i)$  would not be satisfied. Hence, we must doubly orient each edge  $e \in E_{ess}$ , which implies that  $E_{ess} \subseteq E_{min}$ .



Conversely, let  $G' = (V, E', A')$  denote the mixed graph obtained from  $G$  after replacing each  $\mathcal{P}$ -essential edge  $(u, v)$  by the arcs  $uv$  and  $vu$ , i.e.,  $V(G') = V(G)$ ,  $E(G') = E(G) \setminus E_{ess}$  and  $A' = A \cup \{uv, vu : (u, v) \in E_{ess}\}$ . The graph  $G'$  does not contain any  $\mathcal{P}$ -essential edge, then  $G'$  has a  $\mathcal{P}$ -orientation  $G''$  [1]. Thus,  $G''$  is an orientation of  $G$  that satisfies all the pairs in  $\mathcal{P}$  and creates  $|E_{ess}|$  doubly oriented edges, which implies that  $|E_{ess}| \geq |E_{min}|$ . Since we have already shown that  $E_{ess} \subseteq E_{min}$ , we conclude that  $E_{min} = E_{ess}$ .

Now, to solve the MIN-DB-GO problem when  $|\mathcal{P}| = 2$ , we apply the following process.

1. Compute the  $\mathcal{P}$ -essential edges of  $G$  using the polynomial-time algorithm presented in [1];
2. Construct a mixed graph  $G'$  by replacing each  $\mathcal{P}$ -essential edge  $(u, v)$  by two arcs  $uv$  and  $vu$ ;
3. Apply the polynomial-time algorithm presented in [1] to compute a  $\mathcal{P}$ -orientation of  $G'$ .

□

As in the S-GO problem, the MIN-DB-GO problem is polynomial-time solvable when both parameters  $|\mathcal{P}|$  and  $|B|$  are bounded.

**THEOREM 5.** *The MIN-DB-GO problem is polynomial-time solvable when  $G$  is a MAG,  $|\mathcal{P}| = \mathcal{O}(1)$  and  $B = \mathcal{O}(1)$ .*

## 5.2 Difficult cases

Let DB-GO denote the decision version of the MIN-DB-GO problem. We show in the next result that the DB-GO problem is NP-complete even when  $\Delta(G^*) = 3$ .

**THEOREM 6.** *The problem DB-GO is NP-complete when  $G$  is a MAG and  $|\mathcal{P}|$  is unbounded even when  $\Delta(G^*) = 3$  and  $B = 3$ .*

In the remaining of this section, we study the approximability and the parameterized complexity of the MIN-DB-GO problem.

**THEOREM 7.** *Unless  $P = NP$ , the MIN-DB-GO problem is non-approximable when the graph  $G$  is a MAG and  $|\mathcal{P}|$  is unbounded even when  $\Delta(G^*) = 3$  and  $B = 3$ .*


The MIN-DB-GO problem is also APX-hard and W[1]-hard when  $|\mathcal{P}|$  and  $B$  are unbounded even when  $\Delta(G^*) = 3$ .

**THEOREM 8.** *The MIN-DB-GO problem is APX-hard and W[1]-hard (parametrized by the number of doubly oriented edges) when  $G$  is a MAG and  $|\mathcal{P}|$  and  $B$  are unbounded, even when  $\Delta(G^*) = 3$ .*

*Proof.* We propose an L-reduction from the APX-hard problem MINIMUM SET COVER [12,13]: given a ground set  $\mathcal{X} = \{X_1, \dots, X_n\}$ , and a collection of sets  $\mathcal{C} = \{S_1, \dots, S_m\}$  s.t.  $S_i \in 2^{\mathcal{X}}$ , for all  $1 \leq i \leq m$ , the goal is to find a minimum set cover  $\mathcal{C}'$ , i.e., a set  $\mathcal{C}' \subseteq \mathcal{C}$  s.t.  $\mathcal{C} = \bigcup_{S_i \in \mathcal{C}'} S_i$  and  $|\mathcal{C}'|$  is minimum.

We note by  $\alpha_j$ ,  $1 \leq j \leq n$ , the number of the sets containing  $X_j$ . Let us construct an instance  $(G, \mathcal{P})$  of the MIN-DB-GO problem. For each  $X_j \in \mathcal{X}$ , we create the vertex set  $\{x_j\} \cup \{x_j^k, 1 \leq k \leq \alpha_j\}$ , then we create the directed path  $x_j^1 x_j^2 \dots x_j^{\alpha_j} x_j$ . For each  $S_i$ , we add the vertex set  $\{s_i, s'_i\} \cup \{s_i^j, 1 \leq j \leq |S_i|\}$ , and we add an edge  $(s'_i, s_i)$  and a directed path  $s_i s_i^1 s_i^2 \dots s_i^{|S_i|}$ .

Let  $X_l \in \mathcal{X}$  be the  $j$ -th element in a set  $S_i$ , for each  $j$ ,  $1 \leq j \leq |S_i|$ , we add an arc from  $s_i^j$  towards *only one* vertex from the set  $\{x_l^k, 1 \leq k \leq \alpha_l\}$ . Such a vertex is chosen in such a way that the indegree of each vertex  $x_l^k$  is at most two, for all  $1 \leq k \leq \alpha_l$ . To finish the construction of  $G$ , we add a vertex  $r_1$  connected, by two edges, to the vertices  $s'_1$  and  $s'_2$ . Then, we add a new vertex  $r_2$  connected, by two edges, to the vertices  $r_1$  and  $s'_3$ . We continue the creation of vertices  $r_i$  connected, by edges, to  $r_{i-1}$  and  $s'_{i+1}$ , for all  $3 \leq i \leq m-1$ . The set of pairs to satisfy is  $\mathcal{P} = \{(r_{m-1}, x_j), 1 \leq j \leq n\} \cup \{(s_i, s'_i), 1 \leq i \leq m\}$  with  $m = |\mathcal{C}|$ . An example of construction is illustrated in Fig. 3. The degree of each vertex  $r_i$  in  $G$  is at most three and also each vertex  $s_i^j$  is connected to exactly one vertex  $x_{l'}^{j'}$ . Thus, one can easily check that  $\Delta(G^*) = 3$ .



**Figure 3.** (a) Construction of an instance  $(G, \mathcal{P})$  of the S-GO problem from an instance of MINIMUM SET COVER problem. Here,  $\mathcal{X} = \{X_1, X_2, X_3, X_4, X_5, X_6\}$  and  $\mathcal{C} = \{S_1, S_2, S_3, S_4\}$  s.t.  $S_1 = \{X_1, X_3\}$ ,  $S_2 = \{X_2, X_3, X_5\}$ ,  $S_3 = \{X_1, X_4, X_6\}$  and  $S_4 = \{X_1, X_5\}$ . The set of pairs is  $\mathcal{P} = \{(s_1, s'_1), (s_2, s'_2), (s_3, s'_3), (s_4, s'_4), (r_3, x_1), (r_3, x_2), (r_3, x_3), (r_3, x_4), (r_3, x_5), (r_3, x_6)\}$ . (b) The graph  $G'$  is an orientation of  $G$  obtained from the set cover  $\mathcal{C}' = \{S_2, S_3\}$  and satisfying all the pairs in  $\mathcal{P}$ .

We claim that, for every integer  $k \geq 0$ , there is a set cover of  $\mathcal{C}$  of cardinality  $k$  if and only if there is a  $(\mathcal{P}, k)$ -DB-orientation of  $G$ .

$\Rightarrow$ : Given a set cover  $\{S_{i_1}, S_{i_2}, \dots, S_{i_k}\}$ , we doubly orient the edge  $(s_{i_j}, s'_{i_j})$ , for all  $1 \leq j \leq k$ . Then, we replace each edge  $(v_i, v'_i)$  by the arc  $s_i s'_i$ , for all  $i \in \{1, 2, \dots, m\} \setminus \{i_1, i_2, \dots, i_k\}$ . Finally, we orient the tree induced by the vertex set  $\{r_i, 1 \leq i < m\} \cup \{s'_i, 1 \leq i \leq m\}$ , to create a directed tree of root  $r_{m-1}$ .

$\Leftarrow$ : Let  $E_{DB}$  be the set of doubly oriented edges in a  $(\mathcal{P}, k)$ -DB-orientation. Let  $\mathcal{C}' = \{S_i : (s_i, s'_i) \in E_{DB}\}$ . We will show that the set  $\mathcal{C}'$  is a set cover of  $\mathcal{C}$ . Suppose that there is  $X_j \in \mathcal{X}$  s.t.  $X_j \notin S_i$  for all  $S_i \in \mathcal{C}'$ . Let  $\mathcal{C}_j$  denote the collection of the sets containing  $X_j$ , i.e.,  $\mathcal{C}_j = \{S \in \mathcal{C} : X_j \in S\}$ . The graph  $G$  is constructed in such a way that, to satisfy any pair  $(r_{m-1}, x_j)$ , we must replace an edge  $(s'_j, s_i)$  by the arc  $s'_j s_i$  s.t.  $S_i \in \mathcal{C}_j$ . On the other hand, we have to orient each edge  $(s'_j, s_i)$ , from  $s_i$  towards  $s'_j$ , to satisfy the pair  $(s_i, s'_j) \in \mathcal{P}$ . Then the edge  $(s'_j, s_i)$  must be doubly oriented, which implies that  $S_i \in \mathcal{C}'$ . This is a contradiction, because  $\mathcal{C}' \cap \mathcal{C}_j = \emptyset$ .

The above reduction is an L-reduction that preserves the parameter  $k$  (the cardinality of the set cover and the number of doubly oriented edges). Since the problem MINIMUM SET COVER is APX-hard and W[1]-hard when parametrized by  $k$  [12,13], then the MIN-DB-GO problem is also APX-hard and W[1]-hard when parametrized by the number of doubly oriented edges.  $\square$

Now let us show that, unlike the S-GO problem, the MIN-DB-GO problem remains difficult even when  $B = 2$ .

**THEOREM 9.** *The problem MIN-DB-GO is APX-hard when  $G$  is a MAG and  $|\mathcal{P}|$  is unbounded, even when  $B \in \{2, 3\}$  and  $\Delta(G^*) = 3$ .*

*Proof.* Again, we use the above reduction (proof of Theorem 8), but we consider the variant MINIMUM SET COVER- $K$  of the MINIMUM SET COVER problem in which each  $X_j \in \mathcal{X}$  appears in exactly  $K$  sets in  $\mathcal{C}$  s.t.  $K$  is a constant  $\geq 2$ . For each pair of vertices  $(s_i, s'_i) \in \mathcal{P}$  there is a unique path in  $G$ , from  $s_i$  to  $s'_i$  (that

is the edge  $(s_i, s'_i)$ , for all  $1 \leq i \leq m$ . In addition, the fact that each  $X_j \in \mathcal{X}$  appears in exactly  $K$  sets in  $\mathcal{C}$ , implies that for each pair  $(r_{m-1}, x_i)$  there are, in  $G$ ,  $K$  paths from  $r_{m-1}$  to  $s_i$ , for all  $1 \leq i \leq n$ . Thus  $B = K$  and also the graph  $G$  is constructed so that we have  $\Delta(G^*) = 3$ .

As the problem MINIMUM SET COVER- $K$  is APX-hard [11], then we conclude that MIN-DB-GO is APX-hard when  $G$  is a MAG s.t.  $\Delta(G^*) = 3$  and  $|\mathcal{P}|$  is unbounded, even when,  $B$  is a constant  $\geq 2$  (and thus when  $B \in \{2, 3\}$ ).  $\square$

## 6 Conclusion

In this work, we considered two problems that are concerned with the orientation of mixed graphs, both motivated, among others, by biological applications. We studied the complexity of both problems, and in particular we provided polynomial-time algorithms for some restricted instances, and several hardness and inapproximability results. However, some interesting problems remain open so far, such as the following ones:

- Explore the possibility of obtaining fixed-parameterized tractable (FPT) algorithms for MIN-DB-GO.
- Study the approximability of MIN-DB-GO on specific graph classes.
- Study the complexity of S-GO and MIN-DB-GO when  $|\mathcal{P}| \geq 3$  is a constant.

## References

- [1] E. M. Arkin and R. Hassin. A note on orientations of mixed graphs. *Discrete Applied Mathematics*, 116:271–278, 2002.
- [2] B. Aspvall, M. F. Plass, and R. E. Tarjan. A linear-time algorithm for testing the truth of certain quantified boolean formulas. *Inf. Process. Lett.*, 8:121–123, 1979.
- [3] B. Dorn, F. Huffner, D. Kruger, R. Niedermeier, and J. Uhlmann. Exploiting bounded signal flow for graph orientation based on cause–effect pairs. *Algorithms for Molecular Biology*, 6:21, 2011.
- [4] M. Elberfeld, D. Segev, C. R. Davidson, D. Silverbush, and R. Sharan. Approximation algorithms for orienting mixed graphs. *In Proc. CPM*, 6661:416–428, 2011.
- [5] S. Fields and O. Song. A novel genetic system to detect protein-protein interactions. *Nature*, 340(6230):245–246, July 1989.
- [6] I. Gamzu, D. Segev, and R. Sharan. Improved orientations of physical networks. *In Proc. WABI*, 6293:215–225, 2010.
- [7] R. Hassin and N. Megiddo. On orientations and shortest paths. *Linear Algebra and its Applications*, 114:589–602, 1989.
- [8] T. I. Lee, N. J. Rinaldi, F. Robert, D. T. Odom, Z. Bar-Joseph, G. K. Gerber, N. M. Hannett, C. T. Harbison, C. M. Thompson, I. Simon, J. Zeitlinger, E. G. Jennings, H. L. Murray, D. B. Gordon, B. Ren, J. J. Wyrick, J.-B. Tagne, T. L. Volkert, E. Fraenkel, D. K. Gifford, and R. A. Young. Transcriptional regulatory networks in *saccharomyces cerevisiae*. *Science*, 298(5594):799–804, Oct. 2002.
- [9] U. Lucia. Probability, ergodicity, irreversibility and dynamical systems. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 464:1089–1104, 2008.
- [10] A. Medvedovsky, V. Bafna, U. Zwick, and R. Sharan. An algorithm for orienting graphs based on cause-effect pairs and its applications to orienting protein networks. *In Proc. WABI*, 5251:222–232, 2008.
- [11] C. Papadimitriou and M. Yannakakis. Optimization, approximation, and complexity classes. *In Proc. STOC*, 43:229–234, 1988.
- [12] A. Paz and S. Moran. Non deterministic polynomial optimization problems and their approximations. *Theor. Comput. Sci.*, 15:251–277, 1981.
- [13] R. Raz and S. Safra. A sub-constant error-probability low-degree test, and a sub-constant error-probability PCP characterization of NP. *In Proc. STOC*, pages 475–484, 1997.
- [14] D. Silverbush, M. Elberfeld, and R. Sharan. Optimally orienting physical networks. *Journal of Computational Biology*, 18:1437–1448, 2011.
- [15] C. A. Tovey. A simplified NP-complete satisfiability problem. *Discrete Applied Mathematics*, 8:85–89, 1984.
- [16] C.-H. Yeang, T. Ideker, and T. Jaakkola. Physical network models. *Journal of Computational Biology*, 11:243–262, 2004.