Iodine(III)-Mediated Umpolung of Bromide Salts for the Ethoxybromination of Enamides.

Sophie Nocquet-Thibault, Pascal Retailleau, Kevin Cariou, Robert H Dodd

► To cite this version:

Sophie Nocquet-Thibault, Pascal Retailleau, Kevin Cariou, Robert H Dodd. Iodine(III)-Mediated Umpolung of Bromide Salts for the Ethoxybromination of Enamides.. *Organic Letters*, 2013, 15 (8), pp.1842-5. 10.1021/ol400453b . hal-00827755

HAL Id: hal-00827755

<https://hal.science/hal-00827755>

Submitted on 25 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Iodine(III)-Mediated Umpolung of Bromide Salts for the Ethoxybromination of Enamides

Sophie Nocquet-Thibault, Pascal Retailleau,[†] Kevin Cariou and Robert H. Dodd*

Centre de Recherche de Gif-sur-Yvette, Institut de Chimie des Substances Naturelles, CNRS, 91198 Gif-sur-Yvette Cedex, France
robert.dodd@icsn.cnrs-gif.fr

Received Date (will be automatically inserted after manuscript is accepted)

ABSTRACT

Using (diacetoxyiodo)benzene in conjunction with simple bromide salts in ethanol allows the regioselective ethoxybromination of a wide range of enamides, thus yielding highly versatile α -bromo hemiaminals, which can then be engaged in a broad array of transformations.

In recent years, polyvalent iodine compounds have been at the center of tremendous activity and research developments.¹ Of special interest are iodine(III) compounds such as Koser's reagent² and (diacetoxyiodo)benzene (PIDA).^{1,2} One of the particular characteristics of these compounds is that they behave in a similar fashion to transition-metal complexes.³ Indeed, around the central iodine atom, ligands can be exchanged and then transferred through a formal reductive elimination. In this context, halides can be used as ligands, in which case an umpolung of the salt⁴ can occur to give birth to electrophilic halogen species. Several combinations of alkali or metal halide salts and hypervalent iodine(III) derivatives have previously been

studied and reacted with substrates such as carbonyls,⁵ electron-rich arenes,⁶ olefins,⁷ and even aliphatic CH bonds.⁸

We have recently explored the reactivity of enamides **1** toward nitrenoids generated from iodine(III) species, namely iminoiodanes, in the presence of a nucleophile.^{9a} This set of reaction conditions allowed the oxidative difunctionalization of the enamide to give the corresponding amino hemiaminal in a completely regio- and stereoselective manner (Scheme 1, eq 1).

¹ (a) Brown, M.; Farid, U.; Wirth, T. *Synlett* **2013**, 424. (b) Zhdankin, V. V.; Stang, P. J. *Chem. Rev.* **2008**, *108*, 5299. (c) Wirth, T. *Angew. Chem., Int. Ed.* **2005**, *44*, 3656. (d) Zhdankin, V. V.; Stang, P. J. *Chem. Rev.* **2002**, *102*, 2523. (e) Varvoglis, A. *Tetrahedron* **1997**, *53*, 1179. (f) Stang, P. J.; Zhdankin, V. V. *Chem. Rev.* **1996**, *96*, 1123.

² Koser, G. F. *Aldrichimica Acta* **2001**, *34*, 89.

³ Moriarty R. M. *J. Org. Chem.* **2005**, *70*, 2893.

⁴ For another strategy using sulfoxides, see: Klimczyk, S.; Huang, X.; Farès, C.; Maulide, N. *Org. Biomol. Chem.* **2012**, *10*, 4327.

⁵ (a) Liu W. L.; Chen, C.; Zhang, Q.; Zhu Z.-B. *Beilstein J. Org. Chem.* **2012**, *8*, 344. (b) Lee, J. C.; Park, J. Y.; Yoon, S. Y.; Bae, Y. H.; Lee, S. J. *Tetrahedron Lett.* **2004**, *45*, 191. (c) Akula, R.; Galligan, M. Ibrahim, H. *Chem. Commun.* **2009**, *45*, 6991. (d) Coats, S. J.; Wasserman, H. H. *Tetrahedron Lett.* **1995**, *36*, 7735.

⁶ (a) Tu, D.; Ma, L.; Tong, X.; Deng, X.; Xia, C. *Org. Lett.* **2012**, *14*, 4830. (b) Hamamoto, H.; Hattori, S.; Takemaru, K.; Miki Y. *Synlett* **2011**, 1563. (c) Hamamoto, H.; Umemoto, H.; Umemoto, M.; Ohta, C.; Dohshita, M.; Miki Y. *Synlett* **2010**, 2593. (d) Braddock, D. C.; Cansell, G.; Hermitage, S. A. *Synlett* **2004**, 461.

⁷ (a) Fabry, D. C.; Stodulski, M.; Hoerner, S.; Gulder, T. *Chem-Eur. J.* **2012**, *18*, 10834. (b) Pandit, P.; Gayen, K. S.; Khamarui, S.; Chatterjee, N.; Maiti, D. K. *Chem. Commun.* **2011**, *47*, 6933. (c) Yu, L.; Ren, L.; Yi, R.; Wu, Y.; Chen, T.; Guo, R. *Synlett* **2011**, 579. (d) Sanjaya, S.; Chiba, S. *Tetrahedron* **2011**, *67*, 590. (e) Katrun, P.; Chiampanichayakul, S.; Korworapan, K.; Pohmakotr, M.; Reutrakul, V.; Jaipetch, T.; Kuhakarn C. *Eur. J. Org. Chem.* **2010**, 5633. (f) Wu, T.; Yin, G.; Liu, G. *J. Am. Chem. Soc.* **2009**, *131*, 16355. (g) Fan, R.; Wen, F.; Qin, L.; Pu, D.; Wang, B. *Tetrahedron Lett.* **2007**, *48*, 7444. (h) Liu, H.; Tan C.-H. *Tetrahedron Lett.* **2007**, *48*, 8220. (i) Emmanuvel, L.; Shaikh, T. M. A.; Sudala, A. *Org. Lett.* **2005**, *7*, 5071. (j) Braddock, D. C.; Cansell, G.; Hermitage, S. A. *Synlett* **2004**, 461. (k) Hashem, M. A.; Jung, A.; Ries, M.; Kirschning, A. *Synlett* **1998**, 195.

⁸ (a) McMurtrey, K. B.; Racowski, J. M.; Sanford, M. S. *Org. Lett.* **2012**, *14*, 4094. (b) Dohi, T.; Takenaga, N.; Goto, A.; Maruyama, A.; Kita, Y. *Org. Lett.* **2007**, *9*, 3129.

⁹ (a) Nakanishi, M.; Minard, C.; Cariou, K.; Retailleau, P.; Dodd, R. H. *Org. Lett.* **2011**, *13*, 5792. (b) For a related reaction using Rh catalysts, see: Gigant, N.; Dequize, G.; Retailleau, P.; Gillaizeau, I.; Dauban, P. *Chem-Eur. J.* **2012**, *18*, 90.

Extrapolating from this, we reasoned that subjecting the same type of enamide to a combination of a halide source and iodobenzene diacetate,¹⁰ we could in turn

Scheme 1. Initial Endeavor

obtain an α -bromo hemiaminal, which would constitute a highly versatile synthon for further transformations (Scheme 1, eq 2).¹¹

Moreover, the low toxicity of iodine(III) derivatives¹ and their potential for asymmetric transformations¹² led us to consider that this strategy would provide a valuable alternative to the use of anodic oxidation of bromides,¹³ toxic bromine,¹⁴ or NBS,¹⁵ which furthermore, have so far been mostly limited to cyclic enamide derivatives.¹⁶

We started by reacting tosyl enamide **1a** with an excess of iodobenzene diacetate in the presence of a 2-fold amount of lithium bromide in dry acetonitrile. The lithium salt was chosen because, as an alkali halide, it is cheap and widely available and also exhibits a better solubility than other alkali salts in organic solvents. Under these conditions, there is no external nucleophile other than the acetate, which is liberated by reaction of the iodobenzene diacetate and is regioselectively incorporated in the final product (Table 1, entry 1). However, this product could only be isolated in moderate yield (as a mixture of two diastereoisomers in a 4:1 ratio) because of the low stability of the acetoxy hemiaminal moiety.

Table 1. Optimization of Reaction Conditions

entry	X	solvent	Temp (°C)	t (min)	misc	R, yield ^a (%)
1	1.8	MeCN	rt	70	no MS	Ac, 59
2	1.4	MeCN	rt	10	MS	Ac, 72
3	1.4	MeCN	rt	10	5 equiv of EtOH	Et, 85 ^b
4	1.4	MeCN	rt	10	10 equiv of EtOH	Et, 95 ^c
5	1.4	EtOH	rt	10	dr 80/20	Et, 100 ^d
6	1.4	EtOH	0	20	dr 83/17	Et, 91
7	1.1	EtOH	0	90	°	Et, 86
8	1.2	EtOH	0	25	dr 87/13	Et, 97
9	1.2	EtOH	-20	20 h	dr 88/12	Et, 100 ^d

^a Isolated yields after chromatography on silica gel unless otherwise mentioned. ^b Conversion determined by ¹H NMR, along with 15% of R = Ac. ^c Conversion determined by ¹H NMR, along with 5% of R = Ac. ^d Conversion determined by ¹H NMR; ^e 10 mol % of PIDA and 20 mol % of LiBr were added after 60 min.

Adding molecular sieves to the reaction mixture allowed shorter reaction times while improving the yield to some extent (entry 2), but the stability of the adduct remained an issue. Assuming this problem was due to the acetoxy group, ethanol was added as an external nucleophile but competitive addition of the acetate was still observed (entries 3 and 4). Finally, complete chemoselectivity in favor of ethoxy products was obtained by using ethanol as the solvent (entry 5). Lowering the temperature to 0 °C increased the diastereoselectivity, and the optimal amount of reagents was found to be 1.2 equiv of PIDA and 2.4 equiv of lithium bromide, which efficiently provided 97% of the desired product **3a** with a 87:13 diastereomeric ratio (entries 6–8). Diminishing the temperature further to -20 °C only increased the reaction time without any real improvement of the diastereoselectivity (entry 9). The relative configuration of the major diastereoisomer was

¹⁰ For metal-free reactions involving enamides and (diacyloxyiodo)benzenes, see: (a) Zhao, F.; Liu, X.; Qi, R.; Zhang-Negreie, D.; Huang, J.; Du, Y.; Zhao, K. *J. Org. Chem.* **2011**, *76*, 10338. (b) Yu, W.; Du, Y.; Zhao, K. *Org. Lett.* **2009**, *11*, 2417. (c) Huang, X.; Shao, N.; Huryk, R.; Palani, A.; Aslanian, R.; Seidel-Dugan, C. *Org. Lett.* **2009**, *11*, 867. (d) Shao, N.; Huang, X.; Palani, A.; Aslanian, R.; Buevich, A.; Piwinski, J.; Huryk, R.; Seidel-Dugan, C. *Synthesis* **2009**, 2855. (e) Huang, X.; Shao, N.; Palani, A.; Aslanian, R.; Buevich, A. *Org. Lett.* **2007**, *9*, 2597. (f) Huang, X.; Shao, N.; Palani, A.; Aslanian, R. *Tetrahedron Lett.* **2007**, *48*, 1967.

¹¹ An illustration of this versatility, for cyclic α -iodo-hemiaminals, can be found in: (a) Norton Matos, M. R. P.; Afonso, C. A.; Batey, R. A. *Tetrahedron* **2001**, *61*, 1221. (b) Norton Matos, M. R. P.; Afonso, C. A. M.; McGarvey, T.; Lee, P.; Batey, R. A. *Tetrahedron Lett.* **1999**, *40*, 9189.

¹² (a) Farid, U.; Wirth, T. In *Asymmetric Synthesis II*; Christmann, M., Bräse, S., Eds.; Wiley-VCH: Weinheim, 2012; pp 197 – 203. (b) Liang,

H.; Ciufolini M. A. *Angew. Chem. Int. Ed.* **2011**, *50*, 11849. (c) Uyanik, M.; Okamoto, H.; Yasui, T.; Ishihara, K. *Science* **2010**, *328*, 1376.

¹³ (a) Shono, T.; Matsumura, Y.; Onomura, O.; Ogaki, M.; Kanazawa, T. *J. Org. Chem.* **1987**, *52*, 536. (b) Shono, T.; Matsumura, Y.; Ogaki, M.; Onomura, O. *Chem. Lett.* **1987**, 1447.

¹⁴ (a) Kelleher, S.; Quesne, P.-Y.; Evans P. *Beilstein J. Org. Chem.* **2009**, *10*, 3762/bjoc.5.69. (b) Sugiura, M.; Asai, K.; Hamada, Y.; Hatano, K.; Kurono, Y.; Suezawa, H.; Hirota, M. *Chem. Pharm. Bull.* **1997**, *45*, 928. (c) Shono, T.; Matsumura, Y.; Onomura, O.; Yanada, Y. *Tetrahedron Lett.* **1987**, *28*, 4073.

¹⁵ (a) Le Corre, L.; Kizirian, J.-C.; Levraud, C.; Boucher, J.-L.; Bonnet, V.; Dhimane, H. *Org. Biomol. Chem.* **2008**, *6*, 3388. (b) Levraud, C.; Calvet-Vitale, S.; Bertho, G.; Dhimane, H. *Eur. J. Org. Chem.* **2008**, 1901.

¹⁶ For acyclic enamines, see: (a) Alix, A.; Lalli, C.; Retailleau, P.; Masson G. *J. Am. Chem. Soc.* **2012**, *134*, 10389. (b) Jokić, M.; Škarić, V. *Tetrahedron Lett.* **1994**, *35*, 2937.

attributed from the X-ray data obtained for ethoxy-**3a** (Figure 1) which was further confirmed from the X-ray data obtained for compound **3p** (see Table 2).

Figure 1. ORTEP drawings for OEt-**3a** and **3p**.

First, varying the sulfonamide protecting group had only a small influence on the course of the reaction (Table 2, **1a–d**) though the tosyl group was the most satisfying in terms of yield and diastereoselectivity. The slight variations in the diastereoisomeric ratio are consistent with steric effects, while the more electron-withdrawing nosyl group only caused a moderate loss in efficiency. Moreover, comparison between the ^1H NMR spectra of the various α -bromo hemiaminals obtained and **3a** showed that the major diastereoisomer was the same in all cases. Replacement of the phenyl by a *p*-methoxyphenyl, a benzyl, an allyl, or a methyl group (**1e–h**), led to incremental improvements of the dr, probably due to subtle variations of the steric hindrance. It is noteworthy that this reaction proved to be highly chemoselective, as neither the electron-rich anisyl⁶ (**1e**) nor the allyl⁷ (**1g**) moieties were affected in this transformation. On the other hand, switching from a sulfonamide to an amide or a carbamate proved highly detrimental. Indeed, when submitted to the reaction conditions *N*-Ac compounds only gave a complex mixture (**1i,j**). This phenomenon was slightly less pronounced for *N*-Boc substrates, in which case the formation of the desired product could be observed before it decomposed (**1k,l**). When differently functionalized substrates such as (*E*)-2-aminoacrylates were subjected to the same reaction conditions (**1m–p**), the dr were consistently above the 90/10 threshold (as observed for the crude material, the adducts were often isolated as a single diastereoisomer) and the yields remained excellent, with the major isomer corresponding to that obtained for (*E*)-2-aminostyrenes (as evidenced by the X-ray analysis of **3p**, Figure 1). In contrast, the *Z* isomers gave the desired product in lower yields and with almost no diastereoselectivity (**1q,r**). Finally, it should be noted that, despite the oxidative nature of the reaction conditions, free alcohols are tolerated and give the corresponding adducts with moderate yields but excellent diastereoselectivities (**1s,t**).

¹⁷ dr were calculated using ^1H NMR analysis of the crude material. The slightly higher values reported in the Supporting Information correspond to isolated products.

Table 2. Substrate Scope for the Ethoxybromination¹⁷

s.m.	R	PG	R'	<i>t</i> (min)	dr ^a	yield ^b (%)
1a	Ph	Ts	Ph (<i>E</i>)	25	87/13	97
1b	Ph	PhSO ₂	Ph (<i>E</i>)	30	83/17	88
1c	Ph	Ms	Ph (<i>E</i>)	40	76/24	89
1d	Ph	Ns	Ph (<i>E</i>)	25	92/8	81 ^c
1e	PMP	Ts	Ph (<i>E</i>)	60	77/23	71 ^d
1f	Bn	Ts	Ph (<i>E</i>)	20	83/17	88 ^c
1g	Allyl	Ts	Ph (<i>E</i>)	50	90/10	89
1h	Me	PhSO ₂	Ph (<i>E</i>)	20	92/8	89
1i	Ph	Ac	Ph (<i>E</i>)	50	N.D.	c.m.
1j	Bn	Ac	Ph (<i>E</i>)	40	N.D.	c.m.
1k	Ph	Boc	Ph (<i>E</i>)	30	79/21	49 ^e
1l	Bn	Boc	Ph (<i>E</i>)	45	71/29	N.D.
1m	PMP	Ts	CO ₂ Me (<i>E</i>)	120	92/8	79 ^e
1n	Bn	Ts	CO ₂ Me (<i>E</i>)	150	92/8	73 ^e
1o	Bn	Ns	CO ₂ Me (<i>E</i>)	90	92/8	55 ^e
1p	Allyl	Ts	CO ₂ Me (<i>E</i>)	40	91/9	74
1q	Bn	Ts	CO ₂ Me (<i>Z</i>)	60	52/48	52
1r	Allyl	Ts	CO ₂ Me (<i>Z</i>)	45	55/45	65
1s	Bn	Ts	CH ₂ OH (<i>E</i>)	30	92/8	60 ^{e,f}
1t	Allyl	Ts	CH ₂ OH (<i>E</i>)	90	92/8	46 ^f

^a Calculated by ^1H NMR analysis of the crude material. N.D. = not determined. ^b Isolated yields after chromatography on silica gel. c.m. = complex mixture. ^c Reaction run with 1.4 equiv. of PIDA and 2.8 equiv. of LiBr. ^d 10 mol % of PIDA and 20 mol % of LiBr were added after 30 min. ^e Isolated as a single diastereoisomer. ^f Reaction run with 1.2 equiv of PIDA and 4.5 equiv of LiBr on a 1 mmol scale.

The strong discrepancy observed between the reactivity of the *E* and *Z* isomers raised a few questions concerning the mechanism of this transformation. First, at this stage, the exact nature of the electrophile remains conjectural. In the presence of bromide ions a series of equilibria could take place between iodobenzene diacetate and iodobenzene dibromide **5** through the mixed species **4** (Scheme 2).^{5–8,18} For both in situ generated iodine(III)

¹⁸ For the isolation and characterization of such compounds, see: (a) Braddock, D. C.; Cansell, G.; Hermitage, S. A.; White, A. J. P. *Chem.*

reagents a reductive elimination can take place, producing acetyl hypobromite or bromine. Formally, all four species can be considered as Br^+ sources for the reaction.

We then assume that the reaction between the enamide **1** and the electrophilic bromonium species (whatever its exact nature) would stereospecifically lead to bridged bromonium **6**, which would be in equilibrium with iminium **7** (Scheme 3).

Scheme 2. Generation of the Electrophilic Species

Depending on the various stereoelectronic effects at play, one form might be favored over the other, thus influencing the final diastereoselective ratio as the addition of the external nucleophile (ethanol) would proceed through an $\text{S}_{\text{N}}2$ and/or $\text{S}_{\text{N}}1$ type mechanism. Indeed, (*Z*)-enamide would give rise to *cis*-**6** in which unfavorable 1,2 steric interaction would displace the equilibrium toward the iminium and therefore cause a drastic lowering of the diastereoselectivity (as was observed for substrates **1q,r**). Lastly, it should be pointed out that the X-ray data for the major stereoisomer (both for $\text{R}' = \text{Ph}$, **3a** and for $\text{R}' = \text{CO}_2\text{Me}$, **3p**) are consistent with an $\text{S}_{\text{N}}2$ -type opening of bromonium *trans*-**6a** by ethanol.

Scheme 3. Mechanistic Proposal

As already stated, these α -bromo hemiaminals are highly versatile synthons, and so we went on to probe their behavior in several transformations. First, using a Lewis acid such as boron trifluoride etherate allows the regeneration of iminium **7** (and putatively bromonium **6**) and the subsequent addition of soft nucleophiles such as triethylsilane and allyltrimethylsilane (Scheme 4). Both reduction and allylation proceed smoothly to give α -bromo

amines (**8** and **9**) in moderate to good yields (unoptimized) and as a single diastereomer for **9a** and **9b**. Diallylic compound **9b** could be elaborated further by Grubbs' second-generation catalyst-triggered ring-closing metathesis, which yielded the corresponding tetrahydropyridine **10** quantitatively. The bromo moiety could also be exploited. Simple treatment of the free alcohols **3s** and **3t** with sodium hydroxide cleanly afforded α -amino epoxides **11a** and **11b**. *N*-Allyl substrate **8b** could be cyclized under either palladium-catalyzed¹⁹ or radical-initiated conditions to give two types of substituted pyrrolidines **12** and **13**, respectively.

Scheme 4. Exploiting the Versatility of the Adducts^a

^a The relative configuration of compounds **9a**, **9b**, **10**, **11a**, and **11b**, isolated as single diastereoisomers, could not be ascertained.

In summary, using a combination of PIDA and lithium bromide in ethanol enabled the regioselective alkoxybromination of variously substituted enamide derivatives with excellent yield and selectivity. The resulting α -bromo hemiaminals could then be elaborated further using various strategies ranging from radical to transition-metal-catalyzed transformations. Efforts toward the development of an asymmetric variant of this transformation and its use for the synthesis of biologically relevant targets are currently underway in our laboratory.

Acknowledgment. We thank the Institut de Chimie des Substances Naturelles for financial support and fellowship for S. N.-T.. We also thank Dr. Géraldine Masson (ICSN) for helpful discussions.

Supporting Information Available. Experimental procedures and characterization of compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>).²⁰

† P. R. ran the X-ray crystallography experiments.
The authors declare no competing financial interest.