



Local times for functions with finite variation: two versions of Stieltjes change of variables formula

Jean Bertoin, Marc Yor

► To cite this version:

Jean Bertoin, Marc Yor. Local times for functions with finite variation: two versions of Stieltjes change of variables formula. 2013. hal-00835685v2

HAL Id: hal-00835685

<https://hal.science/hal-00835685v2>

Preprint submitted on 4 Jul 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Local times for functions with finite variation: two versions of Stieltjes change of variables formula

Jean Bertoin and Marc Yor

Abstract We introduce two natural notions for the occupation measure of a function V with finite variation. The first yields a signed measure, and the second a positive measure. By comparing two versions of the change-of-variables formula, we show that both measures are absolutely continuous with respect to Lebesgue measure. Occupation densities can be thought of as local times of V , and are described by a Meyer-Tanaka like formula.

Key words: Occupation measure, finite variation, local times.

1 Introduction

Although the setting of the present work is entirely deterministic and does not involve Probability Theory at all, its motivation comes from an important chapter of Stochastic Calculus (see, for instance, Section IV.7 in Protter [7]). Specifically, let $(X_t)_{t \geq 0}$ be a real-valued semimartingale, i.e. $X = M + V$ where M is a local martingale and V a process with finite variation, and the paths of X are right-continuous and possess left-limits (càdlàg) a.s. In turn the local martingale can be expressed as the sum $M = M^c + M^d$ of a continuous local martingale M^c and a purely discontinuous local martingale M^d . The occupation measure of X on a time interval $[0, t]$ is defined as

Jean Bertoin

Institut für Mathematik, Universität Zürich, Winterthurerstrasse 190, CH-8057 Zürich, Switzerland, e-mail: jean.bertoin@math.uzh.ch

Marc Yor

Institut Universitaire de France and Laboratoire de Probabilités et Modèles Aléatoires, UPMC, 4 Place Jussieu, 75252 Paris cedex 05, France e-mail: deaproba@proba.jussieu.fr

$$\int_0^t \mathbf{1}_A(X_s) d\langle M^c \rangle_s, \quad A \in \mathcal{B}(\mathbb{R}), \quad (1)$$

where $\langle M^c \rangle$ denotes the quadratic variation of M^c . A fundamental result in this area is that this occupation measure is a.s. absolutely continuous with respect to Lebesgue measure, i.e. the following occupation density formula holds:

$$\int_0^t \mathbf{1}_A(X_s) d\langle M^c \rangle_s = \int_A L_t^x dx.$$

The occupation densities $\{L_t^x : x \in \mathbb{R} \text{ and } t \geq 0\}$ are known as the local times of X and are described by the Meyer-Tanaka formula

$$\begin{aligned} & (X_t - x)^+ - (X_0 - x)^+ \\ &= \int_0^t \mathbf{1}_{\{X_{s-} > x\}} dX_s + \frac{1}{2} L_t^x + \sum_{0 < s \leq t} (\mathbf{1}_{\{X_{s-} \leq x\}} (X_s - x)^+ + \mathbf{1}_{\{X_{s-} > x\}} (X_s - x)^-). \end{aligned} \quad (2)$$

The purpose of this work is to point out that the above results for semimartingales have natural analogs in the deterministic world of functions with finite variation. The rest of this note is organized as follows. Our main result is stated in Section 2 and proven in Section 4. Section 3 is devoted to two versions of the Stieltjes change-of-variables formula which lie at the heart of the analysis.

2 Local times for functions with finite variation

We consider a càdlàg function $V : [0, \infty) \rightarrow \mathbb{R}$ with finite variation. There is the canonical decomposition of V as the sum of its continuous and discontinuous components,

$$V = V^c + V^d,$$

where

$$V^d(t) = \sum_{0 < s \leq t} \Delta V(s)$$

and the series is absolutely convergent. Plainly, the continuous part V^c has also finite variation; we write $V^c(dt)$ for the corresponding signed Stieltjes measure and $|V^c(dt)|$ for its total variation measure.

We then introduce for every $t > 0$ a signed measure $\theta_t(dx)$ and a positive measure $\vartheta_t(dx)$, which are defined respectively by

$$\theta_t(A) = \int_0^t \mathbf{1}_A(V(s)) V^c(ds) \quad (3)$$

and

$$\vartheta_t(A) = \int_0^t \mathbf{1}_A(V(s)) |V^c(ds)| \quad (4)$$

where $A \subseteq \mathbb{R}$ stands for a generic Borel set. We interpret $\theta_t(dx)$ and $\vartheta_t(dx)$ respectively as the signed and the absolute occupation measure of V on the time interval $[0, t]$.

Our goal is to show that the occupation measures $\theta_t(dx)$ and $\vartheta_t(dx)$ are both absolutely continuous with respect to the Lebesgue measure and to describe explicitly their densities. In this direction, we introduce the following definitions and notations.

For every $x \in \mathbb{R}$ and $t > 0$, we say that V increases through the level x at time t and then write $t \in \mathcal{I}(x)$ if :

- $V(t) = x$ and V is continuous at time t ,
- $V(s) - V(t)$ has the same sign as $s - t$ for all s in some neighborhood of t .

Similarly, we say that V decreases through the level x at time t and then write $t \in \mathcal{D}(x)$ if $-V$ increases through the level $-x$ at time t . We then define

$$\ell^x(t) = \text{Card}\{(0, t] \cap \mathcal{I}(x)\} - \text{Card}\{(0, t] \cap \mathcal{D}(x)\},$$

whenever the two quantities in the difference in the right-hand side are finite. For each such x , $t \rightarrow \ell^x(t)$ is a right-continuous function with integer values, which can only jump at times when V reaches x continuously. We also set

$$\lambda^x(t) = \text{Card}\{(0, t] \cap \mathcal{I}(x)\} + \text{Card}\{(0, t] \cap \mathcal{D}(x)\},$$

so that the point measure $\lambda^x(dt)$ coincides with the total variation measure $|\ell^x(dt)|$.

Our main result identifies $(\ell^x(t) : x \in \mathbb{R} \text{ and } t \geq 0)$ and $(\lambda^x(t) : x \in \mathbb{R} \text{ and } t \geq 0)$ respectively as the signed and absolute local times of V .

Theorem 1. *For every $t > 0$, there are the identities*

$$\theta_t(dx) = \ell^x(t)dx \quad \text{and} \quad \vartheta_t(dx) = \lambda^x(t)dx.$$

Theorem 1 is an immediate consequence of the chain rule when V is piecewise of class \mathcal{C}^1 (with possible jumps); however the general case requires a more delicate analysis. It is of course implicit in the statement that $\ell^x(t)$ and $\lambda^x(t)$ are well-defined for almost all $x \in \mathbb{R}$ and yield measurable functions in $L^1(dx)$, which is not *a priori* obvious.

It is interesting to point out that essentially the same result was established by Fitzsimmons and Port [5] in the special case when V is a Lévy process with finite variation and non-zero drift (then, when for instance the drift is positive, only increasing passages through a level may occur). In the same direction, Theorem 1 applies much more generally to sample paths of a large class of (semi) martingales with finite variation, which arise for instance by predictable compensation of a pure jump process with finite variation. Note that when M is a local martingale with finite variation, then its continuous component in the sense of Martingale Theory is always degenerate (i.e. identically zero), whereas its continuous component in the sense of functions with finite variation may be non-trivial. In that case, the occupa-

tion measure defined by (1) is degenerate, and then definitions (3) and (4) may be sounder.

We further stress that for semi-martingales with non-degenerate continuous martingale component $M^c \not\equiv 0$, the local time $t \rightarrow L_t^x$ defines an increasing process which is always continuous, whereas for functions with finite variation, $t \rightarrow \ell^x(t)$ and $t \rightarrow \lambda^x(t)$ are integer-valued step-functions which only jump at times when the function V crosses continuously the level x .

We also observe the following consequence of Theorem 1. Provided that V is not purely discontinuous, i.e. $V^c \not\equiv 0$, then the set of levels through which V increases or decreases has a positive Lebesgue measure. This contrasts sharply with the case of a typical Brownian path, or also, say, a sample path of a symmetric stable Lévy process with index $\alpha \in (1, 2)$, as then local times exist, but there is no level through which the sample path increases or decreases (cf. Dvoretzky *et al.* [4] and [1]). Of course, in those examples, sample paths have infinite variation.

3 Two versions of the change-of-variables formula

Our strategy for proving Theorem 1 is to compare two change-of-variables formulas. The first version is standard (see, for instance, Dellacherie and Meyer [3] on pages 168-171); a proof will be given for the sake of completeness.

Proposition 1. *Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a function of class \mathcal{C}^1 . Then for every $t > 0$, there is the identity*

$$f(V(t)) - f(V(0)) = \int_0^t f'(V(s))V^c(ds) + \sum_{0 < s \leq t} (f(V(s)) - f(V(s-))) .$$

Proof. With no loss of generality, we may assume that f' has compact support with $|f'(x)| \leq 1$ for all $x \in \mathbb{R}$. We fix $\varepsilon > 0$ arbitrarily small and may find a finite sequence of times $0 = t_0 < t_1 < \dots < t_n = t$ such that

$$\max_{i=1, \dots, n} \max_{t_{i-1} \leq u, v \leq t_i} |V^c(u) - V^c(v)| \leq \varepsilon \quad (5)$$

and

$$\sum_{\substack{0 < s < t \\ s \neq t_1, \dots, t_n}} |\Delta V(s)| \leq \varepsilon \quad (6)$$

where, as usual, $\Delta V(s) = V(s) - V(s-)$. We write

$$f(V(t)) - f(V(0)) = \sum_{i=1}^n (f(V(t_i-)) - f(V(t_{i-1}))) + \sum_{i=1}^n (f(V(t_i)) - f(V(t_i-))) .$$

On the one hand, by the triangle inequality, we have

$$\begin{aligned}
\left| \sum_{\substack{0 < s < t \\ s \neq t_1, \dots, t_n}} (f(V(s)) - f(V(s-))) \right| &\leq \sum_{\substack{0 < s < t \\ s \neq t_1, \dots, t_n}} |f(V(s)) - f(V(s-))| \\
&\leq \sum_{\substack{0 < s < t \\ s \neq t_1, \dots, t_n}} |\Delta V(s)|,
\end{aligned}$$

where in the last line, we applied the mean value theorem and the assumption that $|f'| \leq 1$. Using (6), we have therefore shown that

$$\left| \sum_{0 < s < t} (f(V(s)) - f(V(s-))) - \sum_{i=1}^n (f(V(t_i)) - f(V(t_{i-1}))) \right| \leq \varepsilon.$$

On the other hand, again from the mean value theorem, for each $i = 1, \dots, n$, there is some x_i between $V(t_{i-1})$ and $V(t_i)$ such that

$$f(V(t_i)) - f(V(t_{i-1})) = f'(x_i)(V^c(t_i) - V^c(t_{i-1})) + f'(x_i)(V^d(t_i) - V^d(t_{i-1})).$$

Since $|f'| \leq 1$, it follows readily from (6) and the triangle inequality that

$$\sum_{i=1}^n |f'(x_i)(V^d(t_i) - V^d(t_{i-1}))| \leq \varepsilon.$$

Further, since x_i lies between $V(t_{i-1})$ and $V(t_i)$, we deduce from (5) and (6) that

$$|x_i - V(s)| \leq 2\varepsilon \quad \text{for every } s \in [t_{i-1}, t_i].$$

It follows that

$$\left| \sum_{i=1}^n f'(x_i)(V^c(t_i) - V^c(t_{i-1})) - \int_0^t f'(V(s))V^c(ds) \right| \leq W^c(t) \sup_{|x-y| \leq 2\varepsilon} |f'(x) - f'(y)|,$$

where $W^c(t) < \infty$ stands for the total variation of V^c on $[0, t]$.

Putting the pieces together, we arrive at the inequality

$$\begin{aligned}
&\left| f(V(t)) - f(V(0)) - \int_0^t f'(V(s))V^c(ds) - \sum_{0 < s \leq t} (f(V(s)) - f(V(s-))) \right| \\
&\leq 2\varepsilon + W^c(t) \sup_{|x-y| \leq 2\varepsilon} |f'(x) - f'(y)|.
\end{aligned}$$

Because f' is continuous and has compact support, the upperbound tends to 0 when $\varepsilon \rightarrow 0+$, which establishes the change-of-variables formula. \square

In order to state a second version of the change-of-variables formula, we need first to introduce some further notions. We say that $x \in \mathbb{R}$ is a *simple level* for V if the set

$$\{t > 0 : V(t-) < x < V(t) \text{ or } V(t) < x < V(t-) \text{ or } V(t) = x\}$$

is discrete and if there is no jump of V that starts or ends at x , i.e.

$$\{t > 0 : \Delta V(t) \neq 0 \text{ and either } x = V(t) \text{ or } x = V(t-)\} = \emptyset.$$

Otherwise, we say that x is a *complex level* for V .

Lemma 1. *The set of complex levels for V has zero Lebesgue measure.*

Proof. The set of jump times of V is at most countable. Thus so is

$$\{x \in \mathbb{R} : x = V(t) \text{ or } x = V(t-) \text{ for some } t > 0 \text{ with } \Delta V(t) \neq 0\},$$

and *a fortiori* this set has zero Lebesgue measure.

Consider a sequence of partitions τ_n of $[0, \infty)$ with mesh tending to 0 and such that τ_{n+1} is thinner as τ_n . We write $0 = t_0^n < t_1^n < \dots$ for the elements of τ_n . We set for every $n, i \in \mathbb{N}$

$$J_i^n = [\inf_{t_i^n \leq s \leq t_{i+1}^n} V(s), \sup_{t_i^n \leq s \leq t_{i+1}^n} V(s)],$$

and write as usual $|J_i^n|$ for the length of this interval. If we write $W(t)$ for the total variation of V on $[0, t]$, then for every $t \in \tau_n$, there is the upperbound

$$\sum_{i: t_i^n < t} |J_i^n| \leq W(t).$$

Now introduce

$$N_x(t) = \text{Card}\{0 < s \leq t : V(s-) \leq x \leq V(s) \text{ or } V(s) \leq x \leq V(s-)\}$$

and observe that

$$N_x(t) = \lim_{n \rightarrow \infty} \uparrow \sum_{i: t_i^n < t} \mathbf{1}_{J_i^n}(x).$$

It follows from monotone convergence and Fubini-Tonelli Theorem that

$$\int_{\mathbb{R}} N_x(t) dx \leq W(t) < \infty. \quad (7)$$

A fortiori $N_x(t) < \infty$ for almost every x , which entails that the set

$$\{t > 0 : V(t-) \leq x \leq V(t) \text{ or } V(t) \leq x \leq V(t-)\}$$

is discrete for almost every x . \square

We stress that for every simple level x , we can enumerate the increasing and the decreasing passage times of V through x , and $\ell^x(t)$ and $\lambda^x(t)$ are both well-defined and finite. We are now able to state the second version of the change-of-variables formula.

Proposition 2. *Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a function of class \mathcal{C}^1 . Then for every $t > 0$, there is the identity*

$$f(V(t)) - f(V(0)) = \int_{\mathbb{R}} f'(x) \ell^x(t) dx + \sum_{0 < s \leq t} (f(V(s)) - f(V(s-))) .$$

Remark. Our recent note [2] (see Theorem 2.1(i) there) contains a similar result when V is an increasing process and the function f is merely assumed to be non-decreasing. Of course, when V is non-decreasing, the number of increasing passages through any level is at most 1 and there are no decreasing passages, so $\ell^x(t) = 1$ or 0, depending essentially on whether V hits the level x before time t or not.

Proof. Consider a simple level x . We further suppose that $x \neq V(t)$ and $x \neq V(0)$. For any such x , one can define the consecutive crossings of the level x by the graph of V . Each crossing can be made either upwards or downwards, and occurs either continuously or by a jump. More precisely, an upward crossing of x corresponds either to an increase through the level x , or to a positive jump starting strictly below x and finishing strictly above x ; and there is an analogous alternative for downwards crossings. Plainly upwards and downwards crossings alternate, and the quantity

$$\mathbf{1}_{[x, \infty)} V(t) - \mathbf{1}_{[x, \infty)} V(0)$$

coincides with the difference between upwards and downwards crossings of the level x made before time t . Distinguishing crossings according to whether they occur continuously or by a jump, we arrive at the identity

$$\mathbf{1}_{[x, \infty)} V(t) = \mathbf{1}_{[x, \infty)} V(0) + \ell^x(t) + \sum_{0 < s \leq t} (\mathbf{1}_{[x, \infty)} V(s) - \mathbf{1}_{[x, \infty)} V(s-)) , \quad (8)$$

where the series in the right-hand side accounts for the difference between the number of positive jumps and the number of negative jumps across the level x .

It is convenient to write ν for the point measure on $[0, \infty)$ which has an atom at each jump time of V , viz.

$$\nu(dt) = \sum_{s \geq 0} \mathbf{1}_{\{\Delta V(s) \neq 0\}} \delta_s(dt) .$$

We stress that ν is sigma-finite (because the number of jump times of V is at most countable), and rewrite

$$\sum_{0 < s \leq t} (\mathbf{1}_{[x, \infty)} V(s) - \mathbf{1}_{[x, \infty)} V(s-)) = \int_{[0, t]} (\mathbf{1}_{[x, \infty)} V(s) - \mathbf{1}_{[x, \infty)} V(s-)) \nu(ds) .$$

We also observe that, in the notation of the proof of Lemma 1, there is the inequality

$$\int_{[0, t]} |\mathbf{1}_{[x, \infty)} V(s) - \mathbf{1}_{[x, \infty)} V(s-)| \nu(ds) \leq N_x(t)$$

and recall from (7) that the function $x \rightarrow N_x(t)$ is in $L^1(dx)$. Because the map

$$(s, x) \rightarrow (\mathbf{1}_{[x, \infty)} V(s) - \mathbf{1}_{[x, \infty)} V(s-))$$

is measurable, it follows from Fubini Theorem that the map

$$x \rightarrow \int_{[0, t]} (\mathbf{1}_{[x, \infty)} V(s) - \mathbf{1}_{[x, \infty)} V(s-)) \nu(ds)$$

is also measurable. In particular, we conclude from (8) that $x \rightarrow \ell^x(t)$ is measurable.

We now rewrite the identity (8) in the form

$$\mathbf{1}_{[x, \infty)} V(t) = \mathbf{1}_{[x, \infty)} V(0) + \ell^x(t) + \int_{[0, t]} (\mathbf{1}_{[x, \infty)} V(s) - \mathbf{1}_{[x, \infty)} V(s-)) \nu(ds),$$

multiply both sides by $f'(x)$ and integrate with respect to the Lebesgue measure dx . An application of Fubini Theorem (which is legitimate thanks to the observations above) yields

$$f(V(t)) = f(V(0)) + \int_{\mathbb{R}} f'(x) \ell^x(t) dx + \int_{[0, t]} (f(V(s)) - f(V(s-))) \nu(ds),$$

which is the change-of-variables formula of the statement. \square

4 Proof of Theorem 1

We now turn our attention to the proof of our main result, which will also require the following elementary observation.

Lemma 2. *Let μ be a signed measure on some measurable space, and φ a measurable function with values in $\{-1, 1\}$ such that $\varphi\mu$ is a positive measure. Then $\varphi\mu$ coincides with the total variation measure $|\mu|$.*

Proof. Indeed, $\varphi\mu = |\varphi\mu|$ since $\varphi\mu$ is a positive measure, and on the other hand, we have also $|\varphi\mu| = |\varphi||\mu| = |\mu|$ since $|\varphi| \equiv 1$. \square

We now have all the ingredients needed for establishing Theorem 1.

Proof of Theorem 1 : First, comparing the two change-of-variables formulas in Propositions 1 and 2, we get that for any continuous function $g : \mathbb{R} \rightarrow \mathbb{R}$ (think of $g = f'$ as the derivative of a \mathcal{C}^1 function), there is the identity

$$\int_0^t g(V(s)) V^c(ds) = \int_{\mathbb{R}} g(x) \theta_t(dx) = \int_{\mathbb{R}} g(x) \ell^x(t) dx.$$

Thus $\theta_t(dx) = \ell^x(t) dx$.

Next, we introduce the signed measure $\mu(ds, dx)$ on $[0, t] \times \mathbb{R}$ which is defined by

$$\mu(A) = \int_0^t \mathbf{1}_A(s, V(s)) V^c(ds), \quad A \in \mathcal{B}([0, t] \times \mathbb{R}).$$

The (signed) occupation density formula above yields that for every $f : \mathbb{R} \rightarrow \mathbb{R}$ and $g : [0, t] \rightarrow \mathbb{R}$ measurable and bounded, there is the identity

$$\int_{[0, t] \times \mathbb{R}} f(x) g(s) \mu(ds, dx) = \int_0^t f(V(s)) g(s) V^c(ds) = \int_{\mathbb{R}} \left(\int_{[0, t]} g(s) \ell^x(ds) \right) f(x) dx.$$

More precisely, when g is a step function, this identity follows from the occupation density formula by linearity, and the general case is then derived through a version of the monotone class theorem (cf. Neveu [6]).

Then consider a measurable function $\varphi : [0, \infty) \rightarrow \{1, -1\}$ such that $V^c(dt) = \varphi(t) |V^c(dt)|$. We have

$$\int_{[0, t] \times \mathbb{R}} f(x) g(s) \varphi(s) \mu(ds, dx) = \int_0^t f(V(s)) g(s) |V^c(ds)|,$$

and the right-hand side is nonnegative whenever $f, g \geq 0$. Again by a version of the monotone class theorem, this shows that $\varphi(s) \mu(ds, dx)$ is a positive measure, and more precisely, since $|\varphi| = 1$, Lemma 2 shows that $\varphi(s) \mu(ds, dx) = |\mu(ds, dx)|$ is the total variation measure of $\mu(ds, dx)$. Now we write

$$\int_0^t f(V(s)) g(s) |V^c(ds)| = \int_{\mathbb{R}} \left(\int_{[0, t]} g(s) \varphi(s) \ell^x(ds) \right) f(x) dx.$$

Plainly, whenever $g \geq 0$, we must have $\int_{[0, t]} g(s) \varphi(s) \ell^x(ds) \geq 0$ for almost all $x \in \mathbb{R}$, that is $\varphi(s) \ell^x(ds)$ is a positive measure for almost all $x \in \mathbb{R}$. Again because $|\varphi| = 1$, this entails that

$$\varphi(s) \ell^x(ds) = |\ell^x(ds)| = \lambda^x(ds),$$

where the second equality is merely the definition of λ^x . Putting the pieces together, we have shown that

$$\int_0^t f(V(s)) g(s) |V^c(ds)| = \int_{\mathbb{R}} \left(\int_{[0, t]} g(s) \lambda^x(ds) \right) f(x) dx,$$

which for $g \equiv 1$ simply reads $\vartheta_t(dx) = \lambda^x(t) dx$. \square

We now conclude this note by stressing that the elementary identity (8) for the signed local time $\ell^x(t)$ at a simple level x should be viewed as the analog of the Meyer-Tanaka formula (2) in the semimartingale setting. We also point at the alternative formula

$$\mathbf{1}_{(-\infty, x)} V(t) = \mathbf{1}_{(-\infty, x)} V(0) - \ell^x(t) + \sum_{0 < s \leq t} (\mathbf{1}_{(-\infty, x)} V(s) - \mathbf{1}_{(-\infty, x)} V(s-)). \quad (9)$$

For the absolute local time $\lambda^x(t)$, one sees similarly that for every simple level x , there is the identity

$$H^x(t) = \lambda^x(t) + \sum_{0 < s \leq t} |\mathbf{1}_{[x, \infty)} V(s) - \mathbf{1}_{[x, \infty)} V(s-)|, \quad (10)$$

where $H^x(t)$ denotes the total variation on the time-interval $[0, t]$ of the step function $\mathbf{1}_{[x, \infty)} \circ V$.

References

1. Bertoin, J. Increase of stable processes. *J. Theoretic. Prob.* **7** (1994), 551-563.
2. Bertoin, J. and Yor, M. Pure jump increasing processes and the change of variables formula. *Electron. Commun. Probab.* **18** (2013), paper 41.
3. Dellacherie, C. and Meyer, P.A. *Probabilités et potentiel. Chapitres V à VIII*. Actualités Scientifiques et Industrielles [Current Scientific and Industrial Topics], **1385**, Hermann, Paris, 1980.
4. Dvoretzky, A., Erdős, P. and Kakutani, S. On nonincrease everywhere of the Brownian motion process. *Proc. 4th Berkeley Symp. Math. Stat. Prob. II* (1961), 103-116.
5. Fitzsimmons, P. J. and Port, S. C. Local times, occupation times, and the Lebesgue measure of the range of a Lévy process. In: *Seminar on Stochastic Processes, 1989 (San Diego, CA, 1989)*. Progr. Probab. **18**, pp. 59-73, Birkhäuser Boston 1990.
6. Neveu, J. *Mathematical foundations of the calculus of probability*. Holden-Day, Inc., San Francisco, Calif.-London-Amsterdam, 1965.
7. Protter, Ph. *Stochastic integration and differential equations*. Applications of Mathematics 21 Springer-Verlag, New York, 2004.