

The spinor representation formula in 3 and 4 dimensions

Pascal Romon, Julien Roth

► **To cite this version:**

Pascal Romon, Julien Roth. The spinor representation formula in 3 and 4 dimensions. Conference on Pure and Applied Differential Geometry, Aug 2012, Leuven, Belgium. Shaker Verlag, pp.300, 2013, Berichte aus der Mathematik. <hal-00857015>

HAL Id: hal-00857015

<https://hal.archives-ouvertes.fr/hal-00857015>

Submitted on 2 Sep 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The spinor representation formula in 3 and 4 dimensions

Pascal Romon*, Julien Roth*

Abstract

In the literature, two approaches to the Weierstrass representation formula using spinors are known, one explicit, going back to Kusner & Schmitt, and generalized by Konopelchenko and Taimanov, and one abstract due to Friedrich, Bayard, Lawn and Roth. In this article, we show that both points of view are indeed equivalent, for surfaces in \mathbb{R}^3 , Nil_3 and \mathbb{R}^4 . The correspondence between the equations of both approaches is explicitly given and as a consequence we derive alternate (and simpler) proofs of these previous theorems.

Keywords: Isometric immersions, Weierstrass formula, Dirac equation, Killing spinor.

2010 MSC: Primary 53C27, Secondary 53B25, 53C30

1 Historical background

The starting point of all representation formulae of submanifolds is the celebrated *Weierstrass–Enneper formula* for minimal surfaces in \mathbb{R}^3 : any minimal immersion f is given locally in conformal coordinates $z = x + iy$ by

$$f(z) = \text{Re} \int^z \left(\frac{1-g^2}{2}, i \frac{1+g^2}{2}, g \right) h dz,$$

where g is meromorphic and corresponds to the (stereographic) projection of the Gauss map, h is holomorphic, and $|h|(1+|g|^2)$ is neither 0 nor $+\infty$. This allows to describe locally (and sometimes globally) all minimal surfaces, and plays a key role in understanding and classifying them. However, this formula turns out to characterize many more surfaces than just the minimal ones: namely,

* Université Paris-Est, LAMA (UMR 8050), UPEMLV, UPEC, CNRS, F-77454, Marne-la-Vallée, France.

pascal.romon@univ-mlv.fr, julien.roth@univ-mlv.fr

and up to a cosmetic change, all surfaces, provided they are given in conformal coordinates.

Indeed, in 1979, Kenmotsu [Ke] generalized the formula to any two-dimensional conformal immersion in \mathbb{R}^3 , with prescribed (nonzero) mean curvature H , by proving that the Gauss map and mean curvature actually determine the surface (in the minimal case, an extra function h is required). To that effect, he studied the equation satisfied by the Gauss map.

In 1996, Kusner & Schmitt [KS], following an idea of Sullivan, proposed a similar albeit simpler expression, called the *spinor representation formula*, which gives locally all conformal immersions in \mathbb{R}^3 :

$$f(z) = \operatorname{Re} \int^z \left(i \frac{\bar{\psi}_2^2 + \psi_1^2}{2}, \frac{\bar{\psi}_2^2 - \psi_1^2}{2}, \psi_1 \bar{\psi}_2 \right) dz \quad (\text{W3})$$

where $\psi = (\psi_1, \psi_2)$ is a \mathbb{C}^2 -valued map satisfying the following (nonlinear) ‘‘Dirac equation with potential’’

$$\left(\begin{pmatrix} 0 & \partial/\partial z \\ -\partial/\partial \bar{z} & 0 \end{pmatrix} + \begin{pmatrix} U & 0 \\ 0 & V \end{pmatrix} \right) \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} = 0 \quad (\text{DP})$$

where $U = V = He^\rho/2$ is called the potential, and the metric is $ds^2 = e^\rho |dz|^2 = (|\psi_1|^2 + |\psi_2|^2) |dz|^2$. In the minimal case, $H = 0$, so $\psi_1, \bar{\psi}_2$ are holomorphic and we identify easily the meromorphic data: $h = i\psi_1^{-1}$, $g = -i\bar{\psi}_2/\psi_1$. Note that ψ is defined up to sign and $\psi_1, \bar{\psi}_2$ behave like \sqrt{dz} w.r.t. coordinate change¹.

A few years later, Konopelchenko [Ko] and Taimanov [T1] extended that formulation to conformally parametrized surfaces in $\mathbb{R}^4 \simeq \mathbb{C}^2$ by

$$f(z) = \int^z \begin{pmatrix} s_1 t_1 \\ s_1 t_2 \end{pmatrix} dz + \begin{pmatrix} -\bar{s}_2 \bar{t}_2 \\ \bar{s}_2 \bar{t}_1 \end{pmatrix} d\bar{z} \quad (\text{W4})$$

where s_1, s_2, t_1, t_2 are complex valued functions, obviously defined up to a global sign change, satisfying the following Dirac type equation, with complex potential h

$$\begin{cases} \partial s_1 / \partial \bar{z} = -\bar{h} \bar{s}_2 \\ \partial \bar{s}_2 / \partial z = h s_1 \end{cases} \quad \text{and} \quad \begin{cases} \partial t_1 / \partial \bar{z} = -h \bar{t}_2 \\ \partial \bar{t}_2 / \partial z = \bar{h} t_1 \end{cases} \quad \text{where } |h| = \frac{\|\vec{H}\| e^\rho}{2} \quad (1)$$

(\vec{H} stands for the mean curvature vector and e^ρ is again the conformal factor). The phase of h will be understood later. A particular case of this equation was obtained independently by Hélein & Romon in their study of Lagrangian immersions in euclidean space [HRo]. There, $t_1 = \cos \frac{\beta}{2}$ and $t_2 = \sin \frac{\beta}{2}$, where β is the lagrangian angle $\arg \det df$. This property has led to deep results in the theory of Hamiltonian-stationary lagrangian surfaces (see [LeRo, McRo]

¹A property which indicates that ψ is *not* really a spinor.

and also the surprising parametrization of Aiyama [A]). Finally, Taimanov [T2] expanded the formula (W3) to three dimensional homogeneous manifolds. The formula itself barely changes (see below), as well as the Dirac equation (DP), except for the potential. For example, if the ambient space is Nil_3 endowed with a left-invariant metric, then

$$U = V = \frac{He^\rho}{2} + \frac{i}{4}(|\psi_2|^2 - |\psi_1|^2)$$

However, these representation formulae were purely local and computational in nature, and it was not clear what the relationship they had with the spin bundle until Friedrich bridged the gap in [F1]. He showed that any surface M immersed in \mathbb{R}^3 inherits a spinor φ (given an arbitrary choice of a parallel spinor), which satisfies Dirac's equation with a potential, and can be used to reconstruct the immersion. This property was extended by Morel to \mathbb{S}^3 and \mathbb{H}^3 [Mo], by Roth to homogeneous spaces $E_{\kappa,\tau}$ [R], and most recently by Bayard, Lawn & Roth to \mathbb{R}^4 [BLR].

These abstract approaches all follow Bär's remark [Bä] that an ambient spinor, restricted to an immersed submanifold M , can be identified with an intrinsic spinor (or, in some cases, with a section of a twisted spin bundle). If the ambient spinor field $\tilde{\varphi}$ satisfies some particular PDE, e.g. is parallel or Killing, then the induced spinor φ also satisfies a PDE, typically a Dirac equation with a potential linked to the extrinsic geometry of M . Conversely (and non trivially), the Dirac equation with potential is shown to imply Gauss and Codazzi's equations (and Ricci's when needed), thus guaranteeing the (local) existence of an immersion with prescribed extrinsic data.

Our goal in this paper is to clarify and explicit the relationship between these two point of view, which has not been done before. In particular, we show how the concrete representation formulae and Dirac equations are actually specific gauge choices of the general abstract formulae, explaining their simpler appearance. We give new and more direct proofs of theorems 1 and 2 below, avoiding the cumbersome Gauss, Codazzi and Ricci equations (although they are present implicitly), including their formulation in $E_{\kappa,\tau}$ written in [D]. Note that computations cannot be completely avoided.

We will recall, first in three, then in four dimensions, how one obtains the concrete formulae cited above, then present briefly the induced spinor point of view, and finally write down the correspondence between these two.

Acknowledgements. The authors wish to thank Marie-Amélie Lawn for her interesting suggestions.

2 Conformal immersions in dimension three

We consider in this section the case of surfaces in \mathbb{R}^3 and in metric Lie groups of dimension 3, endowed with a left-invariant metric, and focus on the Heisenberg group $\text{Nil}_3 = \text{Nil}_3(\tau)$ where τ is the fiber torsion (see [D]). When $\tau = 0$ we recover \mathbb{R}^3 . This approach can be extended to other groups and homogeneous spaces without much work, and we direct the reader to [T2].

2.1 Coordinates and the Maurer–Cartan form

The formula (W3) is based on a simple remark: Given an immersion $f : M \rightarrow \mathbb{R}^3$ with coordinate $z = x + iy$, then f is conformal if and only if the derivative $\partial f / \partial z = \frac{1}{2}(\partial f / \partial x - i \partial f / \partial y)$ lies in the quadric

$$Q = \{Z = (Z_1, Z_2, Z_3) \in \mathbb{C}^3, Z_1^2 + Z_2^2 + Z_3^2 = 0\}$$

(note that $\mathbb{P}Q \subset \mathbb{C}\mathbb{P}^2$ is the grassmannian $\text{Gr}_{2,3}$). We can parametrize Q by two complex numbers with what is usually called the Segre map

$$Q \ni Z = \left(\frac{i}{2}(\bar{\psi}_2^2 + \psi_1^2), \frac{1}{2}(\bar{\psi}_2^2 - \psi_1^2), \psi_1 \bar{\psi}_2 \right)$$

and one recovers immediately (W3). If the ambient space \tilde{M} is a Lie group, let (E_1, E_2, E_3) be the orthonormal (moving) frame generated by left translation from any choice of orthonormal frame at the identity e . Then (E_1, E_2, E_3) trivializes TM , in other words, the tangent vectors can be mapped to \mathbb{R}^3 in this frame, or equivalently, one identifies vectors mapped by the differential of f^{-1} to $T_e M$ with vectors in \mathbb{R}^3 (or $T_e M^{\mathbb{C}}$ with \mathbb{C}^3):

$$Z = \begin{pmatrix} Z_1 \\ Z_2 \\ Z_3 \end{pmatrix} \simeq f^{-1} \frac{\partial f}{\partial z}$$

and Z lies again in Q if and only if f is (weakly-)conformal. Hence the same formula (W3) holds, with $f^{-1} \partial f / \partial z$ playing the role of the derivative:

$$f^{-1} \frac{\partial f}{\partial z}(z) = \frac{i}{2}(\bar{\psi}_2^2 + \psi_1^2)E_1 + \frac{1}{2}(\bar{\psi}_2^2 - \psi_1^2)E_2 + \psi_1 \bar{\psi}_2 E_3. \quad (2)$$

Note that integration is less obvious than in the trivial translational group \mathbb{R}^3 .

Conversely, the formula (2) gives an immersion when integrated over a simply connected planar domain, provided the Maurer–Cartan form $\alpha = f^{-1} df$ satisfies the integrability condition $d\alpha + \frac{1}{2}[\alpha \wedge \alpha] = 0$. Let us write down the equation in two explicit cases: \mathbb{R}^3 , where the bracket is identically zero, and

$\text{Nil}_3(\tau)$ where the only nonzero bracket is $[E_1, E_2] = 2\tau E_3$ (and E_3 is the vertical fiber unit tangent vector). We recover the euclidean \mathbb{R}^3 by letting τ go to 0. The integrability equation boils down to, in the (E_1, E_2, E_3) frame at identity,

$$-\frac{\partial}{\partial y} \begin{pmatrix} -\frac{1}{2} \text{Im}(\bar{\psi}_2^2 + \psi_1^2) \\ \frac{1}{2} \text{Re}(\bar{\psi}_2^2 - \psi_1^2) \\ \text{Re}(\bar{\psi}_2 \psi_1) \end{pmatrix} + \frac{\partial}{\partial x} \begin{pmatrix} -\frac{1}{2} \text{Re}(\bar{\psi}_2^2 + \psi_1^2) \\ -\frac{1}{2} \text{Im}(\bar{\psi}_2^2 - \psi_1^2) \\ -\text{Im}(\bar{\psi}_2 \psi_1) \end{pmatrix} + 2\tau \begin{pmatrix} 0 \\ 0 \\ \frac{1}{4}(|\psi_2|^2 + |\psi_1|^2)(|\psi_2|^2 - |\psi_1|^2) \end{pmatrix} = 0 \quad (3)$$

that is

$$\begin{cases} \text{Im}((\bar{\psi}_2)_y \bar{\psi}_2) - \text{Re}((\bar{\psi}_2)_x \bar{\psi}_2) + \text{Im}((\psi_1)_y \psi_1) - \text{Re}((\psi_1)_x \psi_1) = 0 \\ -\text{Re}((\bar{\psi}_2)_y \bar{\psi}_2) - \text{Im}((\bar{\psi}_2)_x \bar{\psi}_2) + \text{Re}((\psi_1)_y \psi_1) + \text{Im}((\psi_1)_x \psi_1) = 0 \\ -\text{Im}[(\bar{\psi}_2)_x + i(\bar{\psi}_2)_y] \psi_1 - \text{Im}[(\psi_1)_x + i(\psi_1)_y] \bar{\psi}_2 \\ \quad + \frac{\tau}{2}(|\psi_2|^2 + |\psi_1|^2)(|\psi_2|^2 - |\psi_1|^2) = 0. \end{cases}$$

The first two lines amount to $(\psi_2)_z \psi_2 + (\psi_1)_{\bar{z}} \psi_1 = 0$, which implies the existence of a function U such that

$$U = \frac{1}{\psi_2} \frac{\partial \psi_1}{\partial \bar{z}} = -\frac{1}{\psi_1} \frac{\partial \psi_2}{\partial z}$$

(at least when ψ_1, ψ_2 do not vanish, and by construction, they do not vanish simultaneously). We obtain the equation (DP) with some complex valued potential U . However the third line yields

$$\begin{aligned} 0 &= 2 \text{Im} \left[\frac{\partial \bar{\psi}_2}{\partial \bar{z}} \psi_1 + \frac{\partial \psi_1}{\partial \bar{z}} \bar{\psi}_2 \right] - \frac{\tau}{2} (|\psi_2|^2 + |\psi_1|^2) (|\psi_2|^2 - |\psi_1|^2) \\ &= 2 \text{Im} [-\bar{U} |\psi_1|^2 + U |\psi_2|^2] - \frac{\tau}{2} (|\psi_2|^2 + |\psi_1|^2) (|\psi_2|^2 - |\psi_1|^2) \\ &= 2 \text{Im} [U (|\psi_1|^2 + |\psi_2|^2)] - \frac{\tau}{2} (|\psi_2|^2 + |\psi_1|^2) (|\psi_2|^2 - |\psi_1|^2) \end{aligned}$$

so that $\text{Im} U = \frac{\tau}{4} (|\psi_2|^2 - |\psi_1|^2)$ in $\text{Nil}_3(\tau)$, and vanishes in \mathbb{R}^3 . Other three-dimensional groups follow the same lines (see [T2] for a full list). Note that the Dirac equation *per se* does not fully determine the potential U , but it can be derived from the representation formula.

2.2 Induced spinors

We assume here a basic acquaintance with spinor and Clifford algebras (see [LM] or [F2] for an introduction). For any spin manifold M^m , we denote by $\text{Spin}(M)$ the spin bundle and $\Sigma(M) = \text{Spin}(M) \times_{\chi_m} \Sigma_n$ the associated spinor bundle,

where χ_m is a representation of $\text{Spin}(M)$ on Σ_m . We recall here briefly the idea in [F1] and [R] in dimension three, and we will give a more explicit formulation in later sections.

Let \tilde{M} be a three dimensional spin manifold², M an oriented immersed surface in \tilde{M} . Because spinor frames in M can be identified with spinor frames of \tilde{M} along M , and both $\text{Spin}(2)$ and $\text{Spin}(3)$ have a canonical representation on $\Sigma_2 = \Sigma_3 = \mathbb{C}^2$, it is possible to identify an ambient spinor $\tilde{\varphi}|_M$ restricted to M with an intrinsic spinor φ of M (when endowed with the induced metric). All identifications are isomorphic (by the universality property) and differ by the corresponding Clifford multiplication. In [R], the identification is such that

$$\forall X \in TM \subset T\tilde{M}, \quad X \cdot_M \varphi = (X \cdot \nu \cdot \tilde{\varphi})|_M$$

where ν is the unit (oriented) normal along M , \cdot_M denotes the Clifford multiplication in M and \cdot the one in \tilde{M} . We will rather use a more direct identification, which will simplify the calculations below, and satisfies the simpler relation

$$\forall X \in TM \subset T\tilde{M}, X \cdot_M \varphi = (X \cdot \tilde{\varphi})|_M. \quad (4)$$

Hence from now on, we will make no difference between Clifford multiplication in M and in \tilde{M} . We will rederive the computations in [R] with this new convention.

A spinor field is the equivalence class of a couple (s, u) , where s is a section of $\text{Spin}(\tilde{M})$ (a moving spinor frame) and u is a Σ_3 -valued function, under the equivalence relation induced by χ_3 :

$$(s, u) \sim (sa^{-1}, \chi_3(a)u)$$

for all $a : \tilde{M} \rightarrow \text{Spin}(3) = \mathbb{S}^3$. When \tilde{M} is a Lie group, we trivialize $\text{Spin}(\tilde{M})$ by a lift \tilde{s} of the left-invariant moving frame defined above, and define $\tilde{\varphi}$ to be *constant* if $\tilde{\varphi}$ is (the equivalence class of) the couple (\tilde{s}, u) and u is constant. (This property is obviously independent of the choice of the reference spinor frame \tilde{s} .) In \mathbb{R}^3 , such a spinor is *parallel*, however that is not the case in general Lie groups. Again, taking as example $\text{Nil}_3(\tau)$, where $\tilde{\nabla}_X E_3 = \tau X \times E_3$, we have, on a constant spinor field $\tilde{\varphi}$

$$\tilde{\nabla}_{E_1} \tilde{\varphi} = \frac{\tau}{2} E_1 \cdot \tilde{\varphi} \quad \tilde{\nabla}_{E_2} \tilde{\varphi} = \frac{\tau}{2} E_2 \cdot \tilde{\varphi} \quad \tilde{\nabla}_{E_3} \tilde{\varphi} = -\frac{\tau}{2} E_3 \cdot \tilde{\varphi}$$

because

$$\tilde{\nabla}_X \tilde{\varphi} = (\tilde{s}, Xu) + \frac{1}{2} \sum_{1 \leq p < q \leq 3} \langle \tilde{\nabla}_X E_p, E_q \rangle E_p \cdot E_q \cdot \tilde{\varphi}$$

where $\tilde{\nabla}$ denotes both the connection on $T\tilde{M}$ and on $\Sigma\tilde{M}$, depending on the context. (Note: The case of \mathbb{R}^3 can be obtained by letting $\tau = 0$, and we recover parallelism.)

²It suffices here to assume that \tilde{M} is orientable.

Let M be an oriented immersed surface, with normal vector ν and let T be the projection of E_3 on TM , so that $E_3 = T + \lambda\nu$. Let $S = -d\nu$ be the shape operator. Then, by the spinorial Gauss relation,

$$\begin{aligned}
\nabla_X \varphi &= \tilde{\nabla}_X \tilde{\varphi}|_M - \frac{1}{2}(SX) \cdot \nu \cdot \varphi \\
&= \frac{\tau}{2}(X^1 E_1 + X^2 E_2 - X^3 E_3) \cdot \varphi - \frac{1}{2}(SX) \cdot \omega_2 \cdot \varphi \\
&= \frac{\tau}{2}(X \cdot \varphi - 2\langle X, T \rangle E_3 \cdot \varphi) - \frac{1}{2}(SX) \cdot \omega_2 \cdot \varphi \\
&= \frac{\tau}{2} \left(-X \cdot \underbrace{\nu \cdot e_1 \cdot e_2}_{\omega_3} \cdot \varphi \right) - \tau \langle X, T \rangle (T \cdot \varphi + \lambda \nu \cdot \varphi) - \frac{1}{2}(SX) \cdot \omega_2 \cdot \varphi \\
&= \frac{\tau}{2} X \cdot \varphi - \tau \langle X, T \rangle (T \cdot \varphi + \lambda \omega_2 \cdot \varphi) - \frac{1}{2}(SX) \cdot \omega_2 \cdot \varphi
\end{aligned}$$

where $\omega_2 = e_1 \cdot e_2 \in Cl_2(M)$ (for any oriented orthonormal frame (e_1, e_2) of M), and $\omega_3 = E_1 \cdot E_2 \cdot E_3 = e_1 \cdot e_2 \cdot \nu \in Cl_3(M)$ are called the *real volume elements* (we used that $\chi_3(\omega_3) = -\text{id}$ to infer $\nu \cdot \varphi = \omega_2 \cdot \varphi$). Finally, $\tilde{\varphi}$ is constant if and only if

$$\nabla_X \varphi = -\frac{1}{2}(SX) \cdot \omega_2 \cdot \varphi + \frac{\tau}{2} X \cdot \varphi - \tau \langle X, T \rangle (T \cdot \varphi + \lambda \omega_2 \cdot \varphi). \quad (\text{C})$$

This implies in particular that $|\varphi|$ is constant. Applying the Dirac operator to φ , we end up with the following Dirac equation

$$D\varphi = H\omega_2 \cdot \varphi + \tau\lambda(\lambda\varphi - T \cdot \omega_2 \cdot \varphi) \quad (5)$$

We now quote the following

Theorem 1 (Friedrich, Roth). *Given a riemannian oriented, simply connected surface M , a field of symmetric endomorphisms S , with trace $2H$ (and in $\text{Nil}_3(\tau)$) a vector field T and function λ , such that $\lambda^2 + \|T\|^2 = 1$). Then the following data are equivalent:*

1. *an isometric immersion from M^2 into \mathbb{R}^3 (resp. $\text{Nil}_3(\tau)$) such that S is the shape operator, and H the mean curvature (and in $\text{Nil}_3(\tau)$, $E_3 = T + \lambda\nu$),*
2. *a non-trivial spinor field φ solution of (C), where the symmetric tensor S and vector field T satisfy, for all X*

$$\nabla_X T = \lambda(SX - \tau JX), \quad X(\lambda) = -\langle SX - \tau JX, T \rangle \quad (6)$$

(J being the complex structure),

3. *a non-trivial spinor field φ of constant norm, solution of the Dirac equation (5), with T , λ satisfying:*

$$d\lambda = -2Q_\varphi(T) - B(T) - \tau JT,$$

where B is the tensor defined in an orthonormal frame $\{e_1, e_2\}$ by the following matrix

$$\begin{pmatrix} 2\tau T_1 T_2 & \tau(T_1^2 - T_2^2) \\ \tau(T_1^2 - T_2^2) & -2\tau T_1 T_2 \end{pmatrix},$$

and Q_φ is the Energy-Momentum tensor associated with the spinor field φ , defined by:

$$Q_\varphi(X, Y) := \frac{1}{2} \operatorname{Re} \langle X \cdot \nabla_Y \varphi + Y \cdot \nabla_X \varphi, \varphi / |\varphi|^2 \rangle.$$

The proof is technical and requires proving that the Gauss & Codazzi equations

$$\begin{aligned} K &= \det A + \tau^2(1 - 4\lambda^2) && \text{(Gauss)} \\ \nabla_X AY - \nabla_Y AX - A[X, Y] &= -4\tau^2 \lambda (\langle Y, T \rangle X - \langle X, T \rangle Y) && \text{(Codazzi)} \end{aligned}$$

are satisfied (with the extra conditions (6) of Daniel on T and λ for $\operatorname{Nil}_3(\tau)$).

2.3 An explicit construction of induced spinors

We will now construct explicitly the bundles involved, using quaternions for shorter notations. Let us identify \mathbb{R}^3 with $\operatorname{Im} \mathbb{H} = \operatorname{Span}(i, j, k)$ and the Clifford algebra with

$$\mathcal{Cl}_3 \simeq \left\{ \begin{pmatrix} a & b \\ b & a \end{pmatrix}, a, b \in \mathbb{H} \right\}, \text{ where } \mathbb{R}^3 \simeq \left\{ \begin{pmatrix} 0 & b \\ b & 0 \end{pmatrix}, b \in \operatorname{Im} \mathbb{H} \right\}.$$

The group $\operatorname{Spin}(3)$ identifies with

$$\operatorname{Spin}(3) \simeq \left\{ \begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix}, a \in \mathbb{S}^3 \subset \mathbb{H} \right\}$$

and acts on \mathbb{R}^3 via matrix multiplication, which amounts to $\theta_3(a) : x \mapsto axa^{-1} = ax\bar{a}$, for any $a \in \mathbb{S}^3$ and $x \in \operatorname{Im} \mathbb{H}$, yielding a twofold cover of $\operatorname{SO}(3)$. From the above action on \mathbb{H}^2 , we also infer an irreducible action χ_3 of $\operatorname{Spin}(3)$ on $\Sigma_3 = \mathbb{H}$, given by the left multiplication by a , $L_a : q \mapsto aq$. The representation is the restriction³ of $\chi_3 : \mathcal{Cl}_3 \rightarrow \operatorname{End}(\mathbb{H})$ such that $\chi_3(x) = L_x$ for $x \in \operatorname{Im} \mathbb{H}$ and $q \in \mathbb{H}$ (since imaginary quaternions have the property that $x^2 = -x\bar{x} = -|x|^2$). Recall that the Clifford multiplication of a spinor by a vector is simply $x \cdot \varphi = \chi_3(x)\varphi$. Finally, we identify Σ_3 with \mathbb{C}^2 by choosing a compatible complex structure; in what follows we let $J = R_i$, the right multiplication by i (so that the complex number i will coincide with the quaternion i).

³The full expression for χ_3 is $\chi_3 \left(\begin{pmatrix} a & b \\ b & a \end{pmatrix} \right) = L_{a+b}$.

The tangent space at identity $e \in \tilde{M}$ identifies with \mathbb{R}^3 and the spinor frame at e with $\text{Spin}(3) \simeq \mathbb{S}^3$. The trivialisation of $\text{Spin}(\mathbb{R}^3)$ as $\mathbb{R}^3 \times \text{Spin}(3)$ holds in greater generality for three dimensional groups \tilde{M} via the group multiplication, where the tangent space $T_x \tilde{M}$ is mapped canonically to $T_e \tilde{M}$ by left multiplication by x^{-1} . Equivalently, a choice of a spinor frame at identity yields by left multiplication a spinor moving frame \tilde{s} on \tilde{M} , which trivializes $\text{Spin}(\tilde{M})$. Such a spinor moving frame will be called *constant*, and is unique up to right multiplication by $\text{Spin}(3)$.

A spinor field $\tilde{\varphi}$ being the (equivalence class of the) couple (s, u) where s is a moving spinor frame and u a map from M to $\Sigma_3 = \mathbb{H}$, the equivalence relation now reads

$$(s, u) \sim (sa^{-1}, \chi_3(a)u) = (sa^{-1}, au)$$

for all $a : \tilde{M} \rightarrow \text{Spin}(3) = \mathbb{S}^3 \subset \mathbb{H}$. Clifford multiplication is well-defined in the following way: since any moving spinor frame s induces by θ_3 an orthonormal moving frame (E_1, E_2, E_3) , and any tangent vector X possesses coordinates (X_1, X_2, X_3) in this frame, we can identify the tangent bundle $T\tilde{M}$ with $\text{Spin}(\tilde{M}) \times_{\theta_3} \mathbb{R}^3$, with the equivalence relation

$$(s, (X_1, X_2, X_3)) \sim (sa^{-1}, \theta_3(a)(X_1, X_2, X_3))$$

or, in quaternionic notations $x = X_1i + X_2j + X_3k$: $(s, x) \sim (sa^{-1}, \theta_3(a)x) = (sa^{-1}, axa^{-1})$. The Clifford multiplication is then

$$X \cdot \varphi = (s, x) \cdot (s, u) = (s, x \cdot u)$$

where $x \cdot u$ is the multiplication in $C\ell_3$. It is immediate that $X \cdot \varphi$ is defined independently of the frame s .

In the following, we pick a constant reference spinor frame \tilde{s} as above, using the group multiplication, and \tilde{s} lies over the orthonormal moving frame (E_1, E_2, E_3) . In $\text{Nil}_3(\tau)$, we choose \tilde{s} such that E_3 lies on the vertical fiber. As a consequence, a spinor field $\tilde{\varphi}$ identifies with a map $u : \tilde{M} \rightarrow \mathbb{H}$. We will then say that $\tilde{\varphi}$ is *constant* if u is, and note that the property does not depend on a particular choice of \tilde{s} .

Writing \mathbb{R}^2 as $\text{Span}(i, j) \in \mathbb{H}$, the two-dimensional Clifford algebra $C\ell_2$ identifies with \mathbb{H} , and $\text{Spin}(2) \simeq U(1) = \{e^{k\alpha} = (\cos \alpha)1 + (\sin \alpha)\omega_2\}$, where $\omega_2 = i \cdot j = k$. The double cover $\theta_2 : \text{Spin}(2) \rightarrow \text{SO}(2)$ is $e^{k\alpha} \mapsto \begin{pmatrix} \cos 2\alpha & -\sin 2\alpha \\ \sin 2\alpha & \cos 2\alpha \end{pmatrix}$. As representation on $\Sigma_2 = \mathbb{H}$, we choose the universal extension of $\chi_2 : \mathbb{R}^2 \rightarrow \text{End}(\mathbb{H})$, $\chi_2(x_1i + x_2j) = L_{x_1i + x_2j}$. In particular, if $a = e^{k\alpha} \in \text{Spin}(2)$, $\chi_2(a)$ is again the left multiplication L_a . Again, we endow Σ_2 with the complex structure $J = R_i$. Finally, the *complex volume element* $\omega_2^{\mathbb{C}} = \sqrt{-1}i \cdot j$, which squares to the identity, acts as $\chi_3(i \cdot j)J = L_k R_i$ on Σ_2 and splits Σ_2 into two eigenspaces $\Sigma_2^{\pm} = (1 \pm j)\mathbb{C}$ of *positive* and *negative* spinors⁴. On a spin manifold

⁴Note for later that this splitting is actually hermitian orthogonal, so that $|v|^2 = |v^{+2}| + |v^{-}|^2$.

M , this allow to view spinors as square roots of forms. Indeed, let s be a spinor frame at a point z , corresponding to the oriented orthonormal frame (e_1, e_2) , and $\varphi = (s, v)$ be a spinor, with $v = \frac{1+j}{\sqrt{2}}v^+ + \frac{1-j}{\sqrt{2}}v^-$. Then the following forms are well defined :

$$T_z M \ni X = X_1 e_1 + X_2 e_2 \mapsto (v^+)^2 (X_1 + iX_2)$$

as well as

$$X \mapsto (\overline{v^-})^2 (X_1 + iX_2) \quad \text{and} \quad X \mapsto (v^+ \overline{v^-}) (X_1 + iX_2).$$

Indeed, the gauge change $s \rightarrow se^{-k\alpha}$ induces $v^+ \rightarrow v^+ e^{-i\alpha}$, $v^- \rightarrow v^- e^{+i\alpha}$ and $X_1 + iX_2 \rightarrow e^{2i\alpha} (X_1 + iX_2)$ (because $e^{k\alpha} (1 \pm j) = (1 \pm j) e^{-i\alpha}$). Obviously these 1-forms are of type $(1, 0)$.

Consider now an immersion $f : M \rightarrow \tilde{M}$. From the embedding of \mathbb{R}^2 into \mathbb{R}^3 as the plane ν^\perp , we deduce an embedding of $\text{Spin}(2)$ into $\text{Spin}(3)$ such that for any $s \in \text{Spin}(2) \subset \text{Spin}(3)$, $\theta_3(s)$ is the rotation of axis ν , whose restriction to the oriented plane ν^\perp is $\theta_2(s)$. This embedding carries over to the spinor frame bundles: In the same way as a tangent frame (e_1, e_2) to M identifies with the tangent frame (e_1, e_2, ν) of \tilde{M} , where ν is the unit normal vector of M , a spinor frame $s \in \text{Spin}(M)$ identifies with some spinor frame in $\text{Spin}(\tilde{M})$, which, given a reference frame \tilde{s} , can be written as $\tilde{s}a^{-1}$, for some $a \in \text{Spin}(3)$. Suppose furthermore that M is parametrized by f in conformal coordinates and that $s(z)$ lies over the frame $(e^{-\rho}\partial f/\partial x, e^{-\rho}\partial f/\partial y)$. Moving all vectors to e by left multiplication by $f(z)^{-1}$, the identification $s(z) \simeq \tilde{s}(f(z))a(z)^{-1}$ reads as⁵

$$\epsilon_1 = e^{-\rho} f^{-1} \frac{\partial f}{\partial x} = a^{-1} i a, \quad \epsilon_2 = e^{-\rho} f^{-1} \frac{\partial f}{\partial y} = a^{-1} j a, \quad \epsilon_3 = f^{-1} \nu = a^{-1} k a \quad (7)$$

where (i, j, k) stand for the frame at e corresponding to $\tilde{s}(e)$ (in quaternionic notations). Clearly a is determined (up to sign) by this property.

The identification of $\Sigma \tilde{M}|_M$ with ΣM used in §2.2 goes as follows: Since, along M , any ambient spinor $\tilde{\varphi} = (\tilde{s}, u)$ can be written as the couple $(\tilde{s}a^{-1}, au)$, and $\tilde{s}a^{-1}$ is identified with the intrinsic spinor frame s , $\tilde{\varphi}$ identifies with the intrinsic spinor $\varphi = (s, v)$ where $v = au$. Of all possible choices, this identification is the one coherent with our choice of Clifford multiplication (4).

Summing up the above, we see that from a ambient spinor field $\tilde{\varphi}$, one can construct $(1, 0)$ -forms along an immersed surface M . We will now replicate the construction of the previous section in order to obtain the representation formula. The principle is the following: Starting with a constant ambient spinor $\tilde{\varphi} = (\tilde{s}, u)$, the induced spinor $\varphi = (s, v) = (\tilde{s}a^{-1}, au)$ determines the spinor

⁵In \mathbb{R}^3 , the group operation is just the vector sum, and one usually identifies (abusively) $T\mathbb{R}^3$ with $T_0\mathbb{R}^3 = \mathbb{R}^3$, so that $f^{-1}df$ is identified with df .

frame a , since $a = vu^{-1}$ and u is fixed. We then deduce $f^{-1}df$ from a via (7). Given a unit quaternion u , and writing $v = \frac{1+j}{\sqrt{2}}v^+ + \frac{1-j}{\sqrt{2}}v^-$,

$$\begin{aligned}
\epsilon_1 &= \bar{a}ia \\
&= \frac{1}{2}u(\overline{v^+}(1-j) + \overline{v^-}(1+j))i((1+j)v^+ + (1-j)v^-)u^{-1} \\
&= u(\overline{v^+}kv^+ - \overline{v^-}kv^- + \overline{v^+}iv^- + \overline{v^-}iv^+)u^{-1} \\
&= u(k((v^+)^2 - (v^-)^2) + i(\overline{v^+}v^- + \overline{v^-}v^+))u^{-1} \\
\epsilon_2 &= \bar{a}ja \\
&= \frac{1}{2}u(\overline{v^+}(1-j) + \overline{v^-}(1+j))j((1+j)v^+ + (1-j)v^-)u^{-1} \\
&= u(\overline{v^+}jv^+ + \overline{v^-}jv^- + \overline{v^+}v^- - \overline{v^-}v^+)u^{-1} \\
&= u(j((v^+)^2 + (v^-)^2) + (\overline{v^+}v^- - \overline{v^-}v^+))u^{-1} \\
\epsilon_3 &= \bar{a}ka \\
&= \frac{1}{2}u(\overline{v^+}(1-j) + \overline{v^-}(1+j))k((1+j)v^+ + (1-j)v^-)u^{-1} \\
&= u(-\overline{v^+}iv^+ + \overline{v^-}iv^- + \overline{v^+}kv^- + \overline{v^-}kv^+)u^{-1} \\
&= u(-i(|v^+|^2 - |v^-|^2) + 2k(v^+v^-))u^{-1}
\end{aligned}$$

Choose $u = (1 + i - j + k)/2$, which satisfies $uk\bar{u} = -j$, $uj\bar{u} = -i$, $ui\bar{u} = k$,

$$\begin{cases}
\epsilon_1 &= -\operatorname{Im}((v^+)^2 - (v^-)^2)i - \operatorname{Re}((v^+)^2 - (v^-)^2)j + 2\operatorname{Re}(\overline{v^-}v^+)k \\
&= -\operatorname{Im}(\overline{(v^-)^2} + (v^+)^2)i + \operatorname{Re}(\overline{(v^-)^2} - (v^+)^2)j + 2\operatorname{Re}(\overline{v^-}v^+)k \\
\epsilon_2 &= -\operatorname{Re}((v^+)^2 + (v^-)^2)i + \operatorname{Im}((v^+)^2 + (v^-)^2)j - 2\operatorname{Im}(\overline{v^-}v^+)k \\
&= -\operatorname{Re}(\overline{(v^-)^2} + (v^+)^2)i - \operatorname{Im}(\overline{(v^-)^2} - (v^+)^2)j - 2\operatorname{Im}(\overline{v^-}v^+)k \\
\epsilon_3 &= -2\operatorname{Im}(v^+v^-)i - 2\operatorname{Re}(v^+v^-)j - (|v^+|^2 - |v^-|^2)k
\end{cases}$$

Defining $\psi_1 = e^{\rho/2}v^+$ and $\psi_2 = e^{\rho/2}v^-$ (so that $|\psi_1|^2 + |\psi_2|^2 = e^\rho$, see footnote 4), i.e.

$$v = e^{-\rho/2} \left(\frac{1+j}{\sqrt{2}}\psi_1 + \frac{1-j}{\sqrt{2}}\psi_2 \right)$$

we have recovered exactly the original representation formula (2) (and (W3) in \mathbb{R}^3).

Note 1. Rigorously speaking, we see that $\psi = (\psi_1, \psi_2)$ is not a spinor, because of the metric factor. But $e^{-\rho/2}\psi$ is. Hence, the integrability equation (DP) reads in terms of v as

$$\begin{cases}
v_z^\dagger = -\frac{\rho_z}{2}v^+ + Uv^- \\
v_{\bar{z}} = -\frac{\rho_z}{2}v^- - Uv^+
\end{cases} \quad (8)$$

Note 2. A different choice of u would yield a different parametrization of the complex curve, or more precisely, a rotation of the formula (2). Our choice of u is the one corresponding to the classical formula. This particular choice will play a role in the next section.

We have thus seen concretely how an intrinsic spinor field induced by a constant ambient spinor is sufficient to reconstruct entirely the immersion (a fact already noted by the previous authors, although without the link to the representation formula). Conversely, given $\psi = (\psi_1, \psi_2)$, then formula (2) defines locally an immersion of M into \tilde{M} , provided the integrability condition (DP) holds. Let $s = \tilde{s}a^{-1}$ be the moving spinor frame along M associated with the conformal coordinate z (s is unique up to sign), and set $\varphi = (s, v)$ with $v = e^{-\rho/2}\psi$. Then along M , the ambient spinor $\tilde{\varphi} = (\tilde{s}, a^{-1}v)$ is constant. Indeed, the translated moving frame $(\epsilon_1, \epsilon_2, \epsilon_3)$ is $(\bar{a}ia, \bar{a}ja, \bar{a}ka)$ as well as $(u\bar{v}iv\bar{u}, u\bar{v}jv\bar{u}, u\bar{v}kv\bar{u})$, by definition of the representation formula. This implies $a = \pm vu^{-1}$, so $a^{-1}v = \pm u$ is constant.

It is worth noting that we used (DP) as an integrability condition and not as a reformulation of $\nabla\tilde{\varphi} = 0$. Rather, the spinor representation formula (2) *per se* implies that the corresponding ambient spinor is constant, *provided the immersion exists*, namely that (DP) holds. In other words (DP) implies (C). There remains to show the converse.

2.4 Dirac: from abstract to explicit

Let us write the constancy equation (C) in terms of quaternions. In the local coordinate z , with associate spinor frame s , the spinor field $\varphi = (s, v)$ identifies with v and we write

$$\begin{aligned}\nabla_X\varphi &= (s, Xv) + \frac{1}{2}\langle\nabla_X\epsilon_1, \epsilon_2\rangle\omega_2 \cdot \varphi = \left(s, Xv + \frac{1}{2}\langle\nabla_X\epsilon_1, \epsilon_2\rangle\chi_2(\omega_2)v\right) \\ &= \left(s, Xv + \frac{1}{2}\langle\nabla_X\epsilon_1, \epsilon_2\rangle kv\right).\end{aligned}$$

From now on, we will abuse notations and write ∇_Xv instead, dropping the spinor frame part s when there is no ambiguity. Using the connection coefficients,

$$\begin{aligned}\nabla_Xv &= e^{-\rho}(X_1v_x + X_2v_y) + \frac{e^{-\rho}}{2}(\rho_xX_2 - \rho_yX_1)kv \\ -\frac{1}{2}(SX) \cdot \omega_2 \cdot v &= \frac{1}{2}((h_{11}X_1 + h_{12}X_2)j - (h_{21}X_1 + h_{22}X_2)i)v \\ X \cdot v &= \chi_2(X)v = (X_1i + X_2j)v \\ T \cdot v + \lambda\omega_2 \cdot v &= (T_1i + T_2j)v + \lambda kv\end{aligned}$$

Finally

$$\begin{aligned}X_1v_x + X_2v_y &= -\frac{1}{2}(\rho_xX_2 - \rho_yX_1)kv \\ &+ \frac{e^\rho}{2}((h_{11}X_1 + h_{12}X_2)j - (h_{21}X_1 + h_{22}X_2)i)v \\ &+ \frac{\tau e^\rho}{2}(X_1i + X_2j)v - \tau e^\rho\langle X, T\rangle((T_1i + T_2j)v + \lambda kv).\end{aligned}$$

Specializing to $\partial/\partial x$ and $\partial/\partial y$

$$\begin{cases} v_x = \left(\frac{\rho y}{2} k + e^\rho \frac{h_{11}}{2} j - e^\rho \frac{h_{21}}{2} i + \frac{\tau e^\rho}{2} i - \tau e^\rho T_1 (T_1 i + T_2 j + \lambda k) \right) v \\ v_y = \left(-\frac{\rho x}{2} k + e^\rho \frac{h_{12}}{2} j - e^\rho \frac{h_{22}}{2} i + \frac{\tau e^\rho}{2} j - \tau e^\rho T_2 (T_1 i + T_2 j + \lambda k) \right) v \end{cases} \quad (\text{C}')$$

Note 3. Since $v_x v^{-1}$ and $v_y v^{-1}$ both lie in $\text{Im } \mathbb{H}$, $|v|$ is clearly constant. From now on, any solution v of (C') will therefore be normalized to norm 1.

Let us now write $v = (v^+, v^-)$ in the $\Sigma_2^+ \oplus \Sigma_2^-$ splitting, so that left quaternionic multiplication (i.e. Clifford multiplication) acts as a skew-hermitian matrix, e.g.

$$(T_1 i + T_2 j + \lambda k)v = i \begin{pmatrix} (T_1 - iT_2)v^- \\ (T_1 + iT_2)v^+ \end{pmatrix} - \lambda i \begin{pmatrix} v^+ \\ -v^- \end{pmatrix}.$$

Then

$$\begin{cases} v_z^+ &= -\frac{\rho z}{2} v^+ + \frac{e^\rho}{2} (H + i\tau\lambda^2)v^- + \frac{i\tau\lambda e^\rho}{2} (T_1 + iT_2)v^+ \\ v_z^- &= -\frac{\rho z}{2} v^- + \frac{e^\rho}{2} (-H + i\tau\lambda^2)v^+ - \frac{i\tau\lambda e^\rho}{2} (T_1 - iT_2)v^- \end{cases} \quad (9)$$

which differs quite a bit from (8), unless $\tau = 0$ (immersions into \mathbb{R}^3). That is not so surprising, since the abstract equation (C) states the existence of an immersion for which φ is induced from *any* constant spinor, while Konopelchenko & Taimanov's equation (DP) implies the same result with φ induced from the *specific* constant spinor field (s, u_0) , namely $u_0 = (1 + i - j + k)/2$. Indeed, because $ku_0 = u_0 i$, we know that the ambient spinor $\tilde{\varphi}$ satisfies

$$E_3 \cdot \tilde{\varphi} = (\tilde{s}, k) \cdot (\tilde{s}, u_0) = (\tilde{s}, ku_0) = (\tilde{s}, u_0 i)$$

so for the induced spinor

$$(E_3 \cdot \tilde{\varphi})|_M = E_3 \cdot \varphi = (T + \lambda\omega_2) \cdot (s, v) = (s, \chi_2(T + \lambda\omega_2)v) = (s, vi),$$

in vector notation,

$$i \begin{pmatrix} (T_1 - iT_2)v^- \\ (T_1 + iT_2)v^+ \end{pmatrix} - \lambda i \begin{pmatrix} v^+ \\ -v^- \end{pmatrix} = i \begin{pmatrix} v^+ \\ v^- \end{pmatrix}$$

or the two (equivalent) colinearity relations

$$\begin{cases} (T_1 - iT_2)v^- = (1 + \lambda)v^+ \\ (T_1 + iT_2)v^+ = (1 - \lambda)v^- \end{cases}$$

Substituting in (9), we obtain

$$\begin{cases} v_z^+ &= -\frac{\rho z}{2} v^+ + \frac{e^\rho}{2} (H + i\tau\lambda)v^- \\ v_z^- &= -\frac{\rho z}{2} v^- - \frac{e^\rho}{2} (H + i\tau\lambda)v^+ \end{cases}$$

exactly the equation (8). In other words, (C) is more general, while (DP) is simpler because it assumes implicitly a given choice of spinor coordinates.

We give now another proof of theorem 1, showing that equation (C') implies the existence of an immersion, for which φ is the restriction of a constant ambient spinor field $\tilde{\varphi} = (\tilde{s}, u)$. We do so without going through the full construction of [R].

The key idea is as follows: if $\varphi = (s, v)$ is the restriction of some constant spinor field $\tilde{\varphi} = (\tilde{s}, u)$, it satisfies the relation

$$E_3 \cdot \tilde{\varphi} = (\tilde{s}, k) \cdot (\tilde{s}, u) = (\tilde{s}, ku) = (\tilde{s}, uq)$$

where $q = u^{-1}ku$. We need the following lemmas, that prepares for the construction of the constants q and u .

Lemma 1. *For any solution v of (C'), where T, λ follow Daniel's equations (6), there exists an imaginary constant q such that*

$$(T_1i + T_2j + \lambda k)v = vq. \quad (10)$$

Proof. The first equation in (6) is written as:

$$\begin{aligned} \nabla_X T &= e^{-\rho}((X_1T_{1x} + X_2T_{1y} + T_2(X_1\rho_y - X_2\rho_x))e_1 \\ &\quad + (X_1T_{2x} + X_2T_{2y} + T_1(-X_1\rho_y + X_2\rho_x))e_2) \\ &= \lambda(SX - \tau JX) = \lambda \left(\begin{pmatrix} h_{11}X_1 + h_{12}X_2 \\ h_{21}X_1 + h_{22}X_2 \end{pmatrix} - \tau \begin{pmatrix} -X_2 \\ X_1 \end{pmatrix} \right) \end{aligned}$$

specializing

$$\begin{cases} T_{1x} = \lambda e^\rho h_{11} - T_2\rho_y \\ T_{1y} = \lambda e^\rho (h_{12} + \tau) + T_2\rho_x \\ T_{2x} = \lambda e^\rho (h_{21} - \tau) + T_1\rho_y \\ T_{2y} = \lambda e^\rho h_{22} - T_1\rho_x \end{cases}$$

The second equation in (6) reads

$$\begin{cases} \lambda_x = -e^\rho \left\langle \begin{pmatrix} h_{11} \\ h_{21} \end{pmatrix} - \tau \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \begin{pmatrix} T_1 \\ T_2 \end{pmatrix} \right\rangle = -e^\rho (T_1 h_{11} + T_2 (h_{21} - \tau)) \\ \lambda_y = -e^\rho \left\langle \begin{pmatrix} h_{12} \\ h_{22} \end{pmatrix} - \tau \begin{pmatrix} -1 \\ 0 \end{pmatrix}, \begin{pmatrix} T_1 \\ T_2 \end{pmatrix} \right\rangle = -e^\rho (T_1 (h_{12} + \tau) + T_2 h_{22}) \end{cases}$$

Set $E = T_1i + T_2j + \lambda k$,

$$\frac{\partial E}{\partial x} = e^\rho (h_{11}(\lambda i - T_1k) + (h_{21} - \tau)(\lambda j - T_2k)) + \rho_y (T_1j - T_2i)$$

Recall from (C') that $v_x = Vv$ for some imaginary quaternion valued function V ; then

$$\frac{\partial \bar{v} E v}{\partial x} = \bar{v} \left(\bar{V} E + \frac{\partial E}{\partial x} + E V \right) v$$

It is long but straightforward to show that $\bar{V}E + \frac{\partial E}{\partial x} + EV$ vanishes identically. The same computations holds for the y variable. Clearly $q \in \text{Im } \mathbb{H}$. \square

Note 4. The equation (C') could accordingly be simplified using (10), eliminating T in the process. However, because right quaternionic multiplication does not commute with the complex structure, the new equation may not be as simple as (8), for which $q = i$.

The mere existence of u implies the following relation for solutions of (C'):

Lemma 2. *For any solution v of (C') assuming (6)*

$$\lambda \bar{v}(T_1 i + T_2 j + \lambda k)v = \langle \bar{v}kv, q \rangle q.$$

Proof.

$$\langle \bar{v}kv, q \rangle = \langle v^{-1}kv, v^{-1}(T_1 i + T_2 j + \lambda k)v \rangle = \langle k, T_1 i + T_2 j + \lambda k \rangle = \lambda$$

so

$$\langle \bar{v}kv, q \rangle q - \lambda \bar{v}(T_1 i + T_2 j + \lambda k)v = \lambda q - \lambda |v|^2 q = 0.$$

\square

We conclude the proof of the theorem by giving an immersion via its Maurer–Cartan form. Using the notations (7) of section 2.3, we need to prove that the $\text{Im } \mathbb{H}$ -valued 1-form

$$\alpha = e^\rho u v^{-1} (i dx + j dy) v u^{-1}$$

is the Maurer–Cartan form $f^{-1}df$ of some immersion into $\text{Nil}_3(\tau)$, where u is a unit quaternion such that $u^{-1}ku = q$, obtained through lemma 1 (u is not unique). The necessary and sufficient condition for local existence is again the integrability equation:

$$\begin{aligned} 0 &= -\frac{\partial}{\partial y}(e^\rho u v^{-1} i v u^{-1}) + \frac{\partial}{\partial x}(e^\rho u v^{-1} j v u^{-1}) + [e^\rho u v^{-1} i v u^{-1}, e^\rho u v^{-1} j v u^{-1}] \\ &= e^\rho u (\rho_x \bar{v} j v + \bar{v}_x j v + \bar{v} j v_x - \rho_y \bar{v} i v - \bar{v}_y i v - \bar{v} i v_y \\ &\quad + 2\tau e^\rho \langle \bar{v}kv, u^{-1}ku \rangle u^{-1}ku) u^{-1}. \end{aligned}$$

(We have used the relation of the Lie bracket in $\mathfrak{nil}_3(\tau) \simeq \text{Im } \mathbb{H}$ with the quaternionic multiplication and scalar product: $[q, q'] = 2\tau \langle qq', k \rangle k$.) Factorizing and

substituting from (C'), and using $T_1^2 + T_2^2 + \lambda^2 = 1$,

$$\begin{aligned}
0 &= \rho_x \bar{v} j v + \bar{v} \left(-\frac{\rho_y}{2} k - e^\rho \frac{h_{11}}{2} j + e^\rho \frac{h_{21}}{2} i - \frac{\tau e^\rho}{2} i + \tau e^\rho T_1 (T_1 i + T_2 j + \lambda k) \right) j v \\
&\quad + \bar{v} j \left(\frac{\rho_y}{2} k + e^\rho \frac{h_{11}}{2} j - e^\rho \frac{h_{21}}{2} i + \frac{\tau e^\rho}{2} i - \tau e^\rho T_1 (T_1 i + T_2 j + \lambda k) \right) v \\
&\quad - \rho_y \bar{v} i v - \bar{v} \left(\frac{\rho_x}{2} k - e^\rho \frac{h_{12}}{2} j + e^\rho \frac{h_{22}}{2} i - \frac{\tau e^\rho}{2} j + \tau e^\rho T_2 (T_1 i + T_2 j + \lambda k) \right) i v \\
&\quad - \bar{v} i \left(-\frac{\rho_x}{2} k + e^\rho \frac{h_{12}}{2} j - e^\rho \frac{h_{22}}{2} i + \frac{\tau e^\rho}{2} j - \tau e^\rho T_2 (T_1 i + T_2 j + \lambda k) \right) v \\
&\quad + 2\tau e^\rho \langle \bar{v} k v, u^{-1} k u \rangle u^{-1} k u \\
&= 2\tau e^\rho (-\lambda \bar{v} (T_1 i + T_2 j + \lambda k) v + \langle \bar{v} k v, u^{-1} k u \rangle u^{-1} k u)
\end{aligned}$$

which vanishes according to lemma 2. This implies the local existence of an immersion in $\text{Nil}_3(\tau)$ (including the special case of \mathbb{R}^3 when $\tau = 0$), for which φ is the restriction of an ambient spinor.

Note 5. A ‘‘colinearity’’ relation similar to (10) appears naturally when using the *spin*^c formalism in Nil_3

$$\xi \cdot \varphi = (T + \lambda v) \cdot \varphi = i\varphi$$

(see [NR] for details).

3 Conformally parametrized surfaces in \mathbb{R}^4

Recently, Bayard, Lawn & Roth [BLR] have characterized isometric immersions of surfaces in \mathbb{R}^4 by means of spinor fields, giving an spinorial analog of the Gauss–Ricci–Codazzi equations. More precisely, they show the following theorem:

Theorem 2 (Bayard, Lawn & Roth). *Given a Killing spinor $\tilde{\varphi}$ with Killing constant λ in $\mathbb{M}^4(4\lambda^2)$ (the real space form of curvature $4\lambda^2$) and an immersed surface M^2 , its restriction φ to M lies in the twisted spinor bundle $\Sigma = \Sigma M \otimes \Sigma E$ (E being the normal bundle) and satisfies the Dirac equation*

$$D\varphi = \vec{H} \cdot \varphi - 2\lambda\varphi \tag{11}$$

for the Dirac operator D along M . Conversely, if (M^2, g) is an oriented riemannian manifold, with given spin structure and rank 2 oriented spin vector bundle E on M , then the following properties are equivalent, where φ is a section of the twisted spinor bundle $\Sigma = \Sigma M \otimes \Sigma E$:

1. φ is a solution of the Dirac equation (11), such that φ^+, φ^- do not vanish and

$$X|\varphi^+| = 2 \text{Re}\langle \lambda X \cdot \varphi^-, \varphi^+ \rangle \quad \text{and} \quad X|\varphi^-| = 2 \text{Re}\langle \lambda X \cdot \varphi^+, \varphi^- \rangle,$$

2. $\nabla_X \varphi = \lambda X \cdot \varphi - \frac{1}{2} \sum_1^2 e_j \cdot \mathcal{B}(X, e_j) \cdot \varphi$ for some symmetric bilinear \mathcal{B} section of E , φ^+, φ^- do not vanish,
3. there exists a local immersion of M with mean curvature $\vec{H} = \frac{1}{2} \text{tr} \mathcal{B}$, second fundamental form \mathcal{B} , normal bundle E into $\mathbb{M}^4(4\lambda^2)$.

The difficult part lies in the converse. Moreover they explicit the reconstruction of f in terms of φ , a procedure now known by the generic term of Weierstrass formula. It must be stressed though that this proof is entirely abstract.

Partially motivating [BLR], in \mathbb{R}^4 only, we have seen in section 1 the *spinor representation formula* (W4) of Konopelchenko and Taimanov, with the condition (1), also named ‘‘Dirac equation’’. We will now bridge the gap between these two approaches, by rewriting [BLR] using concrete spinor coordinates. We will prove directly (without actually resorting to the conditions for isometric immersion of Gauss, Codazzi and Ricci) the existence of an immersion f derived from a spinor φ , and connect equation (11) to (1). Additionally, it will be clear how one recovers a parallel spinor associated to f from the immersion itself. As in the previous sections, we will use quaternions, which play a natural role in \mathbb{R}^4 , and allow for shorter and more elegant formulations. In particular, we express equations (W4) and (1) as follows (a formalism due to [H]). Let f be a conformal immersion $f : M \rightarrow \mathbb{R}^4 \simeq \mathbb{H}$. Since $\text{Spin}(4) \simeq \mathbb{S}^3 \times \mathbb{S}^3 \subset \mathbb{H}^2$ acts on \mathbb{H} via

$$\forall (p, q) \in \mathbb{S}^3 \times \mathbb{S}^3, \forall x \in \mathbb{H}, \quad \theta_4(p, q) : x \mapsto pxq^{-1} = px\bar{q}$$

and $(\partial f/\partial x, \partial f/\partial y)$ is an orthonormal basis, up to a multiplicative factor e^ρ , there exist $A, B : M \rightarrow \mathbb{H}^*$ such that

$$df = \bar{A}dzB = \bar{A}(dx + idy)B = \bar{A}Bdx + \bar{A}iBdy \quad (\text{D})$$

and the conformal factor is $e^\rho = |\bar{A}B|$. Conversely such an expression can be integrated locally into a function f if and only if

$$\begin{aligned} 0 &= d(df) = ((\bar{A}iB)_x - (\bar{A}B)_y)dx \wedge dy \\ &= \bar{A} \left[\bar{A}^{-1} \left(\frac{\partial \bar{A}}{\partial x} + \frac{\partial \bar{A}}{\partial y} i \right) i + i \left(\frac{\partial B}{\partial x} + i \frac{\partial B}{\partial y} \right) B^{-1} \right] B dx \wedge dy \end{aligned} \quad (12)$$

i.e.

$$\bar{A}^{-1}(\partial \bar{A}/\partial \bar{z})i + i(\partial \bar{z} \setminus \partial B)B^{-1} = 0 \quad (13)$$

Note 6. We use here the quaternionic notation for complex derivatives, with $dz = dx + idy$ (and $i \in \mathbb{H}$)

$$df = (\partial f/\partial z)dz + (\partial f/\partial \bar{z})d\bar{z} = dz(\partial z \setminus \partial f) + d\bar{z}(\partial \bar{z} \setminus \partial f)$$

equivalently

$$\begin{aligned}\partial f/\partial z &= \frac{1}{2} \left(\frac{\partial f}{\partial x} - \frac{\partial f}{\partial y} i \right), \quad \partial f/\partial \bar{z} = \frac{1}{2} \left(\frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} i \right), \\ \partial z \backslash \partial f &= \frac{1}{2} \left(\frac{\partial f}{\partial x} - i \frac{\partial f}{\partial y} \right), \quad \partial \bar{z} \backslash \partial f = \frac{1}{2} \left(\frac{\partial f}{\partial x} + i \frac{\partial f}{\partial y} \right).\end{aligned}$$

If we set $A = \bar{s}_1 - js_2$, $B = t_1 + t_2j$ (the reason for which is explained below), then $df = \bar{A}dzB$, yields exactly the ‘‘Weierstrass formula’’ (W4)

$$\bar{A}dzB = (s_1t_1dz - \bar{s}_2\bar{t}_2d\bar{z}) + (s_1t_2dz + \bar{s}_2\bar{t}_1d\bar{z})j$$

when writing $\mathbb{R}^4 \simeq \mathbb{C}^2 \simeq \mathbb{C} \oplus \mathbb{C}j = \mathbb{H}$. Furthermore, the Dirac equation (1) amounts to

$$(\partial z \backslash \partial A)A^{-1} = -hj \quad \text{and} \quad (\partial \bar{z} \backslash \partial B)B^{-1} = hj \quad (\text{I})$$

for some (nonconstant) complex potential h . Whenever the last equation is satisfied, the integrability condition (13) holds immediately:

$$j\bar{h}i + ihj = h(ji + ij) = 0.$$

The converse is false: (13) does not imply (I) in general; however one can use the gauge freedom $(A, B) \mapsto (e^{i\alpha}A, e^{i\alpha}B)$ (already noted in [T1]). Using that freedom, there is always a choice of (A, B) , such that $(\partial z \backslash \partial A)A^{-1}$ lies in $\mathbb{C}j$ (and therefore $(\partial \bar{z} \backslash \partial B)B^{-1}$ as well, because of (13)). They are actually opposite, and (I) holds. This choice is unique up to a constant: if $(\partial \bar{z} \backslash \partial B)B^{-1} \in \text{Span}(j, k)$, then

$$(\partial \bar{z} \backslash \partial (e^{i\alpha}B))(e^{i\alpha}B)^{-1} = e^{i\alpha} \left((\partial \bar{z} \backslash \partial B)B^{-1} + \frac{\partial \alpha}{\partial \bar{z}} \right)$$

lies in $\text{Span}(j, k)$ if and only if $\frac{\partial \alpha}{\partial \bar{z}} = 0$.

3.1 Spinors in two and four dimensions

In two dimensions, we choose as above for the Clifford representation of $\mathbb{R}^2 = \mathbb{R}e_1 \oplus \mathbb{R}e_2$ on $\Sigma_2 = \mathbb{H}$ the map $\chi_2: \chi_2(e_1) = L_i$ and $\chi_2(e_2) = L_j$, where L_i, L_j denote the left quaternionic multiplication by i and j respectively. However, for compatibility purposes with the above equations, we endow \mathbb{H} with the complex structure $J = R_j$, the right multiplication by j . This allows to equate \mathbb{H} with \mathbb{C}^2 , but complex numbers in this setting are linear combinations over \mathbb{R} of quaternions 1 and j . To emphasize this (purely formal) difference and make notations clear, we will write when needed $\hat{\mathbb{C}} = \mathbb{R} \oplus j\mathbb{R} \simeq \mathbb{C}$, so that $\mathbb{H} \simeq \hat{\mathbb{C}}^2$ (6).

⁶Because of the non-commutativity of \mathbb{H} , all vector spaces over $\hat{\mathbb{C}}$ are vector spaces on the right.

As usual, Σ_2 splits into $\Sigma_2^+ \oplus \Sigma_2^-$, eigenspaces of $\chi_2(\omega_2^{\mathbb{C}})$ w.r.t. the eigenvalues $+1$ and -1 ; $\omega_2^{\mathbb{C}} = J e_1 \cdot e_2$ is the complex volume, and is mapped by χ_2 to $L_i L_j R_j = L_k R_j$, whose eigenspaces are

$$\Sigma_2^+ = (1 - i)\hat{\mathbb{C}}, \quad \Sigma_2^- = (1 + i)\hat{\mathbb{C}}.$$

The euclidean space \mathbb{R}^4 is identified with \mathbb{H} (which amounts to taking $1, i, j, k$ as reference basis), on which $\text{Spin}(4) \simeq \mathbb{S}^3 \times \mathbb{S}^3 \subset \mathbb{H} \times \mathbb{H}$ acts as double cover of $SO(4)$ via the classical representation $\theta: \theta(p, q) \cdot x = pxq^{-1} = px\bar{q}$. The Clifford algebra is represented on $\Sigma_4 = \mathbb{H}^2$ via χ_4 , which we define by its restriction to $\mathbb{R}^4 = \mathbb{H}$:

$$\chi_4(x) = \begin{pmatrix} 0 & x \\ -\bar{x} & 0 \end{pmatrix}$$

One checks directly that χ_4 induces the following representation of $\text{Spin}(4)$: $\chi_4((p, q)) = \begin{pmatrix} p & 0 \\ 0 & q \end{pmatrix}$. Note that $\chi_4(x)$ is a complex-linear endomorphism, provided \mathbb{H}^2 is also endowed with J , the right multiplication by j , as above, so that $\mathbb{H}^2 \simeq \hat{\mathbb{C}}^4$. The representation decomposes into two irreducible subspaces, which are here $\Sigma_4^+ = \mathbb{H} \oplus 0$ and $\Sigma_4^- = 0 \oplus \mathbb{H}$. A spinor $\tilde{\varphi}$ on \mathbb{R}^4 is now the equivalence class of couples $((p, q), (a, b))$ where $(p, q) \in \text{Spin}(4) = \mathbb{S}^3 \times \mathbb{S}^3$ and $(a, b) \in \Sigma_4 = \mathbb{H}^2$, under the following equivalence relation: for all $(p_1, q_1) \in \mathbb{S}^3 \times \mathbb{S}^3$

$$((p, q), (a, b)) \sim ((pp_1^{-1}, qq_1^{-1}), \chi_4(p_1, q_1)(a, b)) = ((pp_1^{-1}, qq_1^{-1}), (p_1 a, q_1 b))$$

and a spinor field is a section of the quotient bundle $\text{Spin}(4) \times_{\chi_4} \Sigma_4$. Finally, because \mathbb{R}^4 is globally trivial, one can always write $\tilde{\varphi}$ in a preferred spinor frame, e.g. $(1, 1)$.

The starting point of [BLR] is the following remark: let M be a two-dimensional submanifold, and $\tilde{\varphi}$ be an ambient parallel⁷ spinor, i.e. $\tilde{\nabla}\tilde{\varphi} = 0$, where $\tilde{\nabla}$ denotes the covariant derivative of \mathbb{R}^4 . Define on M the Dirac operator by $D = e_1 \cdot \nabla_{e_1} + e_2 \cdot \nabla_{e_2}$, where now ∇ is the covariant derivative on M (here and in the following, e_1, e_2 will denote any orthonormal frame). Then a simple application of the spinorial Gauss equation

$$\tilde{\nabla}_X \tilde{\varphi} = \nabla_X \tilde{\varphi} + \frac{1}{2} \sum_{j=1}^2 e_j \cdot \mathcal{B}(X, e_j) \cdot \tilde{\varphi}$$

yields (11).

To make sense of the converse result, starting from intrinsic data, one cannot just consider the restriction $\tilde{\varphi}|_M$ as a spinor on M (unlike in the three-dimensional case), since the spinor spaces Σ_2 and Σ_4 have different dimensions.

⁷In \mathbb{R}^4 , a Killing spinor is simply parallel.

Instead, one considers spinors on M and spinors on the normal bundle E . Let us briefly recall the idea of Bär ([Bä], see also [G] for a quick introduction) and consider $\mathbb{R}^2 \oplus \mathbb{R}^2$. Each factor is equipped with its Clifford representation as above, which we denote by χ_2 and χ'_2 on Σ_2 and Σ'_2 (they are identical, the prime just makes it clear which one we are considering). We let γ be the map $\gamma = \chi_2 \otimes \chi'_2(\omega_2^{\mathbb{C}}) \oplus \text{Id}_{\Sigma_2} \otimes \chi'_2$ from $\mathbb{R}^4 = \mathbb{R}^2 + \mathbb{R}^2$ on $\Sigma_2 \otimes \Sigma'_2$. For $u + v \in \mathbb{R}^2 \oplus \mathbb{R}^2$, $\gamma(u + v)^2 = -(|u|^2 + |v|^2)\text{id}$, hence γ extends to a representation of the Clifford algebra $Cl(\mathbb{R}^4)$ on $\Sigma_2 \otimes \Sigma'_2$. by the universal property, γ is equivalent to χ_4 : there exists a isomorphism of algebra $\Psi : \Sigma_2 \otimes \Sigma'_2 \rightarrow \Sigma_4$ making the diagram commutative

$$\begin{array}{ccc} \Sigma_2 \otimes \Sigma'_2 & \xrightarrow{\gamma(x)} & \Sigma_2 \otimes \Sigma'_2 \\ \downarrow \Psi & & \downarrow \Psi \\ \Sigma_4 & \xrightarrow{\chi_4(x)} & \Sigma_4 \end{array}$$

We shall make this correspondence explicit below.

We can now define the *twisted spinor bundle* Σ on a surface M with a rank two vector bundle E , by taking sections in $\Sigma M \otimes \Sigma E$. More precisely, let $\sigma = (s, u)$ be a spinor field on M , i.e. a moving spinor frame s (over an orthogonal frame e_1, e_2 of TM) and a vector field u on Σ_2 , up to the equivalence relation $(s, u) \sim (sa^{-1}, \chi_2(a)u)$; similarly let $\sigma' = (s', u')$ be the equivalence class of a spinor frame s' of E (over an orthogonal frame e_3, e_4 of E) and u' a vector field on Σ'_2 . Then, when $M \subset \mathbb{R}^4$, the couple (s, s') identifies canonically with a spinor frame of \mathbb{R}^4 (over the frame (e_1, \dots, e_4)) and an element $\sigma \otimes \sigma'$ identifies with (the class of) $((s, s'), \Psi(u \otimes u'))$. This allows to identify ambient spinors $\tilde{\varphi}$ restricted to M with sections of $\Sigma M \otimes \Sigma E$, which are intrinsic objects.

The identification $\Sigma_2 \otimes \Sigma'_2 \simeq \Sigma_4$ goes as follows, using the quaternionic notations. From the natural basis of $\Sigma_2 \simeq \mathbb{H}$ over \mathbb{C} : $(1 - i), (1 + i)$, we build the following basis

$$(1 - i) \otimes (1 - i), (1 + i) \otimes (1 + i), (1 - i) \otimes (1 + i), (1 + i) \otimes (1 - i)$$

which actually spans $(\Sigma_2^+ \otimes \Sigma'^+ + \Sigma_2^- \otimes \Sigma'^-) \oplus (\Sigma_2^+ \otimes \Sigma'^-) \oplus (\Sigma_2^- \otimes \Sigma'^+)$. The corresponding basis in $\Sigma_4 \simeq \mathbb{H}^2$ spans $\Sigma_4^{++} \oplus \Sigma_4^{--} \oplus \Sigma_4^{+-} \oplus \Sigma_4^{-+}$, where $(a, b) \in \Sigma_4^{\varepsilon\varepsilon'}$ if and only if $\chi_4(\omega_2^{\mathbb{C}})(a, b) = (\varepsilon a, \varepsilon' b)$. Then a possible (but not unique) choice of basis is

$$\begin{pmatrix} 1+k \\ 0 \end{pmatrix}, \begin{pmatrix} -i-j \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ i+j \end{pmatrix}, \begin{pmatrix} 0 \\ -1-k \end{pmatrix}$$

and Ψ is the map sending on basis to the other:

$$\begin{aligned} & \Psi((1-i) \otimes (1-i)a^{++} + (1+i) \otimes (1+i)a^{--} + (1-i) \otimes (1+i)a^{+-} + (1+i) \otimes (1-i)a^{-+}) \\ &= \begin{pmatrix} (1+k)a^{++} - (i+j)a^{--} \\ (i+j)a^{+-} - (1+k)a^{-+} \end{pmatrix} = \begin{pmatrix} a \\ b \end{pmatrix} \quad (14) \end{aligned}$$

3.2 The abstract Dirac equation

Using this correspondence we can write the Dirac equation, *using only intrinsic data* on M and a given rank two bundle E . We will assume that M is parametrized locally by the conformal coordinate $z = x + iy$, inducing a frame $(e_1, e_2) = e^{-\rho}(\partial/\partial x, \partial/\partial y)$, where ρ is the conformal factor, and we will also denote

$$c_x = \langle \nabla_{\partial/\partial x}^E e_3, e_4 \rangle = e^\rho \langle \nabla_{e_1}^E e_3, e_4 \rangle, \quad c_y = \langle \nabla_{\partial/\partial y}^E e_3, e_4 \rangle = e^\rho \langle \nabla_{e_1}^E e_3, e_4 \rangle$$

the normal connection coefficients, and

$$\text{for } \beta = 1, 2 \quad H_\beta = \frac{1}{2} \langle \mathcal{B}(e_1, e_1) + \mathcal{B}(e_2, e_2), e_\beta \rangle$$

the coefficients which will (eventually) be the mean curvature vector coordinates, in a chosen orthonormal frame (e_3, e_4) of E . Since ∇^M, ∇^E (resp. $\nabla^{\Sigma M}, \nabla^{\Sigma E}$) are the connections (resp. spinorial connections) on M and E , then $\nabla = \nabla^{\Sigma M} \otimes \text{Id}_{\Sigma E} + \text{Id}_{\Sigma M} \otimes \nabla^{\Sigma E}$ is the connection on the twisted spinor bundle Σ . For $\sigma \in \Gamma(\Sigma M)$,

$$\nabla_X^{\Sigma M} \sigma = L_X \sigma + \frac{1}{2} \sum_{\alpha < \beta} \langle \nabla_X^M e_\alpha, e_\beta \rangle e_\alpha \cdot e_\beta \cdot \sigma = L_X \sigma + \frac{1}{2} \langle \nabla_X^M e_1, e_2 \rangle e_1 \cdot e_2 \cdot \sigma$$

so that

$$\begin{aligned} \nabla_{e_1}^{\Sigma M} (1 - \varepsilon i) &= \underbrace{L_{e_1} (1 - \varepsilon i)}_{=0} + \frac{1}{2} \langle \nabla_{e_1}^M e_1, e_2 \rangle i j (1 - \varepsilon i) \\ &= -\frac{e^{-\rho}}{2} \frac{\partial \rho}{\partial y} \varepsilon (1 - \varepsilon i) (-j) \end{aligned}$$

Similarly,

$$\begin{aligned} \nabla_{e_2}^{\Sigma M} (1 - \varepsilon i) &= \frac{e^{-\rho}}{2} \frac{\partial \rho}{\partial x} \varepsilon (1 - \varepsilon i) (-j) \\ \nabla_{e_1}^{\Sigma E} (1 - \varepsilon' i) &= \frac{e^{-\rho}}{2} c_1 \varepsilon' (1 - \varepsilon' i) (-j) \\ \nabla_{e_2}^{\Sigma E} (1 - \varepsilon' i) &= \frac{e^{-\rho}}{2} c_2 \varepsilon' (1 - \varepsilon' i) (-j) \end{aligned}$$

Let $D = e_1 \cdot \nabla_{e_1} + e_2 \cdot \nabla_{e_2}$ be the Dirac operator on $\Sigma M \otimes \Sigma E$, then,

$$D((1 - \varepsilon i) \otimes (1 - \varepsilon' i)) = \frac{e^{-\rho}}{2} (1 + \varepsilon i) \otimes (1 - \varepsilon' i) \left(\varepsilon \varepsilon' \frac{\partial \rho}{\partial x} + c_2 + \left(\varepsilon' \frac{\partial \rho}{\partial y} - \varepsilon c_1 \right) j \right)$$

and for any $\hat{\mathbb{C}}$ -valued a ,

$$\begin{aligned} D((1 - \varepsilon i) \otimes (1 - \varepsilon' i) a) &= \\ (1 + \varepsilon i) \otimes (1 - \varepsilon' i) e^{-\rho} &\left(\frac{c_2 + \varepsilon \varepsilon' \rho_x + (\varepsilon' \rho_y - c_1 \varepsilon) j}{2} a + \varepsilon \varepsilon' \frac{\partial a}{\partial x} + \varepsilon' j \frac{\partial a}{\partial y} \right) \quad (15) \end{aligned}$$

Summing up,

$$\begin{aligned}
D\varphi &= D((1-i) \otimes (1-i)a^{++} + (1+i) \otimes (1+i)a^{--} \\
&\quad + (1-i) \otimes (1+i)a^{+-} + (1+i) \otimes (1-i)a^{-+}) \\
&= (1-i) \otimes (1-i)e^{-\rho} \left(\frac{c_y - \rho_x + (\rho_y + c_x)j}{2} a^{-+} - a_x^{-+} + ja_y^{-+} \right) \\
&\quad + (1+i) \otimes (1+i)e^{-\rho} \left(\frac{c_y - \rho_x + (-\rho_y - c_x)j}{2} a^{+-} - a_x^{+-} - ja_y^{+-} \right) \\
&\quad + (1-i) \otimes (1+i)e^{-\rho} \left(\frac{c_y + \rho_x + (-\rho_y + c_x)j}{2} a^{--} + a_x^{--} - ja_y^{--} \right) \\
&\quad + (1+i) \otimes (1-i)e^{-\rho} \left(\frac{c_y + \rho_x + (\rho_y - c_x)j}{2} a^{++} + a_x^{++} + ja_y^{++} \right)
\end{aligned}$$

Switching to the notation in Σ_4 , through the isomorphism Ψ in (14) (which we will be implicit), we now write $\varphi = (a, b)$, so

$$D\varphi = D \begin{pmatrix} a \\ b \end{pmatrix} = e^{-\rho} \begin{pmatrix} \frac{\partial b}{\partial x} + i \frac{\partial b}{\partial y} + \left(\frac{ic_x - cy}{2} + \frac{\partial \rho}{\partial \bar{z}} \right) b \\ -\frac{\partial a}{\partial x} + i \frac{\partial a}{\partial y} + \left(-\frac{ic_x + cy}{2} - \frac{\partial \rho}{\partial z} \right) a \end{pmatrix}.$$

Since

$$\vec{H} \cdot \varphi = \begin{pmatrix} 0 & H_3j + H_4k \\ H_3j + H_4k & 0 \end{pmatrix} \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} (H_3j + H_4k)b \\ (H_3j + H_4k)a \end{pmatrix}$$

the Dirac equation is written intrinsically as

$$\begin{cases} \left(\frac{\partial b}{\partial x} + i \frac{\partial b}{\partial y} \right) b^{-1} = -\frac{\partial \rho}{\partial \bar{z}} + \frac{c_y - ic_x}{2} + e^\rho (H_3j + H_4k) \\ \left(\frac{\partial a}{\partial x} - i \frac{\partial a}{\partial y} \right) a^{-1} = -\frac{\partial \rho}{\partial z} - \frac{c_y + ic_x}{2} - e^\rho (H_3j + H_4k) \end{cases} \quad (16)$$

Or, using the quaternionic notation for complex derivatives,

$$\begin{cases} 2(\partial z \setminus \partial a)a^{-1} = -\frac{\partial \rho}{\partial z} - \frac{c_y + ic_x}{2} - e^\rho (H_3j + H_4k) \\ 2(\partial \bar{z} \setminus \partial b)b^{-1} = -\frac{\partial \rho}{\partial \bar{z}} + \frac{c_y - ic_x}{2} + e^\rho (H_3j + H_4k) \end{cases} \quad (17)$$

3.3 Building the immersion

To show that (16) (or (17)) implies the local existence of an immersion with the prescribed metric data, let us define the following \mathbb{H} -valued 1-form on M

$$\xi = e^\omega a^{-1}(dx + idy)b$$

for the spinor field $\varphi = (a, b)$. As in [BLR], we prove the following

Theorem 3. *If (a, b) satisfy (16) with a and b of constant norm, then ξ is closed, and the (locally defined) application $f : M \rightarrow \mathbb{H}$ such that $df = \xi$ induces the original metric data. Furthermore, we obtain the restriction of a parallel ambient spinor $\tilde{\varphi}$ by setting $\tilde{\varphi} = ((p, q), (a, b))$ for (p, q) a moving spinor frame on M such that $pq^{-1} = e_1$, $piq^{-1} = e_2$, $pjq^{-1} = e_3$ and $pkq^{-1} = e_4$.*

We can prove closedness directly, however we will choose a slightly different route, in order to connect to the formalism of Konopelchenko and Taimanov.

Proposition 1. *If (a, b) satisfy (17) with a and b of constant norm, then the \mathbb{H} -valued 1-form $\xi = e^\omega a^{-1}(dx + idy)b$ can be written as $\xi = \bar{A} dz B$, with A, B satisfying (I). In particular ξ is closed, and the conclusions of theorem 3 hold.*

Proof. Note first that the couples (a, b) and (A, B) are different in nature: $|a|$ and $|b|$ are constant and equal by hypothesis (we will assume that $|a| = |b| = 1$ for simplicity), while A, B clearly are not, unless the induced metric is flat. Nevertheless, (A, B) can be derived from (a, b) by a process similar to a gauge change, up to the conformal factor. Indeed write

$$A = e^{u+iv+\rho/2}a, \quad B = e^{-u+iv+\rho/2}$$

where u, v are real-valued functions to be determined. Then

$$\bar{A} dz B = a^{-1} e^{u-iv+\rho/2} dz e^{-u+iv+\rho/2} b = e^\rho a^{-1} dz b = \xi.$$

To prove our claim that ξ is closed, we need only show that A, B satisfy the equation (I) for a suitable choice of u and v .

$$\begin{aligned} (\partial z \setminus \partial A) A^{-1} &= \frac{\partial(u+iv+\rho/2)}{\partial z} + e^{u+iv+\rho/2} (\partial z \setminus \partial a) a^{-1} e^{u+iv+\rho/2} \\ &= \frac{\partial u}{\partial z} + i \frac{\partial v}{\partial z} - \frac{c_y + ic_x}{4} - \frac{e^{\rho+2iv}}{2} (H_3 j + H_4 k) \end{aligned}$$

because complex numbers commute, and commute modulo conjugation with j and k . Similarly,

$$(\partial \bar{z} \setminus \partial B) B^{-1} = -\frac{\partial u}{\partial \bar{z}} + i \frac{\partial v}{\partial \bar{z}} + \frac{c_y - ic_x}{4} + \frac{e^{\rho+2iv}}{2} (H_3 j + H_4 k).$$

We want to solve in \mathbb{C}

$$\begin{cases} \frac{\partial u}{\partial z} + i \frac{\partial v}{\partial z} = \frac{c_y + ic_x}{4} \\ \frac{\partial u}{\partial \bar{z}} - i \frac{\partial v}{\partial \bar{z}} = \frac{c_y - ic_x}{4} \end{cases}$$

but the second equation is the conjugate of the first one. Under smoothness assumption (see [Hö, Theorem 4.2.5]), there is always a solution of the $\bar{\partial}$ problem

$$\frac{\partial(u-iv)}{\partial \bar{z}} = \frac{c_y - ic_x}{4}.$$

Given this solution, we remain with

$$(\partial z \setminus \partial A) A^{-1} = -hj \quad \text{and} \quad (\partial \bar{z} \setminus \partial B) B^{-1} = hj$$

where

$$h = \frac{e^{\rho+2iv}}{2}(H_3 + H_4i)$$

depends on the mean curvature vector. We have obtained exactly the Dirac equation of Konopelchenko and Taimanov, which ensures integrability, as we proved earlier.

The map f is an immersion because, by construction, A and B are not allowed to vanish. There remains to show that f induces on M and its normal bundle the same metric data we started with. Because of $df = e^\rho a^{-1} dzb$, the metric on M is clearly the one desired. For the normal bundle connection and second fundamental form, just pick up the constant spinor field $\tilde{\varphi} = ((a^{-1}, b^{-1}), (a, b)) \sim ((1, 1), (1, 1))$. We know from the beginning that $\tilde{\varphi}|_M$ satisfies the Dirac equation (11), which in intrinsic coordinates yields (16) for the induced metric and spinor coefficients a, b . So that a, b satisfy two differential systems, and we can identify the right hand side term by term, using the fact that \mathbb{H} is a vector space (so quaternions split along the basis $(1, i, j, k)$). E.g. $-\frac{\partial \rho}{\partial \bar{z}} + \frac{c_y - ic_x}{2}$ has to coincide with $-\frac{\partial \rho}{\partial \bar{z}} + \frac{\langle \nabla_{\partial/\partial y}^\perp e_3, e_4 \rangle - i \langle \nabla_{\partial/\partial x}^\perp e_3, e_4 \rangle}{2}$ for the induced normal covariant derivative, which proves that both connections are equal. The same holds for the mean curvature. \square

We therefore reach the same conclusion as [BLR]: The Dirac equation (11) on $\Sigma M \otimes \Sigma E$ for a spinor field φ such that $|\varphi^+| = |\varphi^-|$ is equivalent to the equations of Gauss, Codazzi and Ricci for surfaces in \mathbb{R}^4 .

References

- [A] R. Aiyama, Lagrangian surfaces in the complex 2-space, *Proceedings of the Fifth International Workshop on Differential Geometry (Taegu)* 25–29, 2000.
- [Bä] C. Bär, Extrinsic Bounds for Eigenvalues of the Dirac Operator, *Annals of Global Analysis and Geometry* 16(6):573–596, 1998.
- [BLR] P. Bayard, M.-A. Lawn & J. Roth, Spinorial Representation of Surfaces into 4-dimensional Space Forms, [arXiv:1210.7386](https://arxiv.org/abs/1210.7386), to appear in *Annals of Global Analysis and Geometry*.
- [D] B. Daniel, Isometric immersions into 3-dimensional homogeneous manifolds, *Commentarii Mathematici Helvetici* 82(1):87–131, 2007.
- [F1] T. Friedrich, On the spinor representation of surfaces in Euclidean 3-space, *Journal of Geometry and Physics* 28(1-2):143–157, 1998.
- [F2] T. Friedrich, Dirac operators in Riemannian Geometry, *Graduate studies in mathematics*, Volume 25, American Mathematical Society, 2000.

- [G] N. Ginoux, Opérateurs de Dirac sur les sous-variétés, *PhD thesis*, Université Henri Poincaré, Nancy 1, 2002.
- [H] F. Hélein, On Konopelchenko’s representation for surfaces in 4 dimensions, [arXiv:math.DG/0104101](https://arxiv.org/abs/math/0104101), 2001.
- [HRo] F. Hélein & P. Romon, Weierstrass representation of Lagrangian surfaces in four-dimensional space using spinors and quaternions, *Commentarii Mathematici Helvetici* 75(4):668–680, 2000.
- [Hö] L. Hörmander, An introduction to complex analysis in several variables, third edition, *North-Holland Mathematical Library*, 7. North-Holland Publishing Co., Amsterdam, 1990.
- [Ke] K. Kenmotsu, Weierstrass formula for surfaces of prescribed mean curvature, *Mathematische Annalen* 245(2):89–99, 1979.
- [Ko] B. G. Konopelchenko, Weierstrass representations for surfaces in 4D spaces and their integrable deformations via DS hierarchy, *Annals of Global Analysis and Geometry* 18(1):61–74, 2000.
- [KS] R. Kusner & N. Schmitt, The Spinor Representation of Surfaces in Space, [arXiv:dg-ga/9610005](https://arxiv.org/abs/dg-ga/9610005), 1996.
- [LeRo] K. Leschke & P. Romon, Darboux transforms and spectral curves of Hamiltonian stationary Lagrangian tori, *Calculus of Variations and Partial Differential Equations* 38(1-2):45–74, 2010.
- [LM] H.B. Lawson & M.L. Michelson, Spin Geometry, *Princeton University Press*, Princeton, New Jersey, 1989.
- [McRo] I. McIntosh & P. Romon, The spectral data for Hamiltonian stationary Lagrangian tori in \mathbb{R}^4 , *Differential Geometry and its Applications* 29:125–146, 2011.
- [Mo] B. Morel, Surfaces in \mathbb{S}^3 and \mathbb{H}^3 via spinors, *Actes du séminaire de théorie spectrale et géométrie* 23:131–144, 2005.
- [NR] R. Nakad & J. Roth, Hypersurfaces of Spin^c Manifolds and Lawson-Type Correspondence, *Annals of Global Analysis and Geometry* 42(3), 421-442, 2012.
- [R] J. Roth, Spinorial characterizations of surfaces into 3-homogeneous manifolds, *Journal of Geometry and Physics* 60 1045-1061, 2010.
- [T1] I. A. Taimanov, Surfaces in the four-space and the Davey-Stewartson equations, *Journal of Geometry and Physics* 56(8):1235–1256, 2006.
- [T2] I. A. Taimanov, Surfaces in three-dimensional Lie groups in terms of spinors, *RIMS Kokyuroku* 1605, 133-150, 2008.