Robust and asymptotically unbiased estimation of extreme quantiles for heavy tailed distributions

Yuri Goegebeur, Armelle Guillou, Andréhette Verster

► To cite this version:

Yuri Goegebeur, Armelle Guillou, Andréhette Verster. Robust and asymptotically unbiased estimation of extreme quantiles for heavy tailed distributions. *Statistics and Probability Letters*, 2014, 87, pp.108-114. 10.1016/j.spl.2014.01.010 . hal-00880530

HAL Id: hal-00880530

<https://hal.science/hal-00880530>

Submitted on 6 Nov 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Robust and asymptotically unbiased estimation of extreme quantiles for heavy tailed distributions

Yuri Goegebeur ^{*}

Armelle Guillou [†]

Andréhette Verster [‡]

Abstract. A robust and asymptotically unbiased extreme quantile estimator is derived from a second order Pareto-type model and its asymptotic properties are studied under suitable regularity conditions. The finite sample properties of the proposed estimator are investigated with a small simulation experiment.

1 Introduction

In extreme value statistics, the estimation of extreme quantiles of a distribution function is a central topic. Indeed, many important applications in climatology, finance, actuarial science, hydrology and geology, to name but a few, require extrapolations outside the data range, and extreme value theory provides the only realistic framework for such an exercise. In the present paper we shall address this estimation problem, with special focus on asymptotic unbiasedness and robustness against outliers.

^{*}Department of Mathematics and Computer Science, University of Southern Denmark, Campusvej 55, 5230 Odense M, Denmark (email: yuri.goegebeur@imada.sdu.dk).

[†]Institut Recherche Mathématique Avancée, UMR 7501, Université de Strasbourg et CNRS, 7 rue René Descartes, 67084 Strasbourg cedex, France (email: armelle.guillou@math.unistra.fr).

[‡]Department of Mathematical Statistics and Actuarial Science, University of the Free State, 205 Nelson Mandela drive, Park West, 9300 Bloemfontein, South Africa (email: VersterA@ufs.ac.za).

We consider the framework of Pareto-type distributions satisfying a second order condition. In particular, we assume the following (see Beirlant *et al.*, 2009). Let RV_β denote the class of the regularly varying functions at infinity with index β , i.e. Lebesgue measurable ultimately positive functions z satisfying $\lim_{t \rightarrow \infty} z(tx)/z(t) = x^\beta$ for all $x > 0$.

Condition (R). *Let $\gamma > 0$ and $\tau < 0$ be constants. The distribution function F is such that $x^{1/\gamma}\bar{F}(x) \rightarrow C \in (0, \infty)$ as $x \rightarrow \infty$ and the function δ defined via*

$$\bar{F}(x) = Cx^{-1/\gamma}(1 + \gamma^{-1}\delta(x)),$$

is ultimately nonzero, of constant sign and $|\delta| \in RV_\tau$.

Clearly, condition (R) implies that the tail quantile function U , defined as $U(y) := \inf\{x : F(x) \geq 1 - 1/y\}$, $y > 1$, satisfies $y^{-\gamma}U(y) \rightarrow C^\gamma$ as $y \rightarrow \infty$ and the function a implicitly defined by

$$U(y) = C^\gamma y^\gamma (1 + a(y)) \tag{1}$$

satisfies $a(y) = \delta(C^\gamma y^\gamma)(1 + o(1))$ as $y \rightarrow \infty$, so $|a| \in RV_\rho$, with $\rho = \gamma\tau$.

The second order condition (R) can be used to derive the so-called extended Pareto distribution, EPD (Beirlant *et al.*, 2004, Beirlant *et al.*, 2009), with distribution function given by

$$G(y) = \begin{cases} 1 - [y(1 + \delta - \delta y^\tau)]^{-1/\gamma}, & y > 1, \\ 0, & y \leq 1, \end{cases} \tag{2}$$

where $\gamma > 0$, $\tau < 0$, and $\delta > \max\{-1, 1/\tau\}$. As shown in Proposition 2.3 of Beirlant *et al.* (2009), for distribution functions satisfying (R), the distribution function of the relative excess $Y := X/u$ given that $X > u$ can be approximated by (2) with $\delta = \delta(u)$ up to an error that is uniformly $o(\delta(u))$ for $u \rightarrow \infty$. In Dierckx *et al.* (2013), a robust and asymptotically unbiased estimator for γ was introduced by fitting the EPD to a sample of relative excesses by the minimum density power divergence (MDPD) criterion (Basu *et al.*, 1998). In particular, let $X_1, \dots, X_n$ be independent and identically distributed (i.i.d.) random variables with a distribution function

satisfying $(\mathcal{R})$, and denote by $X_{1,n} \leq \dots \leq X_{n,n}$ the corresponding order statistics. The parameters γ and δ of the EPD are then estimated with the minimum density power divergence criterion applied to the relative excesses over the random threshold $u = X_{n-k,n}$, namely $Y_j := X_{n-k+j,n}/X_{n-k,n}$, $j = 1, \dots, k$, i.e. one minimises the empirical divergence

$$\hat{\Delta}_\alpha(\gamma, \delta) := \int_1^\infty g^{1+\alpha}(y)dy - \left(1 + \frac{1}{\alpha}\right) \frac{1}{k} \sum_{j=1}^k g^\alpha(Y_j),$$

in case $\alpha > 0$, and

$$\hat{\Delta}_0(\gamma, \delta) := -\frac{1}{k} \sum_{j=1}^k \log g(Y_j),$$

in case $\alpha = 0$, where g denotes the density function of G . The parameter ρ is estimated externally, e.g. by using one of the estimators proposed in Fraga Alves *et al.* (2003) or Goegebeur *et al.* (2010). Other robust estimators for γ were proposed by e.g. Peng and Welsh (2001), Juárez and Schucany (2004), Vandewalle *et al.* (2007), Kim and Lee (2008).

In the present paper we will consider robust and asymptotically unbiased extreme quantile estimation under model $(\mathcal{R})$, using the MDPD estimator of Dierckx *et al.* (2013). Beirlant *et al.* (2009) studied the asymptotically unbiased estimation of small tail probabilities based on the EPD, fitted by the maximum likelihood method. In Gomes and Pestana (2007) an asymptotically unbiased extreme quantile estimator was introduced for heavy-tailed distributions. These approaches are however not robust against outliers. To the best of our knowledge, robust and asymptotically unbiased extreme quantile estimation has not been considered before.

The remainder of our paper is organised as follows. In the next section we will introduce the robust and asymptotically unbiased estimator for extreme quantiles and study its asymptotic properties under suitable regularity conditions. The finite sample behaviour of the proposed estimator and some alternatives from the literature is illustrated with a small simulation experiment in Section 3.

2 Estimator and asymptotic properties

From the second order condition $(\mathcal{R})$ and using the EPD as approximation to the distribution of X/u_n given $X > u_n$ we can for $\bar{F}(u_n) \rightarrow 0$ and $p_n \rightarrow 0$ such that $p_n/\bar{F}(u_n) \rightarrow c \in [0, \infty)$ introduce

$$U_0\left(\frac{1}{p_n}\right) := u_n \left(\frac{p_n}{\bar{F}(u_n)}\right)^{-\gamma} \left(1 - \delta(u_n) \left(1 - \left(\frac{p_n}{\bar{F}(u_n)}\right)^{-\rho}\right)\right) \quad (3)$$

as approximation for $U(1/p_n)$.

Lemma 1 *Assume $(\mathcal{R})$. If $\bar{F}(u_n) \rightarrow 0$ and $p_n \rightarrow 0$ such that $p_n/\bar{F}(u_n) \rightarrow c \in [0, \infty)$ we have that $U_0(1/p_n)/U(1/p_n) \rightarrow 1$.*

The proof of this lemma is straightforward and therefore it is for brevity omitted from the paper. Now, let $X_1, \dots, X_n$ be i.i.d. random variables with a distribution function satisfying $(\mathcal{R})$, and denote by $X_{1,n} \leq \dots \leq X_{n,n}$ the corresponding order statistics. Taking $u_n = X_{n-k,n}$, replacing F by the empirical distribution function in (3), and using the fact that $e^{-x} \sim 1 - x$ for $x \rightarrow 0$, we can introduce the following extreme quantile estimator

$$\hat{U}\left(\frac{1}{p_n}\right) := X_{n-k,n} \left(\frac{np_n}{k}\right)^{-\hat{\gamma}_n} \exp\left(-\hat{\delta}_n \left(1 - \left(\frac{np_n}{k}\right)^{-\hat{\rho}_n}\right)\right), \quad (4)$$

where $(\hat{\gamma}_n, \hat{\delta}_n)$ is the MDPD estimator for (γ, δ) and $\hat{\rho}_n$ is a consistent estimator sequence for ρ .

In order to study the asymptotic behaviour of $\hat{U}(1/p_n)$, properly normalised, we need some preliminary results. Firstly, we need the limiting distribution of the MDPD estimator for $(\hat{\gamma}_n, \hat{\delta}_n)$. This was already derived in Dierckx *et al.* (2013), but we repeat the result for completeness here. Let the arrow $\rightsquigarrow$ denote convergence in distribution, and let $\xrightarrow{\mathbb{P}}$ denote convergence in probability. From now on we denote by γ_0 and ρ_0 the true values of the parameters γ and ρ , respectively, and $\delta_n := \delta(X_{n-k,n})$.

Theorem 1 *Let $X_1, \dots, X_n$ be a sample of i.i.d. random variables from a distribution function satisfying $(\mathcal{R})$. Then if $k, n \rightarrow \infty$ with $k/n \rightarrow 0$ and $\sqrt{k}a(n/k) \rightarrow \lambda \in \mathbb{R}$, we have that*

$$\sqrt{k} \begin{bmatrix} \hat{\gamma}_n - \gamma_0 \\ \hat{\delta}_n - \delta_n \end{bmatrix} \rightsquigarrow (\Gamma, \Delta)$$

with

$$(\Gamma, \Delta) \sim N_2(\mathbf{0}, \mathbb{C}^{-1}(\rho_0) \mathbb{B}(\rho_0) \boldsymbol{\Sigma}(\rho_0) \mathbb{B}'(\rho_0) \mathbb{C}^{-1}(\rho_0)),$$

where $\boldsymbol{\Sigma}(\rho_0)$ is a symmetric (3×3) matrix with elements

$$\begin{aligned} \sigma_{11}(\rho_0) &:= \frac{\alpha^2(1+\gamma_0)^2}{[1+\alpha(1+\gamma_0)]^2[1+2\alpha(1+\gamma_0)]}, \\ \sigma_{21}(\rho_0) &:= \frac{\alpha(1+\gamma_0)[\alpha(1+\gamma_0)-\rho_0]}{[1+\alpha(1+\gamma_0)][1-\rho_0+\alpha(1+\gamma_0)][1-\rho_0+2\alpha(1+\gamma_0)]}, \\ \sigma_{22}(\rho_0) &:= \frac{[\alpha(1+\gamma_0)-\rho_0]^2}{[1-\rho_0+\alpha(1+\gamma_0)]^2[1-2\rho_0+2\alpha(1+\gamma_0)]}, \\ \sigma_{31}(\rho_0) &:= \gamma_0 \left(\frac{1}{[1+2\alpha(1+\gamma_0)]^2} - \frac{1}{[1+\alpha(1+\gamma_0)]^3} \right), \\ \sigma_{32}(\rho_0) &:= \gamma_0 \left(\frac{1}{[1-\rho_0+2\alpha(1+\gamma_0)]^2} - \frac{1}{[1+\alpha(1+\gamma_0)]^2[1-\rho_0+\alpha(1+\gamma_0)]} \right), \\ \sigma_{33}(\rho_0) &:= \gamma_0^2 \left(\frac{2}{[1+2\alpha(1+\gamma_0)]^3} - \frac{1}{[1+\alpha(1+\gamma_0)]^4} \right), \end{aligned}$$

$\mathbb{C}(\rho_0)$ is a symmetric (2×2) matrix with elements

$$\begin{aligned} c_{11}(\rho_0) &:= \gamma_0^{-\alpha-2} \frac{1+\alpha^2(1+\gamma_0)^2}{[1+\alpha(1+\gamma_0)]^3}, \\ c_{12}(\rho_0) &:= \gamma_0^{-\alpha-2} \frac{\rho_0(1-\rho_0)[1+\alpha(1+\gamma_0)+\alpha^2(1+\gamma_0)^2]+\alpha^3\rho_0(1+\gamma_0)^3}{[1+\alpha(1+\gamma_0)]^2[1-\rho_0+\alpha(1+\gamma_0)]^2}, \\ c_{22}(\rho_0) &:= \gamma_0^{-\alpha-2} \frac{(1-\rho_0)\rho_0^2+\alpha\rho_0^2(1+\gamma_0)[\alpha(1+\gamma_0)-\rho_0]}{[1+\alpha(1+\gamma_0)][1-\rho_0+\alpha(1+\gamma_0)][1-2\rho_0+\alpha(1+\gamma_0)]}, \end{aligned}$$

and

$$\mathbb{B}(\rho_0) := \gamma_0^{-\alpha-2} \begin{bmatrix} \gamma_0 & 0 & -1 \\ \gamma_0 & -\gamma_0(1-\rho_0) & 0 \end{bmatrix}.$$

Secondly, we need the limiting distribution of the intermediate order statistic $X_{n-k,n}$ under $(\mathcal{R})$, properly normalised.

Lemma 2 *Let $X_1, \dots, X_n$ be a sample of i.i.d. random variables from a distribution function satisfying $(\mathcal{R})$. For $k, n \rightarrow \infty$ such that $k = o(n)$ and $\sqrt{k}a(n/k) \rightarrow \lambda \in \mathbb{R}$ we have that*

$$\sqrt{k} \left(\frac{X_{n-k,n}}{U(n/k)} - 1 \right) \rightsquigarrow \mathbb{X}$$

where $\mathbb{X} \sim N(0, \gamma_0^2)$.

In the next theorem we state the limiting distribution of the extreme quantile estimator (4), when properly normalised.

Theorem 2 *Let $X_1, \dots, X_n$ be a sample of i.i.d. random variables from a distribution function satisfying $(\mathcal{R})$. Then if $k \rightarrow \infty$ as $n \rightarrow \infty$ with $k/n \rightarrow 0$, $\sqrt{k}a(n/k) \rightarrow \lambda \in \mathbb{R}$, $np_n/k \rightarrow 0$ and $\ln(np_n)/\sqrt{k} \rightarrow 0$ we have that*

$$\frac{\sqrt{k}}{\ln \frac{k}{np_n}} \left(\frac{\hat{U}\left(\frac{1}{p_n}\right)}{U\left(\frac{1}{p_n}\right)} - 1 \right) \rightsquigarrow \Gamma.$$

Theorem 2 indicates that the normalised extreme quantile estimator inherits the asymptotic distribution of the MDPD estimator for γ_0 . As shown in Dierckx *et al.* (2013), the MDPD estimator for γ_0 based on the EPD is robust against outliers and asymptotically unbiased.

3 Simulation experiment

In this section we investigate the finite sample properties of $\hat{U}(1/p_n)$ as given in (4) with different parameter estimators, in particular the MDPD estimator $\hat{\gamma}_n$ and $\hat{\delta}_n$ with $\alpha = 0.1, 0.5$ and 1 , and the maximum likelihood estimator (corresponding to MDPD with $\alpha = 0$, see also Beirlant *et al.*, 2009). We also consider the Weissman estimator (Weissman, 1978) given by

$$\hat{U}_W(1/p_n) = X_{n-k,n} \left(\frac{np_n}{k} \right)^{-H_{k,n}},$$

with $H_{k,n}$ being Hill's estimator (Hill, 1975). For the parameter ρ we use the estimator of Fraga Alves *et al.* (2003).

Figures 1 to 6 illustrate the results of a small simulation study based on 100 datasets, each of size $n = 200$, simulated from the distributions given below. The same distributions were considered in Dierckx *et al.* (2013).

- Uncontaminated Fréchet distribution (Figure 1): $F(x) = \exp(-x^{-\beta}), x > 0, \beta > 0$, denoted Fréchet(β). For this study β was chosen as 2.
- Contaminated Fréchet distribution: $F_\epsilon(x) = (1 - \epsilon)F(x) + \epsilon\tilde{F}(x)$ where $F(x)$ represents the uncontaminated Fréchet(2) and $\tilde{F}(x) = 1 - (x/x_c)^{-\beta}, x > x_c$ where β is chosen as 0.5

and $x_c = 2$ times the 99.99% quantile of the uncontaminated Fréchet(2). We take $\epsilon = 0.01$ (Figure 2) and $\epsilon = 0.02$ (Figure 3).

- Uncontaminated Burr distribution (Figure 4): $F(x) = 1 - (\eta/(\eta + x^\tau))^\lambda, x > 0, \eta, \tau, \lambda > 0$, denoted $\text{Burr}(\eta, \tau, \lambda)$. For this study we have chosen $\eta = 1, \tau = 1$ and $\lambda = 2$.
- Contaminated Burr distribution: $F_\epsilon(x) = (1 - \epsilon)F(x) + \epsilon\tilde{F}(x)$ where $F(x)$ represents the uncontaminated $\text{Burr}(1,1,2)$ and $\tilde{F}(x) = 1 - (x/x_c)^{-\beta}, x > x_c$ where $\beta = 0.5$ and $x_c = 1.2$ times the 99.99% quantile of the uncontaminated $\text{Burr}(1,1,2)$. We take $\epsilon = 0.01$ (Figure 5) and $\epsilon = 0.02$ (Figure 6).

We report only the results for quantile $1 - 1/500$. The $1 - 1/1000$ quantile was also considered and resulted in similar outcomes.

In Figures 1 to 6, the left panels show the median of the extreme quantile estimators and the right panels the mean squared error (MSE) of $\ln(\hat{U}_*(1/p_n)/U(1/p_n))$, where $\hat{U}_*(1/p_n)$ denotes any of the considered estimators of $U(1/p_n)$, as a function of k . The top panels of the figures illustrate the behavior of $\hat{U}(1/p_n)$ with the MDPD estimator at different levels of robustness, namely $\alpha = 0.1$ (dotted), $\alpha = 0.5$ (solid) and $\alpha = 1$ (dashed). The true quantile is indicated by the horizontal reference line on the left panels. In the uncontaminated cases (Figures 1 and 4) the bias and variability increase as α increases and by judging from the mean squared error of the log ratios the performance weakens as α increases, as expected. Thus, for uncontaminated cases $\alpha = 0.1$ results in the best performance. For the contaminated cases (Figures 2 - 3 and 5 - 6), $\alpha = 0.5$ seems to be less biased compared to the other two and it results in the best MSE of the log ratios (at least for small values of k). The bottom panels of the figures illustrate the behavior of $\hat{U}(1/p_n)$ with the MDPD estimator at $\alpha = 0.1$ for uncontaminated datasets (solid) and $\alpha = 0.5$ for contaminated datasets (solid), as well as with the MLE of the EPD (dotted), and $\hat{U}_W(1/p_n)$ (dashed). For both uncontaminated and contaminated cases the bias corrected and robust MDPD based estimator outperforms the non-robust estimators in terms of bias and MSE. In particular note that compared to the other estimators, the MDPD based estimator has very stable sample paths, especially in presence of outliers.

Figure 1: Fréchet simulation, quantile 1-1/500. Top: median (left) and MSE (right) of the MDPD based estimator with $\alpha = 0.1$ (dotted), $\alpha = 0.5$ (solid) and $\alpha = 1$ (dashed). Bottom: median (left) and MSE (right) of the MDPD based estimator with $\alpha = 0.1$ (solid), MLE based estimator (dotted) and Weissman estimator (dashed).

Appendix

Proof of Lemma 2

Using the inverse probability integral transform we have that $X_{n-k,n} \stackrel{D}{=} U(Y_{n-k,n})$, where $Y_{n-k,n}$ denotes order statistic $n - k$ of a random sample $Y_1, \dots, Y_n$ from the unit Pareto distribution

Figure 2: Fréchet simulation, quantile $1-1/500$, 1% contamination. Top: median (left) and MSE (right) of the MDPD based estimator with $\alpha = 0.1$ (dotted), $\alpha = 0.5$ (solid) and $\alpha = 1$ (dashed). Bottom: median (left) and MSE (right) of the MDPD based estimator with $\alpha = 0.5$ (solid), MLE based estimator (dotted) and Weissman estimator (dashed).

with distribution function $H(y) = 1 - 1/y$, $y > 1$. Thus,

$$\begin{aligned}
 \sqrt{k} \ln \frac{X_{n-k,n}}{U(n/k)} &\stackrel{D}{=} \sqrt{k} \ln \frac{U(Y_{n-k,n})}{U(n/k)} \\
 &= \gamma_0 \sqrt{k} \ln \left(\frac{k}{n} Y_{n-k,n} \right) + \sqrt{k} \ln \frac{1 + a(Y_{n-k,n})}{1 + a(n/k)} \\
 &=: L_1 + L_2.
 \end{aligned}$$

Figure 3: Fréchet simulation, quantile 1-1/500, 2% contamination. Top: median (left) and MSE (right) of the MDPD based estimator with $\alpha = 0.1$ (dotted), $\alpha = 0.5$ (solid) and $\alpha = 1$ (dashed). Bottom: median (left) and MSE (right) of the MDPD based estimator with $\alpha = 0.5$ (solid), MLE based estimator (dotted) and Weissman estimator (dashed).

For L_1 , use the well-known fact that $\sqrt{k}(k/nY_{n-k,n} - 1) \rightsquigarrow Z$ where $Z \sim N(0, 1)$ (see for instance Corollary 2.2.2 in de Haan and Ferreira, 2006) and the delta method to obtain that $L_1 \rightsquigarrow \mathbb{X}$ under the conditions of the lemma. For the term L_2 , a straightforward application of Taylor's theorem gives

$$L_2 = \sqrt{k}a(n/k) \left(\frac{a(Y_{n-k,n})}{a(n/k)} - 1 + o(1) + o\left(\frac{a(Y_{n-k,n})}{a(n/k)}\right) \right).$$

Figure 4: Burr simulation, quantile 1-1/500. Top: median (left) and MSE (right) of the MDPD based estimator with $\alpha = 0.1$ (dotted), $\alpha = 0.5$ (solid) and $\alpha = 1$ (dashed). Bottom: median (left) and MSE (right) of the MDPD based estimator with $\alpha = 0.1$ (solid), MLE based estimator (dotted) and Weissman estimator (dashed).

Since a is regularly varying we have that $a(tx)/a(t) \rightarrow x^\rho$ as $t \rightarrow \infty$, locally uniformly for $x > 0$. Combining this with the fact that $k/nY_{n-k,n} \rightarrow 1$ a.s. and the assumption $\sqrt{k}a(n/k) \rightarrow \lambda \in \mathbb{R}$ we have that $L_2 \xrightarrow{\mathbb{P}} 0$. Lemma 2 follows then by collecting the terms and another application of the delta method.

Figure 5: Burr simulation, quantile $1-1/500$, 1% contamination. Top: median (left) and MSE (right) of the MDPD based estimator with $\alpha = 0.1$ (dotted), $\alpha = 0.5$ (solid) and $\alpha = 1$ (dashed). Bottom: median (left) and MSE (right) of the MDPD based estimator with $\alpha = 0.5$ (solid), MLE based estimator (dotted) and Weissman estimator (dashed).

Proof of Theorem 2

First we comment on the joint convergence in distribution of the random vector

$$(\sqrt{k}(\hat{\gamma}_n - \gamma_0), \sqrt{k}(\hat{\delta}_n - \delta_n), \sqrt{k}(X_{n-k,n}/U(n/k) - 1), \hat{\rho}_n).$$

Figure 6: Burr simulation, quantile 1-1/500, 2% contamination. Top: median (left) and MSE (right) of the MDPD based estimator with $\alpha = 0.1$ (dotted), $\alpha = 0.5$ (solid) and $\alpha = 1$ (dashed). Bottom: median (left) and MSE (right) of the MDPD based estimator with $\alpha = 0.5$ (solid), MLE based estimator (dotted) and Weissman estimator (dashed).

According to the proof of Theorem 2 in Dierckx *et al.* (2013), we have that

$$(\sqrt{k}(\hat{\gamma}_n - \gamma_0), \sqrt{k}\hat{\delta}_n) \rightsquigarrow (\Gamma, \tilde{\Delta}),$$

where $(\Gamma, \tilde{\Delta}) \sim N_2((0, \lambda), \mathbb{C}^{-1}(\rho_0)\mathbb{B}(\rho_0)\Sigma(\rho_0)\mathbb{B}'(\rho_0)\mathbb{C}^{-1}(\rho_0))$. From the proof of Lemma 1 and Theorem 2 in Dierckx *et al.* (2013) we can deduce that $\hat{\gamma}_n$ and $\hat{\delta}_n$ are independent of $X_{n-k,n}$,

and therefore

$$(\sqrt{k}(\hat{\gamma}_n - \gamma_0), \sqrt{k}\hat{\delta}_n, \sqrt{k}(X_{n-k,n}/U(n/k) - 1)) \rightsquigarrow (\Gamma, \tilde{\Delta}, \mathbb{X}),$$

where $(\Gamma, \tilde{\Delta}, \mathbb{X}) \sim N_3((0, \lambda, 0), \Psi)$, with

$$\Psi := \begin{bmatrix} \mathbb{C}^{-1}(\rho_0)\mathbb{B}(\rho_0)\Sigma(\rho_0)\mathbb{B}'(\rho_0)\mathbb{C}^{-1}(\rho_0) & \mathbf{0} \\ \mathbf{0} & \gamma_0^2 \end{bmatrix}.$$

Finally, using the fact that $\sqrt{k}\delta_n \xrightarrow{\mathbb{P}} \lambda$ and $\hat{\rho}_n \xrightarrow{\mathbb{P}} \rho_0$ we have also that

$$(\sqrt{k}(\hat{\gamma}_n - \gamma_0), \sqrt{k}(\hat{\delta}_n - \delta_n), \sqrt{k}(X_{n-k,n}/U(n/k) - 1), \hat{\rho}_n) \rightsquigarrow (\Gamma, \Delta, \mathbb{X}, \rho_0).$$

Now, consider $\ln(\hat{U}(1/p_n)/U(1/p_n))$. Let $d_n := k/(np_n)$. Straightforward calculations give

$$\begin{aligned} \ln \frac{\hat{U}\left(\frac{1}{p_n}\right)}{U\left(\frac{1}{p_n}\right)} &= \ln \frac{X_{n-k,n}}{U\left(\frac{n}{k}\right)} + (\hat{\gamma}_n - \gamma_0) \ln d_n + \ln \frac{1 + a\left(\frac{n}{k}\right)}{1 + a\left(\frac{1}{p_n}\right)} - \hat{\delta}_n \left(1 - d_n^{\hat{\rho}_n}\right) \\ &=: T_1 + T_2 + T_3 - T_4. \end{aligned} \quad (5)$$

Clearly $T_1 = O_{\mathbb{P}}(1/\sqrt{k})$ by Lemma 2 and $T_2 = O_{\mathbb{P}}\left(\frac{\ln d_n}{\sqrt{k}}\right)$ by Theorem 1. From Taylor's theorem we can write

$$T_3 = a(n/k) \left(1 - \frac{a\left(\frac{1}{p_n}\right)}{a\left(\frac{n}{k}\right)} + o(1) + o\left(\frac{a\left(\frac{1}{p_n}\right)}{a\left(\frac{n}{k}\right)}\right) \right).$$

By using the regular variation properties of the function a and the fact that $d_n \rightarrow \infty$ we have that $a(1/p_n)/a(n/k) \rightarrow 0$, and thus under the conditions of the theorem $T_3 = O(a(n/k))$. Finally, for T_4 note that $d_n^{\hat{\rho}_n} = o_{\mathbb{P}}(1)$ and $\hat{\delta}_n = O_{\mathbb{P}}(1/\sqrt{k})$. Collecting all the terms we see thus that the rate of convergence of $\ln(\hat{U}(1/p_n)/U(1/p_n))$ is given by $\frac{\ln d_n}{\sqrt{k}}$. Multiplying both sides of (5) by $\sqrt{k}/\ln d_n$, the result of the theorem follows.

References

- [1] Basu, A., Harris, I.R., Hjort, N.L., Jones, M.C., 1998. Robust and efficient estimation by minimizing a density power divergence. *Biometrika*, 85, 549–559.

- [2] Beirlant, J., Goegebeur, Y., Segers, J., Teugels, J., 2004. *Statistics of Extremes – Theory and Applications*. Wiley Series in Probability and Statistics.
- [3] Beirlant, J., Joossens, E., Segers, J., 2009. Second-order refined peaks-over-threshold modelling for heavy-tailed distributions. *Journal of Statistical Planning and Inference*, 139, 2800–2815.
- [4] de Haan, L., Ferreira, A., 2006. *Extreme Value Theory: An Introduction*. Springer.
- [5] Dierckx, G., Goegebeur, Y., Guillou, A., 2013. An asymptotically unbiased minimum density power divergence estimator for the Pareto-tail index. *Journal of Multivariate Analysis*, to appear.
- [6] Fraga Alves, M.I., Gomes, M.I., de Haan, L., 2003. A new class of semi-parametric estimators of the second order parameter. *Portugaliae Mathematica*, 60, 193–213.
- [7] Goegebeur, Y., Beirlant, J., de Wet, T., 2010. Kernel estimators for the second order parameter in extreme value statistics. *Journal of Statistical Planning and Inference*, 140, 2632–2652.
- [8] Gomes, M.I., Pestana, D., 2007. A sturdy reduced-bias extreme quantile (VaR) estimator. *Journal of the American Statistical Association*, 102, 280–291.
- [9] Hill, B., 1975. A simple general approach to inference about the tail of a distribution. *The Annals of Statistics*, 3, 1163–1174.
- [10] Juárez, S.F., Schucany, W.R., 2004. Robust and efficient estimation for the generalized Pareto distribution. *Extremes*, 7, 237–251.
- [11] Kim, M., Lee, S., 2008. Estimation of a tail index based on minimum density power divergence. *Journal of Multivariate Analysis*, 99, 2453–2471.
- [12] Peng, L., Welsh, A.H., 2001. Robust estimation of the generalized Pareto distribution. *Extremes*, 4, 53–65.

- [13] Vandewalle, B., Beirlant, J., Christmann, A., Hubert, M., 2007. A robust estimator for the tail index of Pareto-type distributions. *Computational Statistics & Data Analysis*, 51, 6252–6268.
- [14] Weissman, I., 1978. Estimation of parameters and large quantiles based on the k largest observations. *Journal of the American Statistical Association*, 73, 812–815.