



HAL
open science

Relationship between CO₂ assimilation and leaf anatomical characteristics of two grapevine cultivars

Angelos Patakas, Dimitrios Stavrakas, Ioannis Fisarakis

► **To cite this version:**

Angelos Patakas, Dimitrios Stavrakas, Ioannis Fisarakis. Relationship between CO₂ assimilation and leaf anatomical characteristics of two grapevine cultivars. *Agronomie*, 2003, 23 (4), pp.293-296. 10.1051/agro:2003002 . hal-00886181

HAL Id: hal-00886181

<https://hal.science/hal-00886181>

Submitted on 11 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Relationship between CO₂ assimilation and leaf anatomical characteristics of two grapevine cultivars

Angelos PATAKAS^a, Dimitrios STAVRAKAS^a, Ioannis FISARAKIS^b

^a Laboratory of Viticulture, Aristotle University of Thessaloniki, PO Box 236, 54006 Thessaloniki, Greece

^b Dept. of Agriculture, Technological Institution of Crete, Estavromenos, 71500 Heraklion, Greece

(Received 18 October 2001; revised 25 July 2002; accepted 27 August 2002)

Abstract – The possible implication of leaf anatomical characteristics on the photosynthetic rate was studied in two grapevine cultivars, Ribier (*Vitis vinifera* L.) and Isabella (*Vitis labrusca*), grown under field conditions. Ribier exhibited higher photosynthetic rates than Isabella, although there were no significant differences in the Rubisco activity and the stomatal conductance. The fraction of mesophyll volume represented by the intercellular spaces as well as the surface area of mesophyll cells exposed to intercellular air spaces were significantly lower in Isabella. Both gaseous CO₂ conductance through intercellular airspaces and liquid phase conductance were significantly higher in Ribier than in Isabella, contributing to a higher photosynthetic rate in this cultivar.

leaf anatomy / mesophyll conductance / photosynthetic rate / *Vitis vinifera* / *Vitis labrusca*

Résumé – Relation entre l'assimilation de CO₂ et les caractéristiques anatomiques des feuilles de deux espèces de vigne. L'influence possible des caractéristiques anatomiques des feuilles sur le taux de photosynthèse a été étudiée pour deux espèces de vigne : *Vitis Labrusca* (Isabella) and *Vitis vinifera* L. (Ribier), cultivées en culture au champ. Ribier a présenté un taux photosynthétique plus élevé que Isabella, bien qu'il n'y ait pas de différences significatives de l'activité de la Rubisco et de la conductance stomatique. La fraction du volume de mésophylle représentée par l'espace intercellulaire, ainsi que la surface des cellules du mésophylle exposée à l'air dans l'espace intercellulaire, étaient significativement plus faibles chez Isabella. Les conductances gazeuses pour le CO₂ à travers l'espace intercellulaire et en phase liquide étaient significativement plus fortes pour Ribier que pour Isabella, contribuant à des taux photosynthétiques plus importants pour cette espèce.

anatomie des feuilles / conductance interne / taux photosynthétique / *Vitis vinifera* / *Vitis labrusca*

1. INTRODUCTION

It is well known that, under non-limiting environmental conditions for gas exchange by leaves, rates of CO₂ assimilation may be regulated by the intrinsic photosynthetic capacity of the mesophyll and by CO₂ transfer conductances from ambient air to carboxylation sites within chloroplasts [11, 12]. These conductances include boundary layer, stomatal (g_s) and internal conductance (g_i) within the mesophyll. The first two conductances could be accurately estimated by measurements of transpiration rates. On the other hand, changes in leaf anatomical characteristics are known to alter the internal CO₂ conductance from the substomatal cavities to sites of carboxylation and thus the photosynthetic rate [7]. A strong correlation between CO₂ internal conductance and photosynthetic

rate has been found both within and between species [11]. However, the relative importance of internal conductance in determining the photosynthetic rate in grapevine cultivars remains unknown.

Differences in photosynthetic capacity are of great importance as they are closely related to plant productivity. Photosynthetic rate is known to vary considerably in different grapevine cultivars [3, 4, 5, 20]. The reasons for the differential photosynthetic behavior in these cultivars remain more or less obscure.

The aim of this study was to determine the leaf anatomical and physiological differences between two grapevine cultivars and evaluate whether photosynthetic rate could be related to their leaf anatomical characteristics.

Communicated by Gérard Guyot (Avignon, France)

* Correspondence and reprints
patak@agro.auth.gr

2. MATERIALS AND METHODS

Fifteen-year-old field-grown grapevines of *Vitis vinifera* L. cv. Ribier and *Vitis labrusca* cv. Isabella were used. The soil was a clay loam, 17% clay, 36% silt, 47% sand, CEC = 14.5 cmol·kg⁻¹, 11% CaCO₃ and a field capacity of 18.5% w/w. The plants were grafted onto 110R (*V. berlandieri* × *V. rupestris*) rootstock and were cordon-trained in a bilateral cordon. Vine spacing was 1.8 m in rows and 2.5 m apart. All plants were uniformly pruned in the winter with a load of 12 buds per plant. Approximately one month after bud break all the plants were thinned to one cluster per shoot. Electronic tensiometers, placed at 25 cm depth, were used for monitoring soil water potential. In order to maintain high water availability in the soil, drip irrigation was applied when soil water potential reached -0.03 MPa (average value from 10 tensiometers), approximately two times every week. Starting in May and continuing throughout the season, predawn and midday leaf water potential (Ψ) as well as osmotic potential (Π) were determined every ten days. These measurements were performed on three mature leaves per treatment, at each sampling time during the day, using the psychrometric technique [15]. Two pairs of leaf discs were obtained from each one of the three leaves. The first pair of discs was used for leaf water potential determination while osmotic potential was measured on the second pair. Turgor potential (P) was estimated using the equation [9]:

$$P = \Psi - \Pi \quad (1)$$

Diurnal measurements of gas exchange parameters were made four times during the growing season, at ten-day intervals, starting 15 days after berry set. These measurements were performed every two hours, from 06:00 to 18:00 h, on ten fully expanded mature exterior leaves per treatment, at saturation light intensity, using a portable gas exchange system (Li-6400, Li-Cor Inc, Nebraska USA). The leaves that were used for these measurements were 40–50 days old. In order to estimate exactly the leaf age, leaf unfolding was systematically recorded throughout the growing season. The values of gas exchange parameters measured early in the morning – approximately at 10:00 pm – were considered as maximum diurnal values. Concomitant measurements of the maximum rate of Rubisco activity (V_c) were also performed on the leaves that were used for the maximum P_n determination. This was calculated using the equation [13, 14]:

$$V_c = (P_n + R_d)(C_i + K_m)/(C_i - \Gamma) \quad (2)$$

where P_n is the maximum diurnal values of photosynthetic rate, K_m is the effective Michaelis-Menten constant for CO₂ (46 Pa according to Farquhar et al. [6]), Γ is the compensation point of CO₂ in the absence of respiration [17], and C_i the intercellular CO₂ concentration. The rate of respiration (R_d) was measured according to the gas exchange measurements protocol with the only exception that the sample cuvette was enclosed in aluminum foil to exclude light [16].

Chlorophyll content was estimated by absorbance at 645 and 663 nm of an 80% acetone extract [1] of ten leaf discs

obtained from each leaf used for gas exchange measurements each sampling time.

For anatomical measurements, ten sections obtained from each of ten mature fully expanded exterior leaves from each treatment were examined under a scanning electron microscope. Five of the best transverse sections without folds were photographed to be used for anatomical measurements. Photographs were taken using a transmission electron microscope attachment and printed in four to eight electron micrographs. These micrographs were pieced together to yield a composite photomicrograph of sufficient enlargement to allow measurement of the surface areas of individual chloroplasts using a BioQuant IV electronic digitizer. Total leaf thickness (L) and mesophyll thickness (l), as well as the width of sections (w) were also measured. The cross-sectional length of mesophyll tissue exposed to intercellular air spaces was measured (L_m), and the surface of mesophyll cells exposed to the intercellular air spaces per unit leaf area (S_m) was calculated as [18]:

$$S_m = 1.34 (L_m/w) \quad (3)$$

The factor 1.34 accounts for cell surfaces that were not uniformly perpendicular to the plane of the section based on the average width to height ratio for palisade and spongy cells [7, 19].

The fraction of mesophyll tissue occupied by the intercellular air spaces (f_{ias}) was determined as:

$$f_{ias} = 1 - A_m/lw \quad (4)$$

where A_m is the total cross-sectional area of mesophyll cells per unit leaf area and l is the mesophyll thickness between the upper and lower leaf epidermises.

An estimate of gaseous conductance through the intercellular air spaces (g_{ias}) was obtained using the equation [18]:

$$G_{ias} = (f_{ias})^{1.55} / lk \quad (5)$$

where the 1.55 power accounts for a modeled tortuosity in the diffusion path through small pores [2], and k is a fitted constant, which for simplicity was considered as equal to one in both cultivars.

Leaf tissue density (T_d) was determined as [18]:

$$T_d = (D/a) / L(1 - f_{ias}) \quad (6)$$

where L is the total leaf thickness and D/a represents the leaf dry weight (D) per unit leaf area (a).

A completely randomized experimental design was used. Statistical analysis was done using the SPSS statistical computer package (SPSS for Windows, standard version, release 6.1). ANOVA and mean separation by LSD test were used to compare the treatments.

3. RESULTS AND DISCUSSION

Seasonal values of predawn and midday leaf water potential (Ψ) decreased continuously from May to August with no significant differences between the two cultivars (Figs. 1 and 2).




Figure 1. Seasonal changes in predawn water potential (-----), osmotic (.....) and turgor potential (solid line) in the two cultivars. Vertical lines indicate the standard error of six replicates for each of the two cultivars.




Figure 2. Seasonal changes in midday water potential (-----) and turgor potential (solid line) in the two cultivars. Vertical lines indicate the standard error of six replicates for each of the two cultivars.

Also, the seasonal changes in predawn and midday osmotic potential (Π) followed the same pattern, resulting in turgor maintenance in both cultivars (Figs. 1 and 2). Diurnal values of gas exchange parameters also did not significantly differ during the period in both cultivars (data not shown) and thus the results presented here (Fig. 3) refer to the means of all the measurements. The maximum values of photosynthetic rate – obtained early in the morning – were significantly higher in Ribier compared with Isabella (Fig. 3). Lower values of photosynthetic rate in Isabella can be attributed to: (i) stomatal limitations; (ii) biochemical constraints on CO_2 assimilation, and/or (iii) differences in internal conductance for CO_2 diffusion from the stomatal cavity to chloroplasts [7, 10]. The relationship between maximum diurnal P_n and stomatal conductance (C_s) revealed that Ribier exhibited higher values of P_n for the same values of C_s (Fig. 4). Since no differences in stomatal conductance between the two cultivars occurred, we can assume that the lower values in the photosynthetic rate in Isabella can




Figure 3. Diurnal changes in photosynthetic rate (P_n) in the two cultivars. Vertical lines indicate the standard error of 40 replicates for each of the two cultivars.




Figure 4. Relationship between leaf stomatal conductance to water vapor and photosynthetic rate in the two cultivars. Vertical lines indicate the standard error of 40 replicates for each of the two cultivars.

be attributed to either biochemical or anatomical limitations. As far as the possible biochemical constraints on CO_2 assimilation are concerned, current biochemical models of photosynthesis consider that CO_2 assimilation is limited by the activity of ribulose 1,5-biphosphate carboxylase/oxygenase (Rubisco) and/or chlorophyll content [8, 11]. However, the estimated values of maximum Rubisco activity and the chlorophyll content did not significantly differ in the two cultivars (Tab. I). Thus, lower values of photosynthetic rate in Isabella do not seem to be related to Rubisco activity. This inability of either stomatal or biochemical limitations to account for the low CO_2 assimilation rate leads us to suggest that probably a low internal

Table I. Changes in maximum photosynthetic rate, Rubisco activity and chlorophyll content in the two grapevine cultivars. Data are means \pm S.E. of 40 replicates.

	Photosynthetic rate (P_n) ($\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$)	Rubisco activity (V_c) ($\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$)	Chlorophyll content ($\text{mg}\cdot\text{dm}^{-2}$)
Ribier	$13.1 \pm 0.7^*$	36.3 ± 5.2	3.51 ± 0.15
Isabella	10.1 ± 0.5	33.9 ± 4.7	3.47 ± 0.09

* $P < 0.05$.

Table II. Leaf anatomical properties of the two grapevine cultivars (f_{ias} = the fraction of mesophyll tissue occupied by the intercellular air spaces, g_{ias} = gaseous conductance through intercellular air spaces, T_d = leaf tissue density and S_m = surface area of mesophyll cells exposed to the intercellular air spaces per unit leaf area). Data are means \pm S.E. of 10 replicates.

	f_{ias}	g_{ias} ($\text{mol} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$) $\times 10^{-4}$	T_d ($\text{g} \cdot \text{m}^{-3}$) $\times 10^5$	S_m ($\text{m}^2 \cdot \text{m}^{-2}$)
Ribier	0.4 \pm 0.06 **	15.39 \pm 2.5 **	7.52 \pm 0.8	15.2 \pm 1.1 **
Isabella	0.28 \pm 0.04	8.96 \pm 1.2	6.27 \pm 0.6	12.5 \pm 0.8

$P < 0.05$, ** $P < 0.01$.

conductance (g_j) for CO_2 diffusion from the stomatal cavity to chloroplasts could be responsible for the lower values of P_n in leaves of Isabella. Internal conductance is considered as the sum of two components: an intercellular gas phase conductance from the stomatal cavity to outer surfaces of mesophyll cell walls (g_{ias}) and a liquid-phase conductance across cell walls to sites of carboxylation within chloroplasts (g_{cw}) [10, 14].

The values of both components of the total internal conductance are known to be closely related to leaf anatomical features [10, 18]. The measurements of leaf anatomical characteristics showed significant differences between the two cultivars (Tab. II). The fraction of mesophyll volume represented by intercellular air spaces (f_{ias}) was significantly greater in Ribier, resulting in almost double values of the calculated gaseous conductance through the intercellular air spaces (g_{ias}) (Tab. II). High g_{ias} means more rapid diffusion of CO_2 from the substomatal cavity to the cell wall surfaces, a fact that may explain the higher photosynthetic rate in Ribier, despite the similar values of stomatal conductance. On the other hand, the liquid face conductance (g_{cw}) includes the CO_2 flow across the cell walls and across cytoplasmic and chloroplastic membranes, and thus it is much more difficult to be estimated [14]. However, reasonable estimates of g_{cw} could be obtained from the average tissue density as well as the surface of the mesophyll cells exposed to intercellular air spaces (S_m) [18]. Our results show no significant differences in leaf tissue density despite the differences in f_{ias} (Tab. II), a fact that might be attributed to: (i) the difference in the number of mesophyll cells per unit of leaf area, and/or (ii) to differences in cell wall thickness between the two cultivars. Since no significant difference in the number of mesophyll cells per unit of leaf area between the two cultivars occurred (data not shown) we can suppose that cell wall thickness would be lower in Ribier compared with Isabella. A less thick cell wall shortens the path from the outer surface of the cell walls to the sites of carboxylation over which the CO_2 must diffuse, resulting in higher g_{cw} . Furthermore, the surface of mesophyll cells exposed to the intercellular air spaces per unit leaf area (S_m) was also significantly greater in Ribier (Tab. II). These data provide evidence that g_{cw} might be greater in Ribier, contributing to the higher photosynthetic rates of this cultivar.

4. CONCLUSION

Our data confirm that differences in photosynthetic rate in the two grapevine cultivars could be attributed to their differ-

ences in leaf anatomical characteristics, which affected the total internal CO_2 conductance.

REFERENCES

- [1] Arnon D.I., Copper enzymes in isolated chloroplasts. Polyphenol oxidase in *Beta vulgaris*, *Plant Physiol.* 24 (1949) 1–15.
- [2] Ball B.C., Modeling of soil pores as tubes using gas permeabilities, gas diffusivities and water release, *Soil Sci.* 32 (1981) 465–481.
- [3] Chaves M.M., Harley P.C., Tenhunen J.D., Lange O.L., Gas exchange in two grapevine cultivars, *Physiol. Plant.* 70 (1987) 639–647.
- [4] During H., CO_2 assimilation and photorespiration of grapevine leaves: Responses to light and drought, *Vitis* 27 (1988) 199–208.
- [5] During H., Photosynthesis of ungrafted and grafted grapevines: effects of rootstock genotype and plant age, *Am. J. Enol. Vitic.* 45 (1994) 297–299.
- [6] Farquhar G.D., Von Caemmerer S., Berry J.A., A biochemical model of photosynthetic CO_2 assimilation in leaves of C_3 species, *Planta* 149 (1980) 78–90.
- [7] Evans J.R., Von Caemmerer S., Setchell B.A., Hudson G.S., The relationship between CO_2 transfer and leaf anatomy in transgenic tobacco with a reduced content of Rubisco, *Aust. J. Plant Physiol.* 21 (1994) 475–495.
- [8] Iacono F., Buccella A., Peterlunger E., Water stress and rootstock influence on gas exchange of grafted and ungrafted grapevines, *Sci. Hortic.* 75 (1998) 27–39.
- [9] Koide R.T., Robichaux R.H., Morse S.R., Smith C.M., Plant water status, hydraulic resistance and capacitance, in: Pearcy R.W., Ehleringer J., Mooney H.A., Runder P.W. (Eds.), *Physiological Ecology*, Chapman and Hall, London-New York, 1991, pp. 161–183.
- [10] Nobel P.S., Leaves and fluxes, in: Nobel P.S. (Ed.), *Physicochemical and Environmental Plant Physiology*, Chap. 8, Academic Press, San Diego, 1991, pp. 393–472.
- [11] Lloyd J., Syvertsen J.P., Kriedemann P.E., Farquhar G.D., Low conductances for CO_2 diffusion from stomata to the sites of carboxylation in leaves of woody species, *Plant Cell Environ.* 15 (1992) 873–899.
- [12] Loreto F., Harley P.C., Di Marco G., Sharkey T.D., Estimation of mesophyll conductance to CO_2 flux by three different methods, *Plant Physiol.* 98 (1992) 1437–1443.
- [13] Pammenter N.W., Loreto F., Sharkey T.D., End product feedback effects on photosynthetic electron transport, *Photosynth. Res.* 35 (1993) 5–14.
- [14] Parkhurst D.F., Diffusion of CO_2 and other gases inside leaves, *New Phytol.* 126 (1994) 449–479.
- [15] Patakas A., Noitsakis B., Cell wall elasticity as a mechanism to maintain favorable water relations during leaf ontogeny in grapevines, *Am. J. Enol. Vitic.* 48 (1997) 352–358.
- [16] Ranney T.G., Ruter J.M., Foliar heat tolerance of three holly species (*Ilex* spp): responses to chlorophyll fluorescence and leaf gas exchange to supraoptimal leaf temperatures, *J. Am. Soc. Hortic. Sci.* 122(1997) 499–503.
- [17] Sharkey T.D., Estimating the rate of photorespiration in leaves, *Physiol. Plant.* 73 (1988) 147–152.
- [18] Syvertsen J.R., Lloyd J., Mc Conchie C., Kriedemann P.E., Farquhar G.D., On the relationship between leaf anatomy and CO_2 diffusion through the mesophyll of hypostomatous leaves, *Plant Cell Environ.* 18 (1995) 149–157.
- [19] Thain J.F., Curvature correction factors in the measurement of cell surface areas, *J. Exp. Bot.* 34 (1983) 87–94.
- [20] Zufferey V., Murisier F., Shultz H.R., A model analysis of the photosynthetic response of *Vitis vinifera* L. cvs. Riesling and Chasselas leaves in the field: I. Interaction of age, light and temperature, *Vitis* 39 (2000) 19–26.