



Amino acid (Lysine) requirement of the Pig: Methods of Approach and Recommendations

W. Wiesemüller

► To cite this version:

W. Wiesemüller. Amino acid (Lysine) requirement of the Pig: Methods of Approach and Recommendations. Annales de zootechnie, 1980, 29 (2), pp.179-193. hal-00887952

HAL Id: hal-00887952

<https://hal.science/hal-00887952>

Submitted on 11 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Amino acid (Lysine) requirement of the Pig: Methods of Approach and Recommendations

W. WIESEMÜLLER

*Wilhelm-Pieck. Universität Rostock — Sektion Tierproduktion. Justus Von Liebig Weg
DDR 25, ROSTOCK*

Summary

Selection for leaner types of pigs is associated with a more sustained daily lean tissue growth and consequently with an increase in the daily lysine requirement. Accordingly, the lysine/energy ratio should be constant for the whole fattening period between 35 and 120 kg live weight.

Similar final carcass characteristics, i.e. the same amount of deposited protein, at a given slaughter weight, may be obtained by distinct energy levels and lysine/energy ratios. For the limiting amino acid (lysine) the lysine/energy ratio was 2.0 g/Mcal D.E. and, taking into account amino acid digestibility, 1.8 g dig. lysine/Mcal D.E.

To optimize energy and protein (lysine) allowances for a given slaughter weight the energy level and profile should be adjusted to the final fattening weight. This is of great importance for producing heavy pigs as compared to lighter ones.

Very high energy consumption in the fattening stage of 35 to 70 kg resulted in a decline of weight and protein deposition at the end of the fattening period from about 90 kg live weight onwards. It cannot be compensated for by a high protein supplementation. At a finishing weight of 120 kg a too intensive nutrient level (more than 650 g gain/day) at the first fattening stage proved to be unfavourable. The resulting depression of protein deposition in the second fattening period accounts for an insufficient expression of the muscle growth potential. It is defined as the upper limit of protein deposition (and consequently lean meat production) in a fattening animal at a given final weight for minimizing both the fattening length and fat deposition. When producing heavy pigs, a reduction of energy supply during the second half of the fattening period is not advisable when using modern crossbred lean pigs. On the contrary, all efforts should be made so that the energy intake during the second half of the fattening period is not limiting for lean tissue growth.

Introduction

The amino acid requirement depends upon performance and age. The effect of protein feeding in fattening pigs was first studied in our laboratory by means of nitrogen balance experiments. Various amounts of different protein sources were added together to a protein-free basal diet (WIESEMÜLLER and POPPE, 1969). The nitrogen balance measurements showed that a smaller quantity of egg protein is required to obtain a given nitrogen balance than from barley or sesame protein. In other words, a given performance is obtained with a considerably lower protein

level if a high-quality protein is used instead of a low quality protein. Thus, for the dietary protein content, considerable differences are observed in growth performance when proteins of different qualities are used. Therefore, formulation of pig diets should not be based only on total protein. In each case the reasons for the differences on performance and nitrogen balance can be explained by the corresponding intakes of the limiting amino acid, lysine for instance. In this connection it does not matter whether the diet involved contains vegetable protein or high value protein of animal origin (POPPE and WIESEMÜLLER, 1968).

The recommended allowances of amino acids were calculated on the basis of the minimum intake required to obtain a nitrogen balance which did not differ statistically from the maximum nitrogen balance as determined mathematically with a curvilinear function. These requirement values were determined within short intervals during the whole fattening period.

In the meantime, the values obtained from nitrogen balance experiments were checked during both fattening and slaughtering experiments using "Edelschwein" and "Landrace" crossbreds (WIESEMÜLLER, POPPE and SIEG, 1974; JANCKER and POPPE, 1975). Recently we also studied to which extent breeding progress and changes in genetic potential may modify the nutritional requirements. Investigations were made to test the variation in the lysine requirement of the growing pig according to the dietary energy level and lean growth potential. The results obtained on animals from the rotational breeding programme have been published (WIESEMÜLLER *et al.*, 1978) and will be discussed here together with results obtained on crossbred pigs.

Materials and Methods

In order to allow a rapid introduction of our research results into the feed industry, these studies were devoted to lysine, the limiting amino acid, all other essential amino acids being present in the experimental rations at such concentrations that requirements could be met.

One hundred and fifteen crossbred *castrated males* (Edelschwein \times Landrace) \times K 250 and 105 crossbred *gilts* (Edelschwein \times Landrace) \times F 150 were fattened from 35 to 120 kg live weight. In addition, in the second group of pigs 35 castrated males were fattened for comparison. All pigs were kept in individual pens.

The line K 250 was bred for rotation crossing in order to improve the fertility performance of the "hybrid" sows. This line is based on the Estonian Bacon Race, the Lithuanian Large White, the Sattel Swine, the Landrace line 05 and the Duroc. The line designated F 150 with rapid meat gain rates has been created for cross-breeding. The breeds originally used, in a variety of combinations, for the crossing scheme were the Pietrain, Lacombe, Belgian Landrace and Hampshire.

Some of the pigs were slaughtered at 40, 70, and 95 kg liveweight and analysis of the chemical composition was performed.

The experiment was designed to test the *effects of lysine intake at 80, 100, and 120 p. 100 of the recommended level at a mean energy intake estimated to be adequate for a daily gain of around 600 g during the whole fattening period*. In addition, a lower level of energy intake with amounts of lysine representing 80 and 100 p. 100 of the recommended level, and a high energy level during the fattening period, were tested. Table 1 gives some precisions about the energy and lysine levels used.

TABLE I

Variation in daily energy and lysine supply according to live weight

Level	40	60	80	100	120 kg live weight
-------	----	----	----	-----	-----------------------

Trial 1 : *Castrated males* (Edelschwein \times Landrace) \times K 250

Energy intake: Mcal D.E. per day

H : 100 %	6.70	8.06	8.84	9.56	9.77
M : 93 %	6.22	7.52	8.48	9.15	9.51
L : 86 %	5.73	6.73	7.66	8.18	8.59

Lysine intake: g per day

80 %	11.3	12.3	13.1	13.5	13.8
100 %	14.0	15.8	16.8	17.3	17.3
120 %	16.8	18.6	19.3	20.1	20.4

Trial 2: *Fattening gilts* (Edelschwein \times Landrace) \times F 150

Energy intake: Mcal D.E. per day

H : 100 %	5.51	6.84	7.97	9.07	10.22
M : 88 %	4.87	6.46	7.59	8.39	9.03
L : 71 %	3.91	5.04	5.84	6.35	6.40

Lysine intake: g per day

80 %	10.5	12.8	13.9	14.2	13.2
100 %	13.6	15.6	17.0	17.8	18.1
120 %	16.4	18.6	20.2	21.1	21.1

The composition of the experimental diets as well as their digestibility values and energy content were published by WIESEMÜLLER *et al.* (1978) and POPPE *et al.* (1979).

Results

I. — *Castrated males* (trial 1)

The daily weight gains and energy expenditure mainly depended on the energy intake. Up to 100 kg liveweight both low and high levels of amino acid supplies produced the same fattening results as a standard amino acid supply.

TABLE 2

Daily gain and feed conversion ratio as related to energy and lysine intakes

Treatment	Energy level %	Lysine level %	Daily gain g	kg dry matter of feed (1) per kg gain
L 80.	86	80	540	3.27
L 100.	86	100	579	3.12
M 80.	93	80	610	3.20
M 100.	93	100	631	3.04
M 120.	93	120	663	2.88
H 100.	100	100	684	3.07
L/H (*)	86/100	100	681	2.85
H/L (*)	100/86	100	589	3.25

(*) Change on the 39th day of the experiment.

(1) 1 kg dry matter of feed = 2 727 kcal NEFs, i.e. approximately 4.04 Mcal D.E.

Within the 100 to 120 kg liveweight range, the animals fed higher levels of protein were slightly superior. As to energy expenditure, the animals subjected to restricted feeding up to 70 kg liveweight and then fed full rations ingested 2.85 kg feed dry matter per kg gain and were significantly better than those fed intensively at the beginning of the experiment and restricted beyond 70 kg liveweight (3.25 kg DM per kg gain), and better than those receiving high levels of energy all the time (3.07 kg DM per kg gain). Using all the gain data within 40 to 120 kg live weight range, a close linear relationship was established between the daily live weight gain (Y , g/kg^{0.75}) and the daily energy consumption (X , F.U./kg^{0.75}) according to the equation: $Y = (-21.06 + 0.7935) \times (R = 0.960, s_b = 0.0315)$. A total of 126 F.U.* or approximately 0.65 Mcal D.E. was needed for every 100 g extra live weight gain.

In order to determine the effects of these levels on the composition of gain, the animals from all treatments were slaughtered and analysed when reaching live weights of 40, 70, 95 and 120 kg. The growth intensities of the animals expressed as daily protein, water and fat gains on four treatments are shown in figure 1.

The daily live weight gain shown by the upper curve remained almost constant throughout the fattening period in pigs with moderate and low energy intakes. The growth intensity exhibited by the pigs on the high energy level (= 100) gradually decreased during the fattening period. The daily protein gain of animals on the levels "Middle-100" and "Middle-120" remained almost constant at 85 g throughout the fattening period. All other levels resulted in a reduced protein deposition with increasing live weight from 70 to 95 kg. The greater the gain performance before this range was reached, the more pronounced was this effect. For example, the high energy level 100 increased the daily protein depo-

* 1 F.U. (feed unit) = 3.5 kcal NEFs, i.e. approximately 5.18 kcal D.E. for the experimental diets.




FIG. 1. — The composition of daily live weight gain of pigs in relation to energy and lysine intakes.

sition to 133 g during the initial development period, i.e. by a factor of 1.5. As the live weight continued to increase protein deposition was obviously influenced by development at an earlier age and the initial high deposition rate was followed by a negative compensatory effect. The daily fat gain was dependent on the energy intake. The average values for the whole fattening period were 260 g/day for the Low energy level, 290 g/day for the Medium level, and 360 g/day for the High level.

Finally, the levels had a decisive effect on the daily water retention of the animals. The results obtained with the M 120 level showed that muscle growth was connected with water deposition.

The protein contents of animals, all slaughtered at 120 kg live weight, with

TABLE 3

Comparison of protein content of the whole body with the content of lean cuts (120 kg live weight)

Treatment	L 100	M 100	M 120
Protein, kg.	15.52	15.46	15.93
% of lean cuts.	49.23	45.51	50.26
Protein + water, kg	68.31	63.05	68.79

the proportion of lean cuts obtained with three treatments are compared in table 3. The latter produced no differences in the protein content, but the animals fed on M 100 had 5 p. 100 less lean cuts per half-carcase than those receiving the other two treatments. This difference could also be shown when the protein and water contents were added together and compared instead of protein alone. The poorest performance exhibited by the animals fed M 100 and slaughtered at 120 kg live weight compared with the animals fed L 100 and M 120 can be related to the lower water retention and greater fat deposition during fattening from 95 to 120 kg live weight.

If the gain is plotted versus time (fig. 2) it is clear how the time taken to reach the final fattening weight of 120 kg can be used to maximise protein production.




FIG. 2. — Body composition changes during the fattening period.

○ H 100. ▲ L 100. ● L 80

The amounts of energy and amino acids supplied to the animals subjected to treatment M 120 fully expressed the protein deposition capacity of the animals while producing a minimum of fat.

This graph shows clearly that protein and fat are produced simultaneously during growth. Although it is possible, by reducing energy intake as in the L 100 treatment, to reduce daily fat deposition and thus produce pigs with carcasses containing the same amounts of protein and fat as those grown to the full expression of their protein deposition ability, this can be done only at the expense of longer fattening periods and higher energy costs.

The capacity or potential for protein deposition is defined as the genetically programmed ability to deposit protein while reaching a given final weight during the shortest possible fattening period and with a minimum fat deposition. The method chosen to show the results gives a good illustration of the negative effects of protein-deficient diets. The shortest fattening period required to reach a defined final weight depends largely on the genetic constitution of the animal.




FIG. 3. — Protein content of the whole body of castrated males ((E \times L) \times K 250) depending on nutrition and live weight.
For each live weight, from left to right: H 100, M 120, L 100 and L 80

In the GDR, pigs for slaughter should have a live weight of 120 kg. For this live weight, the protein retention potential of the genotype tested ((Landrace \times Edelschwein) \times K 250) allowed a final amount of 16 kg protein in the body (fig. 3).

This maximum protein content of 16 kg was achieved by the animals on treatment L 100 and those fed more intensively on M 120. However, the protein deposition potential must be considered as a gain per unit of time. This potential was fully used by the animals on treatment M 120. The forced protein deposition achieved during the first half of the fattening period in animals on H 100 made these animals superior at 70 kg live weight. The subsequent drop in protein deposition accompanied by an increased fat deposition, however, indicated that the protein deposition potential was no longer fully used at 120 kg live weight.

Since, under practical pig fattening conditions, high-energy diets are provided *ad libitum* and the reduced performance during the second part of the fattening period is a cause of general concern, any differences induced by changing the energy input levels at a live weight of about 70 kg would be of interest. For this reason, the L-H level in our studies corresponded to the L 100 treatment up to day 39 of the experiment and to H 100 thereafter, i.e. the animals were on L 100 level up to day 39 (65 kg live weight), whereafter they were fed on treatment H 100. The growth intensity and composition of gain of these animals are shown in figure 4.

The change from a low to a high energy intake level resulted in a substantial increase in the rate of gain first accompanied by an increase in daily fat deposition. However, the situation changed in the course of further growth until the daily protein deposition rate rose to 115 g during the fattening period from 95 to 120 kg live weight. This is also an evidence of the high meat production rate of these animals even towards the end of the fattening period. The high rates of gain




FIG. 4. — Influence of changes in the energy input levels at a live weight of about 65 kg on growth intensity and composition of gain.

LH : Low-High; HL : High-Low. ○ LH; ▲ HL.

shortened the second part of the fattening period to such an extent that these animals had the lowest energy costs.

The lower intake involved in the H-L-programme produced no substantially lower fat deposition, but, due to the reduction in energy intake, the daily protein deposition rate dropped to below the level reached by means of the H 100 treatment alone.

2. — Fattening gilts (trial 2)

More recently GOLDMANN (1978), POPPE *et al.*, (1979) investigated the same problems by using (Edelschwein × Landrace) × F 150 crossbred pigs. Daily gains are shown in table 4. In addition 5 castrated males in each treatment were fattened for comparison.

The weight increase and feed conversion ratio showed a distinct dependence on the energy level of the ration and were mainly determined by the energy intake. Up to a live weight of 90 kg a 20 p. 100 lysine over or undersupply in comparison with the recommended level had no influence on the growth intensity of the animals. On the low and medium energy levels there was a tendency for a superiority of the animals which received protein rich feeds between 90 and

TABLE 4
Daily gains and feed conversion ratios related to energy and lysine intakes in the (Edelschwein × Landrace) × F 150 crossbred pigs

Treatment	Energy level %	Lysine level %	Fattening gilts		Castrated males	
			Daily gain g	kg dry matter of feed (1) per kg gain	Daily gain g	kg dry matter of feed (1) per kg gain
L 80.	71	80	433	3.38	430	3.40
L 100.	71	100	452	3.23	427	3.43
M 80.	88	80	646	2.86	616	3.09
M 100.	88	100	641	2.86	637	2.93
M 120.	88	120	675	2.71	633	2.97
H 100.	100	100	687	2.98	695	2.95
H 120.	100	120	655	3.09	670	3.07

(1) 1 kg dry matter of feed = 2 632 kcal NEFs, i.e. approximately 3.90 Mcal D.E.

120 kg. However, with a high energy level a smaller increase was detected in the last third of the fattening period in connection with the increasing level of lysine. The most suitable result by gilts was obtained with M 120, i.e. a daily increase of 675 g and a feed conversion ratio of 2.71 kg dry matter of feed per kg weight gain.

With all gain data of the 40 to 120 kg live weight range, a close linear relationship was established between daily live weight gain (Y , g/kg^{0.75}) and daily energy consumption (X , F.U./kg^{0.75}), according to the equation: $Y = (-13.97 + 0.7037) \times (R = 0.836; s_b = 0.0295)$. This and the former equation were the basis of the energy requirement data in table 7.

Figure 5 shows the variation in the protein content of gilts.




FIG. 5. — The protein content of the whole body of fattening gilts $((E \times L) \times F 150)$ according to nutrition and live weight.
For each live weight, from left to right: H 100 + 120, M 100 and L 100.

Four treatments were compared with the previously described experiment. In this case a protein deposition potential of around 18 kg at a slaughter weight of 120 kg was also achieved by using a moderate level of energy intake accompanied by an adequate level of lysine intake. The animals fed on the Low 100 treatment with an extremely low energy content reached exactly the same satisfactory carcass characteristics but considerably later with a mean daily gain of 430 g.

It is interesting to note that a high energy intake during the first part of the fattening period also greatly stimulated protein deposition in this case. As in the previous experiment, the daily protein gain was 130 g. However, protein deposition also decreased as fattening proceeded in the case of this meat-producing crossbred pigs, and these losses could not be compensated for by the greater amino acid intake of the animals on the H 120 treatment.

Discussion and Conclusion

To optimize energy and lysine allowances for a given slaughter weight, there is a need to adjust the energy level and profile to the final fattening weight. This is of great importance for producing heavy pigs as compared to lighter ones. It can be concluded that very high energy intakes during the first part of the fattening period are advisable if the animals are to be slaughtered at a live weight of 90 kg. If the final fattening weight is to be 120 kg, a too intensive intake during the period up to a live weight of about 60 kg appears to be unfavourable. The reduced protein deposition observed during the second part of the fattening period appears to result from an insufficient expression of the protein deposition ability of the animal. In the case of crossbred pigs, with a high muscle development, any reduction in the energy intake must be prevented during the second part of the fattening period. Therefore with this type of pigs, the energy intake should be increased during this period, as shown by HENRY (1979).

Now we have to derive the most favourable Lysine/energy ratio since this amino acid was the most limiting in our diets. Pigs on treatment L 100 and M 120 reached the highest protein contents at a final weight of 120 kg. The better results obtained with these treatments justifies the assumption that, in comparison to the energy intake of the pigs on M100 and H100, lysine intake was not quite sufficient to allow maximum utilisation of the protein deposition potential at live weights beyond 70 kg.

At lower weights, reduced lysine intake was compensated for by a greater utilisation, particularly among the animals on the High 100 treatment (table 5). The rate of lysine utilisation was almost 67 p. 100 in the case of these animals. For the whole fattening period, in animals on treatment M 100 it was 36 p. 100, i.e. roughly the same as previously found.

On the basis of a constant lysine requirement of 2.0 g lysine/Mcal digestible energy (table 6), the lysine intake during fattening from 40 to 70 kg is rather smaller than the levels provided in treatment L100 and M100. However, due to the high rate of lysine utilisation, the lower level of lysine intake had no negative effect on protein deposition between these weight limits. This fact must be

TABLE 5

Rate of lysine utilisation for growth (%)

Treatment	Energy/lysine level	Live weight		
		40-70 kg	70-120 kg	40-120 kg
L 80.	86/80	53.8	23.6	34.3
L 100.	86/100	49.9	25.1	33.7
M 80.	93/80	49.8	31.7	38.0
M 100.	93/100	44.0	31.3	35.6
M 120.	93/120	38.5	29.8	32.9
H 100.	100/100	66.8	20.5	35.2

TABLE 6

Lysine/energy ratios in the experimental diets (trial 1)
Amounts in g/Mcal D.E.

Treatment	Live weight (kg)				
	40	60	80	100	120
M 80.	1.84	1.67	1.65	1.62	1.50
H 100	<u>2.09</u>	<u>1.96</u>	1.90	1.81	1.77
L 100.	2.37	2.14	<u>2.09</u>	<u>1.97</u>	1.91
M 100.	2.25	2.10	<u>1.98</u>	1.89	1.82
M 120.	2.66	2.62	2.40	2.30	2.18

TABLE 7

Energy and lysine requirements of growing-fattening pigs as related to growth performance

Daily gain (g)	500		600		700	
	CM	F	CM	F	CM	F
Live weight (kg)	Daily energy needs (Mcal D.E.)					
40.	5.45	5.31	6.10	6.05	6.75	6.75
50.	5.85	5.62	6.50	6.35	7.15	7.09
60.	6.23	5.90	6.88	6.64	7.53	7.37
70.	6.59	6.17	7.24	6.90	7.89	7.64
80.	6.94	6.43	7.59	7.17	8.25	7.90
90.	7.28	6.69	7.94	7.42	8.59	8.16
100.	7.61	6.93	8.26	7.67	8.91	8.40
110.	7.94	7.17	8.59	7.91	9.24	8.65
120.	8.25	7.41	8.90	8.14	9.55	8.88
Live weight (kg)	Daily lysine needs (g)					Recommendations in GDR
40.	11.0		12.4		13.7	14.0
50.	11.9		13.2		14.5	15.0
60.	12.6		14.0		15.2	15.8
70.	13.3		14.7		16.0	16.4
80.	14.1		15.4		16.7	16.8
90.	14.8		16.1		17.4	17.3
100.	15.4		16.7		18.1	17.3
110.	16.1		17.4		18.7	17.3
120.	16.7		18.1		19.3	17.3

CM = Castrated males, F = Females.

checked during further studies. Full utilisation of the protein deposition potential until the end of the fattening period is ensured by the intake of 2.0 g lysine/Mcal D.E. corresponding to the value recommended by HENRY, PION and RERAT (1976) for fattening pigs between 50 and 100 kg live weight. For fattening gilts with smaller needs in the energy intake the lysine/energy ratio could be 2.2 g/Mcal D.E., but for a common fattening of castrated males and gilts until 120 kg the ratio of 2.0 g lysine/Mcal D.E. is sufficient.

Taking into account the lysine requirement of 2.0 g/Mcal D.E. and the energy requirement of growing pigs, the variation with live weight and average daily gain in the daily energy and lysine requirement is shown in table 7.

The current recommendations applied in GDR were determined by using pigs with daily gains of about 700 g (POPPE and WIESEMÜLLER, 1968; WIESEMÜLLER, POPPE and SIEG, 1974). These recommendations are in a good agreement with those presented here for daily gains up to 700 g and up to 90 kg live weight. On account of a constant high rate of protein deposition provided by modern genotypes, the lysine/energy ratio selected here should be maintained up to a final weight of 120 kg. This results in a slightly higher lysine requirement than currently recommended.

Digestibility values are not included in this table. The lysine digestibility of the mixed feed used in our experiments, tested under standardized conditions to secure the amino acid requirements, was 91 p. 100. Our nitrogen balance experiments performed to determine the amino acid requirement were also based on a lysine digestibility of 90 p. 100 and this is why the present requirement values are in good agreement with the recommended ones. However, the protein and amino acid digestibility of the feeds currently used in agricultural practice is far below 90 p. 100 in many instances. For this reason, the figures regarding both requirements and feed contents should refer to the digestible fraction in the case of amino acids. On account of the apparent lysine digestibility of 91 p. 100 determined in our experiments, the calculated digestible lysine requirement for pigs of a modern genotype is 1.8 g/Mcal D.E. Table 8 shows the daily requirements

TABLE 8

Proposals for the supply of digestible lysine to fattening pigs (g per day)

Daily gain (g)	500	600	700	ANDERSEN and JUST (1975)
Live weight (kg)				
40	10.0	11.2	12.4	12.0
50	10.7	12.0	13.1	13.0
60	11.4	12.6	13.8	13.5
70	12.1	13.3	14.4	13.5
80	12.7	14.0	15.1	14.0
90	13.4	14.5	15.8	14.0
100	14.0	15.2	16.3	—
110	14.5	15.8	16.9	—
120	15.1	16.3	17.5	—

for digestible lysine according to live weight and daily gain. The values are in good agreement with the Danish recommendations (ANDERSEN and JUST, 1975).

Accepté pour publication en Juin 1980.

Acknowledgements

The author wishes to acknowledge Mrs. K. Rérat and Mr. Y. Henry for the correction of the manuscript and translation of the summary into French.

Résumé

Besoins en acides aminés (lysine) du porc en croissance Méthodes d'approche et recommandations

La sélection de porcs de type maigre se traduit par un accroissement plus soutenu du dépôt journalier de tissus maigres et, par voie de conséquence, par une augmentation concomitante du besoin journalier de lysine. Il en découle que le rapport lysine/énergie se situe à un niveau relativement constant pendant la durée de l'engraissement entre 35 et 120 kg du poids vif.

Des caractéristiques de carcasse comparables, c'est-à-dire les mêmes quantités de protéines fixées, à un poids d'abattage donné, peuvent être obtenues à l'aide de programmes alimentaires distincts (niveau énergétique et rapport lysine/énergie). Pour l'acide aminé limitant primaire (lysine), le rapport lysine/énergie se situe à 2,0 g/Mcal d'énergie digestible (E.D.), ou 1,8 g de lysine digestible/Mcal E.D., en prenant en compte la digestibilité de l'acide aminé. Afin d'optimiser les apports recommandés d'énergie et de protéines (lysine) pour un poids d'abattage donné, le niveau et le profil de l'apport énergétique devraient être ajustés au poids final d'abattage. Ceci est très important pour la production de porcs lourds comparativement à ceux abattus à un poids plus faible.

Un niveau très élevé d'apport énergétique dans l'intervalle de poids 35-70 kg entraîne une diminution du gain pondéral et du dépôt de protéines à la fin de la période d'engraissement (au delà de 90 kg de poids vif). Cet effet dépressif ne peut pas être compensé par un haut niveau de complémentation azotée. Un niveau alimentaire trop intensif au cours de la première partie de l'engraissement (plus de 650 g de gain/j) aboutit à des résultats défavorables au poids final de 120 kg. La diminution du dépôt de protéines pendant la deuxième partie de l'engraissement est le reflet d'une expression insuffisante du potentiel de développement musculaire. Ce potentiel est défini par la limite supérieure du dépôt azoté (et par conséquent de la production de tissus maigres) chez un porc à l'engraissement, à un poids final donné, permettant d'atteindre le seuil minimum à la fois pour la durée d'engraissement et le dépôt de gras. Lors de la production de porcs lourds, une réduction par l'apport énergétique pendant la deuxième moitié de l'engraissement n'est pas conseillée dans le cas de l'utilisation de porcs de type maigre, provenant des croisements modernes. Au contraire, il importe que le niveau de l'apport énergétique soit suffisant durant la seconde partie de l'engraissement pour ne pas limiter le dépôt de tissus maigres.

References

- ANDERSEN E., JUST A., 1975. Tabeller over fodermidlers sammensætning m.m. Kvaeg-Svin; Det kgl. Danske Landhusholdningsselskab, København.
- GOLDMANN Sylvia, 1978. *Untersuchungen zum Aminosäurenbedarf wachsender weiblicher Mastschweine*. Thesis, W.-Pieck-Universität Rostock.
- HENRY Y., PION R., RÉRAT A., 1976. Protein supply for pigs and possibilities of reducing protein feeding standards. *World Rev. Anim. Prod.* 12, N° 1 : 9-32.
- HENRY Y., 1979. Feed Intake Regulation for growth in the Pig; *Wiss. Z. W.-P.-Univ. Rostock* 28, 129-135.

- JANCKER H., POPPE S., 1975. Untersuchungen zum Lysinbedarf wachsender Mastschweine (männl. Kastrate) im Lebendmasseabschnitt von 50 bis 115 kg. *Jahrb. Tierernährung u. Fütterung*. **9**, 183-192.
- POPPE S., WIESEMÜLLER W., 1968 Untersuchungen über den Aminosäurenbedarf wachsender Schweine. *Arch. Tierernähr.* **18**, 392-416.
- POPPE S., GOLDMANN Sylvia, HACKL W., WIESEMÜLLER W., 1979. Untersuchungen zum leistungsabhängigen Lysinbedarf von Mastsaugen. *Arch. Tierernähr.* **29**, 349-358; 413-418.
- WIESEMÜLLER W., POPPE S., 1969. Die Bewertung der Proteine durch die Bestimmung der N-Bilanz. *Arch. Tierernähr.* **19**, 133-147.
- WIESEMÜLLER W., POPPE S., 1969. Zum biologischen Wert einiger Futterproteine beim Schwein. *Arch. Tierernähr.* **19**, 157-169, 171-185, 273-279.
- WIESEMÜLLER W., POPPE S., SIEG G., 1974. Untersuchungen zum Aminosäurenbedarf von Mastschweinen (Börge). *Arch. Tierernähr.* **24**, 397-406, 407-419, 453-463, 465-477.
- WIESEMÜLLER W., POPPE S., HACKL W., 1978. Untersuchungen zum leistungsabhängigen Lysinbedarf von Mastschweinen. *Arch. Tierernähr.* **28**, 31-43, 165-171.
- WIESEMÜLLER W., OTTO E., POPPE S., BEHM R., HACKL W., 1978. Untersuchungen zum leistungsabhängigen Lysinbedarf von Mastschweinen. *Arch. Tierernähr.* **28**, 305-316, 479-490.
-