Feeding corn and barley based concentrates to grazing dairy cows. I. Responses to insulin and β -adrenergic stimuli

Ga Gagliostro, Se Lavandera, Sc García, Fj Santini

► To cite this version:

Ga Gagliostro, Se Lavandera, Sc García, Fj Santini. Feeding corn and barley based concentrates to grazing dairy cows. I. Responses to insulin and β -adrenergic stimuli. *Annales de zootechnie*, 1995, 44 (Suppl1), pp.278-278. hal-00889437

HAL Id: hal-00889437

<https://hal.science/hal-00889437>

Submitted on 1 Jan 1995

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Feeding corn and barley based concentrates to grazing dairy cows. I. Responses to insulin and β -adrenergic stimuli

GA Gagliostro¹, SE Lavandera², SC García¹, FJ Santini¹

¹INTA EEA Balcarce, CC 276-7620 Balcarce, ²Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Argentine

A high protein (CP) non structural carbohydrate (NSCH) ratio in autumn-winter forages lead to high rumen ammonia (N-NH₃) levels (16-60 mg/dl). Feeding concentrates with different ruminal starch degradability may be a way to synchronize protein/energy availabilities reducing the risk of N-NH₃ excess. Hyperammonemia may induce insulin resistance and enhanced β -adrenergic responses (Fernández *et al*, 1988, J Anim Sci, 66, 3259-3266; Visek, 1984, J Dairy Sci, 67, 481-498; Symonds *et al*, 1981, Br J Nutr, 46,481-486).

Twenty seven Holstein cows (85 $\pm$ 17 days postpartum) grazing an oat sward (CP = 23.9 %, NSCH = 16.9 %) and a pasture of ryegrass (CP = 15.9 %, NSCH = 19.5 %) in winter were assigned to three treatments : PA : only forage, CO : forage plus 7 kg/cow/d of a corn (75 %) based concentrate and BA : forage plus 7 kg/cow/d of a barley (85 %) based concentrate. Preprandial jugular blood samples were taken (5 wks) to evaluate metabolite concentrations. Samples were taken before (INS-0), 30 (INS-30), 60 (INS-60) and 90 (INS-90) minutes after an intravenous bovine insulin challenge (0.12 units/kg body weight) (wk 2) and before (ISO-0) and 15 minutes after (ISO-15) isoproterenol challenge (4 nmol/kg body weight) (wk 3). Urea, triglycerides (TG), glucose and free fatty acids (FFA) were determined using enzymatic kits. Data were analysed using a split-plot design with treatment as main plot and week or sampling time as secondary plot.

Intake of concentrates was 6.3 and 5.3 kg/cow/d in CO and BA. Forage dry matter intake was higher in PA respect to CO and BA (11.93 vs 11.02 and 10.75 kg DM/cow/d, $P < 0.05$). Milk yield was higher in CO (21.5 kg/cow/d) respect to PA (17.0) ($P < 0.05$) but similar between BA (18.4 kg/cow/d) and the other treatments. Milk fat was lower in BA (2.84 %) respect to MA (3.01 %) and PA (3.45 %). Rumen N-NH₃ (six additional fistulated cows) was 15.7, 14.1 and 15.7 mg/dl in PA, CO and BA ($P < 0.15$) (range 5 to 26 mg/dl). Plasma urea was higher in PA (24.2 mg %) respect to BA (21.2 mg %) ($P < 0.05$) and similar between CO (22.3 mg %) and the other treatments. Plasma urea correlated with milk yield (-0.204, $P < 0.05$) and FFA (0.28, $P < 0.01$). Glucose, TG and FFA were similar across treatments. Glucose and FFA concentrations for ISO-0 were 74.7, 71.8 and 70.0 mg/dl and 558, 412 and 364 μ eq/l in PA, CO and BA respectively ($P > 0.05$). ISO-15 values were 84.3, 81.3, 80.0 mg/dl for glucose and 906, 722, 770 μ eq/l for FFA in PA, CO and BA respectively ($P > 0.05$). Changes in plasma glucose and FFA concentrations after INS challenge did not differ.

Higher β -adrenergic responses or resistance to insulin action were not observed in cows grazing pastures rich in CP in winter but in the experiment, the forage NSCH content was high and N-NH₃ and plasma urea were relatively low.

	PA	CO	BA	Mean
<i>Glucose (mg/dl)</i>				
INS-0	78.2	75.6	77.1	76.9 ^a
INS-30	40.1	34.0	40.27	38.1 ^b
INS-60	50.1	48.7	51.5	50.1 ^c
INS-90	65.1	60.5	64.4	63.3 ^d
<i>FFA (μeq/l)</i>				
INS-0	338	309	401	349 ^a
INS-30	271	246	271	262 ^b
INS-60	381	328	334	348 ^a
INS-90	490	359	393	414 ^c

a,b,c,d : means with different letters differ ($P < 0.05$). Neither treatment nor interaction effects were detected