

Performances of dairy cows grazing on alpine pasture under a Leaders-Followers system

P Berzaghi, I Andrighetto, G Cozzi

► **To cite this version:**

P Berzaghi, I Andrighetto, G Cozzi. Performances of dairy cows grazing on alpine pasture under a Leaders-Followers system. *Annales de zootechnie, INRA/EDP Sciences*, 1995, 44 (Suppl1), pp.369-369. <hal-00889523>

HAL Id: hal-00889523

<https://hal.archives-ouvertes.fr/hal-00889523>

Submitted on 1 Jan 1995

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Performances of dairy cows grazing on alpine pasture under a Leaders-Followers system

P Berzaghi, I Andrighetto, G Cozzi

Department of Animal Science, University of Padua, Italy

At the beginning of the grazing season it is usual to observe, in the mountain regions, a severe drop in milk yield in high producing cows (Andrighetto and Ramanzin, 1987, Zoot Nutr Anim, 13, 119). The present study aims to evaluate in an alpine pasture the Leaders-Followers grazing system (LF) as a management tool to limit this decrease in production.

The study was carried out in a commercial farm located in the north-west area of the Italian Alps on an elevation of 1000 m above the sea level. The pasture (23.5 ha), divided into four paddocks, was composed of more than 30 species, mainly grasses (*Festuca rubra*, *Poa pratensis* and *trivialis*) and weeds (*Deschampsia caespitosa*). The milking herd (36 Brown Swiss cows) was divided into two groups (Control and LF) balanced for the milk yield recorded before the beginning of the grazing season. Each group was allowed to graze half of each paddock divided by electric fences. The LF group grazed the assigned areas under a Leaders-Followers system dividing the animals between Leaders and

Followers according to milk produced. During milking all animals received a commercial supplement (CP : 18.6 % DM ; NDF : 16.1 % DM ; UFL : 1.02 /kg DM). Daily milk yield was recorded at wk 4 and 8 of the grazing trial and herbage and milk samples were collected for composition analysis.

Although Leaders had the opportunity to graze each paddock first, the advantage in terms of herbage quality was low and the forage available to the LF and Control had similar composition. Under the Leaders-Followers grazing system high producing animals were able to maintain a milk yield above 20 kg/d but with a decrease due to grazing, similarly to lower producing animals. However, as indicated by the results of milk persistency, on average, the reduction in milk yield for LF was greater than Control cows.

In conclusion, in alpine pastures, the Leader-Followers grazing system is able to sustain milk yield of the more productive animals but it does not offer any advantage compared to the conventional rotational grazing.

	Leaders	Followers	Avg LF	Control	Significance ¹
Herbage composition					
CP	13.4	12.1	12.8	12.2	NS
NDF	63.5	66.1	64.8	64.4	NS
Milk production (kg/d)					
before grazing	27.3	17.4	22.4	22.6	NS
grazing	23.0	12.9	18.0	19.0	NS
Persistency (%)	84.0	72.4	78.2	83.4	NS
Milk composition (%)					
fat	3.6	3.7	3.7	3.7	NS
protein	3.4	3.7	3.6	3.5	NS

¹ Leaders and Followers vs Control