A study of molecular mechanisms for spreading of kanamycin resistance gene among ruminal bacteria

P Pristas, I Vanat, N Kostrabova, P Javorsky

► To cite this version:

P Pristas, I Vanat, N Kostrabova, P Javorsky. A study of molecular mechanisms for spreading of kanamycin resistance gene among ruminal bacteria. *Annales de zootechnie*, 1996, 45 (Suppl1), pp.344-344. hal-00889655

HAL Id: hal-00889655

<https://hal.science/hal-00889655>

Submitted on 1 Jan 1996

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A study of molecular mechanisms for spreading of kanamycin resistance gene among ruminal bacteria

P Pristas, I Vanat, N Kostrabova, P Javorsky

Institute of Animal Physiology, Slovak Academy of Sciences, Soltésovej 4-6, 04001 Kosice, Slovakia

The rumen is one of the most complex microbial ecosystems in the world, and forms an environment especially suited for the study of gene movement and exchange in natural conditions. Antibiotic resistance genes present a good model for studies. In our previous work (Pristas et al, in press) we have detected a *Tn 5* like *aphII* gene among kanamycin-resistant ruminal streptococci and *Selenomonas*. From both microorganisms resistance genes were cloned and shown to confer resistance to kanamycin sensitive *Escherichia coli*. The abundance of *aphII* like sequences in populations of bacteria was analyzed by DNA-DNA hybridization.

It was found that different strategies are used by Gram-negative resp. Gram-positive bacteria:

-from Gram-negative *Selenomonas rumi-*

nantium kanamycin-resistance was transferred to *E. coli* using free DNA transformation. Probably plasmids and classical type II transposons are involved in the spreading of *aphII* resistance gene;

-a different situation occurred for the Gram-positive streptococci. No plasmid DNA was detected in a strain of *Streptococcus bovis*. However, it was shown that a cloned streptococcal kanamycin resistance determinant could be transferred between *E. coli* cells by conjugation. Upon conjugation, site specific integration of the *aphII* determinant into host genome was observed, resembling those found for some integrons (Collins et al, 1993, Mol Microbiol, 9, 41-52).

Detailed analysis both of elements involved in the spreading of kanamycin resistance among ruminal bacteria is under progress.