

Electroporation of G+ host plasmids into *Selenomonas ruminantium*

J Kopecny, V Kostyukovsky, K Fliegerova

► **To cite this version:**

J Kopecny, V Kostyukovsky, K Fliegerova. Electroporation of G+ host plasmids into *Selenomonas ruminantium*. *Annales de zootechnie*, INRA/EDP Sciences, 1996, 45 (Suppl1), pp.356-356. <hal-00889667>

HAL Id: hal-00889667

<https://hal.archives-ouvertes.fr/hal-00889667>

Submitted on 1 Jan 1996

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Electroporation of G⁺ host plasmids into *Selenomonas ruminantium*

J Kopecny, V Kostyukovsky, K Fliegerova

*Institute of Animal Physiology and Genetics, Czech Academy of Sciences, Prague 10,
Uhrineves 104000, Czech Republic*

There are now several systems by which new genetic material can be introduced into rumen bacteria as plasmids (Thomson et al, 1991, *Curr Microbiol*, 24, 49-54). The technique that has proven generally useful for inserting new DNA is that of electroporation. Combining this with plasmids originating from the bacterial species to be modified has led to at least two of the more successful processes reported (Coconcelli et al, 1992, *FEMS Microbiol Lett*, 941, 203-208; Beard et al, 1995, *Curr Microbiol*, 30, 105-109).

All these experiments were done with *Butyrivibrio*, *Bacteroides* and *Ruminococcus* species. With *Selenomonas ruminantium* we have tried to introduce our constructs with ampicillin resistance from *E. coli* with the help of electroporation. Transformants were usually very unstable (Kopecny and Fliegerova, 1994, *Proc Soc Nutr Physiol*, 3, 150).

In this case plasmid pJW1 of *Selenomonas ruminantium* was cut with Pst1 and subcloned

into pG⁺ host 4 in *E. coli* strain SURE. The vector was chosen because of some G⁺ features of rumen selenomonas (Flint, personal communication). The construct pJW1/pG⁺ host4 (Pst1) was isolated and orientation of pJW1 insert verified. For transfection was used apparatus TA 750 (Krüss, Germany). Electroporation buffer (EBc) was composed of 10 mM PIPES pH 7, 15% glycerol and 60 mM CaCl₂. Buffer was prepared anaerobically and autoclaved. Into the transfection chamber 1 ml of culture in EB and 10-50 µl (10-20 ng/µl) of plasmid was added. Treatments were done in range of 1000-15000 V/cm with 100-800 µs duration and 1-9 pulses. Temperature of the electroporation cell and cell culture was 4°C. After electroporation cells were cultivated for 18 hours in M8 medium at 33°C and then selected on M8-agar with erythromycin.

Advantage of these clones is the thermostability of the plasmid pJW1/pG⁺host up to 35°C.