

Varroa jacobsoni as a carrier of bacterial infections to a recipient bee host

Z Gliński, J Jarosz

► **To cite this version:**

Z Gliński, J Jarosz. Varroa jacobsoni as a carrier of bacterial infections to a recipient bee host. Apidologie, Springer Verlag, 1992, 23 (1), pp.25-31. hal-00890967

HAL Id: hal-00890967

<https://hal.archives-ouvertes.fr/hal-00890967>

Submitted on 1 Jan 1992

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

***Varroa jacobsoni* as a carrier of bacterial infections to a recipient bee host**

Z Gliński¹, J Jarosz^{2*}

¹ Agricultural University, Bee Diseases Research Laboratory, Lublin;
² Marie Curie-Skłodowska University, Department of Insect Pathology,
Akademicka 19, Lublin, Poland

(Received 23 December 1988; accepted 20 September 1991)

Summary — Using *Serratia marcescens* as an indicator bacterium and the worker brood of the honey bee, experiments have shown that *Varroa jacobsoni* transmits bacterial infections to a recipient bee host. Integument and haemolymph of brood artificially contaminated with *S. marcescens*, preferably individuals with *Serratia septicaemia*, were the source of bacterial contamination of the mites. The infected parasites during feeding carry *S. marcescens* to about 20% of the healthy brood.

***Varroa jacobsoni* / *Serratia marcescens* / carrier / bacterial infection / septicaemia**

INTRODUCTION

One of the factors that restricts the development of insect diseases caused by bacterial invaders is low capacity of entomopathogenic bacteria to disperse throughout the host environment and ultimately to infect a susceptible host. Bacterial pathogens lack means of locomotion and active penetration *via* the insect integument, and therefore must rely on biotic and abiotic agents for their dispersal in an insect niche and penetration into the host

haemocoel. In the case of biotic agents, bacteria may be transmitted by healthy and infected hosts, parasites and predators. The external stenophagous parasites which feed on infected hosts or on hosts killed by diseases, even though they are not susceptible to the bacteria, are capable of distributing the pathogens to their common hosts.

Bees may become susceptible to viral, bacterial and fungal infections when they are attacked by external parasites that damage their integument by ruptures or

* Correspondence and reprints

wounds. Once infection has been initiated by *V jacobsoni* as a vector or inducer of acute bee paralysis virus (Ball, 1983; Ball and Allan, 1986; Wiegres, 1986), regardless of the initial density of viral pathogen, a causal agent is transmitted to recipient hosts and numbers of sick bees increase in a colony. Data suggests that bacterial infections could start from microbes present on a host body surface and from microbial flora in the bee environment that the *V jacobsoni* mite transmits mechanically or biologically (Smirnov and Kudriavcev, 1977; Horn, 1984; Shabanov, 1984; Strick, 1986; Strick and Madel, 1986).

The transmission of bacteria by *V jacobsoni* to bees has not been established, though the presence of *Hafnia alvei* septicaemia in colonies severely invaded by *V jacobsoni* supports this possibility (Kutzenko, 1975). It has recently been shown that *Serratia marcescens* can be transmitted from artificially contaminated brood and workers of *Apis mellifera* to *V jacobsoni* mites (Gliński and Jarosz, 1990). In this paper we present evidence from microbiological bioassays, using *S marcescens* as an indicator bacterium, that show it is possible to infect healthy brood with mites fed on infected bee larvae.

MATERIALS AND METHODS

Serratia marcescens, a prodigiosin producing strain CCM 2222 obtained from the Culture Collection of Microorganisms, University of Purkinje (Brno, Czechoslovakia), was used as an indicator bacterium. This facultative bee pathogen multiplies after entering the *Apis mellifera* subsp *mellifera* haemocoel and produces a lethal septicaemia. Red colour of *Serratia* colonies on bacteriological media and a characteristic red colour of infected bee cadavers enables identification of *S marcescens* on microbiological media and detection of septicaemia caused by this pathogen.

Females of *V jacobsoni* collected from honeybee prepupae (13 days old) and pupae (15 days old) were starved for 10–12 hours before colonization of brood subcuticle injected or surface contaminated with a log phase of a broth culture of *S marcescens*. Viable cells for inoculation of larvae were prepared by diluting an overnight culture of *S marcescens* (incubated at 28 °C) to a specific concentration in insect Ringer (130 mM NaCl, 5.0 mM KCl and 1.0 mM CaCl₂). Larvae (10 days old) were from capped cells and, after uncapping, were injected in the thoracic region with 2.0 µl of bacterial suspension (about 2 000 cells) or sprayed with about 50 000 cells of the pathogen. Two hours later the cells were individually infested with *V jacobsoni*, 2–3 females per larva, and artificially capped by hand with bee wax. The brood was then reared in an incubator at 34 °C. Bee larvae without any contact with *S marcescens* but infected with the mite, and bees sprayed or injected with a sterile insect Ringer alone, served as controls.

The total number of *S marcescens* associated with the parasite fed on brood was estimated by an agar plate colony count technique after 24 and 72 hours post-colonization. Usually 7 randomly chosen mites were individually blended in sterile insect Ringer solution (1 parasite per 0.2 ml) in a glass tissue grinder. Serial 1/10 dilutions of *Varroa* homogenate were then prepared in the insect Ringer and 10 µl of the solution was streaked in triplicate onto agar medium. The number of *S marcescens* associated with *V jacobsoni* was evaluated after incubation at 28 °C for 2 days.

Multiplication of *S marcescens* in body of bee larvae artificially injected or surface sprayed with bacterial suspension was monitored after 72 hours in larval body homogenates. Specimens were thoroughly rinsed 3 times by dipping them into 10 ml of insect Ringer. After checking the surface sterility of larvae by rolling them on an agar plate, the larvae were blended in 10 ml of Ringer solution. Larval homogenates were then diluted serially in a sterile insect Ringer and inoculated onto agar plates to detect the titer of *S marcescens* present in *Apis mellifera* larvae.

To test for the transmission of bacteria by *V jacobsoni*, female mites that fed 72 hours on injected or surface sprayed *A mellifera* brood were moved onto 100 uncapped healthy prepupae (2–3 mites/specimen). The experiments

were replicated 3 times. Brood capped with bee wax was then reared in an incubator. The symptoms of *S marcescens* septicaemia were observed 4 and 5 days after brood exposure to infected *Varroa* mites. Dead larvae with developing red colour were analyzed bacteriologically to detect multiplication of *S marcescens*.

RESULTS

V jacobsoni females parasitizing honey bee larvae artificially infected with *S marcescens* contained from many thousands to about a million bacterial cells per mite (table I). Only a few parasites carried less than one thousand cells of this pathogen. Interestingly, the number of *S marcescens* associated with the mite increased markedly with parasitization time. Bacterial population density reached approximately 10^6 cells in all mites after 72 hours of feeding of the infected brood. The indicator bacterium attained a dense population in the mite independently of the means of brood infec-

tion (table I), although 24 hours post-infestation the count of bacteria present in *Varroa* fed on injected brood was evidently lower than that in mites fed on brood sprayed with the pathogen.

That the source of the bacterial invader for *Varroa* is the haemolymph of *A mellifera* indicates the development of *S marcescens* bacteraemia in the host sprayed or injected with the pathogen (table II). Our bacteriological assays have shown that bee haemolymph is a suitable medium for multiplication of *S marcescens*. Once infection had been initiated in the bee, regardless of the route of infection and initial level of the bacterial population, pathogens increased rapidly in number, usually $> 5 \times 10^7$ cells/larva and bees died with typical colour symptoms of *S marcescens* septicaemia. Few control bees treated with Ringer alone, by spraying or injection, died during 3 days post-treatment, but neither symptoms of *Serratia* bacteraemia nor *S marcescens* cells were found in dead specimens.

Table I. *Serratia marcescens* artificially contaminating upright larvae of *Apis mellifera* pollutes *Varroa jacobsoni* mite.

Specimen	No of <i>S marcescens</i> per <i>Varroa jacobsoni</i> after :			
	24 h		72 h	
	<i>feeding on brood</i>			
	<i>Sprayed</i>	<i>Injected with the bacterial invader</i>	<i>Sprayed</i>	<i>Injected</i>
V ₁	7185	930	8470	10 ⁶
V ₂	10 ⁶	42	10 ⁶	10 ⁶
V ₃	3150	1655	10 ⁶	10 ⁶
V ₄	5825	110	10 ⁶	10 ⁶
V ₅	1655	230	10 ⁶	10 ⁶
V ₆	2715	1580	10 ⁶	10 ⁶
V ₇	1407	112	10 ⁶	10 ⁶

Table II. The multiplication of *Serratia marcescens* in upright larvae of *Apis mellifera*. Count 72 h after spray or injection of the pathogen.

Specimen	Spray	Injection
A ₁	N	N
A ₂	N	N
A ₃	N	3 x 10 ⁷
A ₄	N	N
A ₅	N	N
A ₆	N	N
A ₇	N	N
A ₈	N	N
A ₉	N	2.5 x 10 ⁷
A ₁₀	N	3 x 10 ⁷

N: more than 5 x 10⁷ cells per larva.

It is generally claimed that predators which feed on injected hosts or on host killed by disease, even though not susceptible to the pathogens, are capable of distributing the pathogens to their common hosts. When *V jacobsoni* females

infected with *S marcescens* were transferred to freshly uncapped healthy bee larvae, *S marcescens* septicaemia developed in many larvae (table III).

DISCUSSION

The *V jacobsoni* mite, because of its damage to honey bees, has received a great deal of interest worldwide (Ritter, 1981; Taber, 1986) and research has elucidated some mechanisms responsible for the deleterious effects of the parasite on the host. *V jacobsoni* has been implicated as a vector of viral infections and inducer of latent infections caused by acute bee paralysis virus (Ball, 1983; Ball and Allen, 1986). In contrast, knowledge of the role of *V jacobsoni* as a carrier of bacterial invader is fragmentary and still contradictory (Gliński and Jarosz, 1988). Obviously, infection *per se* is dependent on the pathogenic properties of microorganisms, their dosage, contact of pathogens with the host and point of entry of the pathogen into a recipient host. One of the factors that restricts the development of diseases caused by entomo-

Table III. The occurrence of *Serratia marcescens* septicaemia in the recipient bee larvae parasitized with *V jacobsoni* contaminated by this bacterium.

Batch	Cumulative percentage of mortality of brood with symptoms of <i>Serratia marcescens</i> septicaemia	
	Days post-treatment	
	4	5
I: 100 worker larvae	18.0	21.0
II: 100 worker larvae	14.0	18.0
III: 100 drone larvae	12.0	13.0
Mean	14.6	17.3

athogenic bacteria is their low capacity to disperse throughout the host environment. Bacteria are generally dispersed by infected hosts, contaminated food and by mechanical or biological vectors.

As parasites, the feeding behaviour of *V jacobsoni* may create conditions favouring transfer of bacteria within bee colonies and their penetration into the bee haemocoel by damaging host integument. Dense populations of *S marcescens* (table I) associated with surface or interior of the mite univocally point to *V jacobsoni* as a carrier of this facultative bee pathogen. Moreover, a luxuriant multiplication of *S marcescens* in the host haemocoel and development of fatal septicaemia in bees fed on the parasite enables this bacteria to be transmitted to subsequent hosts (see tables II and III). Of course, leakage of haemolymph through the host wounds preferably contaminates the mouthparts of *Varroa*, thus creating favourable conditions for infection of healthy bee hosts by a direct introduction of bacteria into fresh wounds.

In the light of the studies on pathogenesis of *Varroa* disease, the impairment of host blood antibacterial factors by parasitizing mites (Gliński and Jarosz, 1984; Gliński and Jarosz, 1987) enables development of infection, multiplication of the pathogen in haemocoel and production of lethal septicaemia in the bee. In the experiments carried out, we observed a high percentage of brood dying of bacterial septicaemia after contact with mites which had been infected with *S marcescens* during parasitization on artificially-infected bee larvae. *Serratia marcescens* is known to be a facultative pathogen of *Apis mellifera* causing the death of brood and bees. Therefore, it is reasonable to postulate that transfer of dangerous bee pathogens such as *Melisococcus pluton* or *Ascosphaera apis* by *V jacobsoni* may constitute a threat to beekeeping. Indeed, the incidence of

European foulbrood (Trubin, 1985) and chalkbrood disease (Chmielewski and Gliński, 1987; Gliński, 1988) has been shown to increase in colonies infested with *V jacobsoni*. We conclude that transmission of bacterial infections by *V jacobsoni* to the honey bee constitutes another detrimental effect of this parasite on its host.

Résumé — *Varroa jacobsoni* en tant que vecteur de maladies bactériennes chez son hôte, l'abeille domestique.

Les mécanismes par lesquels *Varroa jacobsoni*, acarien ectoparasite très spécialisé, provoque des dégâts chez son hôte *Apis mellifera* n'ont pas été déterminés de façon définitive. Parmi les diverses possibilités de dégâts, il existe des preuves selon lesquelles *V jacobsoni* est capable de transmettre des bactéries pathogènes à l'hôte qui l'héberge. On a utilisé comme indicateur la bactérie *Serratia marcescens*, qui provoque chez les larves et les nymphes d'*A m mellifera* une septicémie, identifiable par sa coloration rouge. On a mis en évidence que ce germe pathogène reste attaché à *V jacobsoni*, qui peut ainsi servir de vecteur. *V jacobsoni*, qui s'alimente sur du couvain infesté artificiellement par *S marcescens* (par vaporisation en surface ou injection sous la cuticule), contamine ses pièces buccales et la surface de son corps avec des cellules bactériennes (tableau I). Les tests microbiologiques ont également montré la multiplication de la bactérie dans l'acarien, indiquant là encore son rôle possible de vecteur. En outre, le couvain atteint de septicémie, au sein duquel *S marcescens* se multiplie et atteint des populations élevées (tableau II), est une source de bactéries pour les acariens parasites. Le début d'une septicémie, observée sur du couvain operculé sain mais infesté par *V jacobsoni* (tableau III), prouve de façon indubitable que l'acarien a trans-

mis l'infection bactérienne à la colonie d'abeilles.

***Varroa jacobsoni* / *Serratia marcescens* / vektor / bakteriose / septicémie**

Zusammenfassung — *Varroa jacobsoni* als Überträger bakterieller Infektionen auf ihren Wirt, die Honigbienen. Die Mechanismen, durch die *Varroa jacobsoni*, eine außenparasitische, eng spezialisierte Milbe, ihren Wirt *Apis mellifera* schädigt, sind noch nicht endgültig geklärt. Neben anderen Möglichkeiten der Schädigung gibt es auch Hinweise darauf, daß *Varroa* bakterielle Keime auf ihren Wirt übertragen kann. Es wurde das Bakterium *Serratia marcescens* als Indikator benutzt, das bei Larven und Puppen von *A mellifera mellifera* unter gleichzeitiger Rotfärbung der Tiere eine Septikämie verursacht. Es wurde festgestellt, daß dieser fakultative Krankheitskeim an *Varroa jacobsoni* haften bleibt und daß diese Milbe somit als Überträger von *S marcescens*-Infektionen dienen kann. *V jacobsoni*, die an künstlich (durch Spray oder Injektion in die Körperhöhle) mit *S marcescens* infizierter Brut saugten, verunreinigten ihre Mundwerkzeuge und die Körperoberfläche mit Bakterienzellen (Tab I). Mikrobiologische Tests wiesen darauf hin, daß sich dieses Bakterium in der Milbe vermehrt, so daß *V jacobsoni* auch die Rolle eines biologischen Vektors von Krankheitskeimen spielen könnte. Außerdem ist septikämische Brut, in der *S marcescens* zu einer großen Population anwächst (Tab II), eine Bakterienquelle für parasitierende Milben. Der Ausbruch einer *S marcescens*-Septikämie, beobachtet in gesunder verdeckelter Brut, die aber mit *V jacobsoni* befallen war (Tab III), weist eindeutig auf die Milbe als Überträgerin bakterieller Infektionen im Bienenvolk.

***Varroa jacobsoni* / *Serratia marcescens* / Überträger / bakterielle Infektion / Septikämie**

REFERENCES

- Ball BV (1983) Der Zusammenhang zwischen *Varroa jacobsoni* und Virus – erkrankungen der Honigbiene. *Allg Dtsch Imkerztg* 17, 177-179
- Ball BV, Allen MF (1986) The incidence of acute paralysis virus in honey bee colonies infested with the parasitic mite *Varroa jacobsoni*. In: *Fundamental and Applied Aspects of Invertebrate Pathology* (Samson RA, Vlcek JM, Peters D, eds) 4th Int Colloq Invertebr Pathol, p 151
- Chmielewski M, Gliński Z (1987) Effects of *Varroa jacobsoni* Oud on the incidence and course of chalk-brood disease in *Apis mellifera*. Proc 31st Int Congr Apic August 19-25, Warsaw, Apimondia Publ House, Bucharest, p 104
- Gliński Z (1988) *Varroa jacobsoni* affects the incidence and course of chalk-brood disease in *Apis mellifera* colonies. *Ann Univ Mariae Curie-Sklodowska* DD 43, 23-27
- Gliński Z, Jarosz J (1984) Alterations in haemolymph proteins of drone honey bee larvae parasitized by *Varroa jacobsoni*. *Apidologie* 15, 329-338
- Gliński Z, Jarosz J (1987) *Varroa jacobsoni* parasitization lowers the innate concentration of blood lysozyme of upright larvae of the worker honey bee, *Apis mellifera*. Proc 31st Int Congr Apic August 19-25, Warsaw, Apimondia Publ House, Bucharest, p 121
- Gliński Z, Jarosz J (1988) Deleterious effects of *Varroa jacobsoni* on the honey bee. *Apiacta* 23, 42-55
- Gliński Z, Jarosz J (1990) *Serratia marcescens* artificially contaminating brood and worker honey bees pollutes *Varroa jacobsoni* mite. *J Apic Res* 29, 107-111
- Horna H (1984) Zum Zusammenhang zwischen *Varroa jacobsoni* und Bakteriosen bei der Honigbiene. *Allg Dtsch Imkerztg* 18, 328-329
- Kutzenko DM (1975) On the role of the gamasid mite *Varroa jacobsoni* (Oudemans 1904)

- transmitting hafniosis in bees. *Blagovechtchensk* 2, 50-52 (in Russian)
- Ritter W (1981) *Varroa* disease of the honeybee, *A mellifera*. *Bee World* 62, 141-153
- Shabanov M (1984) The role of the mites *Varroa jacobsoni* as a carrier of microorganisms in the bee hive. *Acta Microbiol Bulgarica* 15, 78-82 (in Bulgarian)
- Smirnov AM, Kudriavcev EA (1977) The *Varroa* mite and diseases from contamination. *Pcelovodstvo* 0(5), 13-14 (in Russian)
- Strick H (1986) Die Rolle der ektoparasitischen Bienenmilbe *Varroa jacobsoni* Oudemans als möglicher Bakterienüberträger bei ihrem Wirt, der Honigbiene *Apis mellifera* Linnaeus. Diplomarbeit an der Naturwissenschaftlichen Fakultät der Rheinischen Friedrich-Wilhelms Universität, Bonn, 118 pp
- Strick H, Madel G (1986) Varroatose und bakterielle Sekundärseuchen. *Allg Dtsch Imkerztg* 20, 321-325
- Taber S (1986) Reports on *Varroa* disease of honey bees. *Am Bee J* 126, 430-431
- Trubin AV (1985) Varroatosis and European foul brood. *Pcelovodstvo* 0(8), 19 (in Russian)
- Wieggers FP (1986) Transmission of acute paralysis virus by the honeybee parasite *Varroa jacobsoni* Oud. In: *Fundamental and Applied Aspects of Invertebrate Pathology* (Samson RA, Vlcek JM, Peters D, eds) 4th Int Colloq Invertebr Pathol, Wageningen, p 152