



Juvenile hormone titer and reproduction of *Varroa jacobsoni* in capped brood stages of *Apis cerana indica* in comparison to *Apis mellifera ligustica*

P Rosenkranz, Nc Tewarson, A Rachinsky, A Strambi, C Strambi, W Engels

► To cite this version:

P Rosenkranz, Nc Tewarson, A Rachinsky, A Strambi, C Strambi, et al.. Juvenile hormone titer and reproduction of *Varroa jacobsoni* in capped brood stages of *Apis cerana indica* in comparison to *Apis mellifera ligustica*. *Apidologie*, 1993, 24 (4), pp.375-382. hal-00891081

HAL Id: hal-00891081

<https://hal.science/hal-00891081>

Submitted on 11 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Juvenile hormone titer and reproduction of *Varroa jacobsoni* in capped brood stages of *Apis cerana indica* in comparison to *Apis mellifera ligustica*

P Rosenkranz ^{1,*}, NC Tewarson ², A Rachinsky ¹,
A Strambi ³, C Strambi ³, W Engels ¹

¹ LS Entwicklungsphysiologie, Zoologisches Institut der Universität, Auf der Morgenstelle 28, D-72076 Tübingen, Germany;

² Department of Zoology, Ewing Christian College, Allahabad-211003, India;

³ Laboratoire de Neurobiologie, CNRS, BP 71, F-13402 Marseille Cedex 9, France

(Received 24 November 1992; accepted 19 February 1993)

Summary — The juvenile hormone (JH) III hemolymph titer of late larval and early pupal stages as determined by radioimmunoassay (RIA) does not differ significantly in *Apis cerana* and *Apis mellifera*, particularly in freshly-sealed worker brood. In drone brood, slightly higher JH III hemolymph concentrations were recorded. These results do not agree with the hypothesis that reproduction of the parasitic bee mite *Varroa jacobsoni* is regulated by host-derived JH. After checking over a thousand capped brood cells containing pupae, it was confirmed that in *Apis cerana* colonies fertility of female mites is restricted to drone hosts.

Apis cerana indica* / *Apis mellifera ligustica* / juvenile hormone titer / reproduction / *Varroa jacobsoni

INTRODUCTION

In honey bees, juvenile hormone (JH) has been studied mostly under aspects of caste development (Hartfelder, 1990; Rembold *et al*, 1992), control of fertility (Engels *et al*, 1990), and recently also polyethism regulation (Robinson, 1992).

Modern analytical techniques allow the determination of hemolymph titer (Rachinsky *et al*, 1990) and rate of synthesis (Rachinsky and Hartfelder, 1990) even in individual bees. Since in arthropods (Downer and Laufer, 1983) and also in acarids, ticks (Pound and Oliver, 1979; Connat *et al*, 1983) as well as mites (Oliver *et al*, 1985)

* Present address: Bayerische Landesanstalt für Bienenzucht, Burgbergstr 70, D-91054 Erlangen, Germany

exogenous JH was generally found to affect reproduction, it was hypothesized that this hormone could also regulate oogenesis in the parasitic bee mite *Varroa jacobsoni*, and, in addition, that host-derived JH could be responsible for initiation of reproduction (Hänel, 1983). Particularly, a high JH III hemolymph titer in drone larvae of the early postcapping phase was considered to stimulate *Varroa* vitellogenesis in the original host species, *Apis cerana*, and likewise in worker larvae of the western honey bee, *Apis mellifera* (Hänel and Koeniger, 1986). Thus a low JH III titer in late 5th instar worker larvae would be a resistant factor, restricting mite reproduction to drone brood.

However, in the only known case of varroa tolerance in *Apis mellifera*, the Africanized honey bees of Brazil (Engels *et al*, 1986), fertility of female mites in worker brood is also reduced (Ritter and De Jong, 1984) but was not found to be correlated with a low JH III hemolymph titer in worker larvae (Rosenkranz *et al*, 1990). These data already contradicted the hypothesis of a possible regulation of *Varroa* reproduction by the host's JH. Because detailed JH III titer measurements were lacking in *Apis cerana* larvae, we analyzed hemolymph collected from Indian eastern honey bees together with *Apis mellifera ligustica* samples at the same study site. The fertility of *Varroa* females on drone and worker hosts of *Apis cerana indica* was also evaluated.

MATERIALS AND METHODS

Hemolymph samples were collected during the spring season, March 1991, from 4 colonies of *Apis cerana indica* kept in Newton hives (Tewarson *et al*, 1992) and 1 colony of *Apis mellifera ligustica* originating from the US in a Langstroth hive. All experimental colonies had 5 brood combs or more and were kept on the campus of Ewing Christian College, Allahabad, north India.

The time of brood cell capping was recorded with the help of transparent sheets. Subsequently larvae were at first taken at 6-h intervals, and later on less frequently. For each sample 40–120 µl hemolymph from non-parasitized specimens were collected from 3–9 individuals, pooled, extracted with hexane, coded and stored at –20°C. After being transported to Europe on dry ice, JH III titer determinations were carried out after diol derivatization by radioimmunoassay (Strambi *et al*, 1981; Rosenkranz *et al*, 1990). In the *Apis mellifera ligustica* colony only little drone brood was present and was insufficient for sampling all required stages, probably due to high ambient temperatures.

In order to evaluate the actual reproduction of *Varroa jacobsoni* in the *Apis cerana indica* colonies, ~ 1 100 capped brood cells containing pupae were opened and checked for female mites and their offspring. Because of severe brood infestation with *Tropilaelaps clareae*, no comparative data could be sampled from the *Apis mellifera ligustica* colony.

RESULTS

Juvenile hormone hemolymph titer in postcapping stages of worker brood

At the time of brood cell sealing in worker larvae, the JH III hemolymph titer was found to be low, ranging < 10 pmol/ml (fig 1a). However, soon after cell capping a steady increase was measured. Peak values of 70–100 pmol/ml were reached ~ 30–48 h post cell operculation. There was not much variation in the different pool data sampled during that period after capping. On average, in 5th instar larvae of *Apis cerana indica* slightly higher JH concentrations were determined than in *Apis mellifera ligustica*, but the differences were statistically insignificant ($P > 0.7$; Friedman test) even around the peak values. The duration of the JH III maximum was ~ 24 h, followed by a decrease in titer reached ~ 96 h after cell capping, and again by another increase in the pupal stages (fig 1a).




Fig 1. Hemolymph titer of juvenile hormone III in postcapping brood stages of eastern (○, — *Apis cerana indica*) and western (■, — *Apis mellifera ligustica*) honey bees. (a) = workers, (b) = drones. pw = white-eyed pupae, pr = red-eyed pupae, pb = brown-eyed pupae. Lines = mean values the differences of which in (a) are not significant ($P > 0.7$; Friedman test). The few data on drone brood of western honey bees (b) are insufficient for calculating a mean titer curve.

Juvenile hormone hemolymph titer in postcapping stages of drone brood

Within the first 48 h after brood cell operculum in drone larvae of *Apis cerana indica* the JH III hemolymph titer varied around 50 pmol/ml (fig 1b) which is clearly more than in worker larvae (fig 1a). An increase was observed with peak values up to 170 pmol/ml only ≈ 84 h after cell capping. The subsequent decrease before the prepupal/pupal moult and later reincrease resemble the titer pattern as determined for workers. The few available data for *Apis mellifera ligustica* drones fall into the range described for *Apis cerana indica* males. Whether in drone L5 larvae of the western honey bee the JH peak is reached a little earlier has not yet been determined.

Fertility of *Varroa jacobsoni* in *Apis cerana indica* brood

Because the first *Varroa* egg is usually laid ≈ 3 d after brood cell capping, an evaluation of the reproductivity of female mites at pupal stages of the host allows a distinct discrimination to be made between fertile and infertile parasite individuals. This was analyzed in several hundred *Apis cerana indica* worker and drone brood cells ≈ 1 wk

Table 1. Brood infestation in *Apis cerana indica* and fertility of *Varroa jacobsoni* females (rate in bold type) on drone and worker hosts.

Apis cerana pupae	Capped brood cells opened (n)	Varroa infested brood cells found		Containing fertile Varroa females	
		(N)	(% n)	(x)	(% N)
Drones	373	76	20	69	90
Workers	710	34	4	0	0

after sealing. The results (table I) demonstrate a much lower level of brood infestation in worker than in drone cells. Furthermore, no fertile *Varroa* female was detected in the worker brood at all, whereas most of the mites found in drone brood were reproducing.

DISCUSSION

From the point of view of development, it is not surprising that the JH III hemolymph titer was found to be almost identical in preimaginal stages of *Apis cerana indica* and *A. mellifera ligustica*. JH III titer data for *Apis cerana* obtained by radioimmunoassay (RIA) are presented here for the first time. Similar results have recently been reported for *Apis mellifera carnica* and Africanized bees (Rachinsky *et al*, 1990; Rosenkranz *et al*, 1990). All these data are in good agreement.

Little is known about JH III in drone development (Hartfelder *et al*, 1993). However, the data available on rates of JH synthesis in late larval and early pupal drones as well as the JH III hemolymph titer values reported here are a little higher but do not differ much from the worker data. Therefore, it is improbable that a host-derived JH stimulus enhances *Varroa* reproduction in drone brood. Particularly the time of the JH peak during the spinning L5 drone instar 3–4 d after capping of the *cerana* brood cells already coincides with the oviposition of the first *Varroa* egg and, consequently, occurs far too late to stimulate oocyte development in the first gonocycle. Furthermore, there is only 1 peak within 8 d after capping in the JH III titer as well as in JH III content (Rembold, 1987) and not 2, with the second occurring after $\approx$ 6–7 d, as reported by Hänel and Koeniger (1986). An account of a RIA-determined course of the JH III hemo-

lymph titer in postcapping stages of drones has not been published previously, whereas such data are already known for workers and queens (Rachinsky *et al*, 1990). In both female castes of *Apis mellifera carnica* the JH III content has also been reported recently for the complete period of preadult development (Rembold *et al*, 1992) with similar worker values for all postcapping stages.

The regulation of metamorphosis and the involvement in sex- and caste-specific control of JH III production (Hartfelder *et al*, 1993) probably existed long before the splitting of the genus *Apis* occurred (Ruttner, 1988, 1992). The control of polymorphic patterns in the duration of the postcapping period in honey bees apparently allows some variation (Moritz, 1985), but this concerns the moment of cell operculum in particular and not the hormone titer peaks correlated with the initiation of metamorphosis and the switch from larval to prepupal–pupal developmental programs.

Consequently, species-specific differences in late 5th instar or prepupal JH III hemolymph titers are rather improbable as an adaptive character related to parasitization by *Varroa jacobsoni*. In fact, in newly capped worker larvae and prepupae of *Apis cerana indica* we found a JH III titer that was even a little higher than in immature *Apis mellifera ligustica* sampled at the same study site, and also in *Apis mellifera carnica* (Rachinsky *et al*, 1990) as well as in Africanized bees from Brazil (Rosenkranz *et al*, 1990). All these RIA-based data differ from the Galleria wax test values according to which the postcapping JH III hemolymph titer was described to be higher in *Apis mellifera* than in *Apis cerana* (Hänel and Koeniger, 1986). There is no explanation for this discrepancy.

Therefore, the oft-quoted hypothesis (Ramirez and Otis, 1986; Camazine, 1988;

Ruttner, 1992) of a regulation of parasite reproduction by host-derived JH in case of the *Varroa*-*Apis* relationship is not supported by the hormone titers as measured here with a sensitive and accurate RIA (Strambi *et al.*, 1981). Of course this does not concern the results obtained by JH application to *Varroa* females (Hänel, 1983; Hänel and Koeniger, 1988) which, however, Milani and Chiesa (1990) were unable to reproduce.

On the other hand, there is no doubt about the dependence of initiation of *Varroa* oogenesis on specific host conditions (Rosenkranz, 1990; Steiner, 1991; Rosenkranz and Stürmer, 1993; Steiner *et al.*, 1993). Hemolymph of L5 instars capped less than 24 h earlier was found to stimulate mite egg development. However, in host larvae at this stage the JH III hemolymph titer is very low (Rachinsky *et al.*, 1990; Rosenkranz *et al.*, 1990; this paper). In any case, all the available evidence points towards the significance of inhibiting the fertility of female mites reproducing in worker brood as a natural resistance factor to varroaosis. This phenomenon is well documented both in the original host species, *Apis cerana* (Koeniger *et al.*, 1981; Tewarson, 1987; Tewarson *et al.*, 1992), and has been confirmed by the present study (table I). In the Africanized *Apis mellifera scutellata* hybrid biotype in Brazil (Ritter and De Jong, 1984; Rosenkranz *et al.*, 1990) and hybrid colonies of *Apis mellifera monticola* and *Apis mellifera ligustica* (Thrybom and Fries, 1991) a similarly low rate of *Varroa* fertility was observed. However, the host factors responsible for influencing reproduction of the *Varroa* mite still remain to be detected.

ACKNOWLEDGMENT

This study was supported by a DAAD overseas research grant to P. Rosenkranz.

Résumé — Teneur en hormone juvénile et reproduction de *Varroa jacobsoni* dans le couvain operculé d'*Apis cerana indica* et d'*Apis mellifera ligustica*. À Allahabad (Inde du Nord), on a prélevé au printemps des échantillons d'hémolymphes dans du couvain operculé d'ouvrières et de mâles issu de 4 colonies d'*Apis cerana indica* et d'une colonie d'*Apis mellifera ligustica*. On a en outre ouvert plus de 1 000 cellules de couvain renfermant des nymphes d'*A. c. indica* pour vérifier la présence de *Varroa jacobsoni*, et on a noté si les femelles de *Varroa* s'étaient reproduites. Les échantillons d'hémolymphes ont été envoyés en Europe et leur teneur en hormone juvénile III (JH) a été déterminée par méthode radio-immunologique.

Jusqu'à la mue nymphale, les teneurs en JH du couvain operculé d'ouvrières diffèrent peu chez les 2 races *indica* et *ligustica*. Celle des larves de l'abeille indienne est en moyenne un peu plus élevée que celle de l'abeille européenne (fig 1a). Dans le couvain de mâles, des teneurs en JH un peu plus élevées dans l'ensemble que chez les ouvrières (fig 1b) ont été mesurées après operculation. Les valeurs maximales se rencontrent le 2^e j après l'operculation dans le couvain d'ouvrières et le 4^e j dans le couvain de mâles. Dans les 2 cas, la présence d'un pic était nette. Le moment où il se produit, ainsi que les différences spécifiques à la race et au sexe, ne rentrent pas en ligne de compte pour déclencher la reproduction de *Varroa*.

Chez *A. c. indica* le parasitisme du couvain par *Varroa* a été contrôlé 1 semaine environ après l'operculation. Les cellules renfermaient déjà des nymphes. Le couvain de mâles était nettement plus parasité que celui d'ouvrières, conformément aux résultats antérieurs. Puisque les femelles de *Varroa* pondent leur premier œuf environ 3 j après l'operculation, il était possible au moment du contrôle de différencier de façon sûre les acariens fertiles des non fer-

tiles. Dans les cellules de mâles, presque toutes les femelles d'acariens s'étaient reproduites. En revanche dans les cellules d'ouvrières, pas un seul *Varroa* n'avait pondu (tableau I).

Ces données, en particulier celles relatives à l'évolution de la teneur en JH du couvain operculé chez l'abeille indienne et l'abeille européenne, sont en contradiction avec les résultats précédents, sur lesquels se fonde l'hypothèse souvent citée de la régulation de la reproduction de *Varroa* par la JH III présente dans l'hémolymph d'abeilles (Hänel et Koeniger, 1986).

Apis cerana indica* / *Apis mellifera ligustica* / hormone juvénile / reproduction / *Varroa jacobsoni

Zusammenfassung — Juvenilhormon-Titer und Fortpflanzung von *Varroa jacobsoni* in verdeckelter Brut von *Apis cerana indica* im Vergleich zu *Apis mellifera ligustica*. In Allahabad (Nord-Indien) wurden im Frühjahr Hämolympheproben von verdeckelter Arbeiterinnen- und Drohnenbrut in 4 Völkern von *Apis cerana indica* und 1 Volk von *Apis mellifera ligustica* gesammelt. Außerdem wurden über 1000 verdeckelte Brutzellen mit Puppen von *Apis cerana indica* geöffnet und auf Befall mit *Varroa jacobsoni* geprüft. Dabei wurde notiert, ob die Milbenweibchen sich fortgepflanzt hatten. Die Hämolympheproben wurden anschließend in Europa mittels Radioimmunoassay (RIA) auf ihren Gehalt an Juvenilhormon III (JH) analysiert und daraus JH-Titer der Hämolymphe berechnet.

Die so ermittelten JH-Titer von verdeckelter Arbeiterinnenbrut unterscheiden sich bei *Apis cerana indica* und *Apis mellifera ligustica* bis zur Puppenhäutung kaum, die Titer-Werte der Larven indischer Honigbienen liegen durchschnittlich sogar etwas über denen der europäischen (Abb

1a). Bei Drohnenbrut wurden nach dem Verdeckeln insgesamt leicht höhere JH-Titer gemessen (Abb 1b) als bei Arbeiterinnen. Die Titer-Maxima traten bei Arbeiterinnenbrut am 2. und bei Drohnenbrut am 4. Tag nach dem Verdeckeln auf, in beiden Fällen war lediglich ein Peak ausgeprägt. Diese Zeitpunkte der JH-Peaks und ebenso die geringen Art- und Geschlechtsspezifischen Titerunterschiede kommen als Auslöser für die Milbenfortpflanzung nicht in Frage. Bei *Apis cerana indica* wurde der Milbenbefall von Brut etwa eine Woche nach dem Verdeckelungszeitpunkt kontrolliert. Die Zellen enthielten dann bereits Puppen. In Übereinstimmung mit früheren Befunden war die Drohnenbrut wesentlich stärker infiziert als die Arbeiterinnenbrut. Da *Varroa*-Weibchen das erste Ei ungefähr 3 Tage nach dem Verdeckeln ablegen, konnte zum gewählten Kontrollzeitpunkt sicher zwischen fruchtbaren und unfruchtbaren Milben unterschieden werden. In Drohnenzellen hatten sich fast alle Milbenweibchen fortgepflanzt. In Arbeiterinnenzellen dagegen hatte keine einzige *Varroa*-Milbe ein Ei gelegt (Tabelle I).

Die hier mitgeteilten Daten, insbesondere über Verlauf des JH-Titers in verdeckelter Brut sowohl bei der östlichen wie bei der westlichen Honigbiene, stehen im Widerspruch zu früheren Befunden, auf denen eine vielzitierte Hypothese über Steuerung der *Varroa*-Fortpflanzung durch JH III aus der Bienenhämolymphe basiert (Hänel und Koeniger, 1986).

Apis cerana indica* / *Apis mellifera ligustica* / Juvenilhormon / Fortpflanzung / *Varroa jacobsoni

REFERENCES

- Camazine S (1988) Factors affecting the severity of *Varroa jacobsoni* infestations on European and Africanized honey bees. In: *Afri-*

- canized Honey Bees and Bee Mites* (Needham GR, Page RE, Delfinado-Baker M, Bowman CE, eds) Ellis Horwood Ltd, Chichester, 444-451
- Connat JL, Ducommun J, Diehl PA (1983) Juvenile hormone-like substances can induce vitellogenesis in the tick *Ornithodoros moubata* (Acarina: Argasidae). *Int J Invertebr Reprod* 6, 285-294
- Downer RGH, Laufer H (eds) (1983) *Invertebrate Endocrinology*. AR Liss, New York
- Engels W, Gonçalves LS, Steiner J, Buriolla AM, Cavichio Issa MR (1986) *Varroa*-Befall von *Carnica*-Völkern in Tropenklimate. *Apidologie* 17, 203-216
- Engels W, Kaatz H, Zillikens A, Simoes ZLP, Trube A, Braun R, Dittrich F (1990) Honey bee reproduction: vitellogenin and caste-specific regulation of fertility. *Adv Invertebr Reprod* 5, 495-502
- Hänel H (1983) Effect of JH III on the reproduction of *Varroa jacobsoni*. *Apidologie* 14, 137-142
- Hänel H, Koeniger N (1986) Possible regulation of the reproduction of the honey bee mite *Varroa jacobsoni* (Mesostigmata: Acari) by a host's hormone: juvenile hormone III. *J Insect Physiol* 32, 791-798
- Hartfelder K (1990) Regulatory steps in caste development of eusocial bees. In: *Social Insects, An Evolutionary Approach to Castes and Reproduction* (Engels W, ed) Springer, Heidelberg, 245-264
- Hartfelder K, Tozetto S de O, Rachinsky A (1993) Sex-specific developmental profiles of juvenile hormone synthesis in honey bee larvae. *Roux's Arch Dev Biol* 202, 176-180
- Koeniger N, Koeniger G, Wijayagunasekara NHP (1981) Beobachtungen über die Anpassung von *Varroa jacobsoni* an ihren natürlichen Wirt *Apis cerana* in Sri Lanka. *Apidologie* 12, 37-40
- Milani N, Chiesa F (1990) Some stimuli inducing oviposition in *Varroa jacobsoni* Oud. In: *Proceedings of the International Symposium on Recent Research on Bee Pathology* (Ritter W, Van Laere O, Jacobs F, De Wael L, eds) Apimondia, Ghent, 27-33
- Moritz RFA (1985) Heritability of the postcapping stage in *Apis mellifera* and its relation to varroaosis resistance. *J Hered* 76, 267-270
- Oliver JH Jr, Pound JM, Severino G (1985) Evidence of a juvenile-hormone-like compound in the reproduction of *Dermanyssus gallinae* (Acari: Dermanyssidae). *J Med Entomol* 22, 281-286
- Pound JM, Oliver JH Jr (1979) Juvenile hormone: evidence of its role in the reproduction of ticks. *Science* 206, 355-357
- Ramirez BW, Otis GW (1986) Developmental phases in the life cycle of *Varroa jacobsoni*, an ectoparasitic mite on honeybees. *Bee World* 67, 92-97
- Rachinsky A, Strambi C, Strambi A, Hartfelder K (1990) Caste and metamorphosis: hemolymph titers of juvenile hormone and ecdysteroids in last instar honeybee larvae. *Gen Comp Endocrinol* 79, 31-38
- Rachinsky A, Hartfelder K (1990) Corpora allata activity, a prime regulating element for caste-specific juvenile hormone titre in honeybee larvae (*Apis mellifera carnica*). *J Insect Physiol* 36, 189-194
- Rembold H (1987) Caste differentiation of the honeybee – fourteen years of biochemical research at Martinsried. In: *Chemistry and Biology of Social Insects* (Eder J, Rembold H, eds) Peperny, Munich, 3-15
- Rembold H, Czoppelt C, Grüne M, Lackner B, Pfeffer J, Woker E (1992) Juvenile hormone titers during honey bee embryogenesis and metamorphosis. In: *Insect Juvenile Hormone Research* (Mauchamp B, Couillaud F, Baehr JC, eds) INRA, Paris, 37-43
- Ritter W, De Jong D (1984) Reproduction of *Varroa jacobsoni* Oud in Europe, the Middle East and tropical South America. *Z Angew Entomol* 98, 55-57
- Robinson GE (1992) Regulation of division of labor in insect societies. *Annu Rev Entomol* 37, 637-665
- Rosenkranz P (1990) Der Einfluß larvaler Bienenhämolymph auf die Fertilität von *Varroa jacobsoni* in europäischen und afrikanisierten Bienenenvölkern. *Apidologie* 21, 370-372
- Rosenkranz P, Rachinsky A, Strambi A, Strambi C, Ropstorff P (1990) Juvenile hormone titer in capped worker brood of *Apis mellifera* and reproduction in the bee mite *Varroa jacobsoni*. *Gen Comp Endocrinol* 78, 189-193
- Rosenkranz P, Stürmer M (1993) Ernährungsabhängige Fertilität der *Varroa*

- Weibchen in Arbeiterinnenbrut von *Apis mellifera carnica* und *Apis mellifera capensis*. In: *Proc 1st Polish-German Symp Progress in Varroa Research*. Stettin, Poland (in press)
- Ruttner F (1988) *Biogeography and Taxonomy of Honeybees*. Springer, Berlin
- Ruttner F (1992) *Naturgeschichte der Honigbienen*. Ehrenwirth, Munich
- Steiner J (1991) Oogenese und Embryogenese während des ersten Gonozyklus von *Varroa jacobsoni*. *Apidologie* 22, 460-462
- Steiner J, Dittmann F, Rosenkranz P, Engels W (1993) Interrelation between reproduction of the honey bee (*Apis mellifera carnica*) parasitic mite (*Varroa jacobsoni*) and preimaginal host development during the first gonocycle. *Invertebr Reprod Dev* 23, in press
- Strambi C, Strambi A, de Reggi ML, Hirn MH, Delaage MA (1981) Radioimmunoassay of insect juvenile hormones and of their diol derivatives. *Eur J Biochem* 118, 401-406
- Tewarson NC (1987) Use of host hemolymph proteins, seasonal reproduction and a hypothesis on nutritional imprinting in the honey bee mite, *Varroa jacobsoni*, on *Apis mellifera* and *Apis cerana*. In: *Chemistry and Biology of Social Insects* (Eder J, Rembold H, eds) J Peperny, Munich, 688-689
- Tewarson NC, Singh A, Engels W (1992) Reproduction of *Varroa jacobsoni* in colonies of *Apis cerana indica* under natural and experimental conditions. *Apidologie* 23, 161-171
- Thrybom B, Fries I (1991) Development of infestations by *Varroa jacobsoni* in hybrid colonies of *Apis mellifera monticola* and *Apis mellifera ligustica*. *J Apicult Res* 30, 151-155