



The nature of dietary fatty acids affects the glycemic and insulinemic responses to carbohydrates in healthy subjects

Jl Joannic, S Auboiron, J Raison, A Basdevant, F Bornet, B Guy-Grand

► To cite this version:

Jl Joannic, S Auboiron, J Raison, A Basdevant, F Bornet, et al.. The nature of dietary fatty acids affects the glycemic and insulinemic responses to carbohydrates in healthy subjects. *Reproduction Nutrition Development*, 1996, 36 (4), pp.418-418. hal-00899863

HAL Id: hal-00899863

<https://hal.science/hal-00899863>

Submitted on 11 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

which parallels an improvement in protein oxidative parameters and abnormal Cu Zn SOD activity. This study highlights the relationships between glucose homeostasis and oxidative stress and the possible utilization of antioxidant agents to increase insulin sensitivity.

The nature of dietary fatty acids affects the glycemic and insulinemic responses to carbohydrates in healthy subjects.

JL Joannic ^{1,2}, S Auboiron ^{1,2}, J Raison ¹, A Basdevant ¹, F Borner ², B Guy-Grand ¹ (¹ Internal Medicine and Nutrition, Hôtel-Dieu; ² Éridania Béghin-Say, 75008 Paris, France)

The effect of a mixed meal composed of different kinds of carbohydrates and fats on postprandial plasma concentrations of glucose, insulin, free fatty acids and triglycerides was investigated in eight young normolipidemic men (24 ± 1 year, body mass index (BMI) 21.5 ± 0.8 kg/m²). Three hours after a standardized breakfast (300 kcal, 18% fats, 70% carbohydrates, 12% proteins), the subjects ingested four test meals (1 200 kcal, 50% fats, 38% carbohydrates, 12% proteins) in 30 min on separate days in random order according to a Latin square design.

The meals contained two kinds of carbohydrates: instant mashed potatoes (high glycemic index 70–90%) or rice (low glycemic index 50–55%) and two mixtures of vegetable oils, with either a high monounsaturated/polyunsaturated fatty acids n-6 ratio (M), or a low one (P). Proteins, saturated and polyunsaturated fatty acids n-3 were comparable in all meals. The plasma parameters were measured every 30 min during 3 h after the beginning of the test meal.

During the postprandial kinetic, the glycemic response was significantly lower with rice-P than potato-M or rice-M ($P < 0.01$) after 30 min. The insulinemic response was

lower for rice-P than with potatoes-M ($P < 0.05$). At 90 min, the average insulin level was similar for rice-P and rice-M and significantly lower than potato-P or potato-M ($P < 0.005$). No significant differences were found between meals in free fatty acid or triglyceride plasma levels.

In conclusion: i) The insulin response was significantly different between rice and instant mashed potatoes only when carbohydrates are associated with a polyunsaturated rich meal. ii) The polyunsaturated rich meals decreased the insulin response to the two kinds of carbohydrates. The same tendency was observed for the glycemic response.

Thus, the postprandial plasma concentrations of glucose and insulin are influenced by the nature of dietary fatty acids present in the meal.

Effect of selenium supplementation on clinical manifestations and plasma biochemical parameters in streptozotocin-diabetic rats mildly balanced by insulin.

C Douillet ¹, M Accominoti ², M Bost ³, F Borson-Chazot ⁴, M Ciavatti ¹ (¹ Inserm 63, 69675 Bron; ² Hôpital Édouard-Herriot, 69008; ³ Trace Element Institute for UNESCO, 69008; ⁴ Hôpital de l'Antiquaille, 69005 Lyon, France)

Seventy-six Sprague-Dawley rats were used in this study. Twelve rats were used as control (group C). Diabetes was induced in 64 rats by iv injection of streptozotocin (30 mg/kg). All rats with glycemia levels > 2.5 g were considered diabetics. All rats received a purified diet (in calories: 30% lipids, 15% proteins, 55% glucides). Three groups of 16 diabetic rats each were supplemented with selenium (Se): a Se-rich yeast (group DSe), or selenomethionine (group DSm) or selenomethionine + vitamin E (group DSmE). The supplementation of Se in all groups corresponded to