



On the ground state energy of the Laplacian with a magnetic field created by a rectilinear current

Vincent Bruneau, Nicolas Popoff

► To cite this version:

Vincent Bruneau, Nicolas Popoff. On the ground state energy of the Laplacian with a magnetic field created by a rectilinear current. *Journal of Functional Analysis*, 2015, 268 (5), pp.1277-1307. 10.1016/j.jfa.2014.11.015 . hal-00911208v3

HAL Id: hal-00911208

<https://hal.science/hal-00911208v3>

Submitted on 19 Feb 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON THE GROUND STATE OF THE LAPLACIAN WITH A MAGNETIC FIELD CREATED BY A RECTILINEAR CURRENT

VINCENT BRUNEAU AND NICOLAS POPOFF

ABSTRACT. We consider the three-dimensional Laplacian with a magnetic field created by an infinite rectilinear current bearing a constant current. The spectrum of the associated hamiltonian is the positive half-axis as the range of an infinity of band functions all decreasing toward 0. We make a precise asymptotics of the band function near the ground energy and we exhibit a semi-classical behavior. We perturb the hamiltonian by an electric potential. Helped by the analysis of the band functions, we show that for slow decaying potential, an infinite number of negative eigenvalues are created whereas only finite number of eigenvalues appears for fast decaying potential. The power-like decaying potential determining the finiteness of the negative spectrum is different than for the free Laplacian.

1. INTRODUCTION

1.1. Motivation and problematic.

- *Physical context.* We consider in \mathbb{R}^3 the magnetic field created by an infinite rectilinear wire bearing a constant current. Let (x, y, z) be the cartesian coordinates of \mathbb{R}^3 and assume that the wire coincides with the z axis. Due to the Biot & Savard law, the generated magnetic field writes

$$\mathbf{B}(x, y, z) = \frac{1}{r^2}(-y, x, 0)$$

where $r := \sqrt{x^2 + y^2}$ is the radial distance corresponding to the distance to the wire. Let $\mathbf{A}(x, y, z) := (0, 0, \log r)$ be a magnetic potential satisfying $\text{curl } \mathbf{A} = \mathbf{B}$. We define the unperturbed magnetic hamiltonian

$$H_{\mathbf{A}} := (-i\nabla - \mathbf{A})^2 = D_x^2 + D_y^2 + (D_z - \log r)^2; \quad D_j := -i\partial_j$$

initially defined on $C_0^\infty(\mathbb{R}^3)$ and then self-adjoint in $L^2(\mathbb{R}^3)$. It is known (see [24], and [25] for a more general setting) that the spectrum of $H_{\mathbf{A}}$ has a band structure with band functions defined on \mathbb{R} and decreasing from $+\infty$ toward 0. Then the spectrum of $H_{\mathbf{A}}$ is absolutely continuous and coincide with $[0, +\infty)$. In that case the presence of the magnetic field does not change the spectrum (i.e. $\mathfrak{S}(H_{\mathbf{A}}) = \mathfrak{S}(-\Delta)$), that may be expected since the magnetic field tends to 0 far from the wire. In this article we study the ground state of $H_{\mathbf{A}}$ and its stability under electric perturbation. These questions are related to the dynamic of spinless quantum particles submitted to the magnetic field \mathbf{B} and perturbed by an electric potential.

- *Comparison with the free hamiltonian.* In general the spectrum of a Laplacian may be higher in the presence of a magnetic field (see [2]). As already said, in our model we still have $\mathfrak{S}(H_{\mathbf{A}}) = \mathbb{R}_+$. However the dynamics are very different from the free motion, see [24] for a description of the classical and quantum dynamics of this model. As we will see, the behavior of the negative spectrum under electrical perturbation is also different that what happens without magnetic field.

If V is a multiplication operator by a real electric potential V such that $V(H_{\mathbf{A}} + 1)^{-1}$ is compact then the operator $H_{A,V} := H_{\mathbf{A}} - V$ is self-adjoint, its essential spectrum coincides with the positive half-axis and discrete spectrum may appear under 0.

Let us recall that, due to the diamagnetic Inequality (see [2, Section 2]), the operator $V(H_{\mathbf{A}} + 1)^{-1}$ is compact as soon as $V(-\Delta + 1)^{-1}$ is compact. Moreover, if $\mathcal{N}_{A,V}(\lambda)$ denotes the number of eigenvalues of $H_{\mathbf{A}} - V$ below $-\lambda < 0$, we have ([2, Theorem 2.15]):

$$(1.1) \quad \mathcal{N}_{A,V}(0^+) \leq C \int_{\mathbb{R}^3} V_+(x, y, z)^{\frac{3}{2}} dx dy dz, \quad V_+ := \max(0, V).$$

In particular, $H_{\mathbf{A}} - V$ has a finite number of negative eigenvalues provided that $V_+ \in L^{\frac{3}{2}}(\mathbb{R}^3)$. But this condition, also valid for $-\Delta - V$, is not optimal in presence of magnetic fields as the results of this article will show.

We will prove that the discrete spectrum of our operator $H_{\mathbf{A}} - V$ below 0 is less dense than for $-\Delta - V$ (see Theorem 1.3 and Corollary 1.4), more precisely for some V the operator $-\Delta - V$ has infinitely many negative eigenvalues whereas $\mathcal{N}_{A,V}(0^+) < +\infty$. In some sense, that means that the absolutely continuous spectrum of $H_{\mathbf{A}}$ near 0 is less dense than the one of the free Laplacian $-\Delta$.

- *Magnetic hamiltonian and band functions.* Several models with constant magnetic field have been studied in the past years. We recall some of them below. In most cases the system has a translation-invariance direction and the magnetic Laplacian is fibered through partial Fourier transform, therefore its study reduces to the study of the band functions that are the spectrum of the fiber operators. The spectrum of the hamiltonian is the range of the band functions (see [9] for a general setting) and the ground state is given by the infimum of the first band function. The number of eigenvalues created under the essential spectrum by a suitable electric perturbation depends strongly on the shape of the band functions near the ground state as shown on the examples below:

For the case of a constant magnetic field in \mathbb{R}^n , the perturbation by electric potential is described for example in [23] or [18]. When $n = 2$, the band functions are constant and equal to the Landau levels. In [20] the authors deal with very fast decaying potential. In that case they prove that the perturbation by an electric potential even compactly supported generates sequences of eigenvalues which converge toward the Landau levels, that is very different from what happens without magnetic field where only a finite number of eigenvalues are created by compactly supported electric perturbation.

In general the band function associated with a Schrödinger operator are not constant. The case where the band functions reach their infimum is described in [19] where the author study the perturbation of a Schrödinger operator with periodic electric potential and no magnetic

field, whose band functions have non-degenerated minima, providing localization in the phase space. Let us come back to the case with constant magnetic field. When adding a boundary, the band functions may not be constant anymore. For example when the domain is a two-dimensional infinite strip of finite width with constant magnetic field, it is proved that all the band functions are even with a non-degenerate minimum, see [8]. In [4], the authors investigate the behavior of the spectral shift function near the minima of the band functions, providing the number of eigenvalue created under the ground state when perturbing by an electric potential. Other examples of such a situation is the case of a half-plane with constant magnetic field and Neumann boundary condition, see [6, Section 4], the case of an Iwatsuka model with an odd discontinuous magnetic field, [15, Section 5] and also the case of the Dirichlet Laplacian on a twisted wave guide, [3].

The case of a half-plane with a constant magnetic field and Dirichlet boundary condition is more intriguing and somehow closer to our model: in that case the bottom of the spectrum of the magnetic Laplacian is the first Landau level, but the associated band function does not reach its infimum. In [6], the authors gives the precise behavior of the counting function when perturbing by a suitable electric potential. Analog situations based on Iwatsuka models are described in [5] or [14].

All the above described situations deal with constant magnetic field. In this article we deal with a three dimensional variable magnetic field going to 0 far from the z -axis and invariant along this axis, therefore the situation is quite different—one may think roughly that the variations of the magnetic field will create non-constant band function as the addition of a boundary does in the case of a constant magnetic field. Moreover in the above described models the band functions are well separated near the ground state in the sense that the infimum of the second band function is larger than the ground state. In our case there are infinitely many band functions that accumulate toward $\inf \mathfrak{S}(H_A)$, see Figure 1, adding a technical challenge when studying the ground state.

In this paper, we give more precise description of the spectrum of H_A near 0 with asymptotic expansion of the band functions. Then, we study the finiteness of the number of the negative eigenvalues of $H_A - V$ for relatively compact perturbations V . On one hand, we display classes of potentials giving rise to an accumulation at 0, of an infinite number of negative eigenvalues, on the other hand, under a decreasing property of V_+ , we prove the finiteness of the discrete spectrum of $H_A - V$ below 0. We obtain a class of polynomially decreasing potentials for which $H_A - V$ has a finite number of negative eigenvalues while the negative spectrum of $-\Delta - V$ is infinite.

1.2. Main results. Using the cylindrical coordinates of \mathbb{R}^3 , we identify $L^2(\mathbb{R}^3)$ with the weighted space $L^2(\mathbb{R}_+ \times (0, 2\pi) \times \mathbb{R}, r dr d\phi dz)$ and the operator H_A writes:

$$H_A = -\frac{1}{r} \partial_r r \partial_r - \frac{\partial_\phi^2}{r^2} + (\log r - D_z)^2$$

acting on functions of $L^2(\mathbb{R}_+ \times (0, 2\pi) \times \mathbb{R}, r dr d\phi dz)$.

Let us recall the fibers decomposition of H_A that can be found with more details in [24]. We denote by \mathcal{F}_3 the Fourier transform with respect to z and Φ the angular Fourier transform.

We have the direct integral decomposition (see [21, Section XIII.16] for the notations about direct decomposition):

$$\Phi \mathcal{F}_3 H_{\mathbf{A}} \mathcal{F}_3^* \Phi^* := \sum_{m \in \mathbb{Z}}^{\oplus} \int_{k \in \mathbb{R}}^{\oplus} g_m(k) dk$$

where the operator

$$(1.2) \quad g_m(k) := -\frac{1}{r} \partial_r r \partial_r + \frac{m^2}{r^2} + (\log r - k)^2$$

is defined as the extension of the quadratic form

$$q_m^k(u) := \int_{\mathbb{R}_+} \left(|u'(r)|^2 + \frac{m^2}{r^2} |u(r)|^2 + (\log r - k)^2 |u(r)|^2 \right) r dr$$

initially defined on $\mathcal{C}_0^\infty(\mathbb{R}_+)$ and closed in $L_r^2(\mathbb{R}_+) := L^2(\mathbb{R}_+, r dr)$.

For all $(m, k) \in \mathbb{Z} \times \mathbb{R}$ the operator $g_m(k)$ has compact resolvent. We denote by $\lambda_{m,n}(k)$ the so-called band functions, i.e. the n -th eigenvalue of $g_m(k)$ associated with a normalized eigenvector $u_{m,n}(k)$.

It is known ([24], see also Section 2.1) that $k \mapsto \lambda_{m,n}(k)$ is decreasing with

$$\lim_{k \rightarrow -\infty} \lambda_{m,n}(k) = +\infty; \quad \lim_{k \rightarrow +\infty} \lambda_{m,n}(k) = 0.$$

Exploiting semi-classical tools (with semi-classical parameter $h = e^{-k}$, $k \gg 1$, see Proposition 2.2), we obtain asymptotic behaviors of the eigenpairs of $g_m(k)$ as k tends to infinity. The main result of Section 2 is the following

Theorem 1.1. *For all $(m, n) \in \mathbb{Z} \times \mathbb{N}^*$, there exist constants $C_{m,n} > 0$ and $k_0 \in \mathbb{R}$ such that for all $k \in (k_0, +\infty)$,*

$$(1.3) \quad |\lambda_{m,n}(k) - (2n-1)e^{-k} + (m^2 - \frac{1}{4} - \frac{n(n-1)}{2})e^{-2k}| \leq C_{m,n} e^{-5k/2}$$

This asymptotics shows that all the band functions tend exponentially to the ground state and cluster according to their energy level, see Figures 1 and 2.

Let us consider V , a multiplication operator such that $V(H_{\mathbf{A}} + 1)^{-1}$ is compact. Considered in $L^2(\mathbb{R}_+ \times (0, 2\pi) \times \mathbb{R}, r dr d\phi dz)$, V is a function of (r, φ, z) and it is said *axisymmetric* when it does not depend of φ .

We want to know how reacts the ground state of $H_{\mathbf{A}}$ under electrical perturbation. For potentials slowly decreasing with respect to r , we have an infinite number of negative eigenvalues of $H_{\mathbf{A}} - V$:

Theorem 1.2. *Suppose V is a potential such that $V(H_{\mathbf{A}} + 1)^{-1}$ is compact and*

$$(1.4) \quad V(x, y, z) \geq \langle (x, y) \rangle^{-\alpha} v_{\perp}(z), \quad \alpha > 0.$$

If α and v_{\perp} satisfy one of the assumptions (i), (ii) below, then, $H_{\mathbf{A}} - V$ have a infinite number of negative eigenvalues which accumulate to 0.

(i) $\alpha < \frac{1}{2}$ and $v_{\perp} \in L^1(\mathbb{R})$ such that

$$\int_{\mathbb{R}} v_{\perp}(z) dz > 0.$$

(ii) $v_\perp \geq C\langle z \rangle^{-\gamma}$ with $\gamma > 0$ and $\alpha + \frac{\gamma}{2} < 1$.

The proof uses a construction of quasi-modes based on the eigenfunctions associated with $\lambda_{m,n}(k)$ that leads to a one-dimensional operator in the z variable. The key point is a projection (in the r variable) of the potential V against the eigenfunctions of $g_m(k)$ that are localized near the wells of the potential $(\log r - k)^2$ for large k .

We also have conditions giving finiteness of the negative spectrum.

Theorem 1.3. *Assume V is a relatively compact perturbation of H_A such that*

$$(1.5) \quad V(x, y, z) \leq \langle (x, y) \rangle^{-\alpha} v_\perp(z),$$

with $\alpha > 1$ and $v_\perp \in L^p(\mathbb{R})$ a non negative function with $p \in [1, 2]$.

Then, $H_A - V$ have, at most, a finite number of negative eigenvalues.

Let us give some comments concerning the above results in comparison with known borderline behavior of perturbations of the Laplacian. It is not true in general that the number of negative eigenvalues of $-\Delta - V$ is larger than when adding a magnetic field, see Exemple 2 after Theorem 2.15 of [2]. Theorem 1.2 is a case where the number of negative eigenvalues in presence of magnetic field is infinite as without magnetic field.

However due to the diamagnetic inequality, one might expect for most cases that the density of negative eigenvalues is more important for $-\Delta - V$ than for $H_A - V$. The above results illustrate this phenomenon, indeed we prove that the borderline behavior of the perturbation determining the finiteness of the negative spectrum of $H_A - V$ is different than for $-\Delta - V$:

Corollary 1.4. *Let V be a measurable function on \mathbb{R}^3 that obeys*

$$c\langle (x, y) \rangle^{-\alpha} \langle z \rangle^{-\gamma} \leq V(x, y, z) \leq C\langle (x, y) \rangle^{-\alpha} \langle z \rangle^{-\gamma},$$

with $\alpha + \gamma < 2$, $\alpha > 1$ and $\gamma > \frac{1}{2}$.

Then the operator $-\Delta - V$ have infinitely many negative eigenvalues while the negative spectrum of $H_A - V$ is finite.

Proof. Since $\langle (x, y) \rangle^{-\alpha} \langle z \rangle^{-\gamma} \geq \langle (x, y, z) \rangle^{-(\alpha+\gamma)}$, according to [21, Theorem XIII.6] we know that for $V(x, y, z) \geq \langle (x, y) \rangle^{-\alpha} \langle z \rangle^{-\gamma}$ with $\alpha + \gamma < 2$, the operator $-\Delta - V$ has infinitely many negative eigenvalues. The corollary is then deduced from Theorem 1.3. □

A natural open question concern the existence of a borderline behavior of V which determine the finiteness of the negative spectrum of $H_A - V$. At the moment we can only say that, if it exists, such borderline potential V_b satisfies:

$$C_- \langle (x, y) \rangle^{-\alpha_-} \langle z \rangle^{-\gamma_-} \leq V_b \leq C_+ \langle (x, y) \rangle^{-\alpha_+} \langle z \rangle^{-\gamma_+},$$

with $0 < \alpha_- \leq \max(1 - \frac{\gamma_-}{2}; \frac{1}{2})$, $\gamma_- > 0$ and $\alpha_+ > 1$, $\gamma_+ > \frac{1}{2}$.

1.3. Organisation of the article. In Section 2 we recall basis on the fibers of the operator H_A and their associated band functions $\lambda_{m,n}(k)$. We give the localization of the associated eigenfunctions for large k and we prove Theorem 1.1. We also provide numerical computations of the band functions. In Section 3, we construct quasi-modes for the perturbed operator $H_A - V$ that leads to study a one-dimensional problem and allows to prove Theorem 1.2. Based on an uniform lower bound of the band functions, Section 4 combines the Birman-Schwinger principle with results of Section 2 to prove Theorem 1.3. The key point is an estimation of the Hilbert-Schmidt norm of Birman-Schwinger type operator associated with the perturbed hamiltonian.

2. DESCRIPTION OF THE 1D PROBLEM ASSOCIATED WITH THE UNPERTURBED HAMILTONIAN

In this section we first recall results from [24] on the behavior of the band functions $k \mapsto \lambda_{m,n}(k)$. Then we give Agmon estimates on the associated eigenfunctions and we perform an asymptotic expansion of $\lambda_{m,n}(k)$ when k goes to $+\infty$. In Section 3 and 4 we will use these expansions to analyse the operator $H_A - V$.

Depending on the context we shall work with different operators all unitarily equivalent to the operator $g_m(k)$ written in (1.2). Table 1 gives a description of these operators and the notations we use.

2.1. Semi-classical point of view.

- *Global behavior of the band functions.* As in [24], we introduce the parameter

$$h := e^{-k}$$

such that $\log r - k = \log(hr)$. The scaling $\rho = hr$ shows that $g_m(k)$ is unitarily equivalent to

$$(2.1) \quad \mathfrak{g}_m(h) := -h^2 \frac{1}{\rho} \partial_\rho \rho \partial_\rho + h^2 \frac{m^2}{\rho^2} + (\log \rho)^2$$

acting on $L^2_\rho(\mathbb{R}_+) := L^2(\mathbb{R}_+, \rho d\rho)$. We denote by $(\mu_{m,n}(h), \mathbf{u}_{m,n}(\cdot, h))_{n \geq 1}$ the normalized eigenpairs of this operator and by \mathfrak{q}_h^m the associated quadratic form. We have $\mu_{m,n}(h) = \lambda_{m,n}(k)$ and

$$\mathbf{u}_{m,n}(\rho, h) = h u_{m,n} \left(\frac{\rho}{h}, -\log h \right)$$

where $u_{m,n}(\cdot, k)$ is a normalized eigenfunction associated with $\lambda_{m,n}(k)$ for $g_m(k)$. Using the min-max principle and the expression (2.1), it is clear that $h \mapsto \mu_{m,n}(h)$ is non decreasing on $(0, +\infty)$ and therefore $k \mapsto \lambda_{m,n}(k)$ is non increasing on \mathbb{R} . It was already used by Yafaev (see [24]) who, moreover, shows (see [24, Lemma 2.2 & 2.3]) that

$$\lim_{h \rightarrow 0} \mu_{m,n}(h) = 0 \quad \text{and} \quad \lim_{h \rightarrow +\infty} \mu_{m,n}(h) = +\infty.$$

Note that these results are extended to more general magnetic fields in [25, Section 3].

- *The fiber operator in an unweighted space.* Sometimes it will be convenient to work in an unweighted Hilbert space on the half-line, therefore we introduce the isometric transformation

$$\begin{aligned}\mathcal{M} &: L^2(\mathbb{R}_+, r dr) \mapsto L^2(\mathbb{R}_+, dr) \\ u(r) &\mapsto \sqrt{r} u(r)\end{aligned}$$

and we define $\tilde{g}_m(k) := \mathcal{M}g_m(k)\mathcal{M}^*$. This operator expressed as

$$(2.2) \quad \tilde{g}_m(k) := -\partial_r^2 + \frac{m^2 - \frac{1}{4}}{r^2} + (\log r - k)^2,$$

acting on $L^2(\mathbb{R}_+)$ and its precise definition can be derived from the natural associated quadratic form initially defined on $C_0^\infty(\mathbb{R}_+)$ and then closed to $L^2(\mathbb{R}_+)$.

2.2. Agmon estimates about the eigenpairs of the fiber operator. We write

$$\mathbf{g}_m(h) = -h^2 \frac{1}{\rho} \partial_\rho \rho \partial_\rho + V_h^m$$

with

$$V_h^m(\rho) := \log(\rho)^2 + h^2 \frac{m^2}{\rho^2}.$$

Let \mathbf{q}_m^h denote the natural associated quadratic form. Assume that μ is an eigenvalue satisfying $\mu \leq E + O(h)$ with $E \geq 0$, the eikonale equation on the Agmon weight ϕ writes

$$h^2 |\phi'|^2 = V_h^m - E$$

that is

$$|\phi'(\rho)|^2 = \frac{(\log \rho)^2 - E}{h^2} + \frac{m^2}{\rho^2}.$$

A solution is given by $\phi_h(\rho)/h$ with

$$(2.3) \quad \phi_h(\rho) := \left| \int_1^\rho \sqrt{\left((\log \rho')^2 - E + h^2 \frac{m^2}{\rho'^2} \right)_+} d\rho' \right|$$

This function provides the general Agmon estimates:

Proposition 2.1. *Let $E \geq 0$ and $C_0 > 0$. For all $\beta \in (0, 1)$ there exist $C(E, \beta) > 0$ and $h_0 > 0$ such that for all eigenpairs (μ, \mathbf{u}_μ) of $\mathbf{g}_m(h)$ with $\mu \leq E + C_0 h$ and \mathbf{u}_μ that is L_ρ^2 -normalized there holds:*

$$(2.4) \quad \forall h \in (0, h_0), \quad \|e^{\beta \frac{\phi_h}{h}} \mathbf{u}_\mu\|_{L_\rho^2(\mathbb{R}_+)} \leq C(E, \beta) \quad \text{and} \quad \mathbf{q}_m^h(e^{\beta \frac{\phi_h}{h}} \mathbf{u}_\mu) \leq C(E, \beta).$$

Proof. This proposition is an application of the well-known Agmon estimates for 1D Schrödinger operators with confining potential. First we have the following identity for any Lipschitz bounded function ϕ , see for example [22], [1] or [12]:

$$(2.5) \quad \langle \mathbf{g}_m(h)u, e^{2\phi}u \rangle_{L_\rho^2(\mathbb{R}_+)} = \mathbf{q}_m^h(e^\phi u) - h^2 \|\phi' e^\phi u\|_{L_\rho^2(\mathbb{R}_+)}^2.$$

In particular when $u = \mathbf{u}_\mu$ is an eigenfunction associated with the eigenvalue μ we get

$$(2.6) \quad \int_{\mathbb{R}_+} (h^2 |\partial_\rho(e^\phi \mathbf{u}_h)|^2 + (V_h^m - h^2 |\phi'|^2 - \mu) |e^\phi \mathbf{u}_h|^2) \rho d\rho = 0.$$

We now use this identity with $\phi = \phi_h/h$ where ϕ_h is defined in (2.3). The remain of the proof is classical and can be found with details in [11, Proposition 3.3.1] for example. \square

Note that

$$\phi_h(\rho) \geq \phi_0(\rho) = \left| \int_1^\rho \sqrt{((\log \rho')^2 - E)_+} d\rho' \right|$$

that does not depend neither on m nor on h . Therefore (2.4) remains true replacing ϕ_h by ϕ_0 and we get L^2 estimates uniformly in m , in particular:

$$(2.7) \quad \forall \beta \in (0, 1), \forall h \in (0, h_0), \quad \|e^{\beta \frac{\phi_0}{h}} \mathbf{u}_{m,n}(\cdot, h)\|_{L^2_\beta(\mathbb{R}_+)} \leq C(E, \beta)$$

for all normalized eigenfunction $\mathbf{u}_{m,n}(\cdot, h)$ of $\mathbf{g}_m(h)$ associated with any eigenvalue $\mu_{m,n}(h)$ satisfying $\mu_{m,n}(h) \leq E + C_0 h$ where $C_0 > 0$ is a set constant.

When $E = 0$ (that means that we are looking at the low-lying energies) the Agmon distance ϕ_0 is explicit:

$$\begin{aligned} \phi_0(\rho) &= \left| \int_1^\rho |\log \rho'| d\rho' \right| \\ &= |[\rho' \log \rho' - \rho']_1^\rho| = |\rho \log \rho - \rho + 1|. \end{aligned}$$

Let us express this in the original cylindrical variable $r = \frac{\rho}{h}$ with the Fourier parameter $k = -\log h$. The associated Agmon distance writes

$$(2.8) \quad \Phi_0(r, k) := \frac{\phi_0(\rho)}{h} = e^k \phi_0(re^{-k}) = r(\log(r) - k) - r + e^k.$$

Writing the previous estimates in these variables we get that for k large enough:

$$(2.9) \quad \|e^{\beta \Phi_0(\cdot, k)} u_{m,n}(\cdot, k)\|_{L^2_\beta(\mathbb{R}_+)} \leq C(0, \beta) \quad \text{and} \quad \|e^{\beta \Phi_0(\cdot, k)} \tilde{u}_{m,n}(\cdot, k)\|_{L^2(\mathbb{R}_+)} \leq C(0, \beta)$$

where $\tilde{u}_{m,n}(r) := \sqrt{r} u_{m,n}(\cdot, k)$ is a normalized eigenvector associated with $\lambda_{m,n}(k)$ for the operator $\tilde{g}_m(k)$ in the unweighted space $L^2(\mathbb{R}_+)$.

The function $r \mapsto \Phi_0(r, k)$ is positive, decreasing on $(0, e^k)$ and increasing on $(e^k, +\infty)$. It vanishes when $r = e^k$, so we find that the eigenfunction of the operator $g_m(k)$ are localized at the minimum of the wells $r = e^k$.

2.3. Asymptotics for the small energy. In this section we provide an asymptotic expansion of $\mu_{m,n}(h)$ for fixed (m, n) when h goes to 0, namely:

Proposition 2.2. *For all $(m, n) \in \mathbb{Z} \times \mathbb{N}^*$ there exists $C_{m,n} > 0$ and $h_0 > 0$ such that*

$$\forall h \in (0, h_0), \quad |\mu_{m,n}(h) - (2n - 1)h - (m^2 - \frac{1}{4} - \frac{n(n-1)}{2})h^2| \leq C_{m,n} h^{5/2}.$$

The operator $\mathbf{g}_m(h)$ written in (2.1) is a semiclassical Schrödinger operator with a potential which has a unique minimum at $\rho = 1$. We will use the technics of the harmonic approximation as described in [7], [22] or [11] to derive the asymptotics of the eigenvalues. The remain of this section is devoted to the proof of Proposition 2.2 which implies Theorem 1.1 because $\lambda_{m,n}(k) = \mu_{m,n}(e^{-k})$.

• *Canonical transformations.* As above we introduce the operator $\tilde{\mathbf{g}}_m(h) := \mathcal{M}\mathbf{g}_m(h)\mathcal{M}^*$ in the unweighted space where $\mathcal{M} : \mathbf{u}(\rho) \mapsto \sqrt{\rho}\mathbf{u}(\rho)$. We get

$$\tilde{\mathbf{g}}_m(h) = -h^2\partial_\rho^2 + h^2\frac{m^2 - \frac{1}{4}}{\rho^2} + \log^2 \rho$$

acting on the unweighted space $L^2(\mathbb{R}_+)$. Apply now the change of variable $t = \frac{\rho-1}{\sqrt{h}}$. We get that $\tilde{\mathbf{g}}_m(h)$ is unitarily equivalent to $h\hat{\mathbf{g}}_m(h)$ where

$$\hat{\mathbf{g}}_m(h) := -\partial_t^2 + \frac{\log^2(1 + \sqrt{ht})}{h} + h\frac{m^2 - \frac{1}{4}}{(1 + \sqrt{ht})^2}$$

acting on $L^2(I_h)$ with $I_h = (-h^{-1/2}, +\infty)$. As we will see below, this operator has a suitable shape to make an asymptotic expansion of its eigenvalues when $h \rightarrow 0$.

• *Asymptotic expansion and formal construction of quasi-modes.* We write a Taylor expansion of the potential near $t = 0$:

$$(2.10) \quad \frac{\log^2(1 + \sqrt{ht})}{h} + h\frac{m^2 - \frac{1}{4}}{1 + \sqrt{ht}} = t^2 - h^{1/2}t^3 + \left(\frac{11}{12}t^4 + m^2 - \frac{1}{4}\right)h + R(t, h)$$

where $R(t, h)$ will later be controlled by $(1 + |t|)^5 h^{3/2}$.

We write

$$\hat{\mathbf{g}}_m(h) = L_0 + h^{1/2}L_1 + hL_2 + R(\cdot, h)$$

where

$$\begin{cases} L_0 := -\partial_t^2 + t^2, \\ L_1 := -t^3, \\ L_2 := \left(\frac{11}{12}t^4 + m^2 - \frac{1}{4}\right). \end{cases}$$

At first we consider these operator as acting on $L^2(\mathbb{R})$ and we look at a quasi-mode for $L_0 + h^{1/2}L_1 + hL_2$ defined on \mathbb{R} . Using a suitable cut-off function this procedure will provide a quasi-mode for $\hat{\mathbf{g}}_m(h)$.

We look for a quasi-mode of the form

$$(E(h), f(\cdot, h)) = (E_0 + h^{1/2}E_1 + hE_2, f_0 + h^{1/2}f_1 + hf_2).$$

We are led to solve the following system:

$$\begin{aligned} (2.11a) \quad & \begin{cases} L_0 f_0 = E_0 f_0, \\ L_1 f_0 + L_0 f_1 = E_0 f_1 + E_1 f_0, \\ L_2 f_0 + L_1 f_1 + L_0 f_2 = E_2 f_0 + E_1 f_1 + E_0 f_2. \end{cases} \\ (2.11b) \quad & \\ (2.11c) \quad & \end{aligned}$$

Since L_0 is the quantum harmonic oscillator, to solve (2.11a) we choose for E_0 the n -th Landau level:

$$(2.12) \quad E_0 := 2n - 1, \quad n \geq 1$$

and

$$f_0 = f_{0,n} := \Psi_n, \quad n \geq 1$$

where Ψ_n is the n -th normalized Hermite's function with the convention that $\Psi_1(t) = (2\pi)^{-1/4} e^{-t^2/2}$.

We take the scalar product of (2.11b) against $f_{0,n}$ and we find

$$E_1 = \langle (L_0 - E_0)f_1, f_{0,n} \rangle + \langle L_1 f_{0,n}, f_{0,n} \rangle = \langle L_1 f_{0,n}, f_{0,n} \rangle.$$

Notice that $f_{0,n}$ is either even or odd and that $L_1 f_{0,n}$ has the opposite parity. Therefore the function $L_1 f_{0,n} \cdot f_{0,n}$ is odd for all $n \geq 1$ and we get

$$(2.13) \quad E_1 = 0.$$

We find f_1 by solving (2.11b):

$$(2.14) \quad (L_0 - E_0)f_1 = -L_1 f_{0,n} = t^3 \Psi_n(t).$$

Using $t\Psi_n(t) = \sqrt{\frac{n-1}{2}}\Psi_{n-1}(t) + \sqrt{\frac{n}{2}}\Psi_{n+1}(t)$, we write $t^3\Psi_n(t)$ on the basis of the Hermite's functions:

$$t^3\Psi_n(t) = a_n\Psi_{n-3}(t) + b_n\Psi_{n-1}(t) + c_n\Psi_{n+1}(t) + d_n\Psi_{n+3}(t)$$

with

$$(2.15) \quad \forall n \geq 1, \quad \begin{cases} a_n = 2^{-3/2} \sqrt{(n-1)(n-2)(n-3)} \\ b_n = 2^{-3/2} 3(n-1)\sqrt{n-1} \\ c_n = 2^{-3/2} 3n\sqrt{n} \\ d_n = 2^{-3/2} \sqrt{n(n+1)(n+2)}. \end{cases}$$

Therefore the unique solution to (2.14) orthogonal to $f_{0,n}$ is:

$$f_1 = f_{1,n} := \left(-\frac{a_n}{6}\Psi_{n-3} - \frac{b_n}{2}\Psi_{n-1} + \frac{c_n}{2}\Psi_{n+1} + \frac{d_n}{6}\Psi_{n+3} \right)$$

with $a_n = 0$ when $n \leq 3$ and $b_n = 0$ when $n = 1$ (see (2.15)).

We now take the scalar product of (2.11c) against $f_{0,n}$:

$$(2.16) \quad E_2 = \langle L_2 f_{0,n}, f_{0,n} \rangle + \langle L_1 f_{1,n}, f_{0,n} \rangle.$$

Computations provides

$$\langle L_2 f_{0,n}, f_{0,n} \rangle = \left(\frac{11}{12} \|t^2 f_{0,n}\|^2 + m^2 - \frac{1}{4} \right) = \left(\frac{11}{16} (2n^2 - 2n + 1) + m^2 - \frac{1}{4} \right)$$

and

$$\langle L_1 f_{1,n}, f_{0,n} \rangle = \left(\frac{a_n^2}{6} + \frac{b_n^2}{2} - \frac{c_n^2}{2} - \frac{d_n^2}{6} \right) = \frac{1}{16} (-30n^2 + 30n - 11),$$

therefore we get

$$(2.17) \quad E_2 = \left(-\frac{n(n-1)}{2} + m^2 - \frac{1}{4} \right).$$

We deduce from (2.11c):

$$(L_0 - E_0)f_2 = E_2f_{0,n} - L_1f_{1,n} - L_2f_{0,n}.$$

Since the compatibility condition is satisfied by the choice of E_2 (see (2.16)), the Fredholm alternative provides a unique solution $f_2 = f_{2,n}$ orthogonal to $f_{0,n}$. As above it may be computed explicitly using the Hermite's functions. Notice that $f_{2,n}$ depends on m as E_2 , see (2.17).

We finally define

$$f_{m,n}(t, h) := f_{0,n}(t) + h^{1/2}f_{1,n}(t) + hf_{2,n}(t)$$

• *Evaluation of the quasi-mode and upper bound.* The above formal construction provides functions on \mathbb{R} and we will now use a cut-off function in order to get quasi-modes for $\hat{\mathbf{g}}_m(h)$. Let $\chi \in C_0^\infty(\mathbb{R}, [0, 1])$ be a cut-off function increasing such that $\chi(t) = 0$ when $t \leq -1/2$ and $\chi(t) = 1$ when $t \geq -1/4$. We define $\chi(t, h) := \chi(h^{1/2}t)$ and

$$\hat{\mathbf{v}}_{m,n}(t, h) := \chi(t, h)f_{m,n}(t, h).$$

Recall that $\hat{\mathbf{g}}_{m,n}(h)$ acts on $L^2(I_h)$ with $I_h = (-h^{-1/2}, +\infty)$. Since $\text{supp}(\hat{\mathbf{v}}_{m,n}(\cdot, h)) \subset (-\frac{1}{2}h^{-1/2}, +\infty)$ and $\hat{\mathbf{v}}_{m,n}(\cdot, h)$ has exponential decay at $+\infty$, we have $\hat{\mathbf{v}}_{m,n} \in \text{dom}(\hat{\mathbf{g}}_m(h))$. Let

$$E_{m,n}(h) := E_0 + h^{1/2}E_1 + hE_2$$

where E_0 , E_1 and E_2 are defined in (2.12), (2.13) and (2.17).

We now evaluate $\|(\hat{\mathbf{g}}_m(h) - E_{m,n}(h))\hat{\mathbf{v}}_{m,n}(\cdot, h)\|_{L^2(I_h)}$. The procedure is rather elementary but for the sake of completeness we provide details below. We have

$$(2.18) \quad \|(\hat{\mathbf{g}}_m(h) - E_{m,n}(h))\hat{\mathbf{v}}_{m,n}(\cdot, h)\|_{L^2(I_h)} \leq \|[\hat{\mathbf{g}}_m(h), \chi(\cdot, h)]f_{m,n}(\cdot, h)\|_{L^2(I_h)} \\ + \|\chi(\cdot, h)R(\cdot, h)f_{m,n}(\cdot, h)\|_{L^2(I_h)} + \|\chi(\cdot, h)(L_0 + h^{1/2}L_1 + hL_2 - E_{m,n}(h))f_{m,n}(\cdot, h)\|_{L^2(I_h)}$$

We have $[\hat{\mathbf{g}}_m(h), \chi(\cdot, h)]f_{m,n}(t, h) = -2h^{1/2}\chi'(h^{1/2}t)f'_{m,n}(t, h) - h\chi''(h^{1/2}t)f_{m,n}(t, h)$ therefore $t \mapsto [\hat{\mathbf{g}}_m(h), \chi(\cdot, h)]f_{m,n}(t, h)$ is supported in $[-\frac{1}{2}h^{-1/2}, -\frac{1}{4}h^{-1/2}]$ and since $f_{m,n}(\cdot, h)$ and $f'_{m,n}(\cdot, h)$ have exponential decay we get

$$(2.19) \quad \|[\hat{\mathbf{g}}_m(h), \chi(\cdot, h)]f_{m,n}(\cdot, h)\|_{L^2(I_h)} = \mathcal{O}(h^\infty).$$

Remind that $R(t, h)$ is defined in (2.10), we get

$$\exists C > 0, \forall h > 0, \forall t \in \text{supp}(\chi(\cdot, h)), \quad |R(t, h)| \leq Ch^{3/2}(1 + |t|)^5.$$

Using the exponential decay of $f_{m,n}$ we get $C_{m,n} > 0$ such that

$$(2.20) \quad \|\chi(\cdot, h)R(\cdot, h)f_{m,n}(\cdot, h)\|_{L^2(I_h)} \leq C_{m,n}h^{3/2}.$$

The last term of (2.18) is easily computed:

$$(L_0 + h^{1/2}L_1 + hL_2 - E_{m,n}(h))f_{m,n}(\cdot, h) = h^{3/2}((L_1 - E_1)f_{2,n} + (L_2 - E_2)f_{1,n}) + h^2L_2f_{2,n}$$

and we get $C_{m,n} > 0$ such that

$$\|\chi(\cdot, h)(L_0 + h^{1/2}L_1 + hL_2 - E_{m,n}(h))f_{m,n}(\cdot, h)\|_{L^2(I_h)} \leq C_{m,n}h^{3/2}.$$

Combining this with (2.19) and (2.20) in (2.18) we get

$$(2.21) \quad \exists C_{m,n}, \exists h_0 > 0, \forall h \in (0, h_0), \quad \|(\hat{\mathbf{g}}_m(h) - E_{m,n}(h))\hat{\mathbf{v}}_{m,n}(\cdot, h)\|_{L^2(I_h)} \leq C_{m,n}h^{3/2}.$$

Moreover we have

$$\begin{aligned}\|\widehat{\mathbf{g}}_{m,n}(\cdot, h)\|_{L^2(I_h)} &= \|f_{m,n}(\cdot, h)\|_{L^2(\mathbb{R})} + \mathcal{O}(h^\infty) \\ &= \|f_{0,n}\|_{L^2(\mathbb{R})} + \mathcal{O}(h^{1/2}) \\ &= 1 + \mathcal{O}(h^{1/2})\end{aligned}$$

where the above estimates depends on (m, n) . Since $\mathbf{g}_m(h)$ is unitarily equivalent to $h\widehat{\mathbf{g}}_m(h)$, $\mu_{m,n}(h)/h$ is the n -th eigenvalue of $\widehat{\mathbf{g}}_m(h)$ and the spectral theorem applied to (2.21) shows that

$$(2.22) \quad \exists C_{m,n}, \exists h_0 > 0, \quad \frac{\mu_{m,n}(h)}{h} \leq E_{m,n}(h) + C_{m,n}h^{3/2}$$

and we have proved the upper bound of Proposition 2.2.

• *Arguments for the lower bound.* The complete procedure for the proof of the lower bound of the eigenvalues of $\widehat{\mathbf{g}}_m(h)$ using the harmonic approximation can be found in [7, Chapter 4] or [11, Chapter 3]. We recall here the main arguments. Let

$$\widehat{\Phi}_0(t, h) := (1 + \sqrt{ht}) \log(1 + \sqrt{ht}) - \sqrt{ht}$$

be the distance of Agmon in the t -variable, the estimates provided in (2.7) becomes:

$$\forall \beta \in (0, 1), \quad \|e^{\beta \frac{\widehat{\Phi}_0}{h}} \widehat{\mathbf{u}}_{m,n}(\cdot, h)\|_{L^2(I_h)} \leq C(E, \beta)$$

where $\widehat{\mathbf{u}}_{m,n}(\cdot, h)$ is the n -th eigenvector associated to $\widehat{\mathbf{g}}_m(h)$. Therefore there holds *a priori* estimates on the eigenfunctions proving that they concentrate near $t = 0$ when h tends to 0. Using a Grushin procedure (see [10]), these eigenfunctions are used as quasi-modes for the first order approximation L_0 and this provides a rough lower bound on the eigenvalues $\frac{\mu_{m,n}(h)}{h}$ of $\widehat{\mathbf{g}}_m(h)$ by the eigenvalues of L_0 that are the Landau levels, modulo some remainders. Combining this with (2.22), we get that there are gaps in the spectrum of $\widehat{\mathbf{g}}_m(h)$ and the spectral theorem applied to (2.21) proved the lower bound on $\frac{\mu_{m,n}(h)}{h}$ and therefore the lower bound of Proposition 2.2.

Notation	Operator	Space	Form	Eigenpairs
$H_{\mathbf{A}}$	$(-i\nabla - \mathbf{A})^2$	$L^2(\mathbb{R}^3)$	—	spectrum = \mathbb{R}_+
$g_m(k)$	$-\frac{1}{r}\partial_r r \partial_r + \frac{m^2}{r^2} + (\log r - k)^2$	$L^2(\mathbb{R}_+, r dr)$	q_m^k	$(\lambda_{m,n}(k), u_{m,n}(r, k))$
$\tilde{g}_m(k)$	$-\partial_r^2 + \frac{m^2 - \frac{1}{4}}{r^2} + (\log r - k)^2$	$L^2(\mathbb{R}_+, dr)$	\tilde{q}_m^k	$(\lambda_{m,n}(k), \tilde{u}_{m,n}(r, k))$
$\mathbf{g}_m(h)$	$-h^2 \frac{1}{\rho} \partial_\rho \rho \partial_\rho + h^2 \frac{m^2}{\rho^2} + (\log \rho)^2$	$L^2(\mathbb{R}_+, \rho d\rho)$	\mathbf{q}_m^h	$(\mu_{m,n}(h), \mathbf{u}_{m,n}(\rho, h))$
$\tilde{\mathbf{g}}_m(h)$	$-h^2 \partial_\rho^2 + h^2 \frac{m^2 - \frac{1}{4}}{\rho^2} + (\log \rho)^2$	$L^2(\mathbb{R}_+, d\rho)$	$\tilde{\mathbf{q}}_m^h$	$(\mu_{m,n}(h), \tilde{\mathbf{u}}_{m,n}(\rho, h))$
$\widehat{\mathbf{g}}_m(h)$	$-\partial_t^2 + h \frac{m^2 - \frac{1}{4}}{(1+h^{1/2}t)^2} + (\log(1+h^{1/2}t))^2$	$L^2(I_h, dt)$	$\widehat{\mathbf{q}}_m^h$	$(h^{-1}\mu_{m,n}(h), \widehat{\mathbf{u}}_{m,n}(t, h))$

TABLE 1. Operators and notations. Remind that $\rho = hr$ with $r = \sqrt{x^2 + y^2}$, $h = e^{-k}$ and $I_h = (-h^{-1/2}, +\infty)$.

2.4. Numerical approximation of the band functions. We use the finite element library Melina ([16]) to compute numerical approximations of the band functions $\lambda_{m,n}(k)$ with $0 \leq m \leq 2$ and $1 \leq n \leq 4$. For $k \in [-2, 6]$, the computations are made on the interval $[0, L]$ with L large enough and an artificial Dirichlet boundary condition at $r = L$. According to the decay of the eigenfunctions provided by the Agmon estimates we have chosen $L = 2e^6$ so that the region $\{r \sim e^k\}$ where are localized the associated eigenfunction is included in the computation domain.

On Figure 1 we have plot the numerical approximation of $\lambda_{m,n}(k)$ for the range of parameters described above. According to the theory, they all decrease from $+\infty$ toward 0. Notice that the band functions may cross for different values of m .

On figure 2 we have zoomed on the lowest energies $\lambda \ll 1$ and we have also plotted the first order asymptotics $k \mapsto (2n - 1)e^{-k}$. We see that for set $1 \leq n \leq 4$, the band functions $\lambda_{m,n}(k)_{0 \leq m \leq 2}$ cluster around the first order asymptotic $(2n - 1)e^{-k}$ according to Theorem 1.1.




FIGURE 1. The band functions $\lambda_{m,n}(k)$ for $0 \leq m \leq 2$ and $1 \leq n \leq 4$ and $k \in [-2, 6]$.




FIGURE 2. Zoom on the lowest energies compared with the first order asymptotics $(2n-1)e^{-k}$. Each cluster corresponds to an energy level n .

3. CONSTRUCTION OF QUASI-MODES AND INFINITENESS OF NEGATIVE EIGENVALUES

In this section we prove Theorem 1.2 giving infinitely many eigenvalues below 0 for a slowly decreasing perturbation.

First, we consider V depending only on (r, z) and we construct quasi-modes which allow to reduce the existence of infinitely many negatives eigenvalues to the existence of sufficiently small eigenvalues of some 1D-effective problems $D_z^2 - V_{m,n}$. Then, we study the effective potential $V_{m,n}$ and conclude the proof of Theorem 1.2.

3.1. Quasi-modes. We construct quasi-modes for the perturbed operator $H_A - V$ where V is axisymmetrical. Let

$$\psi_{m,n}(r, \phi, z, k) := e^{im\phi} e^{ikz} u_{m,n}(r, k) f(z)$$

where $f \in L^2(\mathbb{R})$, (m, n, k) will be chosen later and $u_{m,n}(r, k)$ is a normalized eigenfunction of $g_m(k)$ associated with $\lambda_{m,n}(k)$. We have:

Lemma 3.1. *For any $\epsilon > 0$,*

$$(3.1) \quad \langle (H_{\mathbf{A}} - V)\psi_{m,n}, \psi_{m,n} \rangle \leq (1+\epsilon)\lambda_{m,n}(k)\|f\|_{L^2(\mathbb{R})}^2 + (1+\epsilon^{-1}) \|D_z f\|_{L^2(\mathbb{R})}^2 - \langle V_{m,n}(\cdot, k)f, f \rangle_{L^2(\mathbb{R})}$$

with

$$(3.2) \quad V_{m,n}(z, k) := \int_r |\tilde{u}_{m,n}(r, k)|^2 V(r, z) dr; \quad \tilde{u}_{m,n}(r, k) := \sqrt{r} u_{m,n}(r, k).$$

Proof. We have

$$(3.3) \quad H_{\mathbf{A}}\psi_{m,n}(r, \phi, z, k) = e^{im\phi} e^{ikz} f(z) g_m(k) u_{m,n}(r, k) + e^{im\phi} e^{ikz} u_{m,n}(r, k) (D_z^2 f + 2(\log r - k) D_z f(z)) ,$$

that is

$$(3.4) \quad (H_{\mathbf{A}} - V)\psi_{m,n}(r, \phi, z, k) = \lambda_{m,n}(k)\psi_{m,n}(r, \phi, z, k) + e^{im\phi} e^{ikz} u_{m,n}(r, k) (D_z^2 f + 2(\log r - k) D_z f(z) - V(r, z)f(z)) .$$

$$(3.5) \quad (H_{\mathbf{A}} - V)\psi_{m,n} \cdot \overline{\psi_{m,n}} = \lambda_{m,n}(k) u_{m,n}(r, k)^2 f(z)^2 + u_{m,n}(r, k)^2 \left(D_z^2 f(z) + 2(\log r - k) D_z f(z) - V(r, z)f(z) \right) \overline{f(z)}.$$

Integrating over (r, z) in the weighted space $(\mathbb{R}_+ \times \mathbb{R}, r dr dz)$ we get

$$(3.6) \quad \langle (H_{\mathbf{A}} - V)\psi_{m,n}, \psi_{m,n} \rangle_{L^2(\mathbb{R}_+ \times \mathbb{R}, r dr dz)} = \lambda_{m,n}(k)\|f\|_{L^2(\mathbb{R})}^2 + \|D_z f\|^2 + 2 \int_{r,z} (\log r - k) |u_{m,n}(r, k)|^2 D_z f(z) \overline{f(z)} r dr dz - \int_z V_{m,n}(z, k) |f(z)|^2 dz.$$

Then, using that for any $\epsilon > 0$,

$$|2(\log r - k) D_z f(z) \overline{f(z)}| \leq \epsilon (\log r - k)^2 |f(z)|^2 + \epsilon^{-1} |D_z f|^2,$$

we deduce,

$$\begin{aligned} \langle (H_{\mathbf{A}} - V)\psi_{m,n}, \psi_{m,n} \rangle &\leq \lambda_{m,n}(k)\|f\|_{L^2(\mathbb{R})}^2 + (1 + \epsilon^{-1}) \|D_z f\|_{L^2(\mathbb{R})}^2 \\ &+ \epsilon \int_{r,z} (\log r - k)^2 |u_{m,n}(r, k)|^2 |f(z)|^2 r dr dz - \langle V_{m,n}(\cdot, k)f, f \rangle_{L^2(\mathbb{R})}. \end{aligned}$$

Since in the sense of quadratic form in $L^2(\mathbb{R}_+ \times \mathbb{R}, r dr dz)$, we have $(\log r - k)^2 \leq g_m(k)$, we obtain (3.1) using again that $g_m(k)u_{m,n}(r, k) = \lambda_{m,n}(k)u_{m,n}(r, k)$. \square

Remark 3.2. According to the Feynman-Hellmann formula, the third term in the right hand side of (3.6) is related to the derivative of $\lambda_{m,n}(k)$:

$$\lambda'_{m,n}(k) = -2 \int_{r,z} (\log r - k) |u_{m,n}(r, k)|^2 r dr.$$

This quantity could be studied more carefully as in [13] where it is done for another fibered operator, but here, we need only some rough estimates.

3.2. Estimate on the reduced potential. We are looking at the asymptotic behavior of the 1D potential $z \mapsto V_{m,n}(z, k)$ by using the localization properties of the eigenfunctions $\tilde{u}_{m,n}(\cdot, k)$ when k goes to $+\infty$. In this section all the Landau's notations refers to an asymptotic behavior when k goes to $+\infty$. Set $(m, n) \in \mathbb{Z} \times \mathbb{N}^*$, $C_{m,n} > 2n - 1$ and choose k large enough such that $\lambda_{m,n}(k) \leq C_{m,n}e^{-k}$ (see Theorem 1.1). Write that $\mathbb{R} = I_k \cup \mathbb{C}I_k$ with $I_k = [e^k - a(k), e^k + a(k)]$ and $a(k) = o(e^k)$ will be chosen later. We use (2.9) with $E = 0$:

$$\int_{\mathbb{C}I_k} |\tilde{u}_{m,n}(r, k)|^2 dr \leq C(0, \beta) \sup_{r \in \mathbb{C}I_k} e^{-\beta\Phi_0(r, k)}$$

where the Agmon distance Φ_0 is defined in (2.8). Since $\Phi_0(\cdot, k)$ is decreasing on $(0, e^k)$ and increasing on $(e^k, +\infty)$ we have

$$\inf_{\mathbb{C}I_k} \Phi_0(\cdot, k) = \min(\Phi_0(e^k \pm a(k))).$$

An asymptotic expansion at these points provides

$$\Phi_0(e^k \pm a(k), k) \underset{k \rightarrow +\infty}{=} \frac{1}{2}a^2(k)e^{-k} + O(a(k)^3e^{-2k}).$$

Assume that

$$(3.7) \quad \lim_{k \rightarrow +\infty} a^2(k)e^{-k} = +\infty \quad \text{and} \quad \lim_{k \rightarrow +\infty} a^3(k)e^{-2k} = 0$$

then we have

$$e^{-\beta\Phi_0(e^k \pm a(k), k)} \underset{k \rightarrow +\infty}{\sim} e^{-\frac{\beta}{2}a(k)^2e^{-k}}$$

The condition (3.7) is valid for any $a(k)$ satisfying

$$e^{\frac{k}{2}} \ll a(k) \ll e^{\frac{2k}{3}}$$

and for such an $a(k)$ we get

$$(3.8) \quad \sup_{r \in \mathbb{C}I_k} e^{-\beta\Phi_0(r, k)} \underset{k \rightarrow +\infty}{\sim} e^{-\frac{\beta}{2}a(k)^2e^{-k}}.$$

We have

$$\begin{aligned} V_{m,n}(z, k) &\geq \inf_{r \in I_k} V(r, z) \int_{I_k} |\tilde{u}_{m,n}(r, k)|^2 dr \\ &\geq \inf_{r \in I_k} V(r, z) (1 - C(0, \beta) \sup_{r \in \mathbb{C}I_k} e^{-\beta\Phi_0(r, k)}) \end{aligned}$$

where we have used $\|\tilde{u}_{m,n}(\cdot, k)\|_{L^2(\mathbb{R}_+)} = 1$.

Set $\beta \in (0, 1)$ once for all. Choose $\epsilon > 0$. Then we deduce from the choice of $a(k)$ in (3.7) and (3.8) that there exists k_0 that depends *a priori* of (m, n) such that

$$(3.9) \quad \forall k \geq k_0, \forall z \in \mathbb{R}, \quad V_{m,n}(z, k) \geq (1 - \epsilon) \inf_{r \in I_k} V(r, z)$$

3.3. Proof of Theorem 1.2. According to the min-max principle, since V satisfies (1.4), it is sufficient to prove the infinity of the negative eigenvalues for the axisymmetric potential $V(r, z) = \langle r \rangle^{-\alpha} v_{\perp}(z)$. Let us denote $H_{\mathbf{A}}^m$ the restriction of $H_{\mathbf{A}}$ to $e^{im\phi} L^2(\mathbb{R}_+ \times \mathbb{R}, r dr dz)$. For V axisymmetric, $H_{\mathbf{A}} - V$ is unitarily equivalent to $\oplus_{m \in \mathbb{Z}} (H_{\mathbf{A}}^m - V)$, then $H_{\mathbf{A}} - V$ has infinitely many negative eigenvalues provided that

- Either $H_{\mathbf{A}}^m - V$ has at least one's for all $m \in \mathbb{Z}$,
- or there exists $m \in \mathbb{Z}$ such that $H_{\mathbf{A}}^m - V$ has infinitely many negative eigenvalues.

Thanks to the min-max principle, Lemma 3.1, implies that for each $m \in \mathbb{Z}$ the number of negative eigenvalues of $H_{\mathbf{A}}^m - V$ is at least the number of eigenvalues of $(1 + \epsilon^{-1})D_z^2 - V_{m,n}(\cdot, k)$ below $-(1 + \epsilon)\lambda_{m,n}(k)$, that is the number of eigenvalues of $D_z^2 - \frac{\epsilon}{1+\epsilon}V_{m,n}(\cdot, k)$ below $-\epsilon\lambda_{m,n}(k)$.

For $V(r, z) = \langle r \rangle^{-\alpha} v_{\perp}(z)$, the inequality (3.9) implies:

$$\forall k \geq k_0, \forall z \in \mathbb{R}, \quad V_{m,n}(z, k) \geq C e^{-\alpha k} v_{\perp}(z),$$

and choosing k large enough such that $\lambda_{m,n}(k) \leq C_{m,n} e^{-k}$, we deduce that the number of negative eigenvalues of $H_{\mathbf{A}}^m - V$ is at least the number of eigenvalues of

$$D_z^2 - \frac{C\epsilon}{1+\epsilon} e^{-\alpha k} v_{\perp}$$

below $-\epsilon C_{m,n} e^{-k}$. Then Theorem 1.2 follows by applying the following lemmas (Lemma 3.3 and Lemma 3.4), for k sufficiently large with $\Lambda(k) = e^{-\alpha k}$, $v = \frac{C\epsilon}{1+\epsilon} v_{\perp}$ and $\lambda(k) = \epsilon C_{m,n} e^{-k}$.

3.4. Lemmas on negative eigenvalues for a family of some 1D Schrödinger operators.

Lemma 3.3. Let $h(k) = D_z^2 - \Lambda(k)v(z)$ on \mathbb{R} , $k \in \mathbb{R}$ with:

$$v \in L^1(\mathbb{R}); \quad \int_{\mathbb{R}} v(z) dz > 0, \quad \Lambda(k) > 0.$$

Let $\lambda(k)$ be a positive function of $k \in \mathbb{R}$ such that

$$(3.10) \quad \lim_{k \rightarrow +\infty} \lambda(k) = 0; \quad \lim_{k \rightarrow +\infty} \frac{\lambda(k)}{\Lambda(k)^2} = 0.$$

Then, for k sufficiently large, $h(k) + \lambda(k)$ has at least one negative eigenvalue.

Proof. Let us introduce the L^2 -normalized function

$$v_k(z) := a(k)^{\frac{1}{2}} e^{-a(k)|z|}$$

with $a(k)$ satisfying $\lim_{k \rightarrow +\infty} a(k) = 0$ and to be chosen. We use $v_k(z)$ as a quasi-mode:

$$\langle h(k)v_k, v_k \rangle = a(k)^2 - \Lambda(k)a(k) \int_{\mathbb{R}} v(z) e^{-2a(k)|z|} dz.$$

Since

$$\lim_{k \rightarrow +\infty} \int_{\mathbb{R}} v(z) e^{-2a(k)|z|} dz = \int_{\mathbb{R}} v(z) dz > 0,$$

for k sufficiently large, there exists $C > 0$ such that:

$$\langle h(k)v_k, v_k \rangle \leq a(k)^2 - C\Lambda(k)a(k).$$

By using the min-max principle, it remains to choose $a(k)$ such that $a(k)^2 - C\Lambda(k)a(k) < -\lambda(k)$. Under the assumption (3.10), the polynomial $X^2 - C\Lambda(k)X + \lambda(k)$ has two real roots $a_+(k) > a_-(k) > 0$ with $a_-(k) \leq \frac{2\lambda(k)}{C\Lambda(k)}$ tending to 0 as k tends to infinity, see (3.10). Then, there exists $a(k)$ such that, for k sufficiently large,

$$\langle h(k)v_k, v_k \rangle < -\lambda(k),$$

and Lemma (3.3) holds. \square

Lemma 3.4. *Let $h(k) = D_z^2 - V_k$ on \mathbb{R} , $k \in \mathbb{R}$ with V_k satisfying:*

$$V_k(z) \geq \Lambda(k)\langle z \rangle^{-\gamma}; \quad \gamma \in (0, 2); \quad \Lambda(k) \in (0, 1).$$

Let $\lambda(k)$ be a positive function of $k \in \mathbb{R}$ such that

$$(3.11) \quad \lim_{k \rightarrow +\infty} \frac{\lambda(k)}{\Lambda(k)^{\frac{2}{2-\gamma}}} = 0.$$

Then, for k sufficiently large, $h(k) + \lambda(k)$ has at least one negative eigenvalue and the number of negative eigenvalues tends to infinity, as k tends to infinity.

Proof. Using the change of variable $y = \Lambda(k)^{\frac{1}{2-\gamma}}z$, it is clear that $h(k)$ is unitarily equivalent to $\Lambda(k)^{\frac{2}{2-\gamma}}\tilde{h}(k)$ with

$$\tilde{h}(k) := D_y^2 - \frac{1}{\Lambda(k)^{\frac{2}{2-\gamma}}}V_k\left(\frac{y}{\Lambda(k)^{\frac{1}{2-\gamma}}}\right).$$

By assumption on V_k , we have:

$$\frac{1}{\Lambda(k)^{\frac{2}{2-\gamma}}}V_k\left(\frac{y}{\Lambda(k)^{\frac{1}{2-\gamma}}}\right) \geq (\Lambda^{\frac{2}{2-\gamma}}(k) + y^2)^{-\frac{\gamma}{2}} \geq (1 + y^2)^{-\frac{\gamma}{2}}$$

where we have used $\Lambda(k) \in (0, 1)$. Then the min-max principle implies that the number of negative eigenvalues of $h(k) + \lambda(k)$ is larger than the number of eigenvalues of $D_y^2 - \langle y \rangle^{-\gamma}$ below $-\frac{\lambda(k)}{\Lambda^{\frac{2}{2-\gamma}}(k)}$. Since $\gamma < 2$, it is known (see [21, Theorem XIII.82]) that $D_y^2 - \langle y \rangle^{-\gamma}$ has infinitely many negative eigenvalues and Lemma 3.4 follows from (3.11). \square

4. FINITE NUMBER OF NEGATIVE EIGENVALUE FOR PERTURBATION BY SHORT RANGE POTENTIAL

The aim of this section is to prove Theorem 1.3. In Section 4.2, using the Birman-Schwinger principle, we reduce the proof to the analysis of some compact operator involving the contribution of the small energies ($\lambda_{m,n}(k) \leq \nu \ll 1$). Exploiting that the eigenfunctions associated with $\lambda_{m,n}(k)$ are localized near e^k , we obtain in Section 4.3 an upper bound of the counting function including interactions between the behavior in r and z via a convolution product and the Fourier transform w.r.t. z . Then, exploiting a uniform lower bound of the band functions (see Section 4.1), we are able to prove Theorem 1.3 by computing the Hilbert-Schmidt norm of a canonical operator and by using standard Young inequality (see Section 4.4).

4.1. Uniform estimate for the one-dimensional problem. In order to prove Theorem 1.3 we need an uniform lower bound on the band functions near 0.

Lemma 4.1. *Let $\nu_0 > 0$. There exists $C_0 > 0$ such that for all (m, n, h) satisfying $\mu_{m,n}(h) \leq \nu_0$ we have*

$$\mu_{m,n}(h) \geq C_0 n h$$

• *Sketch of the proof.* For convenience, first we work with the operator

$$\mathbf{g}_m(h) = -h^2 \frac{1}{\rho} \partial_\rho \rho \partial_\rho + V_h^m \quad \text{with} \quad V_h^m(\rho) := \log(\rho)^2 + h^2 \frac{m^2}{\rho^2}.$$

We notice that in the sense of quadratic form we have $\mathbf{g}_m(h) \geq \mathbf{g}_0(h)$ and $\text{dom}(\mathbf{g}_m(h)) \subset \text{dom}(\mathbf{g}_0(h))$, therefore for all $m \in \mathbb{Z}$ there holds $\mu_{m,n}(h) \geq \mu_{0,n}(h)$ and it is sufficient to prove the result for $m = 0$.

We will split the proof depending on which region belongs the parameter h :

- (1) For $h \in (0, h_0)$ with h_0 to be chosen, we will use the semi-classical analysis and the Agmon estimates on the eigenfunctions in order to compare $\mathbf{g}_0(h)$ with more standard operators. The idea is to bound from below the potential $\log^2 \rho$ on a suitable interval by a quadratic potential such that the associated operator has known spectrum.
- (2) Since $h \rightarrow \mu_{0,n}(h)$ is unbounded for large h , there exists h_{ν_0} such that for $h \geq h_{\nu_0}$ the eigenvalues $\mu_{m,n}(h)$ are outside the region $\{\mu \leq \nu_0\}$.
- (3) On the compact $[h_0, h_{\nu_0}]$, since $n \rightarrow \mu_{0,n}(h)$ is unbounded for large n , we may find $N \geq 1$ such that for $n \geq N$ the eigenvalues $\mu_{m,n}(h)$ are outside the region $\{\mu \leq \nu_0\}$. Therefore the Lemma is clear on this region since we have to deal with a finite number of eigenvalues.

• *proof.* Assume $\mu_{m,n}(h) \leq \nu_0$. Denote by $0 < \rho_1 < 1 < \rho_2$ the two real numbers (depending on ν_0) such that

$$\log^2(\rho_1) = \log^2(\rho_2) = \nu_0.$$

Set $\rho'_1 \in (0, \rho_1)$, $\rho'_2 \in (\rho_2, +\infty)$ and $I(\nu_0) := (\rho'_1, \rho'_2)$. Let $M(\nu_0) := \min(\phi_0(\rho'_1), \phi_0(\rho'_2))$ where ϕ_0 is defined by

$$\phi_0(\rho) := \left| \int_1^\rho \sqrt{((\log \rho)^2 - \nu_0)_+} d\rho \right|$$

By construction we have $M(\nu_0) > 0$ and the Agmon estimate (2.7) provides $h_0 > 0$ such that (uniformly in n):

$$\forall h \in (0, h_0), \quad \int_{I(\nu_0)} |\mathbf{u}_{0,n}(\rho, h)|^2 \rho d\rho \leq C(\nu_0, \beta) e^{-\beta M(\nu_0)/h}$$

where $\beta \in (0, 1)$ is set.

Recall that $\tilde{u}_{m,n}(\rho, h) = \sqrt{\rho} u_{m,n}(\rho, h)$ is a normalized eigenfunction of $\tilde{\mathbf{g}}_m(h) = \mathcal{M} \mathbf{g}_m(h) \mathcal{M}^*$ associated with the eigenvalue $\mu_{m,n}(h)$. It satisfies

$$(4.1) \quad \forall h \in (0, h_0), \quad \int_{I(\nu_0)} |\tilde{\mathbf{u}}_{0,n}(\rho, h)|^2 d\rho \leq C(\nu_0, \beta) e^{-\beta M(\nu_0)/h}$$

Remark 4.2. Since $\mathbf{g}_m(h) \geq \mathbf{g}_0(h)$, in the sense of quadratic form, the above estimate (4.1) holds also for $\tilde{\mathbf{u}}_{m,n}$:

$$\forall h \in (0, h_0), \quad \int_{\mathbb{C}I(\nu_0)} |\tilde{\mathbf{u}}_{m,n}(\rho, h)|^2 d\rho = \int_{\mathbb{C}I(\nu_0)} |\mathbf{u}_{m,n}(\rho, h)|^2 \rho d\rho \leq C(\nu_0, \beta) e^{-\beta M(\nu_0)/h}$$

uniformly with respect to (m, n) such that $\mu_{m,n}(h) \leq \nu_0$. This estimate will be used in Section 4.2.

Set $\epsilon_0 \in (0, \rho'_1)$. Let $(\chi_j)_{j=1,2} \in \mathcal{C}^\infty(\mathbb{R}_+, [0, 1])$ be a partition of the unity of \mathbb{R}_+ such that $\chi_1^2 + \chi_2^2 = 1$ with $\chi_2 = 0$ on $I(\nu_0)$ and $\chi_2 = 1$ on $(0, \rho'_1 - \epsilon_0) \cup (\rho'_2 + \epsilon_0, +\infty)$. We may assume that there exists $C > 0$ such that $\sum_j |\nabla \chi_j|^2 \leq C$.

The IMS formula provides for any eigenfunction $\tilde{\mathbf{u}}_{0,n}(\cdot, h)$:

$$\begin{aligned} \tilde{\mathbf{q}}_0^h(\tilde{\mathbf{u}}_{0,n}(\cdot, h)) &= \sum_{j=1,2} \tilde{\mathbf{q}}_0^h(\chi_j \tilde{\mathbf{u}}_{0,n}(\cdot, h)) - \sum_{j=1,2} \|(\nabla \chi_j) \tilde{\mathbf{u}}_{0,n}(\cdot, h)\|_{L^2(\mathbb{R}_+)}^2 \\ &\geq \tilde{\mathbf{q}}_0^h(\chi_1 \tilde{\mathbf{u}}_{0,n}(\cdot, h)) - C \int_{\text{supp}(\chi_2')} |\tilde{\mathbf{u}}_{0,n}(\rho, h)|^2 d\rho \end{aligned}$$

and therefore using (4.1):

$$(4.2) \quad \tilde{\mathbf{q}}_0^h(\tilde{\mathbf{u}}_{0,n}(\cdot, h)) \geq \tilde{\mathbf{q}}_0^h(\chi_1 \tilde{\mathbf{u}}_{0,n}(\cdot, h)) - C(\nu_0, \beta) e^{-\beta M(\nu_0)/h}.$$

We now bound from below $\tilde{\mathbf{q}}_0^h(\chi_1 \tilde{\mathbf{u}}_{0,n}(\cdot, h))$ using a lower bound on the potential. We have

$$(4.3) \quad \exists C(\nu_0) \in (0, 1), \forall \rho \in J(\nu_0), \quad C(\nu_0)(\rho - 1)^2 \leq \log^2 \rho$$

where we have denoted $J(\nu_0) := (\rho'_1 - \epsilon_0, \rho'_2 + \epsilon_0)$.

Assume $n \neq n'$. Since $\langle \tilde{\mathbf{u}}_{0,n}(\cdot, h), \tilde{\mathbf{u}}_{0,n'}(\cdot, h) \rangle_{L^2(\mathbb{R}_+)} = 0$, we deduce from (4.1) that

$$(4.4) \quad |\langle \chi_1 \tilde{\mathbf{u}}_{0,n}(\cdot, h), \chi_1 \tilde{\mathbf{u}}_{0,n'}(\cdot, h) \rangle_{L^2(\mathbb{R}_+)}| \leq C(\nu_0, \beta) e^{-\beta M(\nu_0)/h}.$$

Let us introduce the harmonic oscillator

$$\mathbf{g}^{\text{low}}(h) := -h^2 \partial_\rho^2 + (\rho - 1)^2, \quad \rho \in \mathbb{R}$$

initially defined on $\mathcal{C}_0^\infty(\mathbb{R})$ and close on $L^2(\mathbb{R})$, whose eigenvalues are $\{(2n - 1)h\}_{n \in \mathbb{N}^*}$. Due to (4.3) and since $\text{supp}(\chi_1) = J(\nu_0)$ we have

$$(4.5) \quad \begin{aligned} \tilde{\mathbf{q}}_0^h(\chi_1 \tilde{\mathbf{u}}_{0,n}(\cdot, h)) &\geq C(\nu_0) \langle \mathbf{g}^{\text{low}}(h) \chi_1 \tilde{\mathbf{u}}_{0,n}(\cdot, h), \chi_1 \tilde{\mathbf{u}}_{0,n}(\cdot, h) \rangle_{L^2(\mathbb{R})} \\ &\quad - \frac{h^2}{4(\rho'_1 - \epsilon_0)^2} \|\chi_1 \tilde{\mathbf{u}}_{0,n}(\cdot, h)\|_{L^2(\mathbb{R}_+)}^2 \end{aligned}$$

where in the right hand side, $\chi_1 \tilde{\mathbf{u}}_{0,n}$, extended by 0 on \mathbb{R}_- , is also considered as a function defined on \mathbb{R} .

Recall (4.4), the min-max principle combined with (4.5) provides

$$(4.6) \quad \tilde{\mathbf{q}}_0^h(\chi_1 \tilde{\mathbf{u}}_{0,n}(\cdot, h)) \geq \left(C(\nu_0)(2n - 1)h - \tilde{C}(\nu_0, \beta)h^2 \right) \|\chi_1 \tilde{\mathbf{u}}_{0,n}(\cdot, h)\|_{L^2(\mathbb{R}_+)}^2.$$

Using (4.1) we get $|1 - \|\chi_1 \tilde{\mathbf{u}}_{0,n}(\cdot, h)\|_{L^2(\mathbb{R}_+)}^2| \leq C(\nu_0, \beta) e^{-\beta M(\nu_0)/h}$.

Therefore combining (4.2) and (4.6) we have proved the existence of $h_0 > 0$ and $C_0 > 0$ such that for all $(0, n, h)$ such that $\mu_{0,n}(h) \leq \nu_0$ we have

$$\forall h \in (0, h_0), \quad \mu_{0,n}(h) = \tilde{\mathbf{q}}_0^h(\tilde{\mathbf{u}}_{0,n}(\cdot, h)) \geq C_0 n h.$$

We now have to deal with the region $h \in (h_0, +\infty)$. Since $\mu_{0,n}(h)$ tends to $+\infty$ as h tends to $+\infty$, there exists $h_{\nu_0} > 0$ such that

$$\forall n \in \mathbb{N}^*, \forall h \geq h_{\nu_0}, \quad \mu_{0,n}(h) \geq \nu_0.$$

Therefore we are led to prove the lower bound for $h \in [h_0, h_{\nu_0}]$. Since for all $h > 0$ the sequence $(\mu_{m,n}(h))_{n \geq 1}$ converges toward $+\infty$, there exists $n(h)$ such that for all $n \geq n(h)$ we have $\mu_{m,n}(h) \geq \nu_0$. Due to a compact argument we find $N \in \mathbb{N}^*$ such that

$$\forall n > N, \forall h \in [h_0, h_{\nu_0}], \quad \mu_{0,n}(h) \geq \nu_0.$$

Define $C(h) := \min_{1 \leq n \leq N} \mu_{0,n}(h)/n$ and $C := \min_{h \in [h_0, h_{\nu_0}]} \frac{C(h)}{h}$. We clearly have $C > 0$ and by construction, for all $(n, h) \in \mathbb{N}^* \times [h_0, h_{\nu_0}]$ such that $\mu_{0,n}(h) \leq \nu_0$ we have

$$\mu_{0,n}(h) \geq C n h$$

therefore the lemma is proved for $h \in [h_0, h_{\nu_0}]$.

Remark 4.3. In (4.6), the remainder term of order h^2 involves the contributions of $\frac{-h^2}{4\rho^2}$ and has been controlled on $J(\nu_0)$. Another strategy, which improves the remainder term, would have been to work in the weighted space $L_\rho^2(\mathbb{R}_+)$ and to consider

$$\mathbf{g}^{\text{low}}(h) := -h^2 \frac{1}{\rho} \partial_\rho \rho \partial_\rho + (\rho - 1)^2, \quad \rho > 0.$$

In this case, (4.6) is replaced by

$$\mathbf{q}_0^h(\chi_1 \mathbf{u}_{0,n}(\cdot, h)) \geq C(\nu_0)(\zeta_n(h) - C(\nu_0, \beta)e^{-\beta M(\nu_0)/h}) \|\chi_1 \mathbf{u}_{0,n}(\cdot, h)\|_{L_\rho^2(\mathbb{R}_+)}^2$$

with $\zeta_n(h)$ the n -th eigenvalue of the operator $\mathbf{g}^{\text{low}}(h)$. These eigenvalues have already been studied in [25, Section 4.2] and [17] and they can be bounded from below by $C_1 n h$ by exploiting the results from [17].

4.2. Bring the norm of a canonical operator. Let $\lambda > 0$, for simplicity we denote by $\mathcal{N}(\lambda) := \mathcal{N}_{\mathbf{A}, V}(\lambda)$ the number of negative eigenvalues of $H_{\mathbf{A}} - V$ below $-\lambda$:

$$\mathcal{N}(\lambda) := \# \left(\mathfrak{S}(H_{\mathbf{A}} - V) \cap]-\infty, -\lambda] \right).$$

We want to prove that there exists $C > 0$ independent of λ , such that $\mathcal{N}(\lambda) \leq C$. Let us introduce the axisymmetric non negative potential

$$(4.7) \quad V_0(r, z) := \langle r \rangle^{-\alpha} v_\perp(z).$$

The assumption (1.5) means that $V \leq V_0$. Then the min-max principle gives:

$$(4.8) \quad \mathcal{N}(\lambda) \leq \mathcal{N}_0(\lambda) := \# \left(\mathfrak{S}(H_{\mathbf{A}} - V_0) \cap]-\infty, -\lambda] \right).$$

According to the Birman-Schwinger principle, for $\lambda > 0$,

$$(4.9) \quad \mathcal{N}_0(\lambda) = n_+ \left(1, V_0^{\frac{1}{2}} (H_{\mathbf{A}} + \lambda)^{-1} V_0^{\frac{1}{2}} \right),$$

where for a self-adjoint operator T , $n_+(s, T) := \text{Tr } \mathbb{1}_{(s, \infty)}(T)$; is the counting function of positive eigenvalues of T .

Fix a real number $\nu > 0$ (chosen sufficiently small later) and let us introduce the orthogonal projections $P_\nu := \mathbb{1}_{[0, \nu]}(H_{\mathbf{A}})$ and $\overline{P}_\nu := I - P_\nu = \mathbb{1}_{[\nu, +\infty]}(H_{\mathbf{A}})$.

Since $H_{\mathbf{A}} \overline{P}_\nu \geq \nu$, the compact operator $V_0^{\frac{1}{2}} (H_{\mathbf{A}} + \lambda)^{-1} \overline{P}_\nu V_0^{\frac{1}{2}}$ is uniformly bounded with respect to $\lambda \geq 0$ and from the Weyl inequality, for any $\epsilon > 0$, we have:

$$(4.10) \quad n_+ \left(1, V_0^{\frac{1}{2}} (H_{\mathbf{A}} + \lambda)^{-1} V_0^{\frac{1}{2}} \right) \leq n_+ \left(1 - \epsilon, V_0^{\frac{1}{2}} (H_{\mathbf{A}} + \lambda)^{-1} P_\nu V_0^{\frac{1}{2}} \right) + C_\nu, \quad C_\nu \geq 0.$$

According to the decomposition:

$$H_{\mathbf{A}} = \Phi^* \mathcal{F}_3^* \left(\sum_{(m,n) \in \mathbb{Z} \times \mathbb{N}^*}^{\oplus} \int_{k \in \mathbb{R}}^{\oplus} \lambda_{m,n}(k) P_{m,n}(k) dk \right) \mathcal{F}_3 \Phi,$$

with $P_{m,n}(k) : f \mapsto \langle f, u_{m,n}(\cdot, k) \rangle u_{m,n}(\cdot, k)$, the orthogonal projection onto $u_{m,n}(\cdot, k) \in L^2(\mathbb{R}_+, r dr)$, we have

$$V_0^{\frac{1}{2}} (H_{\mathbf{A}} + \lambda)^{-1} P_\nu V_0^{\frac{1}{2}} = V_0^{\frac{1}{2}} \Phi^* \mathcal{F}_3^* \left(\sum_{(m,n) \in \mathbb{Z} \times \mathbb{N}^*}^{\oplus} \int_{k \in \mathbb{R}}^{\oplus} P_{m,n}(k) \frac{\mathbb{1}_{[0, \nu]}(\lambda_{m,n}(k))}{\lambda_{m,n}(k) + \lambda} dk \right) \mathcal{F}_3 \Phi V_0^{\frac{1}{2}}.$$

Since V_0 is axisymmetric, this operator is unitarily equivalent to the direct sum of

$$K_{\nu, m}(\lambda) := V_0^{\frac{1}{2}} \mathcal{F}_3^* \left(\int_{k \in \mathbb{R}}^{\oplus} \sum_{n \in \mathbb{N}^*}^{\oplus} \tilde{P}_{m,n}(k) \frac{\mathbb{1}_{[0, \nu]}(\lambda_{m,n}(k))}{\lambda_{m,n}(k) + \lambda} dk \right) \mathcal{F}_3 V_0^{\frac{1}{2}},$$

defined in $L^2(\mathbb{R}_+ \times \mathbb{R}, dr dz)$, with $\tilde{P}_{m,n}(k) := \mathcal{M}^* P_{m,n}(k) \mathcal{M} = \langle \cdot, \tilde{u}_{m,n}(k) \rangle \tilde{u}_{m,n}(k, \cdot)$, the orthogonal projection onto $\tilde{u}_{m,n}(\cdot, k) \in L^2(\mathbb{R}_+, dr)$, $\tilde{u}_{m,n}(r, k) = \sqrt{r} u_{m,n}(r, k)$.

Let us prove that for some $s \in]0, 1[$, there exists ν sufficiently small such for any $m \in \mathbb{Z}$ and any $\lambda > 0$

$$(4.11) \quad n_+(s, K_{\nu, m}(\lambda)) = 0.$$

Then Theorem 1.3, is a consequence of (4.8), (4.9), (4.10) and (4.11).

Let us introduce the operator:

$$S_m(\lambda) : L^2(\mathbb{R}, l^2(\mathbb{N}^*)) \longrightarrow L^2(\mathbb{R}_+ \times \mathbb{R}, dr dz),$$

defined, for $(g_n(\cdot))_{n \in \mathbb{N}^*} \in L^2(\mathbb{R}, l^2(\mathbb{N}^*))$ by

$$(4.12) \quad S_m(\lambda)(g_n)(r, z) := \frac{V_0^{\frac{1}{2}}(r, z)}{\sqrt{2\pi}} \sum_{n \in \mathbb{N}^*} \int_{\mathbb{R}} g_n(k) \frac{e^{izk} \mathbb{1}_{[0, \nu]}(\lambda_{m,n}(k))}{(\lambda_{m,n}(k) + \lambda)^{\frac{1}{2}}} \tilde{u}_{m,n}(r, k) dk,$$

and its adjoint defined for $f \in L^2(\mathbb{R}_+ \times \mathbb{R}, dr dz)$, by

$$S_m(\lambda)^*(f)_n(k) = \frac{1}{\sqrt{2\pi}} \frac{\mathbf{1}_{[0,\nu]}(\lambda_{m,n}(k))}{(\lambda_{m,n}(k) + \lambda)^{\frac{1}{2}}} \int_{\mathbb{R}_+ \times \mathbb{R}} e^{-izk} \overline{\tilde{u}_{m,n}(r, k)} (V_0^{\frac{1}{2}} f)(r, z) dr dz.$$

We have:

$$K_{\nu,m}(\lambda) = S_m(\lambda) S_m(\lambda)^*,$$

and since

$$(4.13) \quad n_+(s, K_{\nu,m}(\lambda)) = n_+(s, S_m(\lambda) S_m(\lambda)^*) = n_+(s, S_m(\lambda)^* S_m(\lambda)),$$

we have to prove that for ν sufficiently small, the L^2 -norm of $S_m(\lambda)^* S_m(\lambda)$ admits an upper bound by $s < 1$ uniformly with respect to $m \in \mathbb{Z}$ and $\lambda > 0$.

4.3. Computations on the integral kernel of the canonical operator.

Proposition 4.4. *Let V_0 defined in (4.7) and $S_m(\lambda)$ defined in (4.12). Then there exist $C > 0$ and $\nu_0 > 0$ such that for all $\nu \in (0, \nu_0)$, the following upper bound of the Hilbert-Schmidt norm holds:*

$$(4.14) \quad \|S_m(\lambda)^* S_m(\lambda)\|_2^2 \leq C \sum_{n,n'} \int_k \int_{k'} \iota_{m,n'}(k', \nu) \iota_{m,n}(k, \nu) |\widehat{v}_\perp(k' - k)|^2 dk' dk$$

where we have set

$$\iota_{m,n}(k, \nu) := \frac{\mathbf{1}_{[0,\nu]}(\lambda_{m,n}(k))}{\lambda_{m,n}(k) + \lambda} e^{-\alpha k}.$$

Proof. We check that $S_m(\lambda)^* S_m(\lambda) : L^2(\mathbb{R}, l^2(\mathbb{N}^*)) \longrightarrow L^2(\mathbb{R}, l^2(\mathbb{N}^*))$ corresponds with

$$(4.15) \quad (S_m(\lambda)^* S_m(\lambda)(g_{n'}))_n(k) = \frac{1}{2\pi} L_{m,n}(k) \int_z \int_r \overline{\tilde{u}_{m,n}(r, k)} V_0(r, z) \sum_{n'} \int_{k'} g_{n'}(k') L_{m,n'}(k') \tilde{u}_{m,n'}(r, k') e^{iz(k'-k)} dk' dr dz$$

where we have denoted

$$L_{m,n}(k) := \frac{\mathbf{1}_{[0,\nu]}(\lambda_{m,n}(k))}{\sqrt{\lambda_{m,n}(k) + \lambda}}.$$

The integral kernel of this operator is

$$\begin{aligned} \mathfrak{N}_{m,n,n'}(k, k') &:= L_{m,n}(k) L_{m,n'}(k') \int_r \int_z V_0(r, z) \overline{\tilde{u}_{m,n}(r, k)} \tilde{u}_{m,n'}(r, k') e^{iz(k-k')} dz dr. \\ &= L_{m,n}(k) L_{m,n'}(k') \widehat{v}_\perp(k' - k) \int_r \langle r \rangle^{-\alpha} \overline{\tilde{u}_{m,n}(r, k)} \tilde{u}_{m,n'}(r, k') dr. \end{aligned}$$

Then the Hilbert-Schmidt norm is given by

$$(4.16) \quad 4\pi^2 \|S_m(\lambda)^* S_m(\lambda)\|_2^2 = \sum_{n,n'} \int_k \int_{k'} \int_r L_{m,n}(k)^2 L_{m,n'}(k')^2 |\widehat{v}_\perp(k' - k)|^2 \left| \int_r \langle r \rangle^{-\alpha} \overline{\tilde{u}_{m,n}(r, k)} \tilde{u}_{m,n'}(r, k') dr \right|^2 dk dk'.$$

Set $\nu_0 > 0$ and (m, n, k) such that $\lambda_{m,n}(k) \leq \nu_0$. Applying Remark 4.2 we know that there exists $I_k(\nu_0) := [\rho'_1 e^k, \rho'_2 e^k]$, $\rho'_1 < 1 < \rho'_2$, such that for any $k \geq k_0$ sufficiently large (independent of (m, n)),

$$\int_{\mathbb{C}I_k(\nu_0)} \langle r \rangle^{-\alpha} |\tilde{u}_{m,n}(k, r)|^2 dr \leq \int_{\mathbb{C}I_k(\nu_0)} |\tilde{u}_{m,n}(k, r)|^2 dr \leq C(\nu_0, \beta) e^{-\beta M(\nu_0) e^k}$$

with $\beta \in (0, 1)$ and $M(\nu_0) > 0$. On the other hand, on $I_k(\nu_0)$, we have

$$\int_{I_k(\nu_0)} \langle r \rangle^{-\alpha} |\tilde{u}_{m,n}(k, r)|^2 dr \leq C(\nu_0) e^{-\alpha k} \int_{I_k(\nu_0)} |\tilde{u}_{m,n}(k, r)|^2 dr \leq C(\nu_0) e^{-\alpha k}.$$

Consequently,

$$(4.17) \quad \int_{\mathbb{R}_+} \langle r \rangle^{-\alpha} |\tilde{u}_{m,n}(k, r)|^2 dr = O(e^{-\alpha k}),$$

uniformly with respect to $(m, n, k) \in \mathbb{Z} \times \mathbb{N}^* \times \mathbb{R}$ satisfying $\lambda_{m,n}(k) \leq \nu_0$. Using the Cauchy-Schwarz inequality we deduce from (4.16) that for all $\nu \in (0, \nu_0)$:

$$(4.18) \quad \|S_m(\lambda)^* S_m(\lambda)\|_2^2 \leq C \sum_{n, n'} \int_k \int_{k'} \iota_{m, n'}(k', \nu) \iota_{m, n}(k, \nu) |\widehat{v}_\perp(k' - k)|^2 dk' dk$$

and the lemma is proved \square

We notice that the influence of V appears as an interaction between the behaviors in r and z via a convolution product in the phase space. We now estimate the norm of the function $\iota_{m,n}(k, \nu)$:

Lemma 4.5. *There exists $C > 0$ and $\nu_0 > 0$ such that for all $(m, n, k) \in \mathbb{Z} \times \mathbb{N}^* \times \mathbb{R}$, we have*

$$\forall \nu \in (0, \nu_0), \forall q \geq 1, \quad \|\iota_{m,n}(\cdot, \nu)\|_{L^q} \leq C \frac{\nu^{\alpha-1}}{n^\alpha}.$$

Proof. Set $\nu_0 > 0$ and assume $\lambda_{m,n}(k) \leq \nu_0$. According to Lemma 4.1 there exists $C_0 > 0$ such that

$$(4.19) \quad \lambda_{m,n}(k) \geq C_0 n e^{-k},$$

uniformly with respect to $(m, n, k) \in \mathbb{Z} \times \mathbb{N}^* \times \mathbb{R}$. Then for $\nu \in (0, \nu_0)$ there holds $\mathbf{1}_{[0, \nu]}(\lambda_{m,n}(k)) \leq \mathbf{1}_{[0, \frac{\nu}{C_0}]}(n e^{-k})$ and for any $\lambda > 0$ we have

$$\begin{aligned} \|\iota_{m,n}\|_{L^q}^q &= \int_k \frac{\mathbf{1}_{[0, \nu]}(\lambda_{m,n}(k))}{(\lambda_{m,n}(k) + \lambda)^q} e^{-\alpha q k} dk \leq \int_{k \geq \log \frac{C_0 n}{\nu}} \frac{1}{(\lambda_{m,n}(k) + \lambda)^q} e^{-\alpha q k} dk \\ &\leq \frac{1}{(C_0 n)^q} \int_{k \geq \log \frac{C_0 n}{\nu}} e^{(-\alpha+1)qk} dk \\ &= \frac{1}{q(\alpha-1)(C_0 n)^q} \left(\frac{\nu}{C_0 n} \right)^{(\alpha-1)q} \end{aligned}$$

and the lemma is proved. \square

4.4. Convergence of the series and proof of Theorem 1.3. We notice that the r.h.s of (4.14) coincides with

$$\sum_{n,n'} \int_k \iota_{m,n}(k, \nu) (\iota_{m,n'}(\cdot, \nu) * |\widehat{v_\perp}|^2)(k) dk.$$

Assume that $v_\perp \in L^p$ with $p \in [1, 2]$. Then $|\widehat{v_\perp}|^2 \in L^{p'/2}$ with $p' = \frac{p}{p-1} \geq 2$. Young's inequality provides for all $q \geq 1$:

$$\|\iota_{m,n'} * |\widehat{v_\perp}|^2\|_{L^r} \leq \|\iota_{m,n'}\|_{L^q} \|v_\perp\|_{L^p}$$

where $\frac{2}{p'} + \frac{1}{q} = 1 + \frac{1}{r}$. We now use Holder's inequality combined with lemma 4.5 and we get for all (m, n, n') :

$$\forall \nu \in (0, \nu_0), \quad \int_k \iota_{m,n}(k, \nu) (\iota_{m,n'}(\cdot, \nu) * |\widehat{v_\perp}|^2)(k) dk \leq C \|v_\perp\|_{L^p} \frac{\nu^{2\alpha-2}}{n^\alpha n'^\alpha}$$

Since $\alpha > 1$, we get

$$\sum_{n,n'} \int_k \iota_{m,n}(k, \nu) (\iota_{m,n'}(\cdot, \nu) * |\widehat{v_\perp}|^2)(k) dk = O(\nu^{2\alpha-2}) \|v_\perp\|_{L^p} \sum_{n \geq 1} \frac{1}{n^\alpha} \sum_{n' \geq 1} \frac{1}{(n')^\alpha}$$

and therefore using Proposition 4.4:

$$(4.20) \quad \|S_m(\lambda) * S_m(\lambda)\|_2^2 = O(\nu^{2\alpha-2})$$

which, for $\alpha > 1$, tends to 0 with ν , uniformly with respect to $(m, \lambda) \in \mathbb{Z} \times (0, +\infty)$. Then, (4.11) follows from (4.13). In conclusion the hypotheses we have used on $V(r, z)$ is $V(r, z) \leq \langle r \rangle^{-\alpha} v_\perp(z)$ with $\alpha > 1$ and $v_\perp \in L^p(\mathbb{R})$, $p \in [1, 2]$ and we deduce Theorem 1.3.

REFERENCES

- [1] S. AGMON. *Bounds on exponential decay of eigenfunctions of Schrödinger operators*, volume 1159 of *Lecture Notes in Math.* Springer, Berlin 1985.
- [2] J. AVRON, I. HERBST, B. SIMON. Schrödinger operators with magnetic fields. I. General interactions. *Duke Math. J.* **45**(4) (1978) 847–883.
- [3] P. BRIET, H. KOVAŘÍK, G. RAIKOV, E. SOCCORSI. Eigenvalue asymptotics in a twisted waveguide. *Comm. Partial Differential Equations* **34**(7-9) (2009) 818–836.
- [4] P. BRIET, G. RAIKOV, E. SOCCORSI. Spectral properties of a magnetic quantum Hamiltonian on a strip. *Asymptot. Anal.* **58**(3) (2008) 127–155.
- [5] V. BRUNEAU, P. MIRANDA, G. RAIKOV. Discrete spectrum of quantum Hall effect Hamiltonians I. Monotone edge potentials. *J. Spectr. Theory* **1**(3) (2011) 237–272.
- [6] V. BRUNEAU, P. MIRANDA, G. RAIKOV. Dirichlet and neumann eigenvalues for half-plane magnetic hamiltonians. *Submitted* (2013).
- [7] M. DIMASSI, J. SJÖSTRAND. *Spectral asymptotics in the semi-classical limit*, volume 268 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge 1999.
- [8] V. A. GEĬLER, M. M. SENATOROV. The structure of the spectrum of the Schrödinger operator with a magnetic field in a strip, and finite-gap potentials. *Mat. Sb.* **188**(5) (1997) 21–32.
- [9] C. GÉRARD, F. NIER. The Mourre theory for analytically fibered operators. *J. Funct. Anal.* **152**(1) (1998) 202–219.
- [10] V. V. GRUŠIN. Hypoelliptic differential equations and pseudodifferential operators with operator-valued symbols. *Mat. Sb. (N.S.)* **88**(130) (1972) 504–521.
- [11] B. HELFFER. *Semi-classical analysis for the Schrödinger operator and applications*, volume 1336 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin 1988.

- [12] B. HELFFER, J. SJÖSTRAND. Puits multiples en limite semi-classique. II. Interaction moléculaire. Symétries. Perturbation. *Ann. Inst. H. Poincaré Phys. Théor.* **42**(2) (1985) 127–212.
- [13] P. D. HISLOP, N. POPOFF, E. SOCCORSI. Characterization of currents carried by bulk states in one-edge quantum hall systems. *Ongoing work* (2013).
- [14] P. D. HISLOP, E. SOCCORSI. Edge currents for quantum Hall systems. I. One-edge, unbounded geometries. *Rev. Math. Phys.* **20**(1) (2008) 71–115.
- [15] P. D. HISLOP, E. SOCCORSI. Spectral analysis of Iwatsuka “snake” hamiltonians. *Submitted* (2013).
- [16] D. MARTIN. Mélina, bibliothèque de calculs éléments finis. <http://anum-maths.univ-rennes1.fr/melina> (2010).
- [17] N. POPOFF. On the lowest energy of a 3d magnetic laplacian with axisymmetric potential. *Preprint IRMAR* (2013).
- [18] G. D. RAĬKOV. Eigenvalue asymptotics for the Schrödinger operator with homogeneous magnetic potential and decreasing electric potential. I. Behaviour near the essential spectrum tips. *Comm. Partial Differential Equations* **15**(3) (1990) 407–434.
- [19] G. D. RAĬKOV. Eigenvalue asymptotics for the Schrödinger operator with perturbed periodic potential. *Invent. Math.* **110**(1) (1992) 75–93.
- [20] G. D. RAIKOV, S. WARZEL. Quasi-classical versus non-classical spectral asymptotics for magnetic Schrödinger operators with decreasing electric potentials. *Rev. Math. Phys.* **14**(10) (2002) 1051–1072.
- [21] M. REED, B. SIMON. *Methods of modern mathematical physics. IV. Analysis of operators*. Academic Press [Harcourt Brace Jovanovich Publishers], New York 1978.
- [22] B. SIMON. Semiclassical analysis of low lying eigenvalues. I. Nondegenerate minima: asymptotic expansions. *Ann. Inst. H. Poincaré Sect. A (N.S.)* **38**(3) (1983) 295–308.
- [23] A. V. SOBOLEV. Asymptotic behavior of energy levels of a quantum particle in a homogeneous magnetic field perturbed by an attenuating electric field. I. In *Linear and nonlinear partial differential equations. Spectral asymptotic behavior*, volume 9 of *Probl. Mat. Anal.*, pages 67–84. Leningrad. Univ., Leningrad 1984.
- [24] D. YAFAEV. A particle in a magnetic field of an infinite rectilinear current. *Math. Phys. Anal. Geom.* **6**(3) (2003) 219–230.
- [25] D. YAFAEV. On spectral properties of translationally invariant magnetic Schrödinger operators. *Ann. Henri Poincaré* **9**(1) (2008) 181–207.

UNIVERSITÉ BORDEAUX 1, IMB, UMR CNRS 5251, 351 COURS DE LA LIBÉRATION, 33405 TALENCE CEDEX, *E-mail address*: VINCENT.BRUNEAU@U-BORDEAUX1.FR

LABORATOIRE CPT, UMR 7332 DU CNRS, CAMPUS DE LUMINY, 13288 MARSEILLE CEDEX 9, FRANCE, *E-mail address*: NICOLAS.POPOFF@CPT.UNIV-MRS.FR