Wood ash effects on nutrient dynamics and soil properties under Mediterranean climate

María Gómez-Rey, Madeira, Coutinho

► To cite this version:

María Gómez-Rey, Madeira, Coutinho. Wood ash effects on nutrient dynamics and soil properties under Mediterranean climate. *Annals of Forest Science*, 2012, 69 (5), pp.569-579. 10.1007/s13595-011-0175-y . hal-00930746

HAL Id: hal-00930746

<https://hal.science/hal-00930746>

Submitted on 1 Jan 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Wood ash effects on nutrient dynamics and soil properties under Mediterranean climate

María X. Gómez-Rey · Manuel Madeira · João Coutinho

Received: 11 July 2011 / Accepted: 20 December 2011 / Published online: 13 January 2012

© INRA / Springer-Verlag France 2012

Abstract

• **Aims** This study aims to evaluate the effects of wood ash application on nutrient dynamics and soil properties of an acidic forest soil (Arenosol).

• **Methods** Treatments were loose and pelleted ash application (11 Mg ha^{-1}), alone or together with N fertiliser, and control treatment in a lysimeter experiment. Nutrient leaching was followed during a 2-year period and soil chemical and biological properties were evaluated at the end of the experiment.

• **Results** Wood ash increased leaching of total N, $\text{NH}_4^+\text{-N}$, base cations and P, mainly during the first months, the effect being more pronounced for the loose formulation. At the end of the study period, a positive effect on soil nutrient availability and soil acidity reduction was seen. The application of loose and pelleted ash alone decreased N leaching and increased N microbial biomass at the end of the experiment. The C dynamics was weakly affected.

• **Conclusion** Wood ash can be used to improve nutrient availability and balance nutrient exported by tree harvesting

in acid forest soils, the effects at short-term being stronger for loose than for pelleted ash. However, their application should be carried out when vegetation is established to minimise nutrient losses at short-term and reduce the potential risk for water bodies. In N-limited soils, wood ash should be applied with N fertilisers to counteract N immobilisation.

Keywords Forest soils · Carbon · Basal respiration · N mineralisation · Lysimeter

1 Introduction

The amount of forest biomass available in Portugal for energy production is estimated to be $2.2 \times 10^6 \text{ Mg year}^{-1}$, including residues from wood industry and paper pulp plants (MADRP 2005). Large amounts of wood ash are expected to be generated in the near future as a result of 15 new biomass power stations, and alternatives to its deposition in landfills are necessary. Wood ash application in acidic soils, mainly in forest systems, has received growing interest given their alkaline properties and capacity to balance nutrient removal by tree harvesting (Eriksson 1998). Ash application may, therefore, improve the nutrient status of most Portuguese forest soils which are acidic and poor in base cations.

Wood ash is characterised by high alkalinity and high Ca, Mg, K and, to a lesser extent, P contents; in contrast, it is generally poor in N (Khanna et al. 1994). Wood ash application to forest soils can lead to an increment in soil pH and base cations saturation (Bramryd and Fransman 1995; Eriksson 1998; Khanna et al. 1994; Ludwig et al. 2002). However, application of unstabilised highly soluble ash can cause a dramatic increase in soil pH and salt concentration (Clapham and Zibilske 1992), which may disturb ground vegetation

Handling Editor: Erwin Dreyer

Contribution of the co-authors María X. Gómez-Rey: designing and running the experiment, analysing samples and data and writing the manuscript

Manuel Madeira: designing the experiment and supervising
João Coutinho: analysing samples

M. X. Gómez-Rey (✉) · M. Madeira
Centro de Estudos Florestais, Instituto Superior de Agronomia,
Universidade Técnica de Lisboa,
Tapada da Ajuda,
1349-017 Lisbon, Portugal
e-mail: mxgomez@isa.utl.pt

J. Coutinho
Dpto Biologia e Ambiente, Centro de Química,
Universidade de Trás-os-Montes e Alto Douro,
5001-911 Vila Real, Portugal

(Ferm et al. 1992). Moreover, its application in some soils may increase the solubility of organic C and the CO₂ release (Fritze et al. 1994; Weber et al. 1985; Zimmermann and Frey 2002) and may stimulate nitrification and N leaching (Weber et al. 1985). Therefore, to avoid undesirable effects on vegetation and nutrient leaching, stabilisation of wood ash before its application has been encouraged (Steenari et al. 1998). However, the effects of ash application to forest soils may also depend on the respective formulation: loose, crushed hardened, granulated or pelleted ash (Ring et al. 2006; Arvidsson and Lundkvist 2003).

The effects of wood ash on forest ecosystems have mainly been studied in soils of Northern Europe and Northern America (e.g. Ludwig et al. 2002; Park et al. 2005) where environmental conditions favour organic matter accumulation. However, the consequences of ash application in Southern Europe regions, with higher temperature conditions and soils with lower organic C content, have been scarcely studied (Cabral et al. 2008; Perucci et al. 2008; Solla-Gullón et al. 2006) and the effects on C and N dynamics have not been evaluated. Also, the effects of unstabilised and stabilised ashes under Mediterranean conditions have not been compared yet.

Therefore, a study was developed hypothesising that wood ash application in N-limited soils under Mediterranean conditions might lead to changes on soil N and C dynamics. In addition, we hypothesised that the short-term effects on soil properties will be stronger for unstabilised than for stabilised ashes. Thus, our aims were (1) to evaluate the effect of wood ash application on a Mediterranean forest soil before its use on a larger scale and (2) to compare the effects of unstabilised (loose) and stabilised (pelleted) ashes. To evaluate the capability of wood ash to modify the nutrient status of an acidic soil without plant interference and soil spatial variability, a lysimeter experiment was carried out: loose or pelleted wood ash, alone or together with N, was applied to an acidic forest soil and nutrient leaching and soil characteristics were examined over a 2-year period.

2 Material and methods

2.1 Lysimeter station and materials

The study was carried out in a lysimeter station located at the Instituto Superior de Agronomia, Lisbon. The area has a Mediterranean climate, tempered by an oceanic influence. The 30-year mean (1971–2000) rainfall monitored in the adjacent meteorological station of Lisboa/Ajuda was 675 mm, with a drought period usually extending from late May to September; the mean annual temperature was 16.4°C, ranging from a monthly mean of 11.8°C in January to 21.0°C in August (INMG 2007). During the study period (November

2005–November 2007), rainfall was 209, 857 and 411 mm, respectively, in 2005, 2006 and 2007. The rainiest months were October (182 mm) and November (238 mm) 2006.

The lysimeter station consisted of 20 lysimeters constructed in PVC (29.5 cm inner diameter, 50 cm long, 0.068 m² soil area). Each lysimeter had, at its bottom, a drainage system 10 cm thick constituted, from the bottom to the top, by a double plastic mesh (2 mm), a gravel layer washed with deionised water, a double layer of filtering material (Terram) and a washed sand layer. At the top of these layers were placed 35 kg (corresponding to 32.8 kg, 105°C dry weight) of sieved (<5 mm) homogenised mineral soil of the A horizon (about 30 cm depth), packed at a bulk density of about 1.4 g cm⁻³. Over this material was placed 0.5 kg (corresponding to 240.0 g, 45°C dry weight) of forest floor litter, which corresponded to the amount of that measured in old Portuguese pine plantations (Madeira et al. 2009). At the lysimeters base, a hole was made and a plastic tube was connected to a sampling bottle (5 L) where leachates were continuously collected. Sampling bottles were kept in the dark to avoid biological growth in the sample.

The mineral soil and the forest floor litter (L and F layers) used in the experiment were collected in a permanent monitoring plot for inventory purposes from a 67-year-old maritime pine (*Pinus pinaster*) plantation, in the Leiria National Forest (39°45' N; 8°56' W; 75 m a.s.l.), located in central coastal Portugal, with dry sub-humid climate (30-year mean rainfall of about 800 mm). This plantation is representative of wide areas of Portugal, which are characterised by sandy and acidic soils, with low organic matter content and N availability. Soils are mainly Dystric Arenosols associated with Podzols developed on coastal sand dunes. Soil substrate was collected from the Ah horizon of an Arenosol. The soil was sieved (<2 mm), four subsamples were taken for chemical analysis and the rest of the soil was stored at room temperature until being introduced in the lysimeters, within 4 days after collection. The soil showed a sandy texture (98% of sand) and a low pH (5.1). Exchangeable bases (Ca, Mg, Na and K) and effective cation exchange capacity (NH₄OAc extraction) were 1.09 and 1.49 cmol_c kg⁻¹, respectively. Extractable P and K determined by the Egnér–Riehm method (Egnér et al. 1960) were 4.7 and 20.5 mg kg⁻¹, respectively. Organic C content was 12.4 g kg⁻¹ and C/N ratio was high (45).

Twenty samples (0.25 m² each) from the forest floor litter layer were randomly collected in the area (Arenosols) where soil substrate was sampled. Four composite samples were obtained for the analysis of C, N, Ca, Mg, K and P contents that were 384, 7.71, 8.21, 1.41, 1.05 and 0.42 mg g⁻¹, respectively.

The two forms of wood ash used in this study, obtained from combustion of *Eucalyptus* bark in the CELBI Pulp Company, showed a high pH and low concentrations of

heavy metals (Table 1). Loose ash (diameter <0.2 cm) was collected as fly ash. Fly ash was pelleted (diameter=0.5 cm, length=2 cm) by AB Nordströms Konstruktionsbyrå (Sweden).

2.2 Treatments

The experimental design consisted of five treatments replicated four times in a completely randomised design: CT, absence of wood ash and fertiliser application (control treatment); LA, loose ash application; LAN, as LA, but with N fertiliser; PA, pelleted ash application; PAN, as PA, but with N fertiliser. Except in the control treatment, 74.8 g of wood ash was applied on the surface of the forest floor litter of each lysimeter (without being incorporated into the soil), which corresponded to 11 Mg ha⁻¹ (considering 20 cm of depth). The dose of wood ash applied was assessed considering the lime requirement to achieve a soil pH of 6.5 and the mean CaCO₃ equivalents (CCE) of the wood ash. The added amount of wood ash largely exceeded the export of Ca, Mg and K (670, 110 and 251 kg ha⁻¹, respectively) due to whole-tree harvesting of maritime pine in the Leiria National Forest (Madeira et al. 2009). As wood ash supply low amounts of N and P, in the treatments LAN

Table 1 Chemical composition of the wood ash (C, N, Ca, Mg, K, P, Al, Fe and Mn in grammes per kilogramme; other elements in milligrammes per kilogramme) and amounts of nutrients applied (in kilogrammes per hectare) to the lysimeters

Wood ash	Chemical composition		Amounts applied	
	Loose	Pelleted	Loose	Pelleted
pH	12.9	12.6		
CCE (%)	80	83		
C	16.5	14.2		
N	0.08	0.20		
Ca	103.4	117.3	1,129	1,281
Mg	23.94	26.07	261	285
K	32.15	35.41	351	387
P	4.74	2.27	51.8	24.8
Al	4.45	4.41	48.6	48.2
Fe	3.01	2.90	32.9	31.7
Mn	5.40	6.10	59.0	66.6
B	193	250	2,108	2,73
Cd	0.21	0.13	0.002	0.001
Co	3.26	3.75	0.04	0.04
Cr	15.4	17.0	0.17	0.19
Cu	31.0	41.6	0.34	0.45
Ni	21.9	28.1	0.24	0.31
Pb	18.9	16.6	0.21	0.18
Zn	64.7	54.0	0.71	0.59

CCE calcium carbonate equivalent

and PAN, each lysimeter received 2.27 g (corresponding to 334 kg ha⁻¹) of N fertiliser (25%—8% as ammonium, 16% as urea and 1% as dicyandiamide) with low P content (about 2.25%), considering main nutrient limitations of pine plantations in the Leiria National Forest (e.g. N and P; Madeira et al. 2009). The treatment installation was concluded in 7 November 2005, and the study ended 24 months later (November 2007).

2.3 Sampling

The volume of lysimeter leachates was recorded and subsamples were collected for analysis according to the occurrence of rainfall, approximately once a week in the rainy period and once every 2 weeks or once a month in the summer, so that a total of 39 samplings were carried out. Subsamples were filtered (0.45 µm) and stored at -15°C until chemical analyses. At the end of the experiment (December 2007), the lysimeters were destructively sampled to assess soil chemical changes with depth. After removal of the forest floor litter layer remaining on the soil surface, the soil was divided into four depths (0–5, 5–10, 10–20 and 20–30 cm) and samples were dried (45°C) and sieved (<2 mm) for chemical analyses. For the 0–5 and 5–10 cm layers, subsamples of the field-moist soil were stored at 4°C until processing within 3 days after collection to evaluate N mineralisation, soil basal respiration and microbial biomass.

2.4 Laboratory procedures

Organic residues were dried at 45°C and ground in a laboratory mill to a particle size of 0.5 mm. The mineral elements (Ca, Mg, K and P) were determined after ashing (6 h at 450°C) and dissolving in HCl. Total N was determined using Kjeldhal digestion. The C content was measured by wet oxidation (De Leenheer and Van Hove 1958). The Ca, Mg and K were measured by atomic absorption spectrometry (AAS; Aanalyst 300, Perkin Elmer) and P by the molybdate-blue method (Murphy and Riley 1962).

Leachates were filtered (0.45 µm) and pH was measured. Dissolved organic C (DOC) and total N were determined by an elemental analyser (Skalar FORMAC Combustion TOC/N), using a near-infrared spectroscope and a chemiluminescence detector, respectively. Concentrations of NO₃⁻-N and NH₄⁺-N were determined with a segmented flow autoanalyser, using the α-naphthylamine and sulphanilamide method after reduction with Cd and the modified Berthelot method, respectively (Houba et al. 1994). Calcium, Mg and K were measured by AAS and mineral P by the molybdate-blue method.

The pH in soil and wood ash was determined in water (soil or ash/solution ratio, 1:2.5). The C content and total N were measured as mentioned above. The mineral elements

in wood ash were taken up in HNO_3 . The CCE of ash were determined by boiling 1 g of wood ash with 50 ml of 0.5 M HCl, excess acidity being analysed by titrating with 0.25 M NaOH. The soil exchangeable base cations were extracted by the standard method (1 M NH_4OAc , adjusted at pH 7.0; Chapman 1965). To assess the effective cation exchange capacity, the extractable acidity (Al and H) was determined after extraction by 1 M KCl (Thomas 1982). Extractable P and K were obtained by shaking 1 g of soil with 20 ml of K ($\text{SbO})\text{C}_4\text{H}_4\text{O}_6 \cdot \frac{1}{2}\text{H}_2\text{O}$ solution at pH 3.7 for 2 h (Egnér et al. 1960). The Ca, Mg, Na, K and Al of the extracts and solutions were quantified by AAS, P was determined as above and other elements by ICP-OES.

N mineralisation, soil basal respiration and microbial biomass in the 0–5 and 5–10 cm soil layers were determined. To measure N mineralisation, about 200 g of moist soil from each lysimeter were incubated under aerobic conditions in polythene bags, at 25°C, 60% of water-holding capacity and darkness, during 60 days. Initial soil moisture content was adjusted after a 7-day period by adding distilled water. Fortnightly, NO_3^- -N and NH_4^+ -N were extracted by shaking 10 g of soil for 1 h with 2 M KCl (soil/solution ratio, 1:5) and NO_3^- -N and NH_4^+ -N in extracts were determined as above. Net nitrification and net ammonification were calculated by the difference between NO_3^- -N and NH_4^+ -N at each incubation time and their respective initial levels.

Soil basal respiration was measured by incubating 50 g of moist soil adjusted to 60% water-holding capacity at 25°C in a 1.5-L sealed glass jar. The C-CO_2 produced after the 7-, 15-, 30-, 45- and 60-day period was collected in 30 ml of 0.5 M NaOH, which was titrated with 0.5 M HCl to phenolphthalein endpoint, after the addition of 1.5 M BaCl_2 .

Microbial C (C_{mic}) and N (N_{mic}) were measured by the chloroform fumigation–extraction method (Vance et al. 1987) and organic C and total soluble N were determined by elemental analysis, as previously described. Contents of C_{mic} and N_{mic} were calculated as the difference between fumigated and non-fumigated samples, using a K_{EC} and K_{EN} factor of 2.64 and 1.85, respectively (Brookes et al. 1985; Vance et al. 1987). Metabolic quotient ($q\text{CO}_2$) was calculated as the ratio of hourly basal respiration rate produced after 7 days to the corresponding initial C_{mic} [$\text{mg CO}_2\text{-C (g } C_{\text{mic}})^{-1} \text{ h}^{-1}$].

2.5 Statistical analysis

Treatments effects on nutrient leaching and soil chemical and biological properties were tested by one-way ANOVA and differences among means were determined at $p < 0.05$ using the Tukey honestly significant difference test. Statistical procedures were performed using Statgraphics plus 5.1 (Manugistics Inc.).

3 Results

3.1 Nutrient leaching

Lysimeter drainage volume had a similar trend to that of rainfall, with paralleled amounts in both years. Cumulative volume of leachates during the experimental period was similar in all treatments (Table 2). Mean pH of leachates was increased by loose (about 0.25 units) and pelleted (about 0.10 units) ash application relatively to the control treatment, but differences were not significant (Table 2). At the end of the study period, amounts of DOC leached were not significantly affected by treatments (Table 2). Total N leached was significantly lower in treatments with loose and pelleted ash alone (710 and 762 mg/lysimeter in LA and PA treatments, respectively) than in the control. Application of wood ash with N fertiliser promoted significant NH_4^+ -N losses compared with the control, mainly when applied in loose form (Fig. 1, Table 2). Leaching of NO_3^- -N, negligible during the first year of the experiment, was much greater than that of NH_4^+ -N at the end of the study. Losses of NO_3^- -N were mainly affected by the ash form, with a significant reduction only observed for loose ash (treatment LA), and secondarily by N fertiliser application (Fig. 1).

Leaching of K and Ca was faster and more intense for the loose than the pelleted form. Wood ash application resulted in a fast increment of Ca and K leached amounts, and differences among treatments mostly occurred during the first 4 months of study (Fig. 1). Such differences may be also associated with the rainfall amount, as the leaching of Ca and K was also high in the second winter season of the experiment. A similar trend was observed for Mg and Na leached (data not shown). After the first year of the study, loose ash application significantly increased the amounts of Ca, Mg, Na and K leached in relation to the control; for the ash pelleted formulation, such increase was also significant but less pronounced (Table 2). The effect of wood ash decreased with time and, during the second year, only K leaching showed a significant increment in response to wood ash application. At the end of the study, amounts of Na and K leached were significantly higher in treatments with wood ash than in the control (Table 2). As for base cations, P leached during the first year was significantly higher in treatments with loose ash than in the control. Such differences were still noticeable after 2 years (Table 2, Fig. 1). When pelleted ash was added alone (PA treatment), P leached also increased during the first year, but differences were not observed for the cumulative amount leached after 2 years.

3.2 Soil properties

At the end of the experiment, values of pH- H_2O in the 5-cm top soil layer were 0.91 and 0.83 units higher in treatments with loose ash (LA and LAN treatment, respectively) than in

Table 2 Cumulative volume of leachates (in millimetres), mean pH and cumulative DOC and nutrients (in milligrammes per lysimeter) leached after a 1-year (November 2006) and a 2-year (November 2007) period

Treatment	Volume	pH	DOC	Total N	NO ₃ ⁻ -N	NH ₄ ⁺ -N	Ca	Mg	Na	K	P
One-year period											
CT	344±47	4.57	2,390±400	202±55b	54±17	59±6b	147±20b	70±14b	215±50b	183±31c	8±5c
LA	362±42	4.48	2,799±575	217±27ab	27±9	95±18ab	548±198a	241±90a	401±106a	1,302±366a	19±3ab
LAN	359±20	4.64	2,779±267	303±44ab	40±21	148±23a	643±212a	298±80a	426±48a	1,258±215a	20±6a
PA	322±124	4.84	2,433±746	252±79ab	42±16	84±38b	577±60a	233±71a	367±36a	784±169b	19±7abc
PAN	382±21	4.66	2,494±230	341±69a	65±31	155±43a	508±119a	229±46a	323±26a	645±158b	9±5cb
Two-year period											
CT	1,080±44	4.88	4,896±781	968±93a	541±58a	149±55c	801±99	395±24	470±69b	649±179c	24±3b
LA	1,081±46	5.03	6,159±783	710±51b	343±45b	192±14bc	818±209	385±117	692±105a	2,257±332a	36±6a
LAN	1,072±24	5.15	5,766±556	892±105ab	431±91ab	239±25a	939±267	473±104	737±28a	2,311±114a	34±5a
PA	1,122±28	4.94	5,770±545	762±34b	453±46ab	164±33bc	1,012±120	451±79	667±28a	1,641±131b	28±5ab
PAN	1,108±47	4.89	5,301±669	969±122a	499±91a	229±28ab	959±186	449±38	620±48a	1,460±184b	25±7ab

Values are presented as the mean±SD ($n=4$). Treatments are CT, absence of wood ash and fertiliser application; LA, loose ash application; LAN, as LA, but with N fertiliser; PA, pelleted ash application; PAN, as PA, but with N fertiliser. Different letters in the same column and period denote significant differences ($p<0.05$) among treatments by the Tukey multiple range test

the control, while pH differences for pelleted ash were not significant (Fig. 2). Loose ash application also increased exchangeable Ca and Mg (in 0.87 and 0.12 cmol_c kg⁻¹, respectively) and extractable P (in 2.5 mg kg⁻¹) in the 0–5 cm soil layer, when compared with the control. Treatments with loose or pelleted ash (alone or together with N) showed higher contents of exchangeable K for all soil depths than in the control soil.

In the top soil layer of treatments with loose ash, a significant reduction in extractable acidity was observed (0.10–0.18 cmol_c kg⁻¹), while effective cation exchange capacity significantly increased (0.69–1.43 cmol_c kg⁻¹) relatively to the other treatments (Fig. 2). Total N (data not shown) and total C (Fig. 2) content in soil showed a similar pattern and both were not significantly affected by treatments.

After 60 days of aerobic incubation, treatments with loose ash (LA and LAN) showed a significant reduction on cumulative net ammonification when compared with the control (Fig. 3). In contrast, the resulting cumulative net nitrification was significantly increased by loose ash application. Such differences were stronger in the 0–5 cm surface soil layer. No significant effects were observed for the pelleted formulation.

Soil basal respiration and respective rate were not significantly affected by treatments (Table 3), while initial metabolic quotient ($q\text{CO}_2$) values were lower in treatments with wood ash than in the control.

In both 0–5 and 5–10 cm soil layers, soil microbial C and N contents were significantly higher in treatments with loose and pelleted ash alone (LA and PA treatments) than in the control (Table 3). Similarly, the highest $C_{\text{mic}}/C_{\text{org}}$ ratios were generally observed in treatments with wood ash application, but differences were not significant.

4 Discussion

The methodology used in this study presents some limitations (e.g. disturbance of soil structure and water transfers, absence of root uptake and exudates). Soil disturbance during lysimeter installation might have induced a higher mineralisation flux, while the absence of plants probably increased the volume of percolating water and modified the nutrient leaching. Although experimental conditions have a limited value in assessing the complexity of field situations, they give insight into the effect of wood ash application on soil properties and C and N dynamics.

The influence of wood ash on nutrient dynamics and soil properties depended on the form of ash applied. Wood ash did not exhibit a significant effect on the pH of leachates, while loose ash application increased pH and reduced extractable acidity on the top soil layer at the end of the study period. This finding supports the efficiency of unstabilised

Fig. 1 Mean pH of the leachates and cumulative losses (in milligrammes per lysimeter) of Ca, K, NH₄⁺-N, NO₃⁻-N and P from the lysimeters over the experimental period ($n=4$). Different letters for data at the end of the study denote significant differences ($p<0.05$) among

treatments. Treatments are CT absence of wood ash and fertiliser application, LA loose ash application, LAN as LA, but with N fertiliser, PA pelleted ash application, PAN as PA, but with N fertiliser

wood ash to correct acidity of forest mineral soils at the short-term, as reported in several studies (Ludwig et al. 2002; Park et al. 2005). Stabilisation of wood ash by pelleting, granulation or hardening reduces the reactivity of the ash (Eriksson 1998) and explains the absence of appreciable changes after pelleted ash application during the studied period, although a positive effect on soil pH could be expected in the long-term. Also, in a previous laboratory incubation (Gómez-Rey et al. 2006), a significant and fast soil pH increment was initially observed using loose ash at doses similar to that used in this study, but no significant effect was detected even 20 weeks after pelleted ash addition. Although our data do not allow to predict how long this increase would last, loose ash effect on soil acidity reduction might persist or be increased for several years (6–16 years) after its application, as reported by Augusto et al. (2008) in a recent meta-analysis study.

Leaching of base cations and phosphorus following wood ash application mostly occurred during the first year of the study, when loose ash application enhanced Ca, Mg, Na, K and P amounts in leachates. Given the slower dissolution of pelleted ash (Ring et al. 2006), its effect on nutrient leaching during the study period was of less magnitude when compared with the loose formulation, a trend also observed in a laboratory simulation for long-term leaching of untreated and stabilised wood ash (Steenari et al. 1998). However, the effect of pelleted ash on nutrient leaching might be increased during the following years.

At the end of the study period, the amount of K leached increased by wood ash application; this increase accounted for 34% and 68% of the amount of K supplied to lysimeters by ash (pelleted and loose ash, respectively). In contrast, enhancement of Ca, Mg and P amounts in leachates was only 1.6%, 3.3% and 2.0%, respectively, of the quantity supplied

Fig. 2 Values of pH and contents of total C, exchangeable Ca, Mg and K, extractable P, extractable acidity and effective cationic exchange capacity (ECEC), calculated as the sum of extractable acidity (Al and H) and exchangeable cations (Ca, Mg, Na and K) in the mineral

substrate of lysimeters at the end of the study period, according to depth (*n*=4). Significant differences (*p* < 0.05) among treatments in the same depth were marked with different letters. Treatments as in Fig. 1

by both ashes. These differences are explained by the mineralogical speciation of wood ash in which K occurs in salt forms easily dissolved (Steenari et al. 1999), while Ca and Mg are mainly in carbonate forms and P is bound in compounds with low solubility (Steenari and Lindqvist 1997).

Our results show that loose ash recycling as soil amendment may have a great potential to balance base cations and phosphorus losses due to the harvest of Portuguese eucalypt and pine plantations (Cortez 1996; Madeira et al. 2009). In fact, an enhancement of exchangeable Ca, Mg and K and extractable P was observed, at the end of the study, in the top soil layer (0–5 cm) due to loose ash application,

accounting for low proportions of Ca, Mg, K and P (11%, 4%, 8% and 3%, respectively) initially applied as mineral soil, forest floor, wood ash and fertiliser. This was associated with the displacement of Al³⁺ and H⁺ from the exchangeable complex, which is in conformity with the rise of pH values, especially in the LA and LAN treatments (Fig. 2). However, the increment of base cations in the exchangeable complex was much greater than the reduction of extractable acidity. Given the similar organic C content in all treatments (Fig. 2), such difference may be mostly associated with the enhancement of exchange sites (variable negative charges) in soil colloids (mostly organic matter) as a result of the pH

Fig. 3 Cumulative soil net ammonification and net nitrification during 60 days of aerobic incubation. Significant differences (*p* < 0.05) among means (*n*=4) in the same period were marked with different letters. Treatments as in Fig. 1

Table 3 Cumulative soil basal respiration (in milligrammes of C-CO₂ per kilogramme of soil) during 60 days of incubation, soil basal respiration rate (SBR, in milligrammes of C-CO₂ per gramme of C), metabolic quotient ($q\text{CO}_2$, in milligrammes of C-CO₂ per gramme of C_{mic} per hour) and contents of microbial C and N (C_{mic} and N_{mic}, in milligrammes per kilogramme) for mineral substrate at 0–5 and 5–10 cm depth

Treatment	Basal respiration	SBR	$q\text{CO}_2$	C _{mic}	C _{mic} /C _{org}	N _{mic}
0–5 cm						
CT	150.8±21.5	14.2±3.2	5.8±1.2 ^a	52.6±4.7c	0.18±0.05b	3.7±0.6b
LA	163.8±28.1	18.2±3.7	2.3±0.3bc	93.5±3.6ab	0.25±0.02ab	8.2±2.4a
LAN	171.2±35.0	21.4±3.9	3.2±1.2b	61.6±2.6bc	0.16±0.04b	5.7±0.9bc
PA	122.2±19.9	14.8±2.4	1.1±0.3c	118.2±14.9a	0.35±0.09a	7.6±0.1ac
PAN	138.4±35.1	19.5±4.6	3.4±0.8b	89.7±23.8b	0.24±0.07ab	4.4±1.1b
5–10 cm						
CT	114.3±19.6	11.3±3.9	3.4±0.7 ^a	69.2±17.1bc	0.25±0.05a	5.6±0.9b
LA	91.7±18.0	9.1±2.5	1.3±0.5b	152.7±49.9a	0.33±0.09a	11.1±1.8a
LAN	64.5±9.1	7.9±1.9	3.0±0.8a	36.7±11.6c	0.10±0.04b	9.3±0.3ab
PA	107.4±29.5	13.9±5.2	1.2±0.4b	127.4±41.6a	0.33±0.10a	9.9±2.5a
PAN	77.4±19.0	9.7±3.9	2.6±0.6a	76.3±47.6b	0.23±0.08ab	9.2±2.2ab

Values are presented as the mean±SD ($n=4$). Treatments as in the Table 2. Different letters in the same column and soil layer denote significant differences ($p<0.05$) among treatments by the Tukey multiple range test

increment, as reported by Khanna et al. (1994) for effects of ash derived from *Eucalyptus* litter on forest soils. The absence of significant differences between the pelleted ash treatments (PA and PAN) and the control for extractable base cations, extractable acidity and pH may be a result of the smaller amount of base cations released by the pelleted ash. Similar limited effect on soil nutrient and base cation status was reported by Wang et al. (2010) for granulated wood ash in the short-term (5 years). However, as concluded by these authors, pelleted ashes will likely have positive effects with time and can be used for long-term nutrient compensation, following the findings of Arvidsson and Lundkvist (2003) for hardened crushed ash and those of Callesen et al. (2007) for granulated wood ash.

As observed in other studies (Park et al. 2005), soil enrichment in Ca and Mg was only observed at the upper soil layer while increments in exchangeable K were detected in deeper layers, possibly due to its higher mobility. When pelleted ash was applied, only exchangeable K increments were detected. In contrast, exchangeable Ca and Mg tended to decrease in treatments with pelleted ash relatively to the control. This trend may be ascribed to the lower solubility of Ca and Mg than K in pelleted ash, which may favour their displacement from the exchange sites. We also emphasise that the restriction of Ca and Mg changes mostly to the 0–5 cm soil layer may be associated with the fact that ashes were applied directly on the surface of the forest floor litter (not incorporated), which may reduce effects on the soil exchangeable complex, as reported by Khanna et al. (1994).

As for base cations and phosphorus, an initial increase of total N and NH_4^+ -N leaching was observed during the first

year after wood ash application, mainly when added together with N fertiliser. This suggests that wood ash application to forest plantations growing on sandy soils may affect the quality of water bodies due to the increase in N and P leaching. However, root uptake (trees and understory) was not taken into account in the present study and, therefore, losses may be lower under field conditions. In addition, the dose of N applied in the present study was quite high (83.5 kg ha^{-1}) and, therefore, a reduced risk is expected for lower doses.

At the end of the study period, total N leaching was reduced by the application of loose and pelleted ash alone likely due to the low soil N availability of the studied soil. In contrast to some sites of the Northern European forests, the studied site is N limited due to the low N input from the atmosphere ($4.0 \text{ kg ha}^{-1} \text{ year}^{-1}$; Cortez 1996), low soil organic matter and high C/N ratio of the forest floor layer (50) and of the mineral soil (41), which is above the critical condition for N release (Harmsen and Van Shreven 1955). This pattern suggests that N may have been immobilised in the microbial biomass, which agrees with the higher N microbial contents observed at the end of the study in treatments with wood ash alone (Table 3). In fact, the mean increment on microbial N content observed for the LA or PA treatments (5.0 and 4.1 mg kg^{-1} , respectively), estimated for the whole lysimeter (175 and 143 mg/lysimeter), is relatively close to the reduction on total N leached obtained (258 and 206 mg/lysimeter , respectively). Thus, our results suggest that ash addition may not enhance tree growth of stands established in the studied soils, as also reported by Augusto et al. (2008) for N-limited mineral soils. In fact,

soil N immobilisation following wood ash application may reduce N availability for trees, decreasing tree growth, as suggested by Bramryd and Fransson (1995) for Scots pine growing in poor Scandinavian soils ($C/N > 30$) after liming. Also, Jacobson (2003) reported a stem growth decrease in response to wood ash application at poor Scandinavian coniferous stands, while stem growth increased in more fertile sites. The negligible effect on N leaching observed in treatments where wood ash was applied with N fertiliser suggests that, in N-limited soils as those of the present study, N fertiliser supply is required to promote soil N availability.

In contrast with total N and NO_3^- -N, loose ash application enhanced P leaching. No data on microbial P (P_{mic}) contents were obtained. As the C_{mic}/N_{mic} ratio in our study (8–15) is close to that reported by Achat et al. (2010) for comparable systems, the N_{mic}/P_{mic} ratio reported by those authors (0.6) may be used for the present study. Then, P immobilisation might have increased to about 290 mg/lysimeter. Despite this possible increase of P in microbial biomass, leaching of P was not prevented, suggesting that the soil used in the present study did not avoid P leaching, agreeing with Achat et al. (2009) who measured negligible P buffer capacity in similar soils. As organic P in leachates was not measured, the amount associated with organic matter is unknown. Therefore, further research should be developed to assess deeper understanding on the effects of ash application on P leaching.

At the end of the study, soil incubation under laboratory conditions showed that loose ash application increased net nitrification, agreeing with the trend previously reported for laboratory incubations carried out with soil taken in the same site and with similar ash dose (Gómez-Rey et al. 2006). These results suggest that high doses of untreated ash may favour NO_3^- -N losses such as reported for other forest soils (with soil C/N ratio lower than 35) from regions with cold climate (Ludwig et al. 2002). However, such treatment differences were not observed under the studied lysimetric conditions, as NO_3^- -N amounts in leachates were significantly reduced by loose ash and were not significantly affected by pelleted ash. Discrepancy between both studies may be partially associated with more intense denitrification in the lysimeter system. The increased soil pH may also favour losses of soil N by denitrification as a result of nitrifying microbial populations and nitrification process stimulation (Khanna et al. 1994). The application of N fertiliser combined with wood ash might induce additional N losses by denitrification, following the trend reported by Ozolincius et al. (2006) for an Arenosol amended with wood ash.

Carbon dynamics was weakly affected as DOC in leachates, and soil basal respiration was not significantly different between treatments. This trend does not support the findings of other studies carried out in peat and mineral forest soils in

cold climate conditions where wood ash application promoted increases in DOC (Weber et al. 1985) and soil respiration (Fritze et al. 1994; Khanna et al. 1994; Nohrstedt 2001; Zimmermann and Frey 2002). Also, soil respiration stimulation was observed in ash-amended soils of humid areas of the Iberian Peninsula (Solla-Gullón et al. 2006). The lack of significant effects observed in our study may be related to the low soil organic C content. In fact, by incubating a range of Australian forest soils, Khanna et al. (1994) reported that effects of wood ash addition on organic C dynamics decreased in soils with low organic matter content.

Values of microbial metabolic quotient (qCO_2) were comparable to those reported for Mediterranean soils with similar C content (García et al. 1994). Microbial C increment by wood ash application led to a lower qCO_2 , indicating that use efficiency of substrate C by microorganisms may be higher in amended soils, as reported by Insam and Haselwandter (1989). This pattern does not agree with the higher (Fritze et al. 1994) or similar (Zimmermann and Frey 2002) levels in qCO_2 found in cold regions, as well as with the lack of changes reported for wetter areas of the Iberian Peninsula (Solla-Gullón et al. 2006). Our results suggest that wood ash application could create favourable conditions (e.g. better nutrient availability and soil pH) for soil microbial community in acidic soils of dry sub-humid Mediterranean regions.

At the end of the study, low proportions of Ca, Mg and P (11.9%, 6.0% and 5.8%, respectively) initially applied with the loose ash were measured in leachates and mineral soil layers; these proportions were lower when wood ash was applied in pelleted form. This result suggests that nutrients supplied by ash were mostly retained in the organic layer covering the mineral soil, minimising ash effects on mineral soil layers. This is supported by the fact that dissolution of ash components is expected to be slower when ash is applied to the soil surface than when it is mixed with acidic soils (Khanna et al. 1994). Therefore, greater doses of wood ash, mainly in pelleted form, could be applied in forest systems of sub-humid Mediterranean regions without dramatic changes on soil properties, suggesting that the studied soils may be used for ash disposal. Nevertheless, our results indicate that such application should be carried out some years after tree planting or when understory vegetation is established, mainly in forest plantations growing in sandy soils with low effective cation exchange capacity. Under these circumstances, roots demand would minimise nutrient losses at the short-term, reducing water quality risks and contributing to enhance soil nutrients availability in the long-term. Also, the possible negative impacts of wood ash on seedlings are avoided. However, further studies are needed to evaluate wood ash effects on C and N dynamics in organic soil layers as well as the interactions between these

layers and nutrients released from the ash. Likewise, the interactions with plants will be also evaluated through appropriate field trials to assess how the improvement of soil nutrient availability promoted by wood ash application could affect the nutrient status and growth of the tree stands.

5 Conclusions

Our results showed that, applied at a dose to achieve a soil pH of 6.5, loose ash improved nutrient availability and reduced soil acidity, resulting in an alternative to lime and commercial fertilisers for counteracting nutrient removal by tree harvesting in acidic forest soils of sub-humid Mediterranean regions. Such improvement of base cations availability is mostly associated with the enhancement of the soil effective cation exchange capacity. However, C dynamics was not significantly affected by wood ash application. On the contrary, wood ash should be applied together with N fertiliser to balance soil N immobilisation in N-limited soils as those of the present study. At the short-term, pelleted ash application showed negligible effects, suggesting that it may be used in higher doses, but its effect on Ca and Mg losses from the soil mineral layers should be taken into account. In fact, the increment on nutrient leaching at the short-term suggests that both forms of wood ash show potential risks for water contamination. Our results are consistent with the conclusions of former Nordic experiments, suggesting that guidelines proposed for Northern Europe could be used for Mediterranean regions. However, field studies are necessary to evaluate the long-term effects on soil and plants and to determine the highest dose that can be applied without deleterious effects on soil pH and C and N losses.

Acknowledgements The CELBI Pulp Company is acknowledged for providing the wood ash for the experiment. Laboratory staff of the Departamento de Ciências do Ambiente (Instituto Superior de Agronomia) is acknowledged for their assistance in the analytical work and Luis Hilário for his help in the field work.

Funding The participation of the first author in this research was supported by a post-doctoral grant from the Portuguese Science and Technology Foundation and by the Spanish Foundation “Alfonso Martin Escudero”.

References

- Achat DL, Bakker MR, Augusto L, Saur E, Dousseron L, Morel C (2009) Evaluation of the phosphorus status of P-deficient podzols in temperate pine stands: combining isotopic dilution and extraction methods. *Biogeochem* 92:183–200. doi:10.1007/s10533-008-9283-7
- Achat DL, Bakker MR, Zeller B, Pellerin S, Bienaime S, Morel C (2010) Long-term organic phosphorus mineralization in Spodosols under forests and its relation to carbon and nitrogen mineralization. *Soil Biol Biochem* 2:1479–1490. doi:10.1016/j.soilbio.2010.05.020
- Arvidsson H, Lundkvist H (2003) Effects of crushed wood ash on soil chemistry in young Norway spruce stands. *Forest Ecol Manag* 176:121–132. doi:10.1016/S0378-1127(02)00278-5
- Augusto L, Bakker MR, Meredieu C (2008) Wood ash applications to temperate forest ecosystem—potential benefits and drawbacks. *Plant Soil* 306:181–198. doi:10.1007/s11104-008-9570-z
- Bramryd T, Fransman B (1995) Silvicultural use of wood ashes—effects on the nutrient and heavy metal balance in a pine (*Pinus sylvestris*, L) forest soil. *Water Air Soil Poll* 85:1039–1044. doi:10.1007/BF00476967
- Brookes PC, Landman A, Pruden G, Jenkinson DS (1985) Chloroform fumigation and the release of soil nitrogen: a rapid direct extraction method to measure microbial biomass nitrogen in soil. *Soil Biol Biochem* 17:837–842. doi:10.1016/0038-0717(85)90144-0
- Cabral F, Ribeiro HM, Hilário L, Machado L, Vasconcelos E (2008) Use of pulp mill inorganic wastes as alternative liming materials. *Biores Technol* 99:8294–8298. doi:10.1016/j.biortech.2008.03.001
- Callesen I, Ingerslev M, Raulund-Rasmussen K (2007) Dissolution of granulated wood ash examined by in situ incubation: effects of tree species and soil type. *Biomass Bioenergy* 31:693–699. doi:10.1016/j.biombioe.2007.06.014
- Chapman HD (1965) Cation-exchange capacity. In: Black CA (ed) *Methods of soil analysis—chemical and microbiological properties*. Agronomy 9:891–901
- Clapham WM, Zibilske LM (1992) Wood-ash as a liming amendment. *Comm Soil Sci Plant Anal* 23:1209–1227. doi:10.1080/00103629209368661
- Cortez NRS (1996) Compartimentos e ciclos de nutrientes em plantações de *Eucalyptus globulus* Labill. ssp. *globulus* e *Pinus pinaster* Aiton. Ph.D. dissertation, Universidade Técnica de Lisboa
- De Leenheer L, Van Hove J (1958) Détermination de la teneur en carbone organique des sols. Étude critique des méthodes titrimétriques. *Pédologie* 8:39–77
- Egnér H, Riehm H, Domingo WR (1960) Untersuchungen über die chemische Bodenanalyse als Grundlage für die Beurteilung des Nährstoff-zustandes der Böden. II Chemische Extraktion Method zur Phosphor-und Kaliumbestimmung. *Kungl Lantbr Högsk Ann* 26:199
- Eriksson HM (1998) Short-term effects of granulated wood ash on forest soil chemistry in SW and NE Sweden. *Scand J Forest Res Suppl* 2:43–55
- Ferm A, Hokkanen T, Moilanen M, Issakainen J (1992) Effects of wood bark ash on the growth and nutrition of a Scots pine afforestation in central Finland. *Plant Soil* 147:305–316. doi:10.1007/BF00029082
- Fritze H, Smolander A, Levula T, Kitunen V, Mälkönen E (1994) Wood ash fertilization and fire treatments in a Scots pine forest stand: effects on the organic layer, microbial biomass, and microbial activity. *Biol Fertil Soils* 17:57–63. doi:10.1007/BF00418673
- Garcia C, Hernandez T, Costa F (1994) Microbial activity in soils under mediterranean environmental conditions. *Soil Biol Biochem* 26:1185–1191. doi:10.1016/0038-0717(94)90142-2
- Gómez-Rey MX, Vasconcelos E, Madeira M (2006) Efeito da aplicação de cinza de biomassa florestal na dinâmica do N e nas características químicas do solo de uma plantação de *Pinus pinaster* de Portugal. In: Gallardo Lancho JF (ed) *Medioambiente en Iberoamérica. Visión desde la Física y la Química en los albores del Siglo XXI*, vol 3. Diputación de Badajoz, Badajoz, pp 165–172
- Harmsen GW, Van Shreven DA (1955) Mineralization of organic nitrogen in the soil. *Adv Agron* 7:299–398
- Houba VJS, Novozamsky I, Termminghoff E (1994) Soil analysis procedures. Wageningen Agricultural University, Netherlands

- INMG (2007) Normais climáticas da Tapada da Ajuda, 1971/2000. Available at http://www.isa.utl.pt/dca/Clim_Rec_Hidr/index.html. Accessed 25 November 2009
- Insam H, Haselwandter K (1989) Metabolic quotient of the soil microflora in relation to plant succession. *Oecologia* 79:174–178. doi:10.1007/BF00388474
- Jacobson S (2003) Addition of stabilized wood ashes to Swedish coniferous stands on mineral soils: effects on stem growth and needle nutrient concentrations. *Silva Fenn* 37:437–450
- Khanna PK, Raison RJ, Falkner RA (1994) Chemical properties of ash derived from *Eucalyptus* litter and its effects on forest soils. *Forest Ecol Manag* 66:107–125. doi:10.1016/0378-1127(94)90151-1
- Ludwig B, Rumpf S, Mindrup M, Meiwes KJ, Khanna PK (2002) Effects of lime and wood ash on soil-solution chemistry, soil chemistry and nutritional status of a pine stand in Northern Germany. *Scand J Forest Res* 17:225–237. doi:10.1080/028275802753742891
- Madeira M, Fabião A, Páscoa F, Magalhães MC, Cameira MC, Ribeiro C (2009) Carbono e nutrientes na biomassa aérea arbórea, vegetação sob coberto e solo numa cronossequência de povoamentos de pinhal bravo. *Rev Ciências Agrárias* 32:154–170
- MADRP (2005) Biomassa e Energias Renováveis na Agricultura, Pescas e Florestas. Ministério de Agricultura, do Desenvolvimento Rural e das Pescas, Lisbon
- Murphy J, Riley JP (1962) Modified single solution method for the determination of phosphate in natural waters. *Anal Chim Acta* 27:31–36
- Nohrstedt HÖ (2001) Response of coniferous forest ecosystems on mineral soils to nutrient additions: a review of Swedish experiences. *Scand J For Res* 16:555–573
- Ozolincius R, Armolaitis K, Raguotis A, Varnagiryte I, Zenkovaite J (2006) Influence of wood ash recycling on chemical and biological condition of forest Arenosols. *J For Sci* 52:79–86
- Park BB, Yanai RD, Sahm M, Lee DK, Abrahamson LP (2005) Wood ash effects on plant and soil in a willow bioenergy plantation. *Biomass Bioenergy* 28:355–365. doi:10.1016/j.biombioe.2004.09.001
- Perucci P, Monaci E, Onofri A, Vischetti C, Casucci C (2008) Changes in physico-chemical and biochemical parameters of soil following addition of wood ash: a field experiment. *Eur J Agron* 28:155–161. doi:10.1016/j.eja.2007.06.005
- Ring E, Jacobson S, Nohrstedt HÖ (2006) Soil-solution chemistry in a coniferous stand after adding wood ash and nitrogen. *Can J For Res* 36:153–163. doi:10.1139/X05-242
- Solla-Gullón F, Santalla M, Rodríguez-Soalleiro RJ, Merino A (2006) Nutritional status and growth of a young *Pseudotsuga menziesii* plantation in a temperate region after application of wood-bark ash. *Forest Ecol Manage* 237:312–321. doi:10.1016/j.foreco.2006.09.054
- Steenari BM, Lindqvist O (1997) Stabilisation of biofuel ashes for recycling to forest soil. *Biomass Bioenergy* 13:39–50. doi:10.1016/S0961-9534(97)00024-X
- Steenari B, Marsic N, Karsson L, Tomsic A, Linsdqvist O (1998) Long-term leaching of stabilized wood ash. *Scand J Forest Res Suppl* 2:3–16
- Steenari BM, Karlsson LG, Lindqvist O (1999) Evaluation of the leaching characteristics of wood ash and the influence of ash agglomeration. *Biomass Bioenergy* 16:119–136. doi:10.1016/S0961-9534(98)00070-1
- Thomas GW (1982) Exchangeable cations. In: Page A, Miller R, Keeney D (eds) *Methods of soil analysis, part 2. Chemical and microbiological properties*. ASA, Madison, pp 159–165
- Vance PC, Brookes PC, Jenkinson DS (1987) An extraction method for measuring soil microbial biomass C. *Soil Biol Biochem* 19:703–707
- Wang P, Olsson BA, Arvidsson H, Lundkvist H (2010) Short-term effects of nutrient compensation following whole-tree harvesting on soil and soil water chemistry in a young Norway spruce stand. *Plant Soil* 336:323–336. doi:10.1007/s11104-010-0484-1
- Weber A, Karsisto M, Leppknen R, Sundman V, Skujins J (1985) Microbial activities in a histosol: effects of wood ash and NPK fertilizers. *Soil Biol Biochem* 17:291–296. doi:10.1016/0038-0717(85)90063-X
- Zimmermann S, Frey B (2002) Soil respiration and microbial properties in an acid forest soil: effects of wood ash. *Soil Biol Biochem* 34:1727–1737. doi:10.1016/S0038-0717(02)00160-8