Cyclotomic Units and Class Groups in $\mathbb{Z}_p$ -extensions of real abelian number fields

Filippo Alberto Edoardo Nuccio Mortarino Majno Di Capriglio

► To cite this version:

Filippo Alberto Edoardo Nuccio Mortarino Majno Di Capriglio. Cyclotomic Units and Class Groups in $\mathbb{Z}_p$ -extensions of real abelian number fields. Mathematical Proceedings of the Cambridge Philosophical Society, 2009, 148, pp.93-106. 10.1017/S0305004109990119 . hal-00947151

HAL Id: hal-00947151

<https://hal.science/hal-00947151>

Submitted on 21 Feb 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Cyclotomic units and class groups in $\mathbb{Z}_p$ -extensions of real abelian fields

BY FILIPPO ALBERTO EDOARDO NUCCIO

*Università “La Sapienza”,
P.le Aldo Moro, 5-00186, Rome, Italy.
e-mail: nuccio@mat.uniroma1.it*

(Received 9 December 2008; revised 24 April 2009)

Abstract

For a real abelian number field F and for a prime p we study the relation between the p -parts of the class groups and of the quotients of global units modulo cyclotomic units along the cyclotomic $\mathbb{Z}_p$ -extension of F . Assuming Greenberg’s conjecture about the vanishing of the λ -invariant of the extension, a map between these groups has been constructed by several authors, and shown to be an isomorphism if p does not split in F . We focus in the split case, showing that there are, in general, non-trivial kernels and cokernels.

1. Introduction

Let $F/\mathbb{Q}$ be a real abelian field of conductor f and let Cl_F be its ideal class group. A beautiful formula for the order of this class group comes from the group of cyclotomic units: this is a subgroup of the global units $\mathcal{O}_F^\times$ whose index is linked to the order of Cl_F . To be precise, we give the following definition ([Sin81, section 4]):

Definition 1.1. For integers $n > 1$ and a not divisible by n , let ζ_n be a primitive n th root of unity. Then $Norm_{F \cap \mathbb{Q}(\zeta_n)}^{\mathbb{Q}(\zeta_n)}(1 - \zeta_n^a) \in F$ and we define the cyclotomic numbers D_F to be the subgroup of $F^\times$ generated by -1 and $Norm_{F \cap \mathbb{Q}(\zeta_n)}^{\mathbb{Q}(\zeta_n)}(1 - \zeta_n^a)$ for all $n > 1$ and all a not divisible by n . Then we define the cyclotomic units of F to be

$$Cyc_F := D_F \cap \mathcal{O}_F^\times.$$

Sinnott proved in [Sin81, theorem 4.1, with proposition 5.1], the following theorem:

THEOREM (Sinnott). *There exists an explicit constant κ_F divisible only by 2 and by primes dividing $[F : \mathbb{Q}]$ such that*

$$[\mathcal{O}_F^\times : Cyc_F] = \kappa_F |Cl_F|.$$

Let now p be an odd prime that does not divide $[F : \mathbb{Q}]$: by tensoring $\mathcal{O}_F^\times$, Cyc_F and Cl_F with $\mathbb{Z}_p$ we get an equality

$$[\mathcal{O}_F^\times \otimes \mathbb{Z}_p : Cyc_F \otimes \mathbb{Z}_p] = |Cl_F \otimes \mathbb{Z}_p|$$

and it is natural to ask for an algebraic interpretation of this. Moreover, observe that our assumption $p \nmid [F : \mathbb{Q}]$ makes the Galois group $\Delta := \text{Gal}(F/\mathbb{Q})$ act on the modules appearing above through one-dimensional characters, and we can decompose them accordingly: in the sequel we write $M(\chi)$ for every $\mathbb{Z}[\Delta]$ -module M to mean the submodule of

$M \otimes \mathbb{Z}_p[\text{Im}(\chi)]$ on which Δ acts as χ , where $\chi \in \hat{\Delta}$ (see the end of Section 2 for a precise discussion). Then an even more optimistic question is to hope for a character-by-character version of Sinnott's theorem, namely

$$[\mathcal{O}_F^\times(\chi) : \text{Cyc}_F(\chi)] \stackrel{?}{=} |Cl_F(\chi)| \quad (1.1)$$

and then ask for an algebraic interpretation of this. Although it is easy to see that these Δ -modules are in general not isomorphic (see the example on page 143 of [KS95]), it can be shown that they sit in an exact sequence for a wide class of fields arising in classical Iwasawa theory. More precisely, let F_∞/F be the cyclotomic $\mathbb{Z}_p$ -extension of F and let $\Gamma = \text{Gal}(F_\infty/F) \cong \mathbb{Z}_p$; then

$$F_\infty = \bigcup_{n \geq 0} F_n \supset \cdots \supset F_n \supset F_{n-1} \supset \cdots \supset F_0 = F$$

where F_n/F is a cyclic extension of degree p^n whose Galois group is isomorphic to Γ/Γ^{p^n} . In a celebrated work (see [Iwa73]) Iwasawa gives a formula for the growth of the order of $Cl_{F_n} \otimes \mathbb{Z}_p$: he proves that there are three integers μ , λ and ν , and an index $n_0 \geq 0$, such that

$$|Cl_{F_n} \otimes \mathbb{Z}_p| = p^{\mu p^n + \lambda n + \nu} \quad \text{for every } n \geq n_0.$$

Moreover, Ferrero and Washington proved in [FW79] that the invariant μ vanishes. A long-standing conjecture by Greenberg (see [Gre76], where conditions for this vanishing are studied) predicts that $\lambda = 0$: according to the conjecture the p -part of the class groups should stay bounded in the tower.

Although a proof of this conjecture has never been provided, many computational checks have been performed verifying the conjecture in many cases (see, for instance, [KS95]). Under the assumptions $\lambda = 0$ and $\chi(p) \neq 1$, *i. e.* p does not split in F , some authors (see [BNQD01], [KS95], [Kuz96] and [Oza97]) were able to construct an explicit isomorphism

$$\alpha : Cl_{F_n}(\chi) \cong (\mathcal{O}_{F_n}^\times / \text{Cyc}_{F_n})(\chi) \quad (1.2)$$

if n is big enough. Although the construction of the above morphism works also in the case $\chi(p) = 1$, as detailed in the beginning of Section 5, the split case seems to have never been addressed. We focus then on this case, and study the map in this context, still calling it α . Our main result is the following (see Corollary 5.2):

THEOREM. *With notations as above, assume that χ is a non-trivial character of Δ such that $\chi(p) = 1$ and that $\lambda = 0$. Then, for sufficiently big n , there is an exact sequence*

$$0 \longrightarrow K \longrightarrow Cl_{F_n}(\chi) \xrightarrow{\alpha} (\mathcal{O}_{F_n}^\times / \text{Cyc}_{F_n})(\chi) \longrightarrow C \longrightarrow 0:$$

both the kernel K and the cokernel C of α are finite groups with trivial Γ -action isomorphic to $\mathbb{Z}_p[\text{Im}(\chi)]/L_p(1, \chi)$ where $L_p(s, \chi)$ is the Kubota–Leopoldt p -adic L -function.

2. Some Tate cohomology

In this section we briefly recall some well-known facts that are useful in the sequel. Throughout, L/K is a cyclic extension of number fields, whose Galois group we denote by G . In our application, K and L will usually be layers F_m and F_n of the cyclotomic $\mathbb{Z}_p$ -extension for some $n \geq m$, but we prefer here not to restrict to this special case.

We need to introduce some notation. Let

$$\mathbb{U}_K = \prod_{v \nmid \infty} \mathcal{O}_{K,v}^\times \times \prod_{v \mid \infty} K_v^\times$$

be the idèle units, *i. e.* idèles having valuation 0 at all finite place v (we refer the reader to sections 14 – 19 of Cassels' paper in [CF86] for basic properties of idèles and idèle class group) and let Σ be the set of places of K that ramify in L/K . It is known (see [CF86, Serre, section 1.4]) that the Tate cohomology of local units in an unramified extension of local fields is trivial: therefore Shapiro's Lemma ([CF86, Tate, proposition 7.2]) gives

$$\hat{H}^q(G, \mathbb{U}_L) = \hat{H}^q\left(G, \prod_{v \in \Sigma} \prod_{w|v} \mathcal{O}_{L,w}^\times\right) \cong \prod_{v \in \Sigma} \hat{H}^q(G_v, \mathcal{O}_{L,w}^\times),$$

where we fix a choice of a place w of L above v for every $v \in \Sigma$ and we denote by G_v its decomposition group in G ; we will make this identification throughout. We denote the product of local units at places $v \in \Sigma$ appearing above by U_Σ . Consider the following commutative diagram of G -modules:

$$\begin{array}{ccccccc} & & 0 & & 0 & & 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \mathcal{O}_L^\times & \longrightarrow & L^\times & \longrightarrow & \text{Pr}_L \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \mathbb{U}_L & \longrightarrow & \mathbb{A}_L^\times & \longrightarrow & \text{Id}_L \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \mathcal{Q}_L & \longrightarrow & C_L & \longrightarrow & Cl_L \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ & & 0 & & 0 & & 0 \end{array} \quad (2.1)$$

Here Id_L and Pr_L denote the group of all fractional ideals of L and of principal ideals, respectively; while $C_L = \mathbb{A}_L^\times / L^\times$ is the group of idèle classes and $\mathcal{Q}_L = \mathbb{U}_L / \mathcal{O}_L^\times$.

LEMMA 2.1. *Taking Tate cohomology of (2.1) and exploiting the fact that G is cyclic, we find the diagram:*

$$\begin{array}{ccc} & \hat{H}^{-1}(G, \mathbb{U}_L) & \\ & \downarrow \beta & \\ \hat{H}^0(G, Cl_L) & \xrightarrow{\delta} & \hat{H}^{-1}(G, \mathcal{Q}_L) \end{array}$$

Then $\text{Im}(\beta)$ is generated by the image via δ of classes $[\text{Norm}_{G/G_v}(\mathcal{P}_v)]$ for $\mathcal{P}_v$ running through all primes in L above Σ .

Proof. First of all, the above decomposition

$$\hat{H}^{-1}(G, \mathbb{U}_L) \cong \prod_{v \in \Sigma} \hat{H}^{-1}(G_v, \mathcal{O}_{L,w}^\times)$$

allows to write $\beta = \prod \beta_v$ where β_v is the restriction of β to the v th factor $\hat{H}^{-1}(G_v, \mathcal{O}_{L,w}^\times)$.

We therefore fix a place $v \in \Sigma$ and we show that $\text{Im}(\beta_v)$ is the class of $\delta(\sum_{\sigma \in G/G_v} \sigma \mathfrak{P}_v)$ for some prime $\mathfrak{P}_v$ of L above v ; observe that the sum (written by a small abuse of notation, since we are summing over *representatives* of G/G_v in G – but their choice is harmless) makes sense, since $\mathfrak{P}_v$ is fixed by G_v . Choosing a uniformizer π_w at $\mathfrak{P}_w$ we get a generator $(\tau - 1)\pi_w$ of $\hat{H}^{-1}(G_v, \mathcal{O}_w^\times)$, where τ is a generator of G_v : the isomorphism of Shapiro's Lemma gives

$$\hat{H}^{-1}(G_v, \mathcal{O}_w^\times) \xrightarrow{\cong} \prod_{w|v} \hat{H}^{-1}(G, \mathcal{O}_w^\times)$$

$$(\tau - 1)\pi_w \longmapsto \{\sigma((\tau - 1)\pi_w)\}_{\sigma \in G/G_v}$$

and therefore the image of β_v is generated by $\sum_{\sigma \in G/G_v} \sigma((\tau - 1)\pi_w) := \vartheta_v$.

On the other hand, the map δ is given (see [NSW00, chapter I, section 2]) by $x \mapsto (\gamma - 1)x$, where γ is a generator of G , for all $x \in \hat{H}^0(G, Cl_L)$. As $Cl_L = \text{Id}_L/\text{Pr}_L \cong C_L/Q_L$, we can write the class $[\mathfrak{P}_w] \in Cl_L$ as $[(\dots, 1, \pi_w, 1, \dots)]$ where $(\dots, 1, \pi_w, 1, \dots) \in \mathbb{A}_L^\times$ is the idèle whose w -component is π_w and all other components are 1. Then $\text{Norm}_{G/G_v}[\mathfrak{P}_w] = \sum_{\sigma \in G/G_v} \sigma[(\dots, 1, \pi_w, 1, \dots)] \in H^0(G, Cl_L)$ and the image via δ of its class in $\hat{H}^0(G, Cl_L)$ is clearly ϑ_v (observe that $\sum_{\sigma} \sigma(\tau - 1) = \sum_{\sigma} \sigma(\gamma - 1)$).

PROPOSITION 2.2 (See [Iwa73]). *Let $J : Cl_K \rightarrow Cl_L$ be the map induced by extending fractional ideals of $\mathcal{O}_K$ to $\mathcal{O}_L$. Then*

$$\text{Ker}(J) \cong \text{Ker}(\hat{H}^1(G, \mathcal{O}_L^\times) \rightarrow \hat{H}^1(G, U_\Sigma)).$$

Proof. We simply apply the Snake Lemma twice. First of all, apply it to

$$\begin{array}{ccccccc} 0 & \longrightarrow & Q_K & \longrightarrow & C_K & \longrightarrow & Cl_K \longrightarrow 0 \\ & & \downarrow & & \parallel & & \downarrow J \\ 0 & \longrightarrow & Q_L^G & \longrightarrow & C_L^G & \longrightarrow & Cl_L^G \longrightarrow \hat{H}^1(G, Q_L) : \end{array}$$

it shows $\text{Ker}(J) \cong Q_L^G/Q_K$. Then apply it to

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{O}_K^\times & \longrightarrow & \mathbb{U}_K & \longrightarrow & Q_K \longrightarrow 0 \\ & & \parallel & & \parallel & & \downarrow \\ 0 & \longrightarrow & (\mathcal{O}_L^\times)^G & \longrightarrow & (\mathbb{U}_L)^G & \longrightarrow & Q_L^G \longrightarrow \hat{H}^1(G, \mathcal{O}_L^\times), \end{array}$$

finding $Q_L^G/Q_K \cong \text{Ker}(\hat{H}^1(G, \mathcal{O}_L^\times) \rightarrow \hat{H}^1(G, U_\Sigma))$.

Remark. The above proof does not use the hypothesis that G be cyclic. In fact, we will in the sequel apply the proposition assuming this cyclicity, finding $\text{Ker}(J) \cong \text{Ker}(\hat{H}^{-1}(G, \mathcal{O}_L^\times) \rightarrow \hat{H}^{-1}(G, U_\Sigma))$. We remark that in his thesis [Gre76] Greenberg gives the following criterion: $\mu = \lambda = 0$ if and only if for all $m \geq 0$ the map J relative to the cyclic extension F_n/F_m , restricted to p -Sylow subgroups, becomes 0 for sufficiently large n (possibly depending on m). This will be the starting point for our proof of Theorem 5.1.

We end this section with some remarks on the action of Δ on the cohomology along a $\mathbb{Z}_p$ -extension. Pick a p -adic character χ of Δ ; setting $R_\chi = \mathbb{Z}_p[\text{Im}(\chi)]$, we let $\delta \in \Delta$ act on

R_χ by $x \mapsto \chi(\delta)x$. In this way, R_χ becomes a $\mathbb{Z}_p[\Delta]$ -algebra. For every $\mathbb{Z}[\Delta]$ -module M we denote by $M(\chi)$ the χ -eigenspace of $M \otimes_{\mathbb{Z}_p[\Delta]} R_\chi$ for the action of Δ : it is defined as $M(\chi) = \{m \in M \otimes R_\chi \text{ s. t. } \delta \cdot m = \chi(\delta) \cdot m \forall \delta \in \Delta\} \cong \text{Hom}_{\mathbb{Z}_p[\Delta]}(R_\chi, M \otimes \mathbb{Z}_p)$: for more details, see [Sch88, section 2]. Then we have the following

LEMMA 2.3 [Sch88, lemma 2·1]. *Suppose that G and Δ are abelian groups such that $p \nmid |\Delta|$ and G is a p -group. Let M be a $\mathbb{Z}[\Delta \times G]$ -module, then:*

(i) *for every $q \in \mathbb{Z}$, the natural map*

$$\hat{H}^q(G, M \otimes \mathbb{Z}_p)^\Delta \longrightarrow \hat{H}^q(G, (M \otimes \mathbb{Z}_p)^\Delta)$$

is an isomorphism (of abelian groups with trivial $\Delta \times G$ -action)

(ii) *for every character $\chi : \Delta \rightarrow \overline{\mathbb{Q}}_p^\times$, there is an isomorphism of $\mathbb{Z}_p[\Delta]$ -modules*

$$\hat{H}^q(G, M(\chi)) \cong \hat{H}^q(G, M)(\chi) \quad \forall q \in \mathbb{Z}. \quad (2.2)$$

3. Cyclotomic units

We now go back to the situation described in the introduction: let then F_∞/F be the $\mathbb{Z}_p$ -extension of F and denote by Γ its Galois group. For every n and for every prime ideal $\mathfrak{p} \subseteq \mathcal{O}_{F_n}$ dividing p , we let $\mathcal{O}_{F_n, \mathfrak{p}}^\times$ denote the local units at $\mathfrak{p}$. Also, we let Cyc_{F_n} be the cyclotomic units of F_n as defined by Sinnott in [Sin81]. Fix from now on a non-trivial character $\chi \neq 1$ of Δ and an odd prime $p \nmid [F : \mathbb{Q}]$ such that $\chi(p) = 1$: accordingly, let $R_\chi = \mathbb{Z}_p[(\text{Im}(\chi))]$, seen as a Δ -module as detailed at the end of Section 2. Remark that for every $n \geq 0$, $\text{Gal}(F_n/\mathbb{Q}) \cong \Delta \times \text{Gal}(F_n/F)$, and this turns $F_n^\times$, Cyc_{F_n} , Cl_{F_n} , etc... into Δ -modules.

Consider a contravariant functor

$$\mathcal{M} : \underline{\text{SGr}}(\Delta) \longrightarrow \underline{\mathbb{Z}[\Delta]\text{-Mod}}$$

from the category of subgroups of Δ (morphisms being inclusions) to that of $\mathbb{Z}[\Delta]$ -modules, verifying the condition (compare with [NSW00])

$$\mathcal{M}(H) \cong \mathcal{M}(1)^H \text{ for all } H \triangleleft \Delta \text{ functorially in } H. \quad (3.1)$$

Example. Prototypical examples of such a functor are the additive group $\mathcal{M}(H) := F_n^H$, or the global (resp. cyclotomic, local) units $\mathcal{M}(H) := \mathcal{O}_{F_n^H}^\times$ (resp. $\mathcal{M}(H) := \text{Cyc}_{F_n^H}$, $\mathcal{M}(H) = \mathcal{O}_{F_n^H, \mathfrak{p}}^\times$), together with the natural inclusions $F_n^H \hookrightarrow F_n$, $\mathcal{O}_{F_n^H}^\times \hookrightarrow \mathcal{O}_{F_n}^\times$, etc... Moreover, since $p \nmid |\Delta|$, the p -part of the class group verifies (3.1): indeed, for all H the map $\text{Cl}_{F_n^H} \xrightarrow{j} \text{Cl}_{F_n}^H \xrightarrow{\text{Norm}} \text{Cl}_{F_n^H}$ coincides with the multiplication by $[\Delta : H]$, and is therefore an isomorphism when restricted to p -Sylow (and this for all $n \geq 0$). Similarly, we have $H^1(H, \text{Cyc}_n \otimes \mathbb{Z}_p) = 0$ for all $H \triangleleft \Delta$ because $p \nmid |H|$ (apply Corollary 1 to Proposition 8 of Atiyah and Wall's paper in [CF86]), so

$$((\mathcal{O}_{F_n}^\times / \text{Cyc}_{F_n}) \otimes \mathbb{Z}_p)^H \cong (\mathcal{O}_{F_n^H} / \text{Cyc}_{F_n^H}) \otimes \mathbb{Z}_p$$

and also the functors $\mathcal{M}(H) = (\mathcal{O}_{F_n^H} / \text{Cyc}_{F_n^H}) \otimes \mathbb{Z}_p$ verify condition (3.1) for all $n \geq 0$.

Suppose now that a functor as above is given, and that it verifies (3.1): then for every character χ of Δ , there is a natural identification $\mathcal{M}(1)(\chi) \cong \mathcal{M}(\text{Ker}(\chi))(\chi)$. Indeed, $\mathcal{M}(\text{Ker}(\chi)) \cong \mathcal{M}(1)^{\text{Ker}(\chi)} \subseteq \mathcal{M}(1)$, while $\mathcal{M}(1)(\chi)$ has obviously a trivial action of $\text{Ker}(\chi)$, showing $\mathcal{M}(1)(\chi) \subseteq (\mathcal{M}(1) \otimes R_\chi)^{\text{Ker}(\chi)}$. Since, as the above examples show, we

will be dealing only with functors verifying (3.1), this discussion enables us to replace Δ with $\Delta/\text{Ker}(\chi) \cong \text{Im}(\chi)$ without changing the χ -components we are interested in. We can therefore assume that p splits completely in F , and we assume this throughout.

Then we set

$$U_n := \left(\prod_{\mathfrak{p}|p} \mathcal{O}_{F_n, \mathfrak{p}}^\times \right)$$

and we agree to let $\mathcal{O}_n^\times = \mathcal{O}_{F_n}^\times$ and similarly $\text{Cyc}_n = \text{Cyc}_{F_n}$. Observe that since in a $\mathbb{Z}_p$ -extension only primes above p ramify, the set Σ of Section 2 relative to F_n/F is $\Sigma = \{\mathfrak{p} \subset \mathcal{O}_F, \mathfrak{p}|p\}$ for all n and our notation is consistent with the one introduced there. We finally set

$$\mathcal{B}_n := (\mathcal{O}_n^\times / \text{Cyc}_n)(\chi), \quad \mathcal{B}_n := (U_n / \text{Cyc}_n)(\chi) \quad \text{and} \quad A_n := Cl_{F_n}(\chi).$$

By Sinnott's theorem above, together with [Sin81, theorem 5.3] that guarantees $p \nmid \kappa_{F_n}$ for all n , the groups $Cl_{F_n} \otimes \mathbb{Z}_p$ and $(\mathcal{O}_n^\times / \text{Cyc}_n) \otimes \mathbb{Z}_p$ have the same order. The character-by-character version of this result is much deeper: it is known as Gras' Conjecture, and is a consequence of the (now proven) Main Conjecture of Iwasawa Theory, as detailed in [Gre77] or [BNQD01]. It follows that $|A_n| = |\mathcal{B}_n|$ for all $n \geq 0$.

Remark. The semi-local units considered by Gillard [Gil79b] are products over all $\mathfrak{p}$ above p of local units that are 1 (mod $\mathfrak{p}$). In our situation, all completions $F_{\mathfrak{p}}$ at primes $\mathfrak{p} | p$ are isomorphic to $\mathbb{Q}_p$, so the two definitions coincide and $U_n(\chi)$ is a free R_χ -module of rank 1.

Moreover, since p splits completely in $F/\mathbb{Q}$, all primes above p totally ramify in F_n/F : let $\mathfrak{P}_n$ be one such a prime. Its image in A_n generates a subgroup which is independent of the choice of $\mathfrak{P}_n$: indeed, all primes above p are conjugate by an element in Δ , and the image in A_n of $\delta(\mathfrak{P}_n)$ is $\mathfrak{P}_n^{\chi(\delta)}$, which generates the same subgroup as $\mathfrak{P}_n$. We call this subgroup Π_n and we can restate Lemma 2.1 saying that $\text{Im}(\beta) = \delta(\Pi_n)$ (use also Lemma 2.3).

We now investigate in some detail the structure of Cyc_n . First of all, letting f be the conductor of F , we define

$$\eta_n := \left(\text{Norm}_{F_n}^{\mathbb{Q}(\zeta_{fp^{n+1}})} (1 - \zeta_{fp^{n+1}}) \right) \otimes \left(\sum_{\delta \in \Delta} \chi(\delta^{-1}) \delta \right) \in \text{Cyc}_n(\chi).$$

The left factor is a unit since $p \nmid f$ because p splits completely in F , it is cyclotomic by definition and is acted upon by Δ through χ : moreover, Sinnott's description of cyclotomic units shows that $\text{Cyc}_n(\chi) = \eta_0 R_\chi \times \eta_n R_\chi [G_n]$ (see, for instance, [Gil79a, section 3] for details). In particular, we have an isomorphism of G_n -modules $\text{Cyc}_n(\chi) \cong R_\chi \times I_{G_n}$ where I_{G_n} is the augmentation ideal in $R_\chi [G_n]$, and we find a split exact sequence

$$0 \longrightarrow \langle \eta_n \rangle \longrightarrow \text{Cyc}_n(\chi) \longrightarrow \langle \eta_0 \rangle \longrightarrow 0 \quad (3.2)$$

where we denote, here and in what follows, by $\langle \eta_0 \rangle$ and $\langle \eta_n \rangle$ the $R_\chi [G_n]$ -modules generated by η_0 and η_n respectively: by the above isomorphisms, (3.2) corresponds to the sequence

$$0 \longrightarrow I_{G_n} \longrightarrow I_{G_n} \times R_\chi \longrightarrow R_\chi \longrightarrow 0. \quad (3.3)$$

LEMMA 3.1. *We have $\hat{H}^0(G_{n,m}, \langle \eta_n \rangle) = 0$ and $\hat{H}^{-1}(G_{n,m}, \langle \eta_0 \rangle) = 0$. In particular, the natural map $\hat{H}^q(G_{n,m}, \text{Cyc}_n(\chi)) \cong \hat{H}^q(G_{n,m}, \langle \eta_0 \rangle) \times \hat{H}^q(G_{n,m}, \langle \eta_n \rangle)$ induced by (3.2) gives*

isomorphisms of Δ -modules

$$\hat{H}^0(G_{n,m}, \text{Cyc}_n(\chi)) \cong \hat{H}^0(G_{n,m}, \langle \eta_0 \rangle)$$

and

$$\hat{H}^{-1}(G_{n,m}, \text{Cyc}_n(\chi)) \cong \hat{H}^{-1}(G_{n,m}, \langle \eta_n \rangle).$$

Both are groups of order p^{dn} where $d = \text{rk}_{\mathbb{Z}_p} R_\chi$.

Proof. The exact sequence

$$0 \longrightarrow I_{G_n} \longrightarrow R_\chi[G_n] \longrightarrow R_\chi \longrightarrow 0$$

shows, since $\hat{H}^q(G_{n,m}, R_\chi[G_n]) = 0$ for all q , that $\hat{H}^0(G_{n,m}, I_{G_n}) \cong \hat{H}^{-1}(G_{n,m}, R_\chi) = 0$. The $R_\chi[G_n]$ -isomorphisms $\langle \eta_n \rangle \cong I_{G_n}$ and $\langle \eta_0 \rangle \cong R_\chi$ give the result.

Suppose now that $\lambda = 0$. The extension F_∞/F is totally ramified since p splits in F and the maps $N_m^n: A_n \rightarrow A_m$ (induced by norms on (χ) -parts, and that we keep on calling “Norm maps” by a slight abuse of notation) are surjective by class field theory for all $n \geq m$; assuming $\lambda = 0$ and choosing m big enough, the orders of A_n and A_m coincide, and these norm maps are actually isomorphisms. Therefore the projective limit $X = \varprojlim A_n$ with respect to norms stabilizes to a finite group and $A_n \cong X$ for all $n \geq 0$ (we introduce here the notation $a \geq b$, equivalent to $b \leq a$, to mean that there exists a $b_0 \geq b$ such that what we are stating holds for all $a \geq b_0$). In particular, the action of Γ on X must factor through a certain quotient $G_m = \Gamma / \Gamma^{p^m}$. Therefore $G_{n,m}$ acts trivially on A_n for all $n \geq m$ and the $G_{n,m}$ -norm $N_{G_{n,m}} = \sum_{\tau \in G_{n,m}} \tau$ acts on A_n as multiplication by p^{n-m} . Choosing n big enough so that $p^{n-m} A_n = 0$, we find $N_{G_{n,m}} A_n = 0$ and

$$\begin{aligned} \hat{H}^0(G_{n,m}, A_n) &= A_n^{G_{n,m}} / N_{G_{n,m}} A_n = A_n \\ &= A_n [N_{G_{n,m}}] / I_{G_{n,m}} A_n = \hat{H}^{-1}(G_{n,m}, A_n) \end{aligned} \quad (3.4)$$

where $I_{G_{n,m}}$ is the augmentation ideal of $R_\chi[G_{n,m}]$. Therefore $A_n \cong \hat{H}^{-1}(G_{n,m}, A_n) \cong \hat{H}^0(G_{n,m}, A_n)$, whenever $\lambda = 0$ and $n \geq m \geq 0$. A similar argument leads to the equivalent of (3.4) for B_n , namely $\hat{H}^q(G_{n,m}, B_n) \cong B_n$ for all $q \in \mathbb{Z}$.

LEMMA 3.2. *If $\lambda = 0$ and $m \geq 0$, the natural map*

$$H^1(G_{n,m}, \text{Cyc}_n(\chi)) \longrightarrow H^1(G_{n,m}, \mathcal{O}_n^\times(\chi))$$

is injective for all $n \geq m$.

Proof. Taking $G_{n,m}$ -cohomology in the exact sequence defining B_n gives

$$\begin{aligned} 0 \longrightarrow H^0(G_{n,m}, \text{Cyc}_n(\chi)) \longrightarrow H^0(G_{n,m}, \mathcal{O}_n^\times(\chi)) \longrightarrow H^0(G_{n,m}, B_n) \longrightarrow \\ \longrightarrow H^1(G_{n,m}, \text{Cyc}_n(\chi)) \longrightarrow H^1(G_{n,m}, \mathcal{O}_n^\times(\chi)). \end{aligned} \quad (3.5)$$

Since $G_{n,m}$ -invariants of $\text{Cyc}_n(\chi)$ and $\mathcal{O}_n^\times(\chi)$ are $\text{Cyc}_m(\chi)$ and $\mathcal{O}_m^\times(\chi)$ respectively, we immediately see that $\text{Ker}(H^1(G_{n,m}, \text{Cyc}_n(\chi)) \longrightarrow H^1(G_{n,m}, \mathcal{O}_n^\times(\chi))) = B_n^{G_{n,m}} / B_m$. Assuming that m is big enough and $\lambda = 0$ implies that the orders of B_n and B_m coincide, and the same holds, *a fortiori*, for $B_n^{G_{n,m}}$ and B_m . Thus, the above kernel is trivial.

4. Semi-local units modulo cyclotomic units

We now state a very useful result about semi-local units in our setting; it can already be found in a paper by Iwasawa [Iwa60]. We keep the same notation introduced in the previous section and we make from now on constant use of Lemma 2.3 above, especially in the form of the isomorphism in (2.2).

Definition 4.1. We define U_n^1 to be the kernel $U_n^1 = \text{Ker}(N_0^n: U_n \rightarrow U_0)$ and we set $\mathcal{B}_n^1 = U_n^1(\chi)/\langle \eta_n \rangle$.

PROPOSITION 4.2. *The natural map $U_n^1 \times U_0 \hookrightarrow U_n$ induced by injections is an isomorphism of $\Delta \times G_n$ -modules. It induces a decomposition $\mathcal{B}_n \cong \mathcal{B}_n^1 \times \mathcal{B}_0$.*

Proof. Since all primes above p are totally ramified in F_n/F we can restrict our attention to a single prime $\mathfrak{p} \mid p$ in $\mathcal{O}_F$ and show that

$$\mathcal{O}_{n,\mathfrak{p}}^{\times 1} \times \mathcal{O}_{0,\mathfrak{p}}^{\times} \cong \mathcal{O}_{n,\mathfrak{p}}^{\times}$$

where we denote by $\mathcal{O}_{n,\mathfrak{p}}^{\times 1}$ the local units having norm 1 in $\mathcal{O}_{0,\mathfrak{p}}^{\times}$: taking the product over all primes above p will give the first statement of the Proposition. Consider the exact sequence induced by the norm map N_0^n

$$0 \longrightarrow \mathcal{O}_{n,\mathfrak{p}}^{\times 1} \longrightarrow \mathcal{O}_{n,\mathfrak{p}}^{\times} \longrightarrow N_0^n(\mathcal{O}_{n,\mathfrak{p}}^{\times}) \subseteq \mathcal{O}_{0,\mathfrak{p}}^{\times} \longrightarrow 0.$$

Since the extension $F_{n,\mathfrak{p}}/F_{0,\mathfrak{p}}$ is cyclic and totally ramified, local class field theory shows that $\hat{H}^0(G_n, \mathcal{O}_{n,\mathfrak{p}}^{\times}) = \mathcal{O}_{0,\mathfrak{p}}^{\times}/N_0^n(\mathcal{O}_{n,\mathfrak{p}}^{\times})$ is a cyclic group of order p^n . As $\mathcal{O}_{0,\mathfrak{p}}^{\times} \cong \mathbb{Z}_p^{\times}$ this shows $N_0^n(\mathcal{O}_{n,\mathfrak{p}}^{\times}) = (\mathcal{O}_{0,\mathfrak{p}}^{\times})^{p^n} \times \mu_{p-1}$ and since $\mathcal{O}_{0,\mathfrak{p}}^{\times}$ contains no roots of unity of p -power order we can identify this group with $\mathcal{O}_{0,\mathfrak{p}}^{\times}$ simply by extracting p^n th roots. We find

$$0 \longrightarrow \mathcal{O}_{n,\mathfrak{p}}^{\times 1} \longrightarrow \mathcal{O}_{n,\mathfrak{p}}^{\times} \xrightarrow{p^n \sqrt[p^n]{N_0^n}} \mathcal{O}_{0,\mathfrak{p}}^{\times} \longrightarrow 0. \quad (4.1)$$

Since the natural embedding $\mathcal{O}_{0,\mathfrak{p}}^{\times} \hookrightarrow \mathcal{O}_{n,\mathfrak{p}}^{\times}$ is a G_n -linear section of (4.1), it splits the sequence and therefore gives an isomorphism $\mathcal{O}_{n,\mathfrak{p}}^{\times 1} \times \mathcal{O}_{0,\mathfrak{p}}^{\times} \cong \mathcal{O}_{n,\mathfrak{p}}^{\times}$.

The explicit description $\text{Cyc}_n(\chi) = \langle \eta_0 \rangle \times \langle \eta_n \rangle$ together with $N_0^n(\eta_n) = 0$ (see the discussion before equation (3.2)) shows that $N_0^n(\text{Cyc}_n(\chi)) = N_0^n(\langle \eta_0 \rangle) = \langle \eta_0^{p^n} \rangle$. The argument above thus gives the commutative diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \langle \eta_n \rangle & \longrightarrow & \text{Cyc}_n(\chi) & \xrightarrow{p^n \sqrt[p^n]{N_0^n}} & \langle \eta_0 \rangle \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & U_n^1(\chi) & \longrightarrow & U_n(\chi) & \xrightarrow{p^n \sqrt[p^n]{N_0^n}} & U_0(\chi) \longrightarrow 0 \end{array}$$

and this proves $\mathcal{B}_n \cong \mathcal{B}_n^1 \times \mathcal{B}_0$.

More useful than the splitting itself is the following easy consequence:

COROLLARY 4.3. *For every $n \geq m \geq 0$ and for every $q \in \mathbb{Z}$ the natural maps*

$$\hat{H}^q(G_{n,m}, U_n(\chi)) \longrightarrow \hat{H}^q(G_{n,m}, U_n^1(\chi)) \times \hat{H}^q(G_{n,m}, U_0(\chi))$$

and

$$\hat{H}^q(G_{n,m}, \mathcal{B}_n) \longrightarrow \hat{H}^q(G_{n,m}, \mathcal{B}_n^1) \times \hat{H}^q(G_{n,m}, \mathcal{B}_0)$$

induced by Proposition 4.2 are isomorphisms of abelian groups. In particular, we have $\hat{H}^{-1}(G_{n,m}, U_n(\chi)) = \hat{H}^{-1}(G_{n,m}, U_n^1(\chi))$ and $\hat{H}^0(G_{n,m}, U_n(\chi)) = \hat{H}^0(G_{n,m}, U_0(\chi))$.

Proof. The splitting of the cohomology groups follows immediately from the Proposition. Concerning the cohomology of $U_n(\chi)$, we observe that, since $G_{n,m}$ acts trivially on the torsion-free module U_0 , the group $\hat{H}^0(G_{n,m}, U_0)$ is a group of order $p^{[F:\mathbb{Q}](n-m)}$ while $\hat{H}^{-1}(G_{n,m}, U_0) = 0$. This already implies that $\hat{H}^{-1}(G_{n,m}, U_n) = \hat{H}^{-1}(G_{n,m}, U_n^1)$. It also shows that $\hat{H}^0(G_{n,m}, U_n^1)$ must be trivial because $\hat{H}^0(G_{n,m}, U_n)$ has order $p^{[F:\mathbb{Q}](n-m)}$ by local class field theory; Lemma 2.3 then allows to take (χ) -parts and to get the desired equalities.

LEMMA 4.4. *For all $n \geq m \geq 0$, there are isomorphisms $\hat{H}^q(G_{n,m}, \mathcal{B}_n^1) \cong R_\chi/L_p(1, \chi)$ and $\hat{H}^q(G_{n,m}, \mathcal{B}_0) \cong R_\chi/L_p(1, \chi)$ holding for every $q \in \mathbb{Z}$.*

Proof. Let $\Lambda := R_\chi[[T]]$ and fix an isomorphism $\varpi : \Lambda \cong R_\chi[[\Gamma]]$. This isomorphism fixes a choice of a topological generator $\varpi(1+T) =: \gamma_0$ of Γ and we denote by $\kappa \in \mathbb{Z}_p^\times$ the element $\varepsilon_{\text{cyc}}(\gamma_0)$ where

$$\varepsilon_{\text{cyc}} : \text{Gal}(\bar{F}/F) \longrightarrow \mathbb{Z}_p^\times$$

is the cyclotomic character of F . The main tool of the proof will be [Gil79b, theorem 2], that gives isomorphisms of $R_\chi[[\Gamma]]$ -modules

$$\mathcal{B}_0 \cong R_\chi/L_p(1, \chi) \quad \text{and} \quad \mathcal{B}_n \cong \Lambda/(f(T), \omega_n(T)/T) \quad (4.2)$$

where $\omega_n(T) = (1+T)^{p^n} - 1$ and $f(T) \in \Lambda$ is the power series verifying $f(\kappa^s - 1) = L_p(1-s, \chi)$ for all $s \in \mathbb{Z}_p$. We make Γ act on the modules appearing in (4.2) by $\gamma_0 \cdot x = \varpi^{-1}(\gamma_0)x = (1+T)x$ for all $x \in \mathcal{B}_0$ (resp. all $x \in \mathcal{B}_n$): this induces the action of $G_{n,m}$ we need to compute the cohomology with respect to.

Starting with $\mathcal{B}_0$, observe that the action of Γ , and thus of its subquotient $G_{n,m}$, is trivial on the finite group $\mathcal{B}_0$: as in (3.4) we get

$$\hat{H}^q(G_{n,m}, \mathcal{B}_0) \cong \mathcal{B}_0 \quad \text{for all } n \geq m \geq 0,$$

and we apply (4.2) to get our claim.

Now we compute $\hat{H}^{-1}(G_{n,m}, \mathcal{B}_n^1)$: by definition, $\hat{H}^{-1}(G_{n,m}, \mathcal{B}_n^1) = \mathcal{B}_n^1[N_{G_{n,m}}]/I_{G_{n,m}}\mathcal{B}_n^1$. Applying ϖ we find $I_{G_{n,m}} \cong \omega_m(T)(\Lambda/\omega_n(T))$ and $\varpi(N_{G_{n,m}}) = v_{n,m}(T)$ where $v_{n,m}(T) := \omega_n(T)/\omega_m(T)$. Hence

$$\hat{H}^{-1}(G_{n,m}, \mathcal{B}_n^1) \cong \frac{\{g(T) \in \Lambda \mid g(T)v_{n,m}(T) \in (f(T), \omega_n(T)/T)\}}{(f(T), \omega_n(T)/T, \omega_m(T))}.$$

As observed in [Gil79b, lemma 5], $f(T)$ and $\omega_n(T)/T$ have no common zeroes. Therefore a relation

$$g(T) \frac{\omega_n(T)}{\omega_m(T)} = a(T)f(T) + b(T) \frac{\omega_n(T)}{T}$$

implies $v_{n,m}(T) \mid a(T)$ and we find $g(T) = c(T)f(T) + b(T)\omega_m(T)/T$ for some $c(T) \in \Lambda$: thus,

$$\begin{aligned} \hat{H}^{-1}(G_{n,m}, \mathcal{B}_n^1) &\cong \frac{(f(T), \omega_m(T)/T)}{(f(T), \omega_n(T)/T, \omega_m(T))} \\ &\cong \frac{(\omega_m(T)/T)}{(f(T), \omega_n(T)/T, \omega_m(T))}. \end{aligned}$$

The evaluation map $g(T)\omega_m(T)/T \mapsto g(0)$ gives an isomorphism

$$\frac{(\omega_m(T)/T)}{(\omega_m(T))} \cong R_\chi$$

and we find

$$\frac{(\omega_m(T)/T)}{(f(T), \omega_n(T)/T, \omega_m(T))} \cong R_\chi / (f(0), \omega_n(0)).$$

Since this last module is $R_\chi/f(0)$ as soon as n is big enough and, by definition, $f(0) = L_p(1, \chi)$, we get our claim for $q = -1$. Using now that $\mathcal{B}_n^1$ is finite and therefore has a trivial Herbrand quotient, we know that the order of $\hat{H}^0(G_{n,m}, \mathcal{B}_n^1)$ is the predicted one: the fact that it is cyclic as an R_χ -module comes from the exact sequence

$$\begin{aligned} 0 \longrightarrow \hat{H}^0(G_{n,m}, \mathcal{B}_n^1) \longrightarrow \hat{H}^0(G_{n,m}, U_n^1(\chi)) \longrightarrow \hat{H}^0(G_{n,m}, \langle \eta_n \rangle) \longrightarrow \\ \longrightarrow \hat{H}^1(G_{n,m}, \mathcal{B}_n^1) \longrightarrow 0 \end{aligned}$$

since $\hat{H}^0(G_{n,m}, U_n^1(\chi))$ is itself cyclic, as discussed in Corollary 4.3.

Finally, the fact that $G_{n,m}$ is cyclic gives isomorphisms in Tate cohomology $\hat{H}^{2q}(G_{n,m}, M) \cong \hat{H}^0(G_{n,m}, M)$ for all modules M (and analogously $\hat{H}^{2q+1}(G_{n,m}, M) \cong \hat{H}^{-1}(G_{n,m}, M)$), so the claim for all q 's follows from our computation in the cases $q = 0, -1$.

We want to stress that R_χ is an unramified extension of $\mathbb{Z}_p$. In particular, p is a uniformizer in R_χ and therefore the p -adic valuation of an element in R_χ is an integer: for $a, b \in R_\chi$ we will write $a \stackrel{p}{=} b$ to mean that both sides have the same valuation.

PROPOSITION 4.5. *Recall that $X = \varprojlim A_n$: if $\lambda = 0$ then $|X^\Gamma| \stackrel{p}{=} L_p(1, \chi)^d$ where $d = rk_{\mathbb{Z}_p} R_\chi$.*

Proof. Let L_0 be the maximal pro- p abelian extension of F_∞ everywhere unramified and let M_0 be the maximal pro- p abelian extension of F_∞ unramified outside p . We claim that $L_0 = M_0$. This follows from the fact that for every $\mathfrak{p} \subseteq \mathcal{O}_F$ dividing p , the local field $F_{\mathfrak{p}}$ is $\mathbb{Q}_p$, since p splits completely, and it therefore admits only two independent $\mathbb{Z}_p$ -extensions by local class field theory. In particular, every pro- p extension of $F_{\infty, \mathfrak{p}}$ that is abelian over $F_{\mathfrak{p}}$ must be unramified, so $M_0 = L_0$. Now let $Y := \text{Gal}(L_\infty/F_\infty)$ where L_∞ is the maximal pro- p abelian extension of F_∞ everywhere unramified: then the Artin reciprocity map gives an isomorphism $X \cong Y(\chi)$; also, let M_∞ be the maximal pro- p abelian extension of F_∞ unramified outside p and $\mathcal{Y} := \text{Gal}(M_\infty/F_\infty)$. A classical argument (see [Was97, chapter 13]) shows that $Y_\Gamma = \text{Gal}(L_0/F_\infty)$ and $\mathcal{Y}_\Gamma = \text{Gal}(M_0/F_\infty)$: our claim above implies that $Y_\Gamma = \mathcal{Y}_\Gamma$. Since the actions of Δ and Γ commute with each other, this also shows $X_\Gamma = \mathcal{Y}(\chi)_\Gamma$. Combine this with the following exact sequence induced by multiplication by $\gamma_0 - 1$ where γ_0 is a topological generator of Γ

$$0 \longrightarrow X^\Gamma \longrightarrow X \xrightarrow{\gamma_0 - 1} X \longrightarrow X_\Gamma \longrightarrow 0:$$

since we assumed $\lambda = 0$ all the groups are finite and the sequence gives $|X^\Gamma| = |X_\Gamma| = |\mathcal{Y}(\chi)_\Gamma|$. The Main Conjecture of Iwasawa Theory, as proved by Rubin of [Lan90, appendix], shows that the characteristic polynomial of $\mathcal{Y}(\chi)$ is $f(T)$ (see the proof of Lemma 4.4). Since $\mathcal{Y}$ contains no non-zero finite Γ -submodules (see [NSW00]), we find $\mathcal{Y}^\Gamma = 0$ and the order of $\mathcal{Y}(\chi)_\Gamma$ is $f(0)^d \stackrel{p}{=} L_p(1, \chi)^d$, as detailed in [Was97, exercise 13.12].

COROLLARY 4.6. *If $\lambda = 0$, then Π_n is a cyclic R_χ -module of order $|L_p(1, \chi)|_p^{-d}$ for every $n \geq 0$.*

Proof. Indeed, [Gre76, theorem 2] shows that $\lambda = 0$ if and only if $X^\Gamma = \Pi_n$. The result now follows from the proposition and from the remark of Section 3.

5. Main result

We are now in position of proving our main result. We stick to the notation introduced in Section 3. Let $n \geq 0$ and let $\mathcal{Q}_n := (\mathbb{U}_{F_n}/\mathcal{O}_n^\times)(\chi)$ as in Section 2: consider the exact sequence

$$0 \longrightarrow \mathcal{O}_n^\times(\chi) \longrightarrow \mathbb{U}_{F_n}(\chi) \longrightarrow \mathcal{Q}_n \longrightarrow 0.$$

Since $\text{Cyc}_n(\chi) \subseteq \mathcal{O}_n^\times(\chi)$, it induces an exact sequence

$$0 \longrightarrow B_n \longrightarrow (\mathbb{U}_{F_n}/\text{Cyc}_n)(\chi) \longrightarrow \mathcal{Q}_n \longrightarrow 0,$$

and the Tate cohomology of $(\mathbb{U}_{F_n}/\text{Cyc}_n)(\chi)$ coincides with that of B_n , as discussed in Section 2. For every $m \leq n$ the cyclicity of Tate cohomology for cyclic groups induces an exact square

$$\begin{array}{ccc} \hat{H}^0(G_{n,m}, \mathcal{Q}_n) & \xrightarrow{\alpha_{[0,1]}} & \hat{H}^{-1}(G_{n,m}, B_n) \\ \uparrow & & \downarrow \\ \hat{H}^0(G_{n,m}, B_n) & & \hat{H}^{-1}(G_{n,m}, B_n) \\ \uparrow & & \downarrow \\ \hat{H}^0(G_{n,m}, B_n) & \xleftarrow{\alpha_{[1,0]}} & \hat{H}^{-1}(G_{n,m}, \mathcal{Q}_n). \end{array} \quad (5.1)$$

Consider now the exact sequence

$$0 \longrightarrow \mathcal{Q}_n \longrightarrow C_{F_n}(\chi) \longrightarrow A_n \longrightarrow 0: \quad (5.2)$$

by Lemma 2.3, we have $\hat{H}^q(G_{n,m}, C_n(\chi)) \cong \hat{H}^q(G_{n,m}, C_n)(\chi)$ for all $q \in \mathbb{Z}$; global class field theory (see section 11.3 of Tate's paper in [CF86]) shows $\hat{H}^q(G_{n,m}, C_n)(\chi) \cong \hat{H}^{q+2}(G_{n,m}, \mathbb{Z}(\chi)) = 0$ because we assumed $\chi \neq 1$. Therefore the long exact cohomology sequence of (5.2) induces isomorphisms

$$\hat{H}^q(G_{n,m}, A_n) \cong \hat{H}^{q+1}(G_{n,m}, \mathcal{Q}_n) \quad \text{for every } q \in \mathbb{Z}. \quad (5.3)$$

Remark. Observe that our discussion never uses the assumption $\chi(p) = 1$. Indeed, the maps $\alpha_{[0,1]}$ and $\alpha_{[1,0]}$ are defined whenever $\chi \neq 1$ and their composition with the isomorphisms in (5.3) are indeed the same maps appearing in [KS95, proposition 2.6], where the case $\chi(p) \neq 1$ is treated. As discussed in the introduction, in that case they turned out to be isomorphisms if $\lambda = 0$ (see also [BNQD01], [Kuz96] and [Oza97]). We are going to see this is not the case if $\chi(p) = 1$. Before stating the next theorem we recall that $R_\chi = \mathbb{Z}_p[\text{Im}(\chi)]$ with Δ -action $\delta \cdot x = \chi(\delta)x$ for all $\delta \in \Delta$: its rank over $\mathbb{Z}_p$ is denoted by d .

THEOREM 5.1. *Let F be a real abelian field such that $p \nmid [F : \mathbb{Q}]$ and fix a non-trivial character χ of Δ such that $\chi(p) = 1$. Assume that $\lambda = 0$ and let $n \geq m \geq 0$ be big enough. Then the kernels $\text{Ker}(\alpha_{[0,1]})$, $\text{Ker}(\alpha_{[1,0]})$ and the cokernels $\text{Coker}(\alpha_{[0,1]})$, $\text{Coker}(\alpha_{[1,0]})$ of the maps introduced in (5.1) are cyclic R_χ -modules of order $|L_p(1, \chi)|_p^{-d}$.*

Proof. We start by determining $\text{Ker}(\alpha_{[0,1]})$. Choose m big enough so that $|A_{m+k}| = |A_m|$ for all $k \geq 0$ and $n \geq m$ big enough so that $\hat{H}^q(G_{n,m}, A_n) = A_n$. As [Gre76, remark of section 2, proposition 2] shows that $\lambda = 0$ implies $A_m = \text{Ker}(J_{m,n})$ if n is sufficiently large. Combining Proposition 2.2 with (5.3), this gives an injection (here and in the rest of the proof we make constant use of Lemma 2.3)

$$\hat{H}^0(G_{n,m}, Q_n) \hookrightarrow \hat{H}^{-1}(G_{n,m}, \mathcal{O}_n^\times(\chi)). \quad (5.4)$$

Consider now the following commutative diagram, whose row and column are exact and where the injectivity of the vertical arrow in the middle follows from Lemma 3.2:

$$\begin{array}{ccccc} & & \hat{H}^{-1}(G_{n,m}, B_n) & & \\ & \nearrow \alpha_{[0,1]} & \uparrow & & \\ \hat{H}^0(G_{n,m}, Q_n) & \hookrightarrow & \hat{H}^{-1}(G_{n,m}, \mathcal{O}_n^\times(\chi)) & \longrightarrow & \hat{H}^{-1}(G_{n,m}, U_n(\chi)). \\ & & \uparrow & \searrow \psi & \\ & & \hat{H}^{-1}(G_{n,m}, \text{Cyc}_n(\chi)) & & \end{array} \quad (5.5)$$

An easy diagram chase shows that $\text{Ker}(\alpha_{[0,1]}) \cong \text{Ker}(\psi)$. In order to study $\text{Ker}(\psi)$, observe that ψ appears in the sequence

$$0 \longrightarrow \hat{H}^0(G_{n,m}, \mathcal{B}_n^1) \longrightarrow \hat{H}^{-1}(G_{n,m}, \langle \eta_n \rangle) \xrightarrow{\psi} \hat{H}^{-1}(G_{n,m}, U_n^1(\chi)), \quad (5.6)$$

since $\hat{H}^{-1}(G_{n,m}, \text{Cyc}_n(\chi)) = \hat{H}^{-1}(G_{n,m}, \langle \eta_n \rangle)$ by Lemma 3.1 and $\hat{H}^{-1}(G_{n,m}, U_n(\chi)) = \hat{H}^{-1}(G_{n,m}, U_n^1(\chi))$ by Corollary 4.3. Moreover, again by Corollary 4.3, $\hat{H}^0(G_{n,m}, U_n^1(\chi)) = 0$ and (5.6) is exact, thus giving

$$\text{Ker}(\alpha_{[0,1]}) \cong \hat{H}^0(G_{n,m}, \mathcal{B}_n^1) \cong R_\chi / L_p(1, \chi), \quad (5.7)$$

the last isomorphism being Lemma 4.4.

Having determined $\text{Ker}(\alpha_{[0,1]})$, the exactness of (5.1) together with Corollary 4.3 (and Lemma 4.4) show immediately that

$$\text{Coker}(\alpha_{[1,0]}) \cong \hat{H}^0(G_{n,m}, \mathcal{B}_0) \cong R_\chi / L_p(1, \chi). \quad (5.8)$$

Since the orders of A_n and B_n coincide for $n \geq 0$, and since these groups are isomorphic to $\hat{H}^q(G_{n,m}, Q_n)$ and $\hat{H}^q(G_{n,m}, B_n)$ respectively (see (3.4) and (5.3)), the equalities of orders

$$|\text{Ker}(\alpha_{[0,1]})| = |\text{Coker}(\alpha_{[0,1]})| \quad \text{and} \quad |\text{Ker}(\alpha_{[1,0]})| = |\text{Coker}(\alpha_{[1,0]})|$$

hold. By (5.7) and (5.8), the four groups have the same order, equal to $|L_p(1, \chi)|_p^{-d}$.

We are left with the structure of $\text{Ker}(\alpha_{[1,0]})$ and $\text{Coker}(\alpha_{[0,1]})$. The map $\alpha_{[1,0]}$ is the composition

$$\begin{array}{ccccc} \hat{H}^{-1}(G_{n,m}, Q_n) & \xrightarrow{\tilde{\beta}} & \hat{H}^0(G_{n,m}, \mathcal{O}_n^\times(\chi)) & \longrightarrow & \hat{H}^0(G_{n,m}, B_n) \\ & & \searrow & \nearrow & \\ & & \alpha_{[1,0]} & & \end{array} \quad (5.9)$$

and $\text{Ker}(\alpha_{[1,0]}) \supseteq \text{Ker}(\tilde{\beta}) = \Pi_n$, the last identification coming from Lemma 2.1 as discussed in the remark of Section 3. Combining Corollary 4.6 with the computation of the order of $\text{Ker}(\alpha_{[1,0]})$ performed above, the inclusion cannot be strict, and $\text{Ker}(\alpha_{[1,0]})$ is cyclic

over R_χ of the prescribed order. Looking at $\text{Ker}(\alpha_{[1,0]})$ and at $\text{Coker}(\alpha_{[0,1]})$ as subgroups of $\hat{H}^{-1}(G_{n,m}, \mathcal{B}_n)$ as in (5.1), and knowing the structure of this last module by Lemma 4.3, shows that $\text{Coker}(\alpha_{[0,1]})$ is cyclic, too.

Now we can single out from the proof a precise description of the kernels of the maps $\alpha_{[1,0]}$ and $\alpha_{[0,1]}$ when seen as maps

$$\alpha_{[i,j]} : A_n \longrightarrow B_n,$$

by combining (3.4) and (5.3). Observe that, by Lemma (3.1), there is an isomorphism $\hat{H}^{-1}(G_{n,m}, \text{Cyc}_n(\chi)) \cong \hat{H}^{-1}(G_{n,m}, \langle \eta_n \rangle)$, while $\hat{H}^{-1}(G_{n,m}, \langle \eta_n \rangle) \cong \hat{H}^{-1}(G_{n,m}, I_{G_n}) \cong \hat{H}^0(G_{n,m}, R_\chi)$ by (3.2) and (3.3). It is clear that these isomorphisms are not only $G_{n,m}$ -linear, but also G_n -linear and therefore that $\hat{H}^{-1}(G_{n,m}, \text{Cyc}_n(\chi))$ has trivial G_n -action.

COROLLARY 5.2. *With the same hypothesis as in the theorem,*

$$\text{Ker}(\alpha_{[0,1]}) = \text{Ker}(\alpha_{[1,0]}) = \Pi_n,$$

where Π_n is the subgroup generated in A_n by the class of any prime of F_n above p , as discussed in the remark of Section 3.

Proof. While proving the theorem we found $\text{Ker}(\alpha_{[1,0]}) = \Pi_n$, and we now focus on $\text{Ker}(\alpha_{[0,1]})$. Looking again at (5.5) we find $\text{Ker}(\alpha_{[0,1]}) \subseteq \text{Im}(\hat{H}^{-1}(G_{n,m}, \text{Cyc}_n(\chi))) \subseteq \hat{H}^{-1}(G_{n,m}, \mathcal{O}_n^\times(\chi))^{G_n}$: since the isomorphisms $\hat{H}^0(G_{n,m}, Q_n) \cong A_n$ are G_n -linear, we get $\text{Ker}(\alpha_{[0,1]}) \subseteq A_n^{G_n}$. As in the proof of Corollary (4.6), the assumption $\lambda = 0$ is equivalent to $\Pi_n = X^\Gamma$ and $X^\Gamma = A_n^{G_n}$ if n is big enough; putting all together, we have $\text{Ker}(\alpha_{[0,1]}) \subseteq \Pi_n$. Since they have the same order thanks to Corollary 4.6 together with Theorem 5.1, the inclusion turns into an equality.

Remark. As the above Corollary shows, there are indeed two maps $\alpha_{[0,1]}$ and $\alpha_{[1,0]}$ sitting in an exact sequence

$$0 \longrightarrow \Pi_n \longrightarrow A_n \xrightarrow{\alpha} B_n \longrightarrow \mathcal{B}_0/\eta_0 \longrightarrow 0,$$

where α can be either of them. This is the same as in the non-split case, where both $\alpha_{[0,1]}$ and $\alpha_{[1,0]}$ give an isomorphism $A_n \cong B_n$ for $n \gg 0$ if $\lambda = 0$ (see [KS95]).

Acknowledgements. This work is part of my PhD thesis, written under the supervision of René Schoof. I would like to take this opportunity to thank him not only for proposing me to work on this subject and for the help he gave me in writing this paper, but especially for all the time and patience he put in following me through my PhD and for the viewpoint on Mathematics he suggested to me.

It is also a pleasure to thank the anonymous referee for many useful corrections and suggestions that improved the style and the mathematics of this text.

REFERENCES

- [BNQD01] J.-R. BELLARD and T. NGUYEN QUANG DO. Formules de classes pour les corps abéliens réels. *Ann. Inst. Fourier (Grenoble)* **51** (2001), no. 4, 903–937.
- [CF86] J. W. S. CASSELS and A. FRÖHLICH (eds.). *Algebraic number theory*. (Academic Press Inc. [Harcourt Brace Jovanovich Publishers], 1986.) Reprint of the 1967 original.
- [FW79] B. FERRERO and L. C. WASHINGTON. The Iwasawa invariant μ_p vanishes for abelian number fields. *Ann. of Math.* (2) **109** (1979), no. 2, 377–395.
- [Gil79a] R. GILLARD. Remarques sur les unités cyclotomiques et les unités elliptiques. *J. Number Theory* **11** (1979), no. 1, 21–48.

- [Gil79b] R. GILLARD. Unités cyclotomiques, unités semi-locales et $\mathbf{Z}_l$ -extensions, *II. Ann. Inst. Fourier (Grenoble)* **29** (1979), no. 4, viii, 1–15.
- [Gre76] R. GREENBERG. On the Iwasawa invariants of totally real number fields. *Amer. J. Math.* **98** (1976), no. 1, 263–284.
- [Gre77] R. GREENBERG. On p -adic L -functions and cyclotomic fields, *II. Nagoya Math. J.* **67** (1977), 139–158.
- [Iwa60] K. IWASAWA. On local cyclotomic fields. *J. Math. Soc. Japan* **12** (1960), 16–21.
- [Iwa73] K. IWASAWA. On $\mathbf{Z}_l$ -extensions of algebraic number fields. *Ann. of Math. (2)* **98** (1973), 246–326.
- [KS95] J. S. KRAFT and R. SCHOOF. Computing Iwasawa modules of real quadratic number fields. *Compositio Math.* **97** (1995), no. 1-2, 135–155, Special issue in honour of Frans Oort.
- [Kuz96] L. V. KUZ'MIN. On formulas for the class number of real abelian fields. *Izv. Ross. Akad. Nauk Ser. Mat.* **60** (1996), no. 4, 43–110.
- [Lan90] S. LANG. *Cyclotomic fields I and II*, second ed., Graduate Texts in Mathematics, vol. 121. (Springer-Verlag, 1990) With an appendix by Karl Rubin.
- [NSW00] J. NEUKIRCH, A. SCHMIDT and K. WINGBERG. *Cohomology of number fields*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 323, (Springer-Verlag, 2000).
- [Oza97] M. OZAKI. On the cyclotomic unit group and the ideal class group of a real abelian number field, *I, II. J. Number Theory* **64** (1997), no. 2, 211–222, 223–232.
- [Sch88] R. SCHOOF. Cohomology of class groups of cyclotomic fields: an application to Morse-Smale diffeomorphisms. *J. Pure Appl. Algebra* **53** (1988), no. 1-2, 125–137. MR MR955614 (89j:11111)
- [Sin81] W. SINNOTT. On the Stickelberger ideal and the circular units of an abelian field. *Invent. Math.* **62** (1980/81), no. 2, 181–234.
- [Was97] L. C. WASHINGTON. *Introduction to cyclotomic fields*, second ed. Graduate Texts in Mathematics, vol. 83 (Springer-Verlag, 1997).