



HAL
open science

Convolution of orbital measures on symmetric spaces of type C_p and D_p

Piotr Graczyk, Patrice Sawyer

► **To cite this version:**

Piotr Graczyk, Patrice Sawyer. Convolution of orbital measures on symmetric spaces of type C_p and D_p . 2014. hal-00965263

HAL Id: hal-00965263

<https://hal.science/hal-00965263>

Preprint submitted on 24 Mar 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

CONVOLUTION OF ORBITAL MEASURES ON SYMMETRIC SPACES OF TYPE C_p AND D_p

P. GRACZYK AND P. SAWYER

ABSTRACT. We study the absolute continuity of the convolution $\delta_{e^X}^{\natural} \star \delta_{e^Y}^{\natural}$ of two orbital measures on the symmetric spaces $\mathbf{SO}_0(p, p)/\mathbf{SO}(p) \times \mathbf{SO}(p)$, $\mathbf{SU}(p, p)/\mathbf{S}(\mathbf{U}(p) \times \mathbf{U}(p))$ and $\mathbf{Sp}(p, p)/\mathbf{Sp}(p) \times \mathbf{Sp}(p)$. We prove sharp conditions on $X, Y \in \mathfrak{a}$ for the existence of the density of the convolution measure. This measure intervenes in the product formula for the spherical functions.

1. INTRODUCTION

The spaces $G/K = \mathbf{SO}_0(p, p)/\mathbf{SO}(p) \times \mathbf{SO}(p)$ are Riemannian symmetric spaces of non-compact type corresponding to root systems of type D_p . The spaces $\mathbf{SU}(p, p)/\mathbf{S}(\mathbf{U}(p) \times \mathbf{U}(p))$ and $\mathbf{Sp}(p, p)/\mathbf{Sp}(p) \times \mathbf{Sp}(p)$ correspond to root systems of type C_p .

Consider $X, Y \in \mathfrak{a}$ and let m_K denote the Haar measure of the group K . We define $\delta_{e^X}^{\natural} = m_K \star \delta_{e^X} \star m_K$. It is the uniform measure on the orbit Ke^XK . The problem of the absolute continuity of the convolution

$$m_{X, Y} = \delta_{e^X}^{\natural} \star \delta_{e^Y}^{\natural}$$

of two orbital measures that we address in our paper has important applications in harmonic analysis of spherical functions on G/K and in probability theory. Let λ be a complex-valued linear form on \mathfrak{a} and $\phi_\lambda(e^X)$ be the spherical function, which is the spherical Fourier transform of the measure $\delta_{e^X}^{\natural}$. The product formula for the spherical functions states that

$$\phi_\lambda(e^X) \phi_\lambda(e^Y) = \int_{\mathfrak{a}} \phi_\lambda(e^H) d\mu_{X, Y}(H)$$

where $\mu_{X, Y}$ is the projection of the measure $m_{X, Y}$ on \mathfrak{a} via the Cartan decomposition $G = KAK$. Then the existence of a density of $\mu_{X, Y}$, equivalent to the absolute continuity of $m_{X, Y}$, is of great importance.

It was proven in [3] that as soon as the space G/K is irreducible and one of the elements X, Y is regular and the other nonzero, then the convolution $\delta_{e^X}^{\natural} \star \delta_{e^Y}^{\natural}$ has a density. The density can however still exist when both X and Y are singular. It is a challenging problem to characterize all such pairs X, Y .

This problem was solved for symmetric spaces of type A_n and for the exceptional space $\mathbf{SL}(3, \mathbf{O})/\mathbf{SU}(3, \mathbf{O})$ of type E_6 in [4], and for symmetric spaces of type B_p and BC_p in [6]. In the present paper, we present the solution of the problem for Riemannian symmetric spaces of type C_p and D_p .

For the good comprehension of the methods of this paper, it is useful to know the paper [6]. However, the cases C_p and D_p require many original ideas that did not appear in the case B_p . We refer to S. Helgason's books [7] and [8] for the standard notation and results.

In Section 2, we are reminding the reader of the basic information about the Lie group $\mathbf{SO}(p, p)$ and its Lie algebra $\mathfrak{so}(p, p)$. We also provide the necessary notation to describe the configuration of an element of the Cartan subalgebra \mathfrak{a} of $\mathfrak{so}(p, p)$. This configuration notion allows us to "measure" how singular an element of \mathfrak{a} is and to describe in a precise manner which pairs $X, Y \in \mathfrak{a}$ are "eligible", the sharp criterion that we establish in the paper for the absolute continuity of $\mu_{X, Y}$. The following theorem is the main result of the paper:

Theorem A. *Let $G/K = \mathbf{SO}_0(p, p)/\mathbf{SO}(p) \times \mathbf{SO}(p)$ and $X, Y \in \mathfrak{a}$. The density of the convolution $\delta_{e^X}^{\natural} \star \delta_{e^Y}^{\natural}$ exists if and only if X and Y are eligible (see Definition 2.4).*

In the following Section 3, a series of definitions and accessory results are given to set the stage for the proof of Theorem A. In Section 4, we show that (X, Y) has to be an eligible pair for the measure $\mu_{X, Y}$ to be absolutely continuous. In Section 5, we then show that the eligibility condition is also sufficient.

1991 *Mathematics Subject Classification.* Primary 43A90, Secondary 53C35.

Key words and phrases. root systems, spherical functions, symmetric spaces, non-compact type.

Supported by l'Agence Nationale de la Recherche ANR-09-BLAN-0084-01.

In the last section, as in [4] and [6], we extend our results to the complex and quaternionic cases. Again, the richness of the root structure comes into play: in the table in Remark 6.1, we find that the complex and quaternionic cases have much more in common with the cases $q > p$ than with the real case $\mathbf{SO}(p, p)$.

We conclude the paper with a discussion of the absolute continuity of powers of $\delta_{e^X}^h$ for a nonzero $X \in \mathfrak{a}$.

2. PRELIMINARIES AND DEFINITIONS

We start by reviewing some useful information on the Lie group $\mathbf{SO}_0(p, p)$, its Lie algebra $\mathfrak{so}(p, p)$ and the corresponding root system. Most of this material was given in [9]. For the convenience of the reader, we gather below the properties we will need in the sequel. In this paper, E_{ij} is a rectangular matrix with 0's everywhere except at the position (i, j) where it is 1. Recall that $\mathbf{SO}(p, p)$ is the group of matrices $g \in \mathbf{SL}(2p, \mathbf{R})$ such that $g^T I_{p,p} g = I_{p,p}$ where $I_{p,p} = \begin{bmatrix} -I_p & 0_{p \times p} \\ 0_{p \times p} & I_p \end{bmatrix}$. Unless otherwise specified, all 2×2 block decompositions in this paper follow the same pattern. The group $\mathbf{SO}_0(p, p)$ is the connected component of $\mathbf{SO}(p, p)$ containing the identity. The Lie algebra $\mathfrak{so}(p, p)$ of $\mathbf{SO}_0(p, p)$ consists of the matrices

$$\begin{bmatrix} A & B \\ B^T & D \end{bmatrix}$$

where A and D are skew-symmetric. A very important element in our investigations is the Cartan decomposition of $\mathfrak{so}(p, p)$ and $\mathbf{SO}(p, p)$. The maximal compact subgroup K is the subgroup of $\mathbf{SO}(p, p)$ consisting of the matrices

$$\begin{bmatrix} A & 0 \\ 0 & D \end{bmatrix}$$

of size $(2p) \times (2p)$ such that $A, D \in \mathbf{SO}(p)$ (hence $K \simeq \mathbf{SO}(p) \times \mathbf{SO}(p)$). If \mathfrak{k} is the Lie algebra of K and \mathfrak{p} is the set of matrices

$$(2.1) \quad \begin{bmatrix} 0 & B \\ B^T & 0 \end{bmatrix}$$

then the Cartan decomposition is given by $\mathfrak{so}(p, p) = \mathfrak{k} \oplus \mathfrak{p}$ with corresponding Cartan involution $\theta(X) = -X^T$. To shorten the notation, for $X \in \mathfrak{p}$ as in (2.1), we will write $X^s = B$.

The Cartan space $\mathfrak{a} \subset \mathfrak{p}$ is the set of matrices

$$H = \begin{bmatrix} 0_{p \times p} & \mathcal{D}_H \\ \mathcal{D}_H & 0_{p \times p} \end{bmatrix}$$

where $\mathcal{D}_H = \text{diag}[H_1, \dots, H_p]$. Its canonical basis is given by the matrices

$$A_i := E_{i,p+i} + E_{p+i,i}, 1 \leq i \leq p.$$

The restricted roots and associated root vectors for the Lie algebra $\mathfrak{so}(p, p)$ with respect to \mathfrak{a} are given in Table 1. The positive roots can be chosen as $\alpha(H) = H_i \pm H_j$, $1 \leq i < j \leq p$. The simple roots are given by

root α	multiplicity	root vectors X_α
$\alpha(H) = \pm(H_i - H_j)$ $1 \leq i, j \leq p, i < j$	1	$Y_{i,j}^\pm = \pm(E_{i,j} - E_{j,i} + E_{p+i,p+j} - E_{p+j,p+i}) + E_{i,p+j} + E_{p+j,i}$ $+ E_{j,p+i} + E_{p+i,j}$
$\alpha(H) = \pm(H_i + H_j)$ $1 \leq i, j \leq p, i < j$	1	$Z_{i,j}^\pm = \pm(E_{i,j} - E_{j,i} - E_{p+i,p+j} + E_{p+j,p+i}) - (E_{i,p+j} + E_{p+j,i})$ $+ E_{j,p+i} + E_{p+i,j}$

TABLE 1. Restricted roots and associated root vectors

$\alpha_i(H) = H_i - H_{i+1}$, $i = 1, \dots, p-1$ and $\alpha_p(H) = H_{p-1} + H_p$. We therefore have the positive Weyl chamber

$$\mathfrak{a}^+ = \{H \in \mathfrak{a} : H_1 > H_2 > \dots > H_{p-1} > |H_p|\}.$$

The elements of the Weyl group W act as permutations of the diagonal entries of \mathcal{D}_X with eventual sign changes of any even number of these entries. The Lie algebra \mathfrak{k} is generated by the vectors $X_\alpha + \theta X_\alpha$. We will use the notation

$$k_{X_\alpha}^t = e^{t(X_\alpha + \theta X_\alpha)}.$$

The linear space \mathfrak{p} has a basis formed by $A_i \in \mathfrak{a}$, $1 \leq i \leq p$ and by the symmetric matrices $X_\alpha^s := \frac{1}{2}(X_\alpha - \theta X_\alpha)$ which have the following form

$$\begin{aligned} Y_{i,j} &:= E_{i,p+j} + E_{j,p+i} + E_{p+j,i} + E_{p+i,j}, & 1 \leq i < j \leq p; \\ Z_{i,j} &:= -E_{i,p+j} + E_{j,p+i} - E_{p+j,i} + E_{p+i,j}, & 1 \leq i < j \leq p. \end{aligned}$$

Thus, if $X \in \mathfrak{p}$ is as in (2.1), then the vectors A_i generate the diagonal entries of $B = X^s$ and $Y_{i,j}$ and $Z_{i,j}$ the non-diagonal entries.

We now recall the useful matrix $S \in \mathbf{SO}(p+q)$ which allows us to diagonalize simultaneously all the elements of \mathfrak{a} . Let

$$S = \begin{bmatrix} \frac{\sqrt{2}}{2} I_p & \frac{\sqrt{2}}{2} J_p \\ \frac{\sqrt{2}}{2} I_p & -\frac{\sqrt{2}}{2} J_p \end{bmatrix}$$

where $J_p = (\delta_{i,p+1-i})$ is a matrix of size $p \times p$. If $H = \begin{bmatrix} 0 & \mathcal{D}_H \\ \mathcal{D}_H & 0 \end{bmatrix}$ with $\mathcal{D}_H = \text{diag}[H_1, \dots, H_p]$ then $S^T H S = \text{diag}[H_1, \dots, H_p, -H_p, \dots, -H_1]$. The ‘‘group’’ version of this result is as follows:

$$S^T e^H S = \text{diag}[e^{H_1}, \dots, e^{H_p}, e^{-H_p}, \dots, e^{-H_1}].$$

Remark 2.1. The Cartan projection $a(g)$ on the group $\mathbf{SO}_0(p, p)$, defined as usual by

$$g = k_1 e^{a(g)} k_2, \quad a(g) \in \overline{\mathfrak{a}^+}, k_1, k_2 \in K$$

is related to the singular values of $g \in \mathbf{SO}(p, p)$ in the following way. Recall that the singular values of g are defined as the non-negative square roots of the eigenvalues of $g^T g$. Let us write $H = a(g)$. We have

$$g^T g = k_2^T e^{2H} k_2 = (k_2^T S) (S^T e^{2H} S) (S^T k_2)$$

where $S^T e^{2H} S$ is a diagonal matrix with nonzero entries $e^{2H_1}, \dots, e^{2H_p}, e^{-2H_p}, \dots, e^{-2H_1}$, satisfying $H_1 \geq \dots \geq H_{p-1} \geq |H_p|$. Let us write $a_j = e^{H_j}$ for $j = 1, \dots, p-1$ and $a_p = e^{|H_p|}$. Thus the set of $2p$ singular values of g contains the values $a_1 \geq \dots \geq a_p \geq a_p^{-1} \geq \dots \geq a_1^{-1}$ with $a_1 \geq \dots \geq a_p \geq 1$.

Then

$$a(g) = \begin{bmatrix} 0 & \mathcal{D}_{a(g)} \\ \mathcal{D}_{a(g)} & 0 \end{bmatrix} \quad \text{with} \quad \mathcal{D}_{a(g)} = \text{diag}[\log a_1, \dots, \log a_{p-1}, \text{sgn}(H_p) \log a_p].$$

Note that this method does not allow us to distinguish between the situations where H_p is positive or negative.

Singular elements of \mathfrak{a} . In what follows, we will consider singular elements $X, Y \in \partial \mathfrak{a}^+$. We need to control the irregularity of X and Y , i.e. consider the simple positive roots annihilating X and Y . A special role is played by the last simple root $\alpha_p = H_{p-1} + H_p$, different from the simple roots $\alpha_i(H) = H_i - H_{i+1}$, $i = 1, \dots, p-1$. Note that $\alpha_p(X) = 0$ implies that the last diagonal entry of \mathcal{D}_X is negative or 0.

We introduce the following definition of the configuration of $X \in \overline{\mathfrak{a}^+}$.

Definition 2.2. Let $X \in \overline{\mathfrak{a}^+}$. In what follows, $x_i > 0$, $x_i > x_j$ for $i < j$, $s_i \geq 1$, $u \geq 0$ and $\sum_{i=1}^r s_i + u = p$. Let $\mathbf{s} = (s_1, \dots, s_r)$. We define the configuration of X by:

$$(2.2) \quad [\mathbf{s}; u] \quad \text{if} \quad \mathcal{D}_X = \text{diag} \left[\overbrace{x_1, \dots, x_1}^{s_1}, \overbrace{x_2, \dots, x_2}^{s_2}, \dots, \overbrace{x_r, \dots, x_r}^{s_r}, \overbrace{0, \dots, 0}^u \right]$$

$$(2.3) \quad [\mathbf{s}] \quad \text{if} \quad \mathcal{D}_X = \text{diag} \left[\overbrace{x_1, \dots, x_1}^{s_1}, \overbrace{x_2, \dots, x_2}^{s_2}, \dots, \overbrace{x_{r-1}, \dots, x_{r-1}}^{s_{r-1}}, \overbrace{-x_r}^{s_r=1} \right]$$

$$(2.4) \quad [\mathbf{s}]^- \quad \text{if} \quad \mathcal{D}_X = \text{diag} \left[\overbrace{x_1, \dots, x_1}^{s_1}, \overbrace{x_2, \dots, x_2}^{s_2}, \dots, \overbrace{x_r, \dots, x_r, -x_r}^{s_r} \right].$$

We extend naturally the definition of configuration to any $X \in \mathfrak{a}$, whose configuration is defined as that of the projection $\pi(X)$ of X on $\overline{\mathfrak{a}^+}$.

Remark 2.3. We will often write $X = X[\dots]$ when $[\dots]$ is a configuration of X . For the configuration (2.4), we write the $-$ superscript in $X[\mathbf{s}]^-$ to indicate that X contains nonzero opposite entries. We omit the $-$ superscript in (2.3) and write $X[\mathbf{s}]$ because we are essentially in the same case as in (2.2) with $u = 0$ and $s_r = 1$.

If the number of zero entries $u = 0$, it may be omitted in (2.2). In particular, in the configurations (2.3) and (2.4), $u = 0$. Note that $X = 0$ has configuration $[0; p]$. A regular $X \in \mathfrak{a}^+$ has the configuration $[1^p; 0]$ or $[1^{p-1}; 1]$, where $1^k = (1, \dots, 1)$ with 1 repeated k times.

In what follows, we will write $\max \mathbf{s} = \max_i s_i$ and $\max(\mathbf{s}, u) = \max(\max \mathbf{s}, u)$. We will show that in the case of the symmetric spaces $\mathbf{SO}_0(p, p)/\mathbf{SO}(p) \times \mathbf{SO}(p)$, the criterion for the existence of the density of the convolution $\delta_{e^{\mathbf{h}}_X}^{\mathbf{h}} \star \delta_{e^{\mathbf{h}}_Y}^{\mathbf{h}}$ is given by the following definition of an eligible pair X and Y :

Definition 2.4. Let X and Y be two elements of \mathfrak{a} with configurations $[\mathbf{s}; u]$ or $[\mathbf{s}]^-$ and $[\mathbf{t}; v]$ or $[\mathbf{t}]^-$ respectively. We say that X and Y are eligible if one of the two following cases holds

$$(2.5) \quad u \leq 1, v \leq 1 \quad \text{and} \quad \max(\mathbf{s}) + \max(\mathbf{t}) \leq 2p - 2,$$

$$(2.6) \quad u \geq 2 \text{ or } v \geq 2 \quad \text{and} \quad \max(\mathbf{s}, 2u) + \max(\mathbf{t}, 2v) \leq 2p$$

and, for $p = 4$,

$$(2.7) \quad \{\mathcal{D}_X, \mathcal{D}_Y\} \neq \{\text{diag}[a, a, a, a], \text{diag}[b, b, c, c]\} \text{ nor } \{\text{diag}[a, a, a, -a], \text{diag}[b, b, c, -c]\},$$

for any $a \neq 0, b \neq 0, |b| \neq |c|$.

Observe that if X and Y are eligible, then $X \neq 0$ and $Y \neq 0$. The non-eligible pairs given by (2.7) are $\{[4], [(2, 2)]\}, \{[4]^-, [(2, 2)]^-\}$ with $u = v = 0$ or $\{[4], [2; 2]\}$ and $\{[4]^-, [2; 2]\}$ with $u = 0$ and $v = 2$.

Remark 2.5. It is interesting to note that the definition of eligible pairs is more complicated for the space $\mathbf{SO}(p, p)$ than for the spaces $\mathbf{SO}(p, q)$ with $p < q$ (recall that the latter spaces have a much richer root structure). As for the spaces $\mathbf{SO}(p, q)$ with $p < q$, the number of zeroes on the diagonal of \mathcal{D}_X is important. Unlike in the case $\mathbf{SO}(p, q)$ with $p < q$, this only becomes a factor when the number of zeroes is greater than 1 (as opposed to greater than 0).

In [6], we showed that if $p < q$ and X and $Y \in \mathfrak{a}$ were such that the \mathcal{D}_X and \mathcal{D}_Y have no zero diagonal elements on the diagonal then $\mu_{X, Y}$ was absolutely continuous. This is no longer the case when $p = q$ and this is one of main differences between the $\mathbf{SO}(p, p)$ and $\mathbf{SO}(p, q)$ cases. Another difference is the anomalous case when $p = 4$ seen in (2.7) and the fact that lower dimensional cases require different proofs. On the other hand, when the number of zeroes on the diagonal of either \mathcal{D}_X or \mathcal{D}_Y is at least 2, then the proof of Theorem A is similar to the one found in [6] but requires considering separately a low dimension case $X[5], Y[3; 2]$.

3. BASIC TOOLS AND REDUCTIONS

Definition 3.1. For $Z \in \mathfrak{a}$, let V_Z be the subspace of \mathfrak{p} defined by

$$V_Z = \text{span}\{X_\alpha - \theta(X_\alpha) \mid \alpha(Z) \neq 0\} \subset \mathfrak{p}.$$

We denote by $|V_Z|$ the dimension of V_Z . It equals the number of positive roots α such that $\alpha(Z) \neq 0$.

The following definition and lemmas will help reduce the number of cases of configurations of (X, Y) to verify.

Definition 3.2. We will say that X and $X' \in \mathfrak{a}$ are relatives if exactly one of the diagonal entries of \mathcal{D}_X and $\mathcal{D}_{X'}$ differs by sign. If X is a relative to X' and Y is a relative to Y' then we will say that (X, Y) is a relative pair of (X', Y') .

The properties listed in the following lemma are straightforward.

Lemma 3.3.

- (1) X is a relative of X' if and only if \mathcal{D}_X has at least one diagonal entry equal to 0, i.e. $u \geq 1$ in $X[\mathbf{s}; u]$. Thus $X \in \overline{\mathfrak{a}^+}$ has no other relatives in $\overline{\mathfrak{a}^+}$.
- (2) If X and X' are relatives then $|V_X| = |V_{X'}|$.
- (3) If (X, Y) and (X', Y') are relative pairs then (X, Y) is an eligible pair if and only if (X', Y') is an eligible pair.
- (4) If $X = X[\mathbf{s}; u] \in \overline{\mathfrak{a}^+}$ with $u > 0$ then all X_i 's are non-negative.
- (5) If $X = X[\mathbf{s}; u], Y[\mathbf{t}; v] \in \overline{\mathfrak{a}^+}$ are such that $u > 0$ or $v > 0$ then either $\mathcal{D}_X, \mathcal{D}_Y$ have no negative entries or we can choose a relative pair $X', Y' \in \overline{\mathfrak{a}^+}$ in such a way that $\mathcal{D}_{X'}, \mathcal{D}_{Y'}$ have no negative entries.
- (6) If X is a relative of X' and X' is a relative of X'' then X and X'' are in the same Weyl-group orbit.

Lemma 3.4. If (X, Y) and (X', Y') are relative pairs then either both sets $Ke^X Ke^Y K$ and $Ke^{X'} Ke^{Y'} K$ have nonempty interiors or neither has.

Proof. Let $J_0 = \text{diag}\{\overbrace{1, \dots, 1}^{2p-1}, -1\}$ and note that J_0 is orthogonal and that $J_0^2 \in \mathbf{SO}(p) \times \mathbf{SO}(p)$. Suppose that X, Y, X' and Y' are as in the statement of the lemma. Suppose that $w_1, w_2 \in W \subset K$ are such that the element of $\mathcal{D}_{w_1 X'}$ (resp. $\mathcal{D}_{w_2 Y'}$) that differs by a sign from the corresponding element of \mathcal{D}_X (resp. \mathcal{D}_Y) is placed at the end. Then

$$\begin{aligned} Ke^X Ke^Y K &= J_0 K J_0 w_1 e^X w_1^{-1} J_0 K J_0 w_2 e^Y w_2^{-1} J_0 K J_0 \\ &= J_0 K (J_0 w_1 e^X w_1^{-1} J_0) K (J_0 w_2 e^Y w_2^{-1} J_0) K J_0 = J_0 K e^{X'} K e^{Y'} K J_0 \end{aligned}$$

which allows us to conclude. \square

In the sequel we use some ideas, results and notations of [4, Section 3], that we strengthen and complete.

Proposition 3.5.

- (i) *The density of the measure $m_{X,Y}$ exists if and only if its support Ke^XKe^YK has nonempty interior.*
- (ii) *Consider the analytic map $T: K \times K \times K \rightarrow G$ defined by*

$$T(k_1, k_2, k_3) = k_1 e^X k_2 e^Y k_3.$$

The set $T(K \times K \times K) = Ke^XKe^YK$ contains an open set if and only if the derivative of T is surjective for some choice of $\mathbf{k} = (k_1, k_2, k_3)$.

Proof. Part (i) follows from arguments explained in [3] in the case of the support of the measure $\mu_{X,Y}$, equal to $a(e^XKe^Y)$. Part (ii) is justified for example in [8, p. 479]. \square

Corollary 3.6. *Let (X, Y) and (X', Y') be relative pairs. The measure $m_{X,Y}$ is absolutely continuous if and only if the measure $m_{X',Y'}$ is absolutely continuous.*

Proof. We use Proposition 3.5(i) and Lemma 3.4. \square

Proposition 3.7. *Let $U_Z = \mathfrak{k} + \text{Ad}(e^Z)\mathfrak{k}$. The measure $m_{X,Y}$ is absolutely continuous if and only if there exists $k \in K$ such that*

$$(3.1) \quad U_{-X} + \text{Ad}(k)U_Y = \mathfrak{g}.$$

Proof. We want to show that this condition is equivalent to the derivative of T at \mathbf{k} being surjective. We have

$$(3.2) \quad \begin{aligned} dT_{\mathbf{k}}(A, B, C) &= \left. \frac{d}{dt} \right|_{t=0} e^{tA} k_1 e^X e^{tB} k_2 e^Y e^{tC} k_3 \\ &= A k_1 e^X k_2 e^Y k_3 + k_1 e^X B k_2 e^Y k_3 + k_1 e^X k_2 e^Y C k_3 \end{aligned}$$

We now transform the space of all matrices of the form (3.2) without modifying its dimension:

$$\begin{aligned} &\dim\{A k_1 e^X k_2 e^Y k_3 + k_1 e^X B k_2 e^Y k_3 + k_1 e^X k_2 e^Y C k_3 : A, B, C \in \mathfrak{k}\} \\ &= \dim\{k_1^{-1} A k_1 e^X k_2 e^Y + e^X B k_2 e^Y + e^X k_2 e^Y C : A, B, C \in \mathfrak{k}\} \\ &= \dim\{A e^X k_2 e^Y + e^X B k_2 e^Y + e^X k_2 e^Y C : A, B, C \in \mathfrak{k}\} \\ &= \dim\{e^{-X} A e^X + B + k_2 e^Y C e^{-Y} k_2^{-1} : A, B, C \in \mathfrak{k}\} \end{aligned}$$

The space in the last line equals $\mathfrak{k} + \text{Ad}(e^{-X})(\mathfrak{k}) + \text{Ad}(k_2)(\text{Ad}(e^Y)(\mathfrak{k})) = U_{-X} + \text{Ad}(k_2)U_Y$. \square

In order to apply the condition (3.1), we will consider convenient symmetrized root vectors and the spaces V_Z generated by them.

Lemma 3.8. *Let $Z \in \mathfrak{a}$. Then $U_Z = \mathfrak{k} + V_Z = U_{-Z}$.*

Proof. Clearly $V_Z = V_{-Z}$. We show that $V_Z \subset U_Z$ and therefore that $\mathfrak{k} + V_Z \subset U_Z$: let α be a root such that $\alpha(Z) \neq 0$. Note that $[Z, X_\alpha] = \alpha(Z) X_\alpha$ and $[Z, \theta(X_\alpha)] = -\alpha(Z) \theta(X_\alpha)$. Let $U = X_\alpha + \theta(X_\alpha) \in \mathfrak{k}$. Now,

$$\text{Ad}(e^Z)U = e^{\text{ad}Z}(X_\alpha + \theta(X_\alpha)) = e^{\alpha(Z)} X_\alpha + e^{-\alpha(Z)} \theta(X_\alpha).$$

Therefore $X_\alpha = (e^{\alpha(Z)} - e^{-\alpha(Z)})^{-1} (-e^{-\alpha(Z)} U + \text{Ad}(e^Z)U) \in \mathfrak{k} + \text{Ad}(e^Z)(\mathfrak{k}) = U_Z$. The vector θX_α is a root vector for the root $-\alpha$, so we also have $\theta X_\alpha \in U_Z$.

It remains to show that $U_Z \subset \mathfrak{k} + V_Z$. It suffices to show that $\text{Ad}(e^Z)\mathfrak{k} \subset \mathfrak{k} + V_Z$: for every α , $\text{Ad}(e^Z)(X_\alpha + \theta(X_\alpha)) = e^{\alpha(Z)} X_\alpha + e^{-\alpha(Z)} \theta(X_\alpha) = \frac{e^{\alpha(Z)} + e^{-\alpha(Z)}}{2} (X_\alpha + \theta(X_\alpha)) + \frac{e^{\alpha(Z)} - e^{-\alpha(Z)}}{2} (X_\alpha - \theta(X_\alpha)) \in \mathfrak{k} + V_Z$. \square

The following corollary is then straightforward:

Corollary 3.9. *The measure $m_{X,Y}$ is absolutely continuous if and only if there exists $k \in K$ such that*

$$(3.3) \quad V_X + \text{Ad}(k)V_Y = \mathfrak{p}.$$

Corollary 3.10. *The measure $m_{X,Y}$ is absolutely continuous if and only if there exists a dense open subset $U \subset K$ such that for every $k \in U$*

- (1) $V_{w_1 \cdot X} + \text{Ad}(k)V_{w_2 \cdot Y} = \mathfrak{p}$ for every $w_1, w_2 \in W$,
- (2) For every $r < 2p$, the matrix obtained by removing the first r rows and r columns of k is non-singular.

(1) For the root vectors $Z_{i,j}^+$ and $Y_{i,j}^+$, we have

$$\begin{aligned} \text{Ad}(e^{t(Y_{i,j}^+ + \theta(Y_{i,j}^+))})(Y_{i,j}) &= \cos(4t) Y_{i,j} + 2 \sin(4t) (A_i - A_j), \\ \text{Ad}(e^{t(Z_{i,j}^+ + \theta(Z_{i,j}^+))})(Z_{i,j}) &= \cos(4t) Z_{i,j} + 2 \sin(4t) (A_i + A_j). \end{aligned}$$

(2) The operators $\text{Ad}(e^{t(Y_{i,j}^+ + \theta(Y_{i,j}^+))})$ and $\text{Ad}(e^{t(Z_{i,j}^+ + \theta(Z_{i,j}^+))})$ applied to the other symmetrized root vectors do not produce any components in \mathfrak{a} .

In the proof of the existence of the density for the pairs $X[4], Y[2, 2]^-$ and $X[5], Y[3, 2]$ without predecessors, we will need the following elementary lemma. Recall that $Z_{k,l} = \frac{1}{2}(Z_{k,l}^+ - \theta(Z_{k,l}^+)) \in \mathfrak{p}$.

Lemma 3.20. *Let $1 \leq i < j \leq p$ and $1 \leq k < l \leq p$. We have*

$$[Z_{i,j}^+ + \theta(Z_{i,j}^+), Z_{k,l}] = \begin{cases} 0 & \text{if } \{i, j\} \cap \{k, l\} = \emptyset, \\ 4(A_i + A_j) & \text{if } \{i, j\} = \{k, l\}, \\ 2Y_{\min(j,l), \max(j,l)} & \text{if } i = k, j \neq l, \\ 2Y_{\min(i,k), \max(i,k)} & \text{if } i \neq k, j = l, \\ -2Y_{i,l} & \text{if } j = k, \\ -2Y_{k,j} & \text{if } i = l. \end{cases}$$

Proof. We apply the well known fact that $[\mathfrak{g}_\alpha, \mathfrak{g}_\beta] \subset \mathfrak{g}_{\alpha+\beta}$ when $\alpha + \beta$ is a root and $[\mathfrak{g}_\alpha, \mathfrak{g}_\beta] = 0$ otherwise. For the computation of exact coefficients in the formulas, we use Table 1. \square

4. NECESSITY OF THE ELIGIBILITY CONDITION

Proposition 4.1. *If $X = X[\mathbf{s}; u]$ and $Y = Y[\mathbf{t}; v] \in \mathfrak{a}$ with $u \leq 1$ and $v \leq 1$ are such that $\max \mathbf{s} + \max \mathbf{t} > 2p - 2$ then $|V_X| + |V_Y| < p^2$.*

Proof. Assume that $X = X[\mathbf{s}; u]$ and $Y = Y[\mathbf{t}; v] \in \bar{\mathfrak{a}}^+$. Without loss of generality, assume that $\max \mathbf{s} \geq \max \mathbf{t}$. We then have $\max \mathbf{s} = p$ and $\max \mathbf{t} \geq p - 1$. The only possible pairs are

$$(4.1) \quad \begin{aligned} &X[p], Y[p] \text{ and the relative pair } X'[p]^-, Y'[p]^- \\ &X[p], Y[p-1, 1] \text{ and the relative pair } X'[p]^-, Y'[1, p-1]^- \\ &X[p], Y[p]^- \text{ and the relative pair } (X', Y') = (Y, X) \\ &X[p], Y[1, p-1]^- \text{ and the relative pair } X'[p]^-, Y'[p-1, 1]^- \end{aligned}$$

By Remark 3.11 we do not need to consider the configuration $[1, p-1]$. We have $|V_X| = \frac{p(p-1)}{2}$ and $|V_{Y[p-1, 1]}| = \frac{p(p-1)}{2} + p - 1$. We apply Lemma 3.3 and we find by examination that $|V_X| + |V_Y| \leq p^2 - 1$ in all cases. \square

Corollary 4.2. *Let $p \geq 2$. Consider a pair $X = X[\mathbf{s}; u]$ and $Y = Y[\mathbf{t}; v]$ with $u \leq 1$ and $v \leq 1$. Then $|V_X| + |V_Y| \geq p^2$ if and only if*

$$(4.2) \quad \max(\mathbf{s}) + \max(\mathbf{t}) \leq 2p - 2.$$

Proof. By Proposition 4 only the sufficiency of condition (4.2) needs to be proven. Suppose $\max(\mathbf{s}) + \max(\mathbf{t}) \leq 2p - 2$ and that $\max \mathbf{s} \geq \max \mathbf{t}$. If $\max \mathbf{s} = p$ then $|V_X| = p(p-1)/2$ and $\max \mathbf{t} \leq p-2$ implies $|V_Y| \geq p(p-1)/2 + 2(p-2)$. If both $\max \mathbf{s} \leq p-1$ and $\max \mathbf{t} \leq p-1$ then $|V_X| \geq p(p-1)/2 + p-1$ and $|V_Y| \geq p(p-1)/2 + p-1$. In both cases, the result follows. \square

Definition 4.3. *We will call **exceptional** the set of configurations listed in (4.1) and denote it by \mathcal{E} .*

Proposition 4.4. *Let $X, Y \in \mathfrak{a}$ be such that $\mathcal{D}_X = \text{diag}[b, b, c, c]$ with $b > c > 0$ and $\mathcal{D}_Y = \text{diag}[a, a, a, a]$ with $a > 0$. Then $\delta_{e_X}^{\natural} \star \delta_{e_Y}^{\natural}$ has no density.*

Proof. According to Corollary 3.16, we can write

$$\begin{aligned} K e^X K e^Y K &= K e^X \begin{bmatrix} A & 0 \\ 0 & A \end{bmatrix} k_{Z_{1,2}^+}^{t_1} k_{Z_{3,4}^+}^{t_2} \begin{bmatrix} C & 0 \\ 0 & C \end{bmatrix} e^Y K \\ &= K e^X \exp \left(\begin{bmatrix} 0 & R & 0 & 0 \\ -R & 0 & 0 & 0 \\ 0 & 0 & 0 & R \\ 0 & 0 & -R & 0 \end{bmatrix} \right) k_{Z_{1,2}^+}^{t_1} k_{Z_{3,4}^+}^{t_2} e^Y K = K e^X k_{Y_{1,3}^+}^{r_1} k_{Y_{2,4}^+}^{r_2} k_{Z_{1,2}^+}^{t_1} k_{Z_{3,4}^+}^{t_2} e^Y K \end{aligned}$$

(here $Z_{1,2}^+, Z_{3,4}^+, Y_{1,3}^+, Y_{2,4}^+$ are exactly as in Table 1). We used the fact that e^Y and $\begin{bmatrix} C & 0 \\ 0 & C \end{bmatrix}$ commute, the Cartan decomposition $A = \begin{bmatrix} A_1 & 0 \\ 0 & A_2 \end{bmatrix} \exp\left(\begin{bmatrix} 0 & R \\ -R & 0 \end{bmatrix}\right) \begin{bmatrix} C_1 & 0 \\ 0 & C_2 \end{bmatrix}$ ($R = \text{diag}[r_1, r_2]$, $A_i, C_i \in \mathbf{SO}(2)$), and the facts that e^X , $\begin{bmatrix} A_1 & 0 & 0 & 0 \\ 0 & A_2 & 0 & 0 \\ 0 & 0 & A_1 & 0 \\ 0 & 0 & 0 & A_2 \end{bmatrix}$ commute and $\begin{bmatrix} C_1 & 0 & 0 & 0 \\ 0 & C_2 & 0 & 0 \\ 0 & 0 & C_1 & 0 \\ 0 & 0 & 0 & C_2 \end{bmatrix}$ commutes with $k_{Z_{1,2}^+}^{t_1} k_{Z_{3,4}^+}^{t_2}$ and with e^Y .

Now it is easy to see by considering the proof of Proposition 3.5(ii) that for these particular X and Y , the condition $V_X + \text{Ad}(k)V_Y = \mathfrak{p}$ must be satisfied by k of the form $k_0 = k_{Y_{1,3}^+}^{r_1} k_{Y_{2,4}^+}^{r_2} k_{Z_{1,2}^+}^{t_1} k_{Z_{3,4}^+}^{t_2}$. On the other hand, $V_Y = \langle Z_{i,j}, i < j \rangle$ and $\text{Ad}(k_0)(Z_{i,j})$, $i < j$, can only produce diagonal elements which satisfy $H_1 + H_3 = H_2 + H_4$ as can be checked by Lemma 3.19 and using the fact that $\text{Ad}(k_0) = \text{Ad}(k_{Y_{1,3}^+}^{r_1}) \text{Ad}(k_{Y_{2,4}^+}^{r_2}) \text{Ad}(k_{Z_{1,2}^+}^{t_1}) \text{Ad}(k_{Z_{3,4}^+}^{t_2})$. Consequently, $\mathfrak{a} \not\subset V_X + \text{Ad}(k_0)V_Y$ and the density cannot exist. \square

Proposition 4.5. *If X and Y are not eligible then the density does not exist.*

Proof. Let the configuration of X be $[\mathfrak{s}; u]$ or $[\mathfrak{s}]^-$ and the configuration of Y be $[\mathfrak{t}; v]$ or $[\mathfrak{t}]^-$. Proposition 4.1, Proposition 4.4 and Corollary 3.6 imply the non-existence of density when X and Y are not eligible and $u \leq 1$ and $v \leq 1$.

Suppose then that $u \geq 2$ or $v \geq 2$ and $\max(\mathfrak{s}, 2u) + \max(\mathfrak{t}, 2v) > 2p$ and consider the matrices $a(e^X k e^Y)$, $k \in \mathbf{SO}(p) \times \mathbf{SO}(p)$. Using (5) of Lemma 3.3 and Lemma 3.4, we may assume that the diagonal entries of \mathcal{D}_X and \mathcal{D}_Y are non-negative. Applying Remark 2.1, we have

$$\tilde{a}(e^X k e^Y) = \tilde{a}\left(\overbrace{(S^T e^X S)}^{e^{S^T X S}} \overbrace{(S^T k S)}^{\in \mathbf{SO}(p+q)} \overbrace{(S^T e^Y S)}^{e^{S^T Y S}}\right)$$

where $\tilde{a}(g)$ is the diagonal matrix with the singular values of g on the diagonal, ordered decreasingly (see the explanation before Lemma 3.18).

If $u + v > p$ then there are $r - s = r + (N - s) - N = 2u + 2v - 2p = 2(u + v - p)$ repetitions of $0 + 0 = 0$ in coefficients of $\tilde{a}(e^X k e^Y)$. Therefore 0 occurs at least $u + v - p > 0$ times as a diagonal entry of \mathcal{D}_H for every $H \in a(e^X K e^Y)$ which implies that $a(e^X K e^Y)$ has empty interior. If $2u + \max(\mathfrak{t}) > 2p$ denote $t = \max(\mathfrak{t})$. Let $Y_i \neq 0$ be repeated t times in \mathcal{D}_Y (or, if $t = t_r$ and $Y = Y[t]^-$, we have $t - 1$ times Y_r and once $-Y_r$ in \mathcal{D}_Y). Then there are $r - s = r + (N - s) - N = 2u + t - 2p$ repetitions of $Y_i + 0$ in coefficients of $\tilde{a}(e^X k e^Y)$. Therefore Y_i occurs at least $2u + t - 2p > 0$ times as a diagonal entry of \mathcal{D}_H for every $H \in a(e^X K e^Y)$ which implies that $a(e^X K e^Y)$ has empty interior. \square

5. SUFFICIENCY OF THE ELIGIBILITY CONDITION

5.1. Case $u \leq 1$ and $v \leq 1$.

Remark 5.1. In our proof, the case $u \leq 1$ and $v \leq 1$ is equivalent to the case $u = v = 0$. Indeed, for $H \in \overline{\mathfrak{a}^+}$, if the sole diagonal entry 0 in \mathcal{D}_H is replaced by a positive entry different from the existing diagonal entries of \mathcal{D}_H , then V_H is unchanged. We will therefore assume in this section that $u = 0$ and $v = 0$.

Definition 5.2. Let $\mathfrak{s} = (s_1, s_2, \dots, s_m)$ and $\mathfrak{t} = (t_1, t_2, \dots, t_n)$ be two partitions of p ($\sum_i s_i = p = \sum_i t_j$). We will say that \mathfrak{s} is finer than \mathfrak{t} if the t_i 's are sums of disjoint subsets of the s_j 's (for example, $\mathfrak{s} = [3, 2, 2, 1, 1, 1]$ is finer than $\mathfrak{t} = [5, 3, 2] = [3 + 2, 2 + 1, 1 + 1]$).

Remark 5.3. If $X = X[\mathfrak{s}]$ and $Y = Y[\mathfrak{t}]$ and \mathfrak{s} is finer than \mathfrak{t} then $V_Y \subset V_X$.

In the following lemma, we reduce in a significant way the number of elements for which we must prove the existence of the density.

Lemma 5.4. *For $p \geq 5$, it is sufficient to prove the existence of the density in the following cases:*

- S_1 : $X[p], Y[p - k, k]$ for $p - k \geq k \geq 2$
- S_2 : $X[p]^-, Y[p - k, k]$ for $p - k \geq k \geq 2$
- S_3 : $X[1, p - 1]^-, Y[p - 1, 1]$
- S_4 : $X[p - 1, 1], Y[p - 1, 1]$.

For $p = 4$ the same is true provided the case S_1 is replaced by the cases $X[4], Y[2, 1, 1]$ and $X[3, 1], Y[2, 2]$.

Proof. Suppose $p \geq 5$. Let us call A_0 the configurations of the form $[\mathfrak{s}]$ and A_1 all the others, i.e. the configurations of the form $[\mathfrak{s}]^-$.

- (a) We first observe that if the density exists for S_1 then it follows that it exists for all pairs $\{X, Y\}$ such that $X, Y \in A_0$, except when X or Y have configurations $[p]$ or $[p-1, 1]$. This comes from the fact that all these X, Y have structures that are finer and, consequently, the corresponding V_X and V_Y are larger. Thus, existence of the density in the cases S_1 together with S_4 will imply the existence of the density in all the cases when $X, Y \in A_0$, except when $\{X, Y\} \in \mathcal{E}$.
- (b) By switching to relatives and changing the order of X and Y , we see that it implies the existence of the density in all the cases when $X, Y \in A_1$, except when $\{X, Y\} \in \mathcal{E}$.
- (c) It remains to show that the cases S_2 and S_3 imply the existence of the density in all the cases when $X \in A_1$ and $Y \in A_0$, except when $\{X, Y\} \in \mathcal{E}$. Note first that if $X = [s]^-$ then either $[s]^- = [p]^-$ or $V_{X'} \subset V_X$ with $X' = X'[1, p-1]^-$. In the first case, we observe that the case S_2 implies the cases $X[p]^-$ and $Y \in A_0 \setminus \{[p], [p-1, 1]\}$ for the same reason as in (a). The only cases that remain with $Y[p-1, 1]$ are covered by S_3 . Finally, switching to relatives, we get the pairs $X \in A_1, Y[p]$ which are not in \mathcal{E} .

We illustrate the proof of the Lemma in the case $p = 5$.

$[2, 1^3]$	$[2, 2, 1]$	$[3, 1, 1]$	$[3, 2]$	$[4, 1]$	$[5]$	$[1^3, 2]^-$	$[2, 1, 2]^-$	$[1, 1, 3]^-$	$[3, 2]^-$	$[2, 3]^-$	$[1, 4]^-$	$[5]^-$	
✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	$[2, 1^3]$
	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	$[2, 2, 1]$
		✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	$[3, 1, 1]$
			✓	✓	S_1	✓	✓	✓	✓	✓	✓	S_2	$[3, 2]$
				S_4	X	✓	✓	✓	✓	✓	S_3	X	$[4, 1]$
					X	✓	✓	✓	✓	✓	X	X	$[4, 1]$
						✓	✓	✓	✓	✓	✓	✓	$[5]$
							✓	✓	✓	✓	✓	✓	$[1^3, 2]^-$
								✓	✓	✓	✓	✓	$[2, 1, 2]^-$
									✓	✓	✓	✓	$[1, 1, 3]^-$
										✓	✓	✓	$[3, 2]^-$
											✓	✓	$[2, 3]^-$
												✓	$[1, 4]^-$
													$[5]^-$

In the above table, \checkmark indicates that the pair is eligible, X indicates that the pair is not eligible and therefore that the density does not exist (the cases identified in (4.1)) and the S_i 's correspond to the notation above (the pair is eligible where the S_i 's appear). We use the reduction from Remark 3.11. \square

Theorem 5.5. *Let $p \geq 2$ and suppose that $X = X[s; u]$ or $[s]^-$ and $Y = Y[t; v]$ or $[t]^-$ are such that $u \leq 1$ and $v \leq 1$. If the pair $\{X, Y\}$ does not belong to the set $\mathcal{E} \cup \{[4], [2, 2]\} \cup \{[4]^- , [2, 2]^- \}$ then the density exists.*

Proof. The proof proceeds by induction similarly as in [6], but with a different ‘‘asymmetric’’ technique of executing Steps 2 and 3. Also, for small values of p the proofs must be led separately, due the lack of available good predecessors. With some exceptions in the starting phase of the induction, and in the case S_3 of the Lemma 5.4, the elements X and Y will be in \bar{a}^+ and their ‘‘usual’’ predecessors will be obtained by skipping the first diagonal terms of \mathcal{D}_X and \mathcal{D}_Y .

The only case of existence of the density for $p = 2$ is for regular $X[1, 1]$ and $Y[1, 1]$ (according to (4.1), only the pair $X[1, 1], Y[1, 1]$ of two regular elements verifies $|V_X| + |V_Y| \geq p^2$).

For $p = 3$, we have four possible configurations of nonzero singular elements: $[2, 1], [3], [1, 2]^- , [3]^-$.

All exceptional cases listed in (4.1) appear and only three pairs of singular configurations, namely $(X[2, 1], Y[2, 1]), (X[2, 1], Y[1, 2]^-)$ and $(X[1, 2]^- , Y[1, 2]^-)$, verify $|V_X| + |V_Y| \geq p^2$. Given that the pairs $(X[2, 1], Y[2, 1])$ and $(X[1, 2]^- , Y[1, 2]^-)$ are relatives, we only have to check the cases $(X[2, 1], Y[2, 1])$ and $(X[2, 1], Y[1, 2]^-)$.

In the case $X[2, 1], Y[2, 1]$, we write $\mathcal{D}_X = \text{diag}[a, a, b]$, $\mathcal{D}_Y = \text{diag}[c, c, d]$ and the predecessors $\mathcal{D}_{X'} = \text{diag}[a, b]$, $\mathcal{D}_{Y'} = \text{diag}[c, d]$, obtained by skipping the first coordinates, are regular. In the case $X[2, 1], Y[1, 2]^-$ we consider $\mathcal{D}_X = \text{diag}[a, a, b]$, $\mathcal{D}_{w \cdot y} = \text{diag}[-d, c, d]$ and only now go to regular predecessors $\mathcal{D}_{X'} = \text{diag}[a, b]$, $\mathcal{D}_{(w \cdot Y)'} = \text{diag}[c, d]$. The general proof given below applies in these cases.

When $p = 4$, by Lemma 5.4, we must show the existence of the density for:

- (1) $X[4], Y[2, 1, 1]$ and $X[2, 2], Y[3, 1]$,
- (2) $X[4]^- , Y[2, 2]$ or equivalently the relative pair $X[4], Y[2, 2]^-$,
- (3) $X[1, 3]^- , Y[3, 1]$,
- (4) $X[3, 1], Y[3, 1]$.

In the cases (1), (3) and (4), the usual predecessors have density when $p = 3$. The general proof given below applies in these cases.

For the case (2), observe that when $p = 3$, the configuration $X'[3]$ never gives the existence of density, when Y' is singular. That is why the second case $X[4], Y[2, 2]^-$ has no good predecessors and this case must be proved separately. We will do it after the general proof.

Starting from $p = 5$, the **general proof** by induction applies, the exceptions due to small values of p being taken care of. We present this proof now.

Step 1. Let $Y \in \overline{\mathfrak{a}^+}$ be such that $\mathcal{D}_Y = \text{diag}[\overbrace{a, \dots, a}^{p-k}, \overbrace{b, \dots, b}^k]$ and let its predecessor Y' be such that $\mathcal{D}_{Y'} = \text{diag}[\overbrace{a, \dots, a}^{p-k-1}, \overbrace{b, \dots, b}^k]$. The space V_Y is generated by completing a basis of $V_{Y'}$ with

$$N_Y = \{Y_{1,p-k+1}, \dots, Y_{1,p}, Z_{1,2}, \dots, Z_{1,p}\}.$$

We choose the predecessor X' of X in the same manner, except in the case S_3 , where we first write $\mathcal{D}_X = \text{diag}[b, b, \dots, b, a, -b]$ where $a > b > 0$ and skip the first term b in \mathcal{D}_X .

It is easy to see that the predecessors of X and Y are in the corresponding classes S_i for $p - 1$, so with density, except for $X[5], Y[3, 2]$, due to the non-eligible case $X[4], Y[2, 2]$. In this last case we arrange $\mathcal{D}_X = \text{diag}[a, a, a, a, a]$ and $\mathcal{D}_{wY} = \text{diag}[-b, b, b, c, -c]$ and go down to good predecessors $X'[4], (wY)'[2, 2]^-$. The proof described below leads to the existence of the density.

By the induction hypothesis and considering Corollary 3.10, there exists an open dense subset D' of $\mathbf{SO}(p - 1) \times \mathbf{SO}(p - 1)$ such that for all $w' \in W'$ and $k_0 \in D'$,

$$(5.1) \quad V_{w'.X'} + \text{Ad}(k_0)V_{Y'} = \mathfrak{p}'$$

and k_0 verifies condition (2) of Corollary 3.10.

We embed $K' = \mathbf{SO}(p - 1) \times \mathbf{SO}(p - 1)$ in $\mathbf{SO}(p) \times \mathbf{SO}(p)$ in the following manner:

$$K' \ni k' = \begin{bmatrix} 1 & & & \\ & k_{0,1} & & \\ & & 1 & \\ & & & k_{0,2} \end{bmatrix} \in \begin{bmatrix} \mathbf{SO}(p) & \\ & \mathbf{SO}(p) \end{bmatrix}, \quad k_{0,1}, k_{0,2} \in \mathbf{SO}(p - 1).$$

Hence, we have (identifying \mathfrak{p}' with its natural embedding into \mathfrak{p})

$$(5.2) \quad V_1 := V_{w'.X'} + \text{Ad}(k_0)V_{Y'} = \mathfrak{p}' = \begin{bmatrix} 0 & B' \\ B'^T & 0 \end{bmatrix}$$

for any $w' \in W'$, where $B' = \left[\begin{array}{c|c} 0_{1 \times 1} & 0_{1 \times (p-1)} \\ \hline 0_{(p-1) \times 1} & B''_{(p-1) \times (p-1)} \end{array} \right]$, and the matrix B'' is arbitrary (note that \mathfrak{p}' is of dimension $(p - 1)^2$). We must show that for some $k \in K$, the space $V_X + \text{Ad}(k)V_Y = \mathfrak{p}$.

Step 2. The element Y is always of the same form, so the next step of the proof is common for all the 4 cases. Similarly as in [6], Step 2 of the proof of Theorem 4.8, case (i), we prove that for $k_0 \in D' \subset \mathbf{SO}(p - 1) \times \mathbf{SO}(p - 1)$ the following property holds:

The space $\text{Ad}(k_0)\text{span}(N_Y)$ is of dimension $p + k - 1$ and its elements can be written in the form

$$\left[\begin{array}{c|ccc} 0 & a_1 & \dots & a_{p-1} \\ \hline \tau_1 & & & \\ \vdots & & & \\ \tau_{p-1-k} & & & 0 \\ a_{p+k-1} & & & \\ \vdots & & & \\ a_p & & & \end{array} \right]^s$$

with $a_i \in \mathbf{R}$ arbitrary and $\tau_j = \tau_j(a_1, \dots, a_{p+k-1})$. We will not need to write explicitly the functions τ_j . For the sake of completeness, we give a proof of Step 2.

Step 2 comes from the fact that the action of $\text{Ad}(k_0)$ on the elements of N_Y gives the linearly independent matrices

$$(5.3) \quad \text{Ad}(k_0)Y_{1,i} = \left[\begin{array}{c|c} 0 & \beta_{i-1}^T \\ \hline \alpha_{i-1} & 0 \end{array} \right]^s, \quad i = p - k + 1, \dots, p, \quad \text{Ad}(k_0)Z_{1,i} = \left[\begin{array}{c|c} 0 & -\beta_{i-1}^T \\ \hline \alpha_{i-1} & 0 \end{array} \right]^s, \quad i = 2, \dots, p$$

where the α_i 's are the columns of $k_{0,1}$ and the β_i 's are the columns of $k_{0,2}$. Let us write α'_i for a column α_i with the first $p - 1 - k$ entries omitted. In order to prove the statement of Step 2, we must show that the matrices

obtained by replacing α_i by α'_i in (5.3) are still linearly independent. This is equivalent to the linear independence of the matrices

$$(5.4) \quad \left[\begin{array}{c|c} 0 & -\beta_i^T \\ \hline \alpha'_i & 0 \end{array} \right]^s, \quad i = 1, \dots, p-k-1, \quad \left[\begin{array}{c|c} 0 & \beta_i^T \\ \hline 0 & 0 \end{array} \right]^s, \quad i = p-k, \dots, p-1, \quad \left[\begin{array}{c|c} 0 & 0 \\ \hline \alpha'_i & 0 \end{array} \right]^s, \quad i = p-k, \dots, p-1.$$

The matrices in (5.4) are linearly independent given that the matrix k_0 was assumed to satisfy condition (2) of Corollary 3.10.

Observe that contrary to [6], we have filled the zero margins of the matrix B' asymmetrically, which is why we call this method ‘‘asymmetric’’. The reason for doing this will be clear from the structure of the set N_X that we study now.

Step 3. Let us write the set N_X in the four cases S_i :

- (1) $S_1: N_X = \{Z_{1,2}, \dots, Z_{1,p}\}$
- (2) $S_2: N_X = \{Z_{1,2}, \dots, Z_{1,p-1}, Y_{1,p}\}$
- (3) $S_3: N_X = \{Z_{1,2}, \dots, Z_{1,p-1}, Y_{1,p-1}, Y_{1,p}\}$
- (4) $S_4: N_X = \{Z_{1,2}, \dots, Z_{1,p}, Y_{1,p}\}$.

We will now use the elements of N_X in order to generate the missing $p-k-1$ dimensions τ_j in the margins of B' . We use for this the vectors $Z_{1,2}, \dots, Z_{1,p-k}$ available in all four cases for $k \geq 1$. We proceed as follows:

If $\tau_1(1, 0, \dots, 0) = -1$, the vector $Z_{1,2} \in N_X$ is unhelpful. We change X' into X'' by putting the sign $-$ before the second and the last term of X' . We obtain $X'' = w'' \cdot X'$ such that N_X contains $Y_{1,2}$ instead of $Z_{1,2}$ and $w'' \in W'$ changes two signs of X' . This manipulation is justified by the fact that (5.1) holds for any $w'' \in W'$. We repeat this procedure, if needed, whenever $\tau_j(e_j) = -1$ and obtain, from elements of $N_{w \cdot X}$ and $\text{Ad}(k_0)(N_Y)$,

$$(5.5) \quad \left[\begin{array}{c|ccc} 0 & a_1 & \dots & a_{p-1} \\ \hline a_{2p-2} & & & \\ \vdots & & & \\ a_{p+k-1} & & 0 & \\ \vdots & & & \\ a_p & & & \end{array} \right]^s$$

for $w \cdot X$ with $w \in W$ and where the a_i 's are arbitrary.

Step 4. Noting that we have at least one element of N_X that has not been used, either $Y_{1,p}$ or $Z_{1,p}$, combining (5.1) and (5.5), we have

$$(5.6) \quad V_0 := \widetilde{V_{w \cdot X}} + \text{Ad}(k_0)(V_Y) = \left[\begin{array}{c|ccc} 0 & * & \dots & * \\ \hline * & * & \dots & * \\ \vdots & \vdots & \ddots & \vdots \\ * & * & \dots & * \end{array} \right]$$

where $\widetilde{V_{w \cdot X}}$ corresponds to the all of V_X without using the remaining $Y_{1,p}$ or $Z_{1,p}$. To fix things, let us assume that the unused element is $Z_{1,p}$, the reasoning being similar if it is $Y_{1,p}$ instead.

The end of the proof is similar to the final step of the proof in [6]. Refer to Lemma 3.19 and note that for t small, $\text{Ad}(e^{t(Z_{i,j}^+ + \theta Z_{i,j}^+)}) (\widetilde{V_{w \cdot X}}) + \text{Ad}(k_0)(V_Y) = V_0$. Indeed, the lemma shows that no new element is introduced and, for t small, the dimension is unchanged. On the other hand, V_0 is strictly included in $\text{Ad}(e^{t(Z_{i,j}^+ + \theta Z_{i,j}^+)}) (\langle Z_{1,p} \rangle \cup \widetilde{V_{w \cdot X}}) + \text{Ad}(k_0)(V_Y)$ since, still by Lemma 3.19, a new diagonal element is introduced. We conclude that for t small enough, $\text{Ad}(e^{t(Z_{i,j}^+ + \theta Z_{i,j}^+)}) (\langle Z_{1,p} \rangle \cup \widetilde{V_{w \cdot X}}) + \text{Ad}(k_0)(V_Y) = \mathfrak{p}$. Finally,

$$V_{w \cdot X} + \text{Ad}(e^{-t(Z_{i,j}^+ + \theta Z_{i,j}^+)}) k_0(V_Y) = \mathfrak{p}.$$

The case $X[2, 2]^-$, $Y[4]$: This case is awkward since (X, Y) do not have eligible predecessors. We select X and Y such that $\mathcal{D}_X = \text{diag}[a, a, b, -b]$ and $\mathcal{D}_Y = \text{diag}[c, c, c, c]$ (assuming $a, b, c \neq 0$ and $a \neq b$). Then V_Y is generated by the basis B_Y composed of all the 6 vectors $Z_{i,j}$, $i < j$, while the basis B_X of V_X contains the vectors $Y_{1,3}, Y_{1,4}, Y_{2,3}, Y_{2,4}, Y_{3,4}$ and all the vectors $Z_{i,j}$ except $Z_{3,4}$. Note that $|V_X| = 10$.

For a root vector $Z_{i,j}^+$ denote $Z_{i,j}^\mathfrak{k} = Z_{i,j}^+ + \theta(Z_{i,j}^+) \in \mathfrak{k}$. Define $Z_0 = Z_{1,2}^\mathfrak{k} + Z_{2,3}^\mathfrak{k} + Z_{1,4}^\mathfrak{k} + Z_{2,4}^\mathfrak{k} \in \mathfrak{k}$. We denote

$$F_t = V_X + \text{Ad}(e^{tZ_0})V_Y = V_X + e^{t \text{ad } Z_0}(V_Y),$$

$$E_t = V_X + \langle \{v + t[Z_0, v] : v \in V_Y\} \rangle.$$

We will write

$$f_t = \det(B_X, \text{Ad}(e^{tZ_0})B_Y)$$

where the elements of \mathfrak{p} are seen as column vectors in \mathbf{R}^{p^2} . Analogously, we denote by e_t the determinant constructed in a similar way from the vectors of B_X and the vectors $v + t[Z_0, v]$, $v \in B_Y$, belonging to E_t . We write $f_t = e_t + r_t$ and we analyse now e_t and r_t in order to show that $f_t \neq 0$ for some small nonzero t .

Using Lemma 3.20, we check that $e_t = ct^5$ with $c = \det(B_X, Z_{3,4}, [Z_0, Z_{1,2}], \dots, [Z_0, Z_{2,4}]) \neq 0$. The coefficient of t^6 in e_t equals zero since $\det(B_X, [Z_0, B_Y]) = 0$. On the other hand it is easy to see in a similar way that the remainder r_t in the analytic expansion $f_t = e_t + r_t$ does not have terms in t^n for $n < 6$. We conclude that $f_t \neq 0$ for small nonzero t . \square

5.2. Case $u \geq 2$ or $v \geq 2$. This case is handled in much of the same way as the case $u > 0$ or $v > 0$ in [6]. A notable difference is that the basis for induction is the previous case ($u \leq 1$ and $v \leq 1$). For $p = 3$, we only need to consider the pair $X[1; 2], Y[2, 1]$ which has regular predecessors. Similarly, for $p = 4$, we see that all eligible pairs with $u \geq 2$ or $v \geq 2$ have eligible predecessors.

In the case $p = 5$, because of (2.7), there are eligible pairs with no eligible predecessors. It suffices to consider the pair $X[5], Y[3; 2]$. In order to show that the density exists in this case, we use the same technique as for the case $X[2, 2]^- , Y[4]$. We take $Z_0 = Z_{1,2}^{\mathfrak{k}} + Z_{2,3}^{\mathfrak{k}} + Z_{3,4}^{\mathfrak{k}} + Z_{1,5}^{\mathfrak{k}} + Z_{2,5}^{\mathfrak{k}}$. In order to prove that $e_t = ct^9$ with $c \neq 0$, we check using Lemma 3.20 that the 9 vectors $[Z_0, Z_{1,2}], \dots, [Z_0, Z_{3,5}]$ produce the missing vectors $Y_{1,2}, Y_{1,3}, Y_{2,3}, Y_{4,5}$ and the diagonal.

Starting from $p = 6$, the induction proof works. The fact that the roots α_i defined by $\alpha_i(X) = X_i$ are absent in the case $\mathbf{SO}(p, p)$ does not influence the proof (in [6] the roots α_i were not used in Step 4 of the proof). These two differences being overcome, the proof is sufficiently similar that it should not be repeated here.

6. APPLICATIONS

We now extend our results to the symmetric spaces of type C_p , i.e. to the complex and quaternion cases. Recall that $\mathbf{SU}(p, p)$ is the subgroup of $\mathbf{SL}(2p, \mathbf{C})$ such that $g^* I_{p,p} g = I_{p,p}$ while $\mathbf{Sp}(p, p)$ is the subgroup of $\mathbf{SL}(2p, \mathbf{H})$ such that $g^* I_{p,p} g = I_{p,p}$. Their respective maximal compact subgroups are $\mathbf{S}(\mathbf{U}(p) \times \mathbf{U}(p))$ and $\mathbf{Sp}(p) \times \mathbf{Sp}(p) \equiv \mathbf{SU}(p, \mathbf{H}) \times \mathbf{SU}(p, \mathbf{H})$. Their subspaces \mathfrak{p} can be described as $\begin{bmatrix} 0 & B \\ B^* & 0 \end{bmatrix}$ where B is an arbitrary complex (respectively quaternionic) matrix of size $p \times p$. The Cartan subalgebra \mathfrak{a} is chosen in the same way as for $\mathfrak{so}(p, p)$, with real entries in the diagonal.

Remark 6.1. The following table is helpful in showing the differences and similarities between $\mathbf{SO}(p, p)/\mathbf{SO}(p) \times \mathbf{SO}(p)$, $\mathbf{SU}(p, p)/\mathbf{S}(\mathbf{U}(p) \times \mathbf{U}(p))$ and $\mathbf{Sp}(p, p)/\mathbf{Sp}(p) \times \mathbf{Sp}(p)$ (the ‘‘real’’, ‘‘complex’’ and ‘‘quaternionic’’ cases).

	$\mathbf{SO}(p, p)/\mathbf{SO}(p) \times \mathbf{SO}(p)$	$\mathbf{SU}(p, p)/\mathbf{S}(\mathbf{U}(p) \times \mathbf{U}(p))$	$\mathbf{Sp}(p, p)/\mathbf{Sp}(p) \times \mathbf{Sp}(p)$
Root system	D_p	C_p	C_p
m_α where $\alpha(X) = X_i - X_j, i < j$	1	2	4
m_α where $\alpha(X) = X_i + X_j, i < j$	1	2	4
m_α where $\alpha(X) = 2X_i,$ $i = 1, \dots, p$	0	1	3
Dimension of \mathfrak{p}	p^2	$2p^2$	$4p^2$
Action of the Weyl group on $X \in \mathfrak{a}$	Permutes the diagonal entries of $\mathcal{D}(X)$ and changes any pair of signs	Permutes the diagonal entries of $\mathcal{D}(X)$ and changes any sign	Permutes the diagonal entries of $\mathcal{D}(X)$ and changes any sign

Theorem 6.2. Consider the symmetric spaces $\mathbf{SU}(p, p)/\mathbf{S}(\mathbf{U}(p) \times \mathbf{U}(p))$ and $\mathbf{Sp}(p, p)/\mathbf{Sp}(p) \times \mathbf{Sp}(p)$. Let $X = X[\mathbf{s}; u]$ and $Y = Y[\mathbf{t}; v] \in \mathfrak{a}$. Then the measure $\delta_{e^X}^{\mathfrak{a}} * \delta_{e^Y}^{\mathfrak{a}}$ is absolutely continuous if and only if $\max(\mathbf{s}, 2u) + \max(\mathbf{t}, 2v) \leq 2p$.

Proof. Let $X, Y \in \mathfrak{a}$. Note that since

$$a(e^X \mathbf{S}(\mathbf{U}(p) \times \mathbf{U}(p)) e^Y) \subset a(e^X (\mathbf{Sp}(p) \times \mathbf{Sp}(p)) e^Y),$$

if the density exists in the complex case, it also exists in the quaternionic case. On the other hand, given Lemma 3.18, one can reproduce Proposition 4.5 using $\mathbf{F} = \mathbf{C}$ and $\mathbf{F} = \mathbf{H}$ to show that the condition is necessary in the complex and quaternionic cases.

However, the root structure is richer in the complex and quaternionic cases compared to the real cases. The existence of the roots $\alpha(X) = 2X_k$ makes the complex and quaternionic cases very similar to the case $q > p$.

It clearly suffices to prove the result in the complex case. The involution θ is given by $\theta(X) = -X^*$ and the positive root vectors are generated by $X_k^+ = \left[\begin{array}{c|c} -iE_{k,k} & iE_{k,k} \\ \hline -iE_{k,k} & iE_{k,k} \end{array} \right]$ for the root $\alpha(H) = 2H_k$, by $Y_{r,s}^+ = \left[\begin{array}{c|c} E_{r,s} - E_{s,r} & E_{r,s} + E_{s,r} \\ \hline E_{r,s} + E_{s,r} & E_{r,s} - E_{s,r} \end{array} \right]$, $Y_{r,s,\mathbf{C}}^+ = \left[\begin{array}{c|c} i(E_{r,s} + E_{s,r}) & i(E_{r,s} - E_{s,r}) \\ \hline i(E_{r,s} - E_{s,r}) & i(E_{r,s} + E_{s,r}) \end{array} \right]$ for the root $\alpha(H) = H_r - H_s$ and by $Z_{r,s}^+ = \left[\begin{array}{c|c} E_{r,s} - E_{s,r} & E_{s,r} - E_{r,s} \\ \hline E_{r,s} - E_{s,r} & E_{s,r} - E_{r,s} \end{array} \right]$, $Z_{r,s,\mathbf{C}}^+ = \left[\begin{array}{c|c} -i(E_{r,s} + E_{s,r}) & i(E_{r,s} + E_{s,r}) \\ \hline -i(E_{r,s} - E_{s,r}) & i(E_{r,s} + E_{s,r}) \end{array} \right]$ for the root $\alpha(H) = H_r + H_s$ (here the matrices $E_{r,s}$ are of size $p \times p$).

Taking into account the fact that if the density exists in the real case, it also exists in the complex case, we only have a few cases to verify. Given that changing any sign of a diagonal element of \mathcal{D}_X , $X \in \mathfrak{a}$, is a Weyl group action, we can always assume that all entries of \mathcal{D}_X are non-negative. The configuration $[s]^-$ thus disappears.

We will need to show that the cases $(X[p], Y[p])$, $(X[p], Y[p-1; 1])$ and $(X[4], Y[2; 2])$ all have a density. We will use the case $p = 1$ which is of rank 1 as the inductive step (there is nothing to prove for that case).

For the case $X[p], Y[p]$, $p > 1$, we proceed much as in [6]. We note here a few differences. If $k_0 \in \mathbf{S}(\mathbf{U}(p) \times \mathbf{U}(p))$

with $k_0 = \begin{bmatrix} 1 & & & \\ & k_{0,1} & & \\ & & 1 & \\ & & & k_{0,2} \end{bmatrix} \in \mathbf{S}(\mathbf{U}(p) \times \mathbf{U}(p))$ then

$$(6.1) \quad \text{Ad}(k_0)(Z_{1,k}) = \left[\begin{array}{c|c} 0 & -\beta_{k-1}^* \\ \hline \alpha_{k-1} & 0 \end{array} \right]^s, \text{Ad}(k_0)(Z_{1,k,\mathbf{C}}) = \left[\begin{array}{c|c} 0 & i\beta_{k-1}^* \\ \hline i\alpha_{k-1} & 0 \end{array} \right]^s, \quad k = 1, \dots, p-1 \quad \text{and}$$

$$\text{Ad}(k_0)(X_1) = \left[\begin{array}{c|c} i & 0 \\ \hline 0 & 0 \end{array} \right]^s.$$

Given that

$$(6.2) \quad Z_{1,j} = \left[\begin{array}{c|c} 0 & -\mathbf{e}_{j-1}^T \\ \hline \mathbf{e}_{j-1} & 0 \end{array} \right]^s, \quad Z_{1,j,\mathbf{C}} = \left[\begin{array}{c|c} 0 & i\mathbf{e}_{j-1}^T \\ \hline i\mathbf{e}_{j-1} & 0 \end{array} \right]^s, \quad j = 2, \dots, p,$$

we want to show that the matrices in (6.1) together with those of (6.2) are linearly independent for a $k_0 \in \mathbf{S}(\mathbf{U}(p-1) \times \mathbf{U}(p-1))$ for which the equality (5.1) holds. Note that if $k_{0,1} = iI_{p-1}$, $k_{0,2} = -iI_{p-1}$ then the matrices in (6.1) and (6.2) are linearly independent. Since the linear independence is based on a determinant being nonzero, this implies that the set of matrices k_0 for which this is true is open and dense in $\mathbf{S}(\mathbf{U}(p-1) \times \mathbf{U}(p-1))$.

We conclude that if $N'_X = N_X \setminus \{X_1\}$ then $\text{span}(N'_X + V_{X'}) + \text{Ad}(k'_0)V_Y$ has the form $\begin{bmatrix} ia & * & \dots & * \\ * & * & \dots & * \\ \vdots & \vdots & \ddots & \vdots \\ * & * & \dots & * \end{bmatrix}$ where

the $*$'s represent arbitrary complex numbers and a is an arbitrary real number.

We finish the proof as in the case $\mathbf{SO}(p, p)$ using the vector X_1^+ .

The case $(X[p], Y[p-1; 1])$ has eligible predecessors $(X'[p-1], Y'[p-1])$. We then have $N_X = \{Z_{1,k}, Z_{1,k,\mathbf{C}}, X_1\}$ and $N_Y = \{Z_{1,k}, Z_{1,k,\mathbf{C}}, Y_{1,k}, Y_{1,k,\mathbf{C}}\}$. The rest follows easily.

Finally, the case $(X[4], Y[2; 2])$ has predecessors $(X'[3], Y'[2; 1])$ which are eligible. \square

We will conclude this paper with two further applications.

Proposition 6.3. *Let X and $Y \in \mathfrak{a}$ be such that $(\delta_{e^X}^\natural)^{*2}$ and $(\delta_{e^Y}^\natural)^{*2}$ are absolutely continuous. Then $\delta_{e^X}^\natural * \delta_{e^Y}^\natural$ is absolutely continuous.*

Proof. The proof is very similar to the one found in [6]. \square

In previous papers, we have studied a related question: if $X \in \mathfrak{a}$ and $X \neq 0$, for which convolution powers l is the measure $(\delta_{e^X}^\natural)^l$ absolutely continuous? This problem is equivalent to the study of the absolute continuity of convolution powers of uniform orbital measures $\delta_g^\natural = m_K * \delta_g * m_K$ for $g \notin K$. It was proved in [5, Corollary 7] that it is always the case for $l \geq r + 1$, where r is the rank of the symmetric space G/K . It was also shown in [6]

that $r + 1$ is optimal for this property for symmetric spaces of type A_n ([5, Corollary 18]) but this is not the case for the symmetric spaces of type B_p where r was shown to be sufficient in [6].

Proposition 6.4. *If $p = 3$ and $\mathcal{D}_X = \text{diag}[a, a, a]$, $a > 0$, then $(\delta_{e^X}^{\natural})^3$ is not absolutely continuous in $\mathbf{SO}(p, p)/\mathbf{SO}(p) \times \mathbf{SO}(p)$ while $(\delta_{e^X}^{\natural})^4$ is absolutely continuous.*

Proof. Computing the derivative of the map $T(k_1, k_2, k_3, k_4) = k_1 e^X k_2 e^X k_3 e^X k_4$ at (k_1, k_2, k_3, k_4) as in (3.2), we obtain

$$k_1 e^X k_2 (\text{Ad}(k_2^{-1}) U_{-X} + \text{Ad}(e^X) U_X) e^X k_3 k_4 = k_1 e^X k_2 (\mathfrak{k} + \text{Ad}(k_2^{-1}) V_{-X} + \text{Ad}(e^X) V_X) e^X k_3 e^X k_4.$$

The dimension of this space is at most $|\mathfrak{k}| + |V_{-X}| + |V_X| = |\mathfrak{k}| + 3 + 3 < |\mathfrak{k}| + |\mathfrak{p}| = |\mathfrak{g}|$ so the map cannot be surjective.

On the other hand, X' such $\mathcal{D}_{X'} = \text{diag}[2a, a, a]$ belongs to $a(e^X K e^X)$ from what precedes (taking $x = a$) and the pair (X', X') is eligible. From there, we conclude that $a(e^X K e^X K e^X K e^X)$ has nonempty interior. \square

Proposition 6.5. *If $p = 4$ and $\mathcal{D}_X = \text{diag}[a, a, a, a]$, $a > 0$, then $(\delta_{e^X}^{\natural})^3$ is not absolutely continuous in $\mathbf{SO}(p, p)/\mathbf{SO}(p) \times \mathbf{SO}(p)$ while $(\delta_{e^X}^{\natural})^4$ is absolutely continuous. Consequently, if $X = X[\mathfrak{s}; u]$ with $u \geq 1$ or $X = X[\mathfrak{s}]^-$ then $(\delta_{e^X}^{\natural})^4$ is absolutely continuous.*

Proof. We know that the elements of $e^X K e^X$ have the form $k_a e^Z k_b$ where $\mathcal{D}_Z = \text{diag}[c, c, d, d]$, $c \geq d$. From the end of the proof of Proposition 4.4, we know that for all $H \in a(e^X K e^X K e^X)$, $\mathcal{D}_H = \text{diag}[H_1, H_2, H_3, H_4]$ will satisfy $H_1 + H_3 = H_2 + H_4$. We conclude therefore that $a(e^X K e^X K e^X)$ has empty interior. On the other hand, since there exists $Z \in a(e^X K e^X)$ with $\mathcal{D}_Z = \text{diag}[c, c, d, d]$ with $c > d > 0$ and (Z, Z) forms an eligible pair, it follows that $a(e^X K e^X K e^X K e^X)$ has nonempty interior since it contains $a(e^Z K e^Z)$. \square

Proposition 6.6. *If $p \geq 5$ and $\mathcal{D}_X = \text{diag}[a, \dots, a]$, $a > 0$, then $(\delta_{e^X}^{\natural})^3$ is absolutely continuous in $\mathbf{SO}(p, p)/\mathbf{SO}(p) \times \mathbf{SO}(p)$. Consequently, if $X = X[\mathfrak{s}; u]$ with $u \geq 1$ or $X = X[\mathfrak{s}]^-$ then $(\delta_{e^X}^{\natural})^3$ is absolutely continuous.*

Proof. Note that for $t > 0$ small enough, $Z = a(e^X k_t^{Z_{p-1,1}^+} e^X) \in a(e^X K e^X)$ is such that $\mathcal{D}_Z = \text{diag}[\overbrace{a, \dots, a}^{p-2}, x, x]$ with $a > x > 0$. Given that (Z, X) form an eligible pair and that $a(e^Z K e^X) \subset a(e^X K e^X K e^X)$, the result follows. \square

Theorem 6.7. *On symmetric spaces $\mathbf{SO}_0(p, p)/\mathbf{SO}(p) \times \mathbf{SO}(p)$, ($p \geq 4$), $\mathbf{SU}(p, p)/\mathbf{S}(\mathbf{U}(p) \times \mathbf{U}(p))$ and $\mathbf{Sp}(p, p)/\mathbf{Sp}(p) \times \mathbf{Sp}(p)$, $p \geq 2$, for every nonzero $X \in \mathfrak{a}$, the measure $(\delta_{e^X}^{\natural})^p$ is absolutely continuous. Moreover, p is the smallest value for which this is true: if X has a configuration $[1; p-1]$ then the measure $(\delta_{e^X}^{\natural})^{p-1}$ is singular.*

Proof. We used the three preceding propositions and [6, Theorem 5.3]. \square

7. CONCLUSION

With this paper and with [4, 6], we have now obtained sharp criteria on singular X and Y for the existence of the density of $\delta_{e^X}^{\natural} \star \delta_{e^Y}^{\natural}$ for the root systems of types A_n , B_p , C_p , D_p and E_6 . Thanks to [5, 6] and Theorem 6.7 of the present paper, sharp criteria are now given for the l -th convolution powers $(\delta_{e^X}^{\natural})^l$ to be absolutely continuous for any $X \neq 0$, $X \in \mathfrak{a}$. It is interesting to note that the eligibility criterion depends strongly on the geometry of the root system. Consequently, a characterization of eligibility that would be applicable for all Riemannian symmetric spaces of non-compact type is unlikely to exist.

REFERENCES

- [1] J.-P. Anker, *Le noyau de la chaleur sur les espaces symétriques $\mathbf{U}(p, q)/\mathbf{U}(p) \times \mathbf{U}(q)$* , Lectures Notes in Mathematics 1359, (Springer-Verlag, New-York), 1988, 60-82.
- [2] Friedberg, Insel, Spence. *Linear Algebra*. Prentice-Hall, 1997.
- [3] P. Graczyk, P. Sawyer, *The product formula for the spherical functions on symmetric spaces of noncompact type*, J. Lie Theory 13 (2003), 247–261.
- [4] P. Graczyk, P. Sawyer, *A sharp criterion for the existence of the density in the product formula on symmetric spaces of Type A_n* , J. Lie Theory 20 (2010), 751-766.
- [5] P. Graczyk, P. Sawyer, *Absolute continuity of convolutions of orbital measures on Riemannian symmetric spaces*, J. Funct. Anal. 259 (2010), 1759-1770.
- [6] P. Graczyk, P. Sawyer, *On the product formula on non-compact Grassmannians*, Colloq. Math. 133 (2013), 145-167
- [7] S. Helgason, *Differential geometry, Lie groups, and symmetric spaces*, Academic Press, 1978.
- [8] S. Helgason, *Groups and geometric analysis*, Academic Press, 1984.
- [9] P. Sawyer, *Spherical functions on $\mathbf{SO}_0(p, q)/\mathbf{SO}(p) \times \mathbf{SO}(q)$* , Canad. Math. Bull. 42 (1999), no. 4, 486-498.

LABORATOIRE DE MATHÉMATIQUES, LAREMA, UNIVERSITÉ D'ANGERS, 49045 ANGERS CEDEX 01
E-mail address: piotr.graczyk@univ-angers.fr

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, LAURENTIAN UNIVERSITY, SUDBURY, ONTARIO
E-mail address: psawyer@laurentian.ca