Comment on "Phase transition of a two-dimensional generalized XY model"

Aernout van Enter, Silvano Romano, Valentin Zagrebnov

► To cite this version:

Aernout van Enter, Silvano Romano, Valentin Zagrebnov. Comment on "Phase transition of a two-dimensional generalized XY model". Journal of Physics A: Mathematical and Theoretical, 2011, 44 (20), pp.208001. 10.1088/1751-8113/44/20/208001 . hal-00968544

HAL Id: hal-00968544

<https://hal.science/hal-00968544>

Submitted on 2 Apr 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Comment on the paper by
Y. Komura and Y. Okabe, *J.Phys. A* **44**,
015002, 2011

Aernout C.D. van Enter ^{*}, Silvano Romano [†]
and Valentin A. Zagrebnov [‡]

March 27, 2011

Abstract

We point out that the claim of strong universality of [1] is incorrect, as it contradicts known rigorous results.

AMS 2000 subject classification: 82B20, 82B26, 60K35.

Keywords: Generalised XY-model, Kosterlitz-Thouless versus first-order transition, nonuniversality.

^{*}University of Groningen, Johann Bernoulli Institute of Mathematics and Computing Science,
Postbus 407, 9700 AK Groningen, The Netherlands,
A.C.D.v.Enter@math.rug.nl,
<http://www.math.rug.nl/~aenter/>

[†]University of Pavia, Dept of Physics, via A. Bassi 6, Pavia, Italy,
Silvano.Romano@pv.infn.it,
<http://www.pv.infn.it/~romano/>

[‡]Université de la Méditerranée and Centre de Physique Théorique - UMR 6207 Luminy Case
977, 13288 Marseille, Cedex09, France
zagrebnov@cpt.univ-mrs.fr,
<http://www.cpt.univ-mrs.fr/~zagrebno/zagrebnov.htm>

The generalised lattice ferromagnetic XY -model was introduced by two of us in [2]. Its simplest version involves 3-component unit spins on a D -dimensional lattice, parameterized by standard angles in the spherical coordinates and interacting according to the ferromagnetic nearest-neighbour Hamiltonian

$$H = -J \sum_{\langle i,j \rangle} (\sin\theta_i \sin\theta_j)^q \cos(\varphi_i - \varphi_j) , \quad J > 0 .$$

When $D = 2$, the model yields orientational disorder at all finite temperatures, and produces a transition to a low-temperature Berezinskiĭ–Kosterlitz–Thouless (BKT) phase, possessing slow decay of magnetic correlations and infinite susceptibility [2]. In turn, the transition is expected, and was found by simulation [3], to exhibit the usual BKT scenario, at least for small values of q . On the other hand, this scenario does not hold for *all* values of q , i.e. the BKT transitions are *not universal*.

Indeed, we *proved* in [4] that the named transition turns first-order for sufficiently large q . Recall that a hint of this behaviour was already suggested by numerical analysis [3].

Notice that the above *rigorous* results entail that both existence and the type of transition *depend* on the parameter q , i.e. they exclude universality.

The above-mentioned generalized XY -model was recently addressed by simulation for various values of q in Ref. [1]. Although the authors found only a certain evidence of the BKT criticality, they claim that this behaviour holds for *all values* of q . This claim can not be correct since it contradicts to our rigorous results [4], which seem to have been unknown to the authors of the Ref. [1].

References

- [1] Y. Komura and Y. Okabe, *J.Phys. A* **44**, 015002, 2011.
- [2] S. Romano and V. A. Zagrebnov, *Phys. Lett. A* **301**, 201, 2002.
- [3] L. A. S. Mol, A. R. Pereira, H. Chamati and S. Romano, *Eur.Phys. J. B* **50**, 541, 2006.
- [4] A. C. D. van Enter, S. Romano and V. A. Zagrebnov, *J. Phys. . A* **39**, L439, 2006.