

Approximation of the Ventcel problem, numerical results

Marc Dambrine ^{*1} and Charles Pierre ^{†1}

¹ Laboratoire de Mathématiques et de leurs Applications, , UMR CNRS 5142,
Université de Pau et des Pays de l'Adour, France.

December, 2017

Abstract

Report on the numerical approximation of the Ventcel problem. The Ventcel problem is a 3D eigenvalue problem involving a surface differential operator on the domain boundary: the Laplace-Beltrami operator. We present in the first section the problem statement together with its finite element approximation, the code machinery used for its resolution is also presented here.

The last section presents the obtained numerical results. These results are quite unexpected for us. Either super-converging for P^1 Lagrange finite elements or under converging for P^2 and P^3 .

The remaining sections 2 and 3 provide numerical results either for the classical Laplace or for the Laplace-Beltrami operator numerical approximation. These examples being aimed to validate the code implementation.

1 Introduction

We present in this report an algorithm for solving the Ventcel problem [8, 9]. The code is detailed and validated considering various classical test cases. The resolution of the Ventcel problem on the unit ball is used as a test case to provide a convergence numerical analysis of the method.

1.1 The Ventcel problem

The Ventcel problem is the following eigenvalue problem. Let Ω denote some bounded smooth domain in \mathbb{R}^3 . We search for the eigenvalues λ and for the

*marc.dambrine@univ-pau.fr

†charles.pierre@univ-pau.fr

associated eigenfunctions u satisfying,

$$\Delta u = 0 \quad \text{on } \Omega, \quad \Delta_B u - \partial_n u + \lambda u = 0 \quad \text{on } \partial\Omega. \quad (1)$$

Here \mathbf{n} denotes the outward unit normal to $\partial\Omega$ and so $\partial_n u$ the derivative of u on the normal direction to the boundary and $\Delta_B u$ stands for the surface Laplacian of u on the boundary $\partial\Omega$, i.e. the Laplace-Beltrami operator.

This problem has a simple weak formulation (2). Multiplying Δu by a test function v and integrating over Ω and assuming sufficient regularity we get by integration by part,

$$\int_{\Omega} \nabla u \cdot \nabla v \, dx - \int_{\partial\Omega} \partial_n u \, v \, ds = 0.$$

By substituting the boundary condition in (1),

$$\int_{\Omega} \nabla u \cdot \nabla v \, dx - \int_{\partial\Omega} (\Delta_B u + \lambda u) \, v \, ds.$$

We obtain by a second integration by part on the boundary $\partial\Omega$,

$$\int_{\Omega} \nabla u \cdot \nabla v \, dx + \int_{\partial\Omega} \nabla_T u \cdot \nabla_T v \, ds = \lambda \int_{\partial\Omega} u v \, ds, \quad (2)$$

with $\nabla_T u$ the tangential gradient of u on $\partial\Omega$, i.e. $\nabla_T u = \nabla u - \partial_n u \, \mathbf{n}$.

On the Hilbert space $V = \{u \in H^1(\Omega), \nabla_T u \in L^2(\partial\Omega)\}$ the classical theory applies as exposed in [3] providing an orthogonal set of eigenfunctions associated with the eigenvalues $0 = \lambda_0 < \lambda_1 \leq \dots$. The first eigenvalue $\lambda_0 = 0$ is associated with the (one dimensional) eigenspace of the constants. We will focus on the approximation of λ_1 .

The following discretisation of (2) will be used. Let \mathcal{M} be some tetrahedral mesh of Ω . We will denote by Ω_h the computation domain (made of all tetrahedra in the mesh \mathcal{M}). We consider the classical Lagrange finite element spaces $V_h = P^k(\mathcal{M})$. We search for $u \in V_h$ and $\lambda \in \mathbb{R}$ so that, for all $v \in V_h$ we have:

$$\int_{\Omega_h} \nabla u \cdot \nabla v \, dx + \int_{\partial\Omega_h} \nabla_T u \cdot \nabla_T v \, ds = \lambda \int_{\partial\Omega_h} u v \, ds. \quad (3)$$

Considering a canonical bases of V_h we identify u_h with its vectorial representation $U \in \mathbb{R}^N$, with N the number of degrees of freedom (the dimension of V_h). We then consider the matrices S_3 , S_2 and M_2 (mass and stiffness matrices) representing the products on $V_h \times V_h$ in (3):

$$(u, v) \mapsto \int_{\Omega_h} \nabla u \cdot \nabla v \, dx = U^T S_3 V, \quad (4)$$

$$(u, v) \mapsto \int_{\partial\Omega_h} \nabla_T u \cdot \nabla_T v \, ds = U^T S_2 V, \quad (5)$$

$$(u, v) \mapsto \int_{\partial\Omega_h} u v \, ds = U^T M_2 V. \quad (6)$$

Problem (3) under matricial form is the following generalized eigenvalue problem,

$$(S_3 + S_2)U = \lambda M_2 U. \quad (7)$$

The matrix $S_3 + S_2$ is symmetric positive semi definite (with a one dimensional kernel made of the constant vectors $U = c$ representing the constant functions). The matrix M_2 also is symmetric positive semi definite but with a high dimensional kernel made of all functions in V_h vanishing on the boundary $\partial\Omega_h$.

1.2 Implementation

A *python* code is used for the numerical resolution of (7). The assembling of the finite element matrices is made using *Getfem++*¹.

The iterative Lanczos method (see e.g. Saad [6]) is used to solve the eigenvalue problem. Because we are interested in the smallest eigenvalues and since the matrix $S_3 + S_2$ is only semi definite, we consider the shifted invert variant of this method. Actually the shifted matrix $S_3 + S_2 - \sigma M_2$ is symmetric positive definite for any negative value of the shift parameter σ , fixed to $\sigma = -1$ here. In practice the ARPACK library² has been used for this.

Each Lanczos algorithm iteration requires two kind of advanced operations: solving the linear system $(S_3 + S_2 - M_2)X = Y$ (shifted system matrix) and performing matrix-vector multiplication $X \mapsto M_2 X$. These two operations are executed using the *PETSc* library³.

The largest amount of CPU consumption is devoted to the linear system resolution. A conjugate gradient algorithm is used with an incomplete Cholesky (fill in 3) preconditioning, see e.g. Saad [7] for precisions. The system is solved with a tolerance of 10^{-11} .

The residual at end of the Lanczos algorithm is asked to be smaller than 10^{-10} . This residual is algebraic, we additionally re-computed an L^2 residual: denoting λ_i the i^{th} eigenvalue and U_i the associated eigenvector, the L^2 residual is defined as,

$$\frac{\|(S_3 + S_2)U_i - \lambda_i M_2 U_i\|_{L^2}}{\|U_i\|_{L^2}},$$

assuming the identification $V_h \simeq \mathbb{R}^N$. This functional residual is also controlled to remain below 10^{-9} .

Eventually all meshes have been built with the software *Gmsh*⁴.

¹Getfem++: an open-source finite element library, <http://download.gna.org/getfem/html/homepage/index.html>

²ARPACK, Arnoldi Package, <http://www.caam.rice.edu/software/ARPACK/>

³PETSc, Portable, Extensible Toolkit for Scientific Computation, <http://www.mcs.anl.gov/petsc/>

⁴Gmsh: a three-dimensional finite element mesh generator, <http://geuz.org/gmsh/>

2 Code validation

We consider here the resolution of several classical problems aimed to validate the code implementation: matrix assembling, eigenproblem solver and error analysis. On each example our purpose is to recover the correct method's convergence rate as predicted by classical theories.

2.1 Matrix assembling validation

The three matrices S_3 , S_2 and M_2 associated with the products (4), (5) and (6) need to be computed. The matrix S_3 is a classical stiffness matrix on the three dimensional domain Ω_h . Conversely the matrices S_2 and M_2 are not as classical. They correspond to two dimensional stiffness and mass matrices but associated with the non flat domain $\partial\Omega_h$.

Remark nevertheless that the assembling of S_2 (and similarly for M_2) consist of a loop over all element faces E of the mesh \mathcal{M} included in $\partial\Omega_h$. Precisely the elements of \mathcal{M} are tetrahedra, their faces E thus are triangles and these triangles subsets of $\partial\Omega_h$ form a partition of $\partial\Omega_h$. On every such triangle E is assembled the local matrix corresponding to the product,

$$(u, v) \mapsto \int_E \nabla_T u \cdot \nabla_T v \, ds = \int_E \nabla u|_E \cdot \nabla v|_E \, ds.$$

On one hand E is an affine deformation of the reference triangle in dimension 2 and on the other hand $u|_E, v|_E \in P^k(E)$. Therefore these local matrices indeed are local matrices for a classical 2 dimensional stiffness matrix. As a result the assembling of S_2 and similarly of M_2 do not need particular software but only a 2D finite element library.

Taking the previous commentary into account, the validation for the assembling of S_3 , S_2 and M_2 can be performed on flat domains $\omega \subset \mathbb{R}^d$, $d = 2, 3$. We fix $\omega = (0, 1)^d$ and solve the classical elliptic problem,

$$-\Delta u + u = f \quad \text{on } \omega \quad \text{and} \quad \partial_n u = 0 \quad \text{on } \partial\omega,$$

for the right hand side $f = (2\pi^2+1) \cos(\pi x) \cos(\pi y)$ and $f = (3\pi^2+1) \cos(\pi x) \cos(\pi y) \cos(\pi z)$ in dimension 2 and 3 respectively. The associated exact solutions are $u = \cos(\pi x) \cos(\pi y)$ and $u = \cos(\pi x) \cos(\pi y) \cos(\pi z)$ respectively. The numerical solution u_h is given by,

$$S_d U_h + M_d U_h = M_d F, \quad d = 2, 3.$$

We analyze the relative errors between u and u_h both in L^2 -norm and H^1 -semi norm,

$$e_{L^2} = \frac{\|u - u_h\|_{L^2(\omega)}}{\|u\|_{L^2(\omega)}}, \quad e_{H^1} = \frac{\|u - u_h\|_{H^1(\omega)}}{\|u\|_{H^1(\omega)}}.$$

considering a series of refined meshes and using the $P^k(\omega)$ finite element method for $k = 1, 2, 3$, we recovered a k (resp. $k + 1$) order of convergence

in H^1 (resp. L^2) norm as presented in table 1. These results are in complete agreement with the classical theory (see e.g. Ciarlet [2]) and this test fully validates a correct assembling of the desired matrices.

$d = 2$	P^1	P^2	P^3	$d = 3$	P^1	P^2	P^3
e_{H^1}	1.0	2.1	3.0	e_{H^1}	1.0	1.9	2.8
e_{L^2}	2.0	3.1	4.0	e_{L^2}	1.9	3.0	4.0

Table 1: Computed orders of convergence for problem (2.1)

2.2 Eigenvalue problem solver validation

We test the eigenvalue problem solver described in section 1.2 considering the Laplace eigenproblem,

$$-\Delta u = \lambda u \quad \text{on } \omega \quad \text{and} \quad \partial_n u = 0 \quad \text{on } \partial\omega, \quad (8)$$

on the same square or cubic geometry ω as in the previous section.

The first eigenvalue is $\lambda_0 = 0$ with eigenspace the constant functions. The first non-zero eigenvalue is $\lambda_1 = \pi^2$ with multiplicity d and with eigenspace $E_1 = \text{Span}(\cos(\pi x), \cos(\pi y))$ or $E_1 = \text{Span}(\cos(\pi x), \cos(\pi y), \cos(\pi z))$ for $d = 2$ or $d = 3$.

According to the dimension d the numerical approximation for (8) is,

$$S_d U_h = \lambda_h M_d U_h.$$

The first non-zero numerical eigenvalue $\lambda_{h,1}$ is computed to analyze the relative error,

$$e_\lambda = \frac{|\lambda_{h,1} - \lambda_1|}{\lambda_1}.$$

One associated eigenvector $U_{h,1}$ of L^2 -norm equal to 1 (for normalization) is considered to compute the following H^1 and L^2 errors,

$$e_{L^2} = \|u_{h,1} - pu_{h,1}\|_{L^2(\omega)} \quad e_{H^1} = \|u_{h,1} - pu_{h,1}\|_{H^1(\omega)},$$

where p is the L^2 -orthogonal projection onto the eigenspace E_1 .

$d = 2$	P^1	P^2	P^3	$d = 3$	P^1	P^2	P^3
e_{H^1}	1.0	2.0	3.1	e_{H^1}	1.0	2.0	2.7
e_{L^2}	2.0	3.0	4.1	e_{L^2}	1.9	3.0	3.8
e_λ	2.0	4.2	5.9	e_λ	1.95	4.0	5.9

Table 2: Computed orders of convergence for problem (8)

The order of convergence is computed considering a series of refined meshes and several Lagrange finite element spaces $P^k(\omega)$, they are reported in table 2. They are in full agreement with the theoretical orders presented e.g. in Babuřhka and Osborn [1] : using P^k finite elements provides a convergence of order k for e_{H^1} , $k + 1$ for e_{L^2} and $2k$ for e_λ .

3 Curved surface effects

In this section we illustrate the influence of computing the two dimensional stiffness and mass matrices S_2 and M_2 . We consider again the geometrical situation of the Ventcel problem in section 1.1: Ω is an open smooth domain in \mathbb{R}^3 with boundary Γ . \mathcal{M} is a tetrahedral mesh of Ω , Ω_h is the domain of the mesh \mathcal{M} , i.e. the union of its element. The matrices S_2 and M_2 are computed on $\Gamma_h = \partial\Omega_h$. Note that Γ_h itself is the domain of a triangular mesh \mathcal{M}_2 of Γ , the element of \mathcal{M}_2 are the boundary faces of \mathcal{M} .

As developed in section 2.1, the assembling of S_2 and M_2 on the curved domain Γ_h (i.e. embedded in \mathbb{R}^3) is essentially the same as their assembling in a flat domain (a two dimensional domain in \mathbb{R}^2). We however check here that the third dimension z is taken into account for this assembling.

3.1 Laplace-Beltrami problem

We will consider the elliptic problem

$$-\Delta_B u + u = f \quad \text{on } \Gamma, \quad (9)$$

no boundary condition are needed here (Γ has no boundary!). This problem is well posed : existence and uniqueness of a solution $u \in H^1(\Gamma)$ for any $f \in L^2(\Gamma)$.

The finite element approximation of (9) has been theorized by Demlow in [5]. We need some notations. To $x \in \Gamma$ is associated its unit outer normal $\mathbf{n}(x)$. We consider a tubular neighborhood ω of Γ so that for all $x \in \omega$, there exists a unique $p(x) \in \Gamma$ so that $x - p(x) = \lambda(x)\mathbf{n}(p(x))$ ($p(x)$ is an orthogonal projection of x on Γ) and so that the segment $[x, p(x)] \subset \omega$. A function $f : \Gamma \rightarrow \mathbb{R}$ can be extended to a function $f^e : \omega \rightarrow \mathbb{R}$ by: $f^e(x) = f(p(x))$. We assume that $\Gamma_h \subset \omega$. A function $f : \Gamma \rightarrow \mathbb{R}$ can then be lifted to a function $f^l : \Gamma_h \rightarrow \mathbb{R}$ by $f^l = f^e|_{\Gamma_h}$.

The lift operation allows to compare the exact solution u defined on Γ with a numerical approximation u_h defined on Γ_h . Note that it would have also been possible to lift u_h to a function u_h^l on Γ by $u_h^l(p(x)) = u_h(x)$. Demlow showed in [5] that analyzing the error in terms of $u^l - u_h$ (lift of u to Γ_h) or in terms of $u - u_h^l$ (lift of u_h to Γ) are equivalent, we choose the first strategy for its practical convenience.

Equation (9) is discretised on $V_h = P^k(\mathcal{M}_2)$ as,

$$(S_2 + M_2)U_h = M_2 F^l. \quad (10)$$

The L^2 and H^1 approximation errors are alternatively defined as,

$$e_{L^2} = \|u^l - u_h\|_{L^2(\Gamma_h)}, \quad e_{H^1} = \|u^l - u_h\|_{H^1(\Gamma_h)}.$$

The convergence properties of this scheme are quite different from those on flat domains illustrated in section 2.1. The reason for this is analyzed in Demlow

[5]: the curved surface Γ is approximated by the surface Γ_h that is the boundary of a polyhedral and thus piecewise flat (made of triangles). Approximating a curved surface by a (piecewise) linear one induces upper bound errors for (9),

$$e_{H^1} = O(h^k + h^2), \quad e_{L^2} = O(h^{k+1} + h^2),$$

for P^k finite elements. Therefore a saturation of the convergence order to 2 is predicted.

The numerical scheme (10) has been implemented for the sphere. In this case the projection p is very simple, $p(x) = x/\|x\|$ on $\omega = \mathbb{R}^3 - \{0\}$. The right hand side to (9) is set to $f = x(2 + x) \exp(x)$ so that the exact solution is $u = \exp(x)$. The results are reported in table 3. The saturation of the convergence order to h^2 is clearly seen, in agreement with Demlow convergence analysis [5].

	P^1	P^2	P^3		P^1	P^2	P^3
e_{H^1}	1.1	2.0	2.0	e_{H^1}	1	2	2
e_{L^2}	2.0	2.0	2.0	e_{L^2}	2	2	2
				e_λ	2	2	2

Table 3: Computed orders of convergence for the Laplace-Beltrami problem (9) (left) and for the Laplace-Beltrami eigenvalue problem (11).

3.2 Laplace-Beltrami eigenvalue problem

The Laplace-Beltrami eigenproblem is considered on the sphere,

$$-\Delta_B u = \lambda u \quad \text{on } \Gamma. \quad (11)$$

Its discretisation takes the form: find $U_h \in V_h = P^k(\Gamma_h)$ and $\lambda_h \in \mathbb{R}$ so that,

$$S_2 U_h = \lambda_h M_2 U_h.$$

Problem (11) has $\lambda_0 = 0$ for eigenvalue associated to the eigenspace of constant functions. The first non-zero eigenvalue is $\lambda_1 = 2$ of multiplicity 3 with eigenspace $E_1 = \text{Span}(x, y, z)$ the restriction of the linear functions in \mathbb{R}^3 to the sphere.

The space E_1 is lifted to Γ_h as in the previous section. The lifted space E_i^l is the vector space of functions of the form $X = (x, y, z) \mapsto (\alpha x + \beta y + \delta z)/\|X\|$ with $\|X\| = (x^2 + y^2 + z^2)^{1/2}$. The orthogonal projector p from V_h onto E_1^l is considered to define the numerical errors,

$$e_\lambda = \frac{|\lambda_{h,1} - \lambda_1|}{\lambda_1}, \quad e_{L^2} = \frac{\|u_{h,1} - pu_{h,1}\|_{L^2(\Gamma_h)}}{\|u_{h,1}\|_{L^2(\Gamma_h)}} \quad \text{and} \quad e_{H^1} = \frac{\|u_{h,1} - pu_{h,1}\|_{H^1(\Gamma_h)}}{\|u_{h,1}\|_{L^2(\Gamma_h)}},$$

on the first computed non zero eigenvalue $\lambda_{h,1}$ and associated eigenfunction $u_{h,1}/\|u_{h,1}\|_{L^2(\Gamma_h)}$ normalized in L^2 -norm.

The results are reported in table 3 on the right. The convergence in H^1 -norm is in $O(h^k + h^2)$ for P^k Lagrange finite element which is consistent with the previous subsection assertions. Order 2 convergence in L^2 -norm is observed in all cases P^k , $k = 1, 3$.

The eigenvalue convergence clearly shows a saturation to order 2 convergence. In section 2.2 we quote the analysis of Babuška and Osborn [1] who showed an order $2k$ convergence for the eigenvalues with k the convergence order of the eigenfunctions in H^1 -norm. This analysis does not apply here since it is restricted to Galerkin type approximations with $V_h \subset V$. In the present case this is no longer true because V and V_h are function spaces associated to different domains, Γ and Γ_h respectively.

Conclusion. The numerical results showed in this section together with the theoretical analysis of Demlow [5] show that one cannot get better than an order 2 convergence when considering a piecewise affine mesh Γ_h of the curved boundary Γ of the domain Ω . This point also was addressed in [5] where piecewise polynomial interpolation of order p Γ_h^p of Γ are considered providing now a saturation of the convergence order to $p + 1$.

4 The Ventcel problem numerical convergence

	P^1	P^2	P^3
e_{H^1}	1.	1.5	1.5
e_{L^2}	2.	2.5	2.5
e_λ	2.	2.	2.

Table 4: Convergence orders for the Ventcel problem

We analyze here the numerical approximation of the Ventcel problem exposed in section 1.1 using the numerical scheme (7).

The domain Ω is set to the unit ball. In this particular case the eigenvalues/eigenfunctions have been computed in [4]. The n^{th} eigenvalue λ_n for $n \geq 0$ is given by $\lambda_n = n^2 + 2n$, with multiplicity $2n + 1$ and with eigenspace E_n the space of harmonic functions of order n .

The numerical domain Ω_h here is a subset (by convexity) of Ω . Therefore functions f on Ω simply are lifted to functions f^l on Ω_h by restriction: $f^l = f|_{\Omega_h}$. We consider a numerical approximation $\lambda_{n,h}$ of the n^{th} non zero eigenvalue λ_n (since λ_n is of multiplicity $2n + 1$, we numerically get $2n + 1$ approximations of λ_n , we fix one of them). We denote $U_{n,h}$ an associated eigenfunction, so that $\|U_{n,h}\|_{L^2(\Omega_h)} = 1$ (L^2 normalisation). We introduce the orthogonal projection (for the L^2 -scalar product on Ω_h) $p : V_h \rightarrow E_n^l$, where here E_n^l is the space made of the restrictions of all functions in E_n to Ω_h .

with these notations we define the numerical errors,

$$e_{\lambda_n} = \frac{|\lambda_{n,h} - \lambda_n|}{\lambda_n}, \quad e_{L^2} = \|U_{n,h} - pU_{n,h}\|_{L^2(\Omega_h)} \quad \text{and} \quad e_{H^1} = \|\nabla U_{n,h} - \nabla pU_{n,h}\|_{L^2(\Omega_h)}$$