

Two-component equations modelling water waves with constant vorticity

Joachim Escher, David Henry, Boris Kolev, Tony Lyons

► **To cite this version:**

Joachim Escher, David Henry, Boris Kolev, Tony Lyons. Two-component equations modelling water waves with constant vorticity. *Annali di Matematica Pura ed Applicata*, Springer Verlag, 2016, 195 (1), pp.249-271. <http://link.springer.com/article/10.1007/s10231-014-0461-z> . 10.1007/s10231-014-0461-z . hal-01002338v2

HAL Id: hal-01002338

<https://hal.archives-ouvertes.fr/hal-01002338v2>

Submitted on 30 Sep 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

TWO-COMPONENT EQUATIONS MODELLING WATER WAVES WITH CONSTANT VORTICITY

JOACHIM ESCHER, DAVID HENRY, BORIS KOLEV, AND TONY LYONS

ABSTRACT. In this paper we derive a two-component system of nonlinear equations which models two-dimensional shallow water waves with constant vorticity. Then we prove well-posedness of this equation using a geometrical framework which allows us to recast this equation as a geodesic flow on an infinite dimensional manifold. Finally, we provide a criteria for global existence.

CONTENTS

1. Introduction	1
2. Derivation of the model equation	3
2.1. Nondimensionalisation and scaling	4
3. The two component system	6
3.1. The ρ -component	6
3.2. The m -component	7
4. Geometric reformulation as a geodesic flow	10
5. Well-posedness of the equation	14
6. Global solutions	18
Acknowledgements	22
References	22

1. INTRODUCTION

The focus of this paper is the following nonlinear two-component system of equations which model two-dimensional shallow water waves with constant vorticity:

$$(1.1) \quad \begin{cases} m_t = \alpha u_x - a u_x m - u m_x - \kappa \rho \rho_x, \\ \rho_t = -u \rho_x - (a - 1) u_x \rho, \end{cases}$$

where $m = u - u_{xx}$. Here $a \neq 1$ is a real parameter, α is a constant which represents the vorticity of the underlying flow, and $\kappa > 0$ is an arbitrary real parameter. In Sections 2 and 3 we present a derivation of system (1.1) in the hydrodynamical setting using formal asymptotic expansions and perturbation theory [35] applied to the full governing equations for two-dimensional

Date: September 30, 2014.

2010 Mathematics Subject Classification. 35Q35; 76B15; 35Q53; 58D05.

Key words and phrases. Water waves; vorticity; model equations; Euler equation; diffeomorphism group.

water waves with constant vorticity. The above system generalises and incorporates a number of celebrated nonlinear partial differential equations which have recently been derived as approximate models in hydrodynamics.

When $\alpha = 0$ (which in our considerations corresponds to irrotational fluid flow) and $\rho \equiv 0$ we obtain a one-component family of equations which are parameterised by $a \neq 1$. This family of so-called b -equations possess a number of structural phenomena which are shared by solutions of the family of equations [23, 26, 28]. However, there are just two members of this family which are integrable [32]: the Camassa-Holm (CH) [3, 4] equation, when $a = 2$, and the Degasperis-Procesi (DP) [13] equation, when $a = 3$. The CH equation is a remarkable equation which is both integrable and which possesses both global solutions and solutions which exhibit wave-breaking in finite time [6, 9, 8], a feature which is also shared by the DP equation [22, 21]. Both the CH and DP equations can be derived in the hydrodynamic setting of shallow water waves [7, 12, 33], and hence they represent the first examples of integrable equations which possess wave-breaking solutions [7].

In [38] the authors presented a number of integrable multi-component generalisations of the CH equation, the most popular of which corresponds to $\alpha = 0, a = 2, \kappa = \pm 1$ in (1.1) [10, 5, 20, 31, 27]. It was shown in [10] that this two-component generalisation of CH can be derived from a hydrodynamical setting, in which case $\kappa = 1$. The question as to whether these hydrodynamical model equations could be adapted so as to incorporate an underlying vorticity in the flow is a natural one. Physically, vorticity is vital for incorporating the ubiquitous effects of currents and wave-current interactions in fluid motion [39], and furthermore the past decade has seen a burgeoning in the mathematical analysis of the full-governing equations for water waves with vorticity— cf. [7] for an overview of much of this work. From a model equation viewpoint, following [34], Ivanov [30] derived a number of two-component equations for shallow water waves with underlying constant vorticity, including an integrable generalisation of the two-component CH equation. In Sections 2 and 3 of this paper we extend this work to derive (1.1), a family of two-component systems parameterised by a , where $a = 2$ generalises the CH equation, and $a = 3$ generalises the DP equation.

In Section 4 we present a geometrical interpretation of (1.1), which is shown to correspond (when $a = 2$) to the *Arnold–Euler equation* of a right-invariant metric on the infinite dimensional Lie group

$$(\text{Diff}^\infty(\mathbb{S}^1) \otimes \text{C}^\infty(\mathbb{S}^1)) \times \mathbb{R}.$$

This geometrical approach goes back to the pioneering work of V. Arnold [1], published in 1966, who recast the equation of motion of a perfect fluid (with fixed boundary) as the geodesic flow on the volume-preserving diffeomorphisms group of the domain. For the little history, Arnold’s paper was written (in French) for the bi-centenary of the 1765’s paper of Euler [24] (also written in French) who recast the equation of motion of a free rigid body as the geodesic flow on the rotation group. As acknowledged by Arnold himself, his paper concentrated on the *geometrical ideas* and not on the difficult analytical technicalities that are inherent when one works with an *infinite dimensional diffeomorphisms group* rather than a *finite dimensional Lie group*. In 1970, Ebin & Marsden [14] reconsidered this geometric approach

from the analytical point of view. They proposed to consider the group of smooth diffeomorphisms as an *inverse limit of Hilbert manifolds*. The remarkable observation is that, in this framework, the Euler equation (a PDE) can be recast as an ODE (the geodesic equation) on these Hilbert manifolds. Furthermore, following their approach, if we can prove *local existence and uniqueness of the geodesics* (ODE), then the PDE is *well-posed*. This technique has been used in [37, 11] for the periodic Camassa–Holm equation. It was extended to (non metric) geodesic flows such as the Degasperis–Procesi equation in [17] and to right-invariant metrics induced by fractional Sobolev norm (non-local inertia operators) in [18]. It is used in Section 5 to establish the well-posedness of the system (1.1). Finally, in Section 6, we derive some *a priori* estimates on (1.1) which lead to a criteria for global existence of solutions.

2. DERIVATION OF THE MODEL EQUATION

In this section we apply formal asymptotic methods to the full-governing equations for two-dimensional water waves with an underlying constant vorticity α to derive the two-component system (1.1). We choose Cartesian-coordinates with the z -axis pointing vertically upwards and the x -axis perpendicular to the crest-lines of the waves, the flow being in the positive x -direction. We let $z = 0$ denote the location of the flat bed and assume that the mean-depth, or the depth of the undisturbed water domain, is given by h_0 . The velocity field of the two dimensional water flow is given by $(u(t, x, z), w(t, x, z))$ with the water's free surface $z = h_0 + \eta(t, x)$ — we denote by Ω_η the fluid domain with free boundary. For water waves which are large in scale it is reasonable to make the simplifying assumptions of homogeneity (constant density) and non-viscosity (no internal friction forces). We decompose the pressure

$$P(x, z, t) = P_0 + \rho g(h_0 - z) + p(x, z, t)$$

into the hydrostatic pressure term (where ρ is the fluid density and P_0 is the constant atmospheric pressure) and the deviation from hydrostatic pressure p . The equations governing the motion of the fluid comprise Euler's equation

$$(2.1a) \quad u_t + uu_x + ww_z = -\frac{1}{\rho}p_x \quad \text{in } \Omega_\eta,$$

$$(2.1b) \quad w_t + uw_x + ww_z = -\frac{1}{\rho}p_z \quad \text{in } \Omega_\eta,$$

together with the continuity equation

$$(2.1c) \quad u_x + w_z = 0 \quad \text{in } \Omega_\eta,$$

and the dynamic and kinematic boundary conditions

$$(2.1d) \quad w = \eta_t + u\eta_x, \quad p = \eta\rho g \quad \text{on } z = h_0 + \eta,$$

$$(2.1e) \quad w = 0 \quad \text{on } z = 0.$$

2.1. Nondimensionalisation and scaling. The process of nondimensionalisation enables us to reexpress the governing equations (2.1) in terms of dimensionless variables and functions, and so the resulting equations are purely mathematical. The benefit of this nondimensionalisation procedure is that it naturally introduces dimensionless parameters into the mathematical problem, which measure the relative size and significance of the physical terms characteristic to the water wave problem we consider. These parameters enable the formal asymptotic expansion procedures which we engage in when we derive (1.1) below. If a is the typical amplitude of the waves we consider, and λ represents a characteristic horizontal length scale (e.g. the wavelength) for the waves, we define the dimensionless parameters $\epsilon = a/h_0$, $\delta = h_0/\lambda$. Then, performing the change of variables

$$(2.2) \quad \begin{aligned} x &\rightarrow \lambda x, & z &\rightarrow zh_0, & t &\rightarrow \frac{\lambda}{\sqrt{gh_0}}t, & \eta &\rightarrow a\eta, \\ u &\rightarrow \epsilon\sqrt{gh_0}u, & w &\rightarrow \epsilon\delta\sqrt{gh_0}w, & p &\rightarrow \epsilon\rho gh_0p, \end{aligned}$$

the new scaled variables are nondimensionalised. In terms of the new variables the system (2.1) assumes the form

$$\begin{aligned} u_t + \epsilon(uu_x + wu_z) &= -p_x, \\ \delta^2(w_t + \epsilon(uw_x + ww_z)) &= -p_z, \\ u_x + w_z &= 0, \\ w = \eta_t + \epsilon u\eta_x, \quad p &= \eta\rho g && \text{on } z = 1 + \epsilon\eta, \\ w = 0 &&& \text{on } z = 0. \end{aligned}$$

We note that $(u, w, p, \eta) \equiv (U(z), 0, 0, 0)$ gives a solution of the above system, which represents laminar flows with a flat free surface and with an arbitrary shear. To incorporate undulating waves in the presence of a shear flow, we make the transformation $u = (U(z) + \epsilon u)$. If $\epsilon = 0$ (flat surface), then we are simply left with a shear flow as above. The system which describes waves in the presence of a shear flow is given by

$$(2.3a) \quad u_t + Uu_x + wU' + \epsilon(uu_x + wu_z) = -p_x \quad \text{in } \Omega_\epsilon,$$

$$(2.3b) \quad \delta^2(w_t + Uw_x + \epsilon(uw_x + ww_z)) = -p_z \quad \text{in } \Omega_\epsilon,$$

$$(2.3c) \quad u_x + w_z = 0 \quad \text{in } \Omega_\epsilon,$$

$$(2.3d) \quad w = \eta_t + (U + \epsilon u)\eta_x, \quad p = \eta\rho g \quad \text{on } z = 1 + \epsilon\eta,$$

$$(2.3e) \quad w = 0 \quad \text{on } z = 0.$$

For the simplest nontrivial case of a laminar shear flow, we let $U(z) = \alpha z$, α constant and $0 \leq z \leq 1$. We choose $\alpha > 0$ whereby the underlying current is in the positive x -direction. The Burns condition [2, 25] for the non-dimensionalised scaled variables becomes

$$\int_0^1 \frac{dz}{(U(z) - c)^2} = 1,$$

and so

$$(2.4) \quad c^2 - \alpha c - 1 = 0,$$

giving us the nondimensionalised speed

$$c = \frac{1}{2}(\alpha \pm \sqrt{\alpha^2 + 4})$$

of the travelling waves in linear approximation. We mention here that the Burns condition arises as a local bifurcation condition for shallow water waves with constant vorticity, cf. [7]. In physical coordinates the vorticity is given by $\omega = (U - u)_z - w_x$. Scaling the vorticity by $\omega = \sqrt{h_0/g}\omega$ we get

$$\omega = \alpha + \epsilon(u_z - \delta^2 w_x).$$

As we seek a solution with constant vorticity we must have

$$(2.5) \quad u_z - \delta^2 w_x = 0.$$

From (2.5), (2.3c) and (2.3e) we get

$$(2.6a) \quad u = u_0 - \delta^2 \frac{z^2}{2} u_{0,xx} + \mathcal{O}(\epsilon^2, \delta^4, \epsilon\delta^2),$$

$$(2.6b) \quad w = -z u_{0,x} + \delta^2 \frac{z^3}{6} u_{0,xxx} + \mathcal{O}(\epsilon^2, \delta^4, \epsilon\delta^2).$$

Here $u_0(x, t)$ is the leading order approximation for u in an asymptotic expansion, and we see from (2.5) that $u_z = 0$ when $\delta \rightarrow 0$ and hence u_0 is independent of z . From (2.3d), together with (2.6) we get

$$(2.7a) \quad \eta_t + \alpha\eta_x + \left[(1 + \epsilon\eta)u_0 + \epsilon\frac{\alpha}{2}\eta^2 \right]_x - \delta^2 \frac{1}{6} u_{0,xxx} = 0,$$

ignoring terms of $\mathcal{O}(\epsilon^2, \delta^4, \epsilon\delta^2)$. From (2.3b), (2.3d), (2.6) we have, to the same order

$$p = \eta - \delta^2 \left[\frac{1 - z^2}{2} u_{0,xt} + \frac{1 - z^3}{3} \alpha u_{0,xx} \right].$$

Then from (2.3a) we have

$$(2.7b) \quad \left(u_0 - \delta^2 \frac{1}{2} u_{0,xx} \right)_t + \epsilon u_0 u_{0,x} + \eta_x - \delta^2 \frac{\alpha}{3} u_{0,xxx} = 0.$$

Letting $\delta, \epsilon \rightarrow 0$ in (2.7) we get

$$(2.8a) \quad \eta_t + \alpha\eta_x + u_{0,x} = 0,$$

$$(2.8b) \quad u_{0,t} + \eta_x = 0,$$

giving us

$$(2.9) \quad \eta_{tt} + \alpha\eta_{tx} - \eta_{xx} = 0.$$

Travelling wave solutions $\eta(x - ct)$ of (2.9) have a speed c which satisfies (2.4). There are two possible speeds c , corresponding to a left and right running wave respectively. In particular, from (2.8a) we get

$$(2.10) \quad \eta = cu_0 + \mathcal{O}(\epsilon, \delta^2),$$

when we choose a specific value of the wavespeed c . We note here that while ϵ and δ are generally not related, it is known [12] that the small-amplitude long-wave scaling in the absence of vorticity is $\epsilon = O(\delta^2)$, while $\epsilon = O(\delta)$ is the long-wave scaling for waves of moderate amplitude.

3. THE TWO COMPONENT SYSTEM

We introduce the auxiliary variable θ , defined in terms of η and u_0 as follows

$$(3.1) \quad \theta = 1 + \epsilon k_1 \eta + \epsilon^2 k_2 \eta^2 + \epsilon \delta^2 k_3 u_{0,xx},$$

which upon squaring and retaining terms to order $\mathcal{O}(\epsilon^2, \epsilon \delta^2)$ gives

$$(3.2) \quad \theta^2 = 1 + \epsilon(2k_1)\eta + \epsilon^2(k_1^2 + 2k_2)\eta^2 + \epsilon\delta^2(2k_3)u_{0,xx} + \mathcal{O}(\epsilon^3, \epsilon^2\delta^2).$$

The coefficients k_1 , k_2 and k_3 are as yet undetermined, but as we proceed it will be seen that specific values must be assigned to all three if the resulting system is to be integrable. At a later stage, θ will be rewritten in terms of the physical variable ρ , which satisfies one member of the two component system (1.1) we intend to analyse in the following sections.

3.1. The ρ -component. Rewriting η in terms of θ using (3.1) we get

$$(3.3) \quad \begin{aligned} \eta_t &= \frac{1}{k_1\epsilon}\theta_\xi - \epsilon\frac{k_2}{k_1}\eta\eta_t + \delta^2\frac{k_3}{k_1}u_{0,xt}, \\ \eta_x &= \frac{1}{k_1\epsilon}\theta_x - \epsilon\frac{k_2}{k_1}(\eta^2)_x + \delta^2\frac{k_3}{k_1}u_{0,xxx}, \end{aligned}$$

and using (2.8a), (2.8b) and (2.10), we write to leading order

$$(3.4) \quad \begin{aligned} u_{0,xt} &= -\eta_{xxx} = -cu_{0,xxx} + \mathcal{O}(\epsilon, \delta^2), \\ \eta_t &= -\alpha\eta_x - u_{0,x} + \mathcal{O}(\epsilon, \delta^2). \end{aligned}$$

Substituting the expressions (3.4) into equation (3.3) and keeping terms to order $\mathcal{O}(\epsilon, \delta^2)$ we find

$$(3.5) \quad \begin{aligned} \eta_t &= \frac{1}{k_1\epsilon}\theta_t + \epsilon\frac{2k_2(1+\alpha c)}{k_1}\eta u_{0,x} + \delta^2\frac{k_3c}{k_1}u_{0,xxx} + \mathcal{O}(\epsilon^2, \epsilon\delta^2), \\ \eta_x &= \frac{1}{k_1\epsilon}\theta_x - \epsilon\frac{2k_2c}{k_1}\eta u_{0,x} - \delta^2\frac{k_3}{k_1}u_{0,xxx} + \mathcal{O}(\epsilon^2, \epsilon\delta^2). \end{aligned}$$

Then forming the linear combination $\eta_t + \alpha\eta_x$ using the expressions in (3.5) we obtain

$$(3.6) \quad \eta_t + \alpha\eta_x = \frac{1}{\epsilon k_1}(\theta_t + \alpha\theta_x) + \epsilon\frac{2k_2}{k_1}\eta u_{0,x} + \delta^2\frac{k_3(c-\alpha)}{k_1}u_{0,xxx} + \mathcal{O}(\epsilon^2, \epsilon\delta^2).$$

Replacing the $\eta_t + \alpha\eta_x$ term in equation (2.7a) with corresponding the expression given by relation (3.6) yields

$$(3.7) \quad \begin{aligned} \frac{1}{\epsilon k_1}(\theta_t + \alpha\theta_x) + \left((1 + \epsilon\eta)u_0 + \epsilon\frac{\alpha}{2}\eta^2 + \epsilon\frac{k_2}{k_1}\eta u_0 \right)_x = \\ -\delta^2 \left(\frac{k_3(c-\alpha)}{k_1} - \frac{1}{6} \right) u_{0,xxx} + \mathcal{O}(\epsilon^2, \epsilon\delta^2) = 0. \end{aligned}$$

We impose the constraint

$$(3.8) \quad \frac{k_3}{k_1} = \frac{1}{6(c-\alpha)},$$

thereby eliminating terms of order δ^2 which would otherwise introduce dispersion to this particular member of the two component system. Using

the approximation (2.10) and neglecting terms of order $\mathcal{O}(\epsilon^2, \epsilon\delta^2)$, we may rewrite equation (3.7) as

$$(3.9) \quad \frac{1}{\epsilon k_1} (\theta_t + \alpha\theta_x) + \left((1 + \epsilon \left(1 + \frac{\alpha c}{2} + \frac{k_2}{k_1} \right) \eta) u_0 \right)_x + \mathcal{O}(\epsilon^2, \epsilon\delta^2) = 0.$$

Having imposed only one constraint on the coefficients k_1 and k_3 so far, we may also choose k_1 and k_2 such that

$$(3.10) \quad k_1 = 1 + \frac{\alpha c}{2} + \frac{k_2}{k_1},$$

in which case (3.9) becomes

$$(3.11) \quad \frac{1}{\epsilon k_1} (\theta_t + \alpha\theta_x) + ((1 + \epsilon k_1 \eta) u_0)_x + \mathcal{O}(\epsilon^2, \epsilon\delta^2) = 0.$$

Keeping terms of order $\mathcal{O}(\epsilon, \delta^2)$ we approximate the factor $1 + \epsilon k_1 \eta$ in (3.11) by θ , resulting in

$$(3.12) \quad \frac{1}{\epsilon k_1} (\theta_t + \alpha\theta_x) + (\theta u_0)_x = 0.$$

Under the Galilean transformation $x \rightarrow x - \alpha t$, $t \rightarrow t$, equation (3.12) becomes

$$(3.13) \quad \frac{1}{\epsilon k_1} \theta_t + (\theta u_0)_x = 0.$$

We now redefine the auxiliary function θ in terms of the physical variable ρ as follows

$$(3.14) \quad \rho^{\frac{1}{a-1}} = \theta, \quad a \neq 1.$$

We substitute (3.14) into (3.13), and upon multiplying the resulting expression by $a - 1$ we find

$$(3.15) \quad \frac{1}{\epsilon k_1} \rho_t + u_0 \rho_x = (1 - a) \rho u_{0,x},$$

which up to a rescaling is the first member (3.36a) of our two component system (3.36).

3.2. The m -component. Using (3.2), we express η_x in terms of the auxiliary function θ to find

$$(3.16) \quad \eta_x = \frac{1}{2k_1\epsilon} (\theta^2)_x - \epsilon \left(\frac{2k_2 + k_1^2}{2k_1} \right) (\eta^2)_x - \delta^2 \frac{k_3}{k_1} u_{0,xxx} + \mathcal{O}(\epsilon^2, \epsilon\delta^2),$$

which when substituted into (2.7b) gives

$$\begin{aligned} (u_0 - \delta^2 \beta_0^2 u_{0xx})_t + \mathcal{O}(\epsilon^2, \epsilon\delta^2) = & -\epsilon \left[\frac{u_0^2}{2} - \frac{k_1^2 + 2k_2}{2k_1} (\eta^2) \right]_x \\ & - \delta^2 \left[\left(\frac{k_3}{k_1} + \frac{\alpha}{3} \right) u_{0,x} - k_0 u_{0,t} \right]_{xx} - \frac{1}{2\epsilon k_1} (\theta^2)_x. \end{aligned}$$

Here $\beta_0^2 = 1/2 + k_0$ and k_0 is a constant yet to be determined. Using relations (2.10) and (3.4) above, and neglecting terms of order $\mathcal{O}(\epsilon, \delta^2)$, yields

$$(3.17) \quad \left(u_0 - \delta^2 \left(\frac{1}{2} + k_0 \right) u_{0xx} \right)_t + \mathcal{O}(\epsilon^2, \epsilon\delta^2) = \\ - \epsilon \left(1 - c^2 \frac{2k_2 + k_1^2}{k_1} \right) \left(\frac{u_0^2}{2} \right)_x \\ - \delta^2 \left(\frac{k_3}{k_1} + \frac{\alpha}{3} + k_0 c \right) u_{0,xxx} - \frac{1}{2\epsilon k_1} (\theta^2)_x.$$

At this point we reexpress the θ^2 term in (3.17) in terms of ρ^2 using

$$(3.18) \quad \theta = 1 + \epsilon\theta_0,$$

where θ_0 is of order $\mathcal{O}(1, \epsilon, \delta^2)$. Retaining terms to order $\mathcal{O}(\epsilon^2, \epsilon\delta^2)$, we may use a Taylor expansion to write

$$(3.19) \quad \theta^{2(a-1)} = 1 + 2(a-1)\epsilon\theta_0 + (a-1)(2a-3)\epsilon^2\theta_0^2 + \mathcal{O}(\epsilon^3, \epsilon^2\delta^2) = \rho^2$$

and as such

$$(3.20) \quad \epsilon 2\theta_0 = \frac{1}{a-1}\rho^2 - \epsilon^2(2a-3)\theta_0^2 - \frac{1}{a-1} + \mathcal{O}(\epsilon^3, \epsilon^2\delta^2).$$

In addition it follows from (3.2) that

$$(3.21) \quad \epsilon 2\theta_0 = \theta^2 - 1 - \epsilon^2\theta_0^2.$$

Comparing (3.20) and (3.21), we find

$$(3.22) \quad \theta^2 = \frac{1}{a-1}\rho^2 - \epsilon^2 2(a-2)\theta_0^2 + \frac{a-2}{a-1} + \mathcal{O}(\epsilon^3, \epsilon^2\delta^2).$$

Moreover comparing (3.1) and (3.18), we find

$$(3.23) \quad \theta_0 = k_1\eta + \epsilon k_2\eta^2 + \delta^2 k_3 u_{0,xx},$$

which we may replace in (3.22) to obtain

$$(3.24) \quad \theta^2 = \frac{1}{a-1}\rho^2 - \epsilon^2 2c^2 k_1^2 (a-2)u_0^2 + \frac{a-2}{a-1} + \mathcal{O}(\epsilon^3, \epsilon^2\delta^2),$$

where we have used (2.10) to write $\eta^2 = c^2 u_0^2 + \mathcal{O}(\epsilon, \delta^2)$ in the term of order $\mathcal{O}(\epsilon^2)$ above. Equation (3.24) allows us to replace the auxiliary function θ^2 appearing in (3.17) with the physical variable ρ^2 , in which case we find

$$(3.25) \quad m_t + \frac{1}{2\epsilon k_1(a-1)} (\rho^2)_x + \mathcal{O}(\epsilon^2, \epsilon\delta^2) = \\ - \epsilon \left(1 - c^2 \left(\frac{k_1^2 + 2k_2}{k_1} + 2(a-2)k_1 \right) \right) \left(\frac{u_0^2}{2} \right)_x \\ - \delta^2 \left(\frac{k_3}{k_1} + \frac{\alpha}{3} + k_0 c \right) u_{0,xxx},$$

where we define,

$$(3.26) \quad m = u_0 - \delta^2 \left(\frac{1}{2} + k_0 \right) u_{0,xx}.$$

Adding the term $\alpha m_{0,x}$ to both sides of (3.25) we find

$$(3.27) \quad m_t + \alpha m_{0,x} + \frac{1}{2\epsilon k_1(a-1)}(\rho^2)_x + \mathcal{O}(\epsilon^2, \epsilon\delta^2) = \\ \alpha u_{0,x} + \epsilon \left(1 - c^2 \left(\frac{k_1^2 + 2k_2}{k_1} + (a-2)k_1 \right) \right) \left(\frac{u_0^2}{2} \right)_x \\ + \delta^2 \left(\frac{k_3}{k_1} - \frac{\alpha}{6} + k_0(c-\alpha) \right) u_{0,xxx},$$

where on the right hand side we use $m_{0,x}$ as it appears in (3.26). It was previously stated that k_0 is an undetermined coefficient, and so we may choose its value such that

$$(3.28) \quad \frac{k_3}{k_1} - \frac{\alpha}{6} + k_0(c-\alpha) = 0.$$

Combined with the Burns condition imposed on c , as determined by (2.4), along with the first constraint on k_1 and k_3 in (3.8), equation (3.28) gives

$$\beta_0^2 = k_0 + \frac{1}{2} = \frac{\alpha c - \alpha^2 - 1}{6(c-\alpha)^2} + \frac{1}{2} = \frac{1}{3c^2(c-\alpha)^2},$$

thus fixing k_0 uniquely in terms of α and c . With k_0 fixed, (3.27) becomes

$$(3.29) \quad m_t + \alpha m_x + \frac{1}{2\epsilon k_1(a-1)}(\rho^2)_x + \mathcal{O}(\epsilon^2, \epsilon\delta^2) = \\ \alpha u_{0,x} + \epsilon \left(1 - c^2 \left(\frac{k_1^2 + 2k_2}{k_1} + (a-2)k_1 \right) \right) \left(\frac{u_0^2}{2} \right)_x.$$

So far we have only imposed two constraints on the coefficients k_1 , k_2 and k_3 and so we may choose k_1 and k_2 such that,

$$(3.30) \quad k_1(1+a) = 1 - c^2 \left(\frac{k_1^2 + 2k_2}{k_1} + 2(a-2)k_1 \right),$$

which upon substitution into (3.29) gives,

$$(3.31) \quad m_t + \alpha m_x - \alpha u_{0,x} + \epsilon k_1(1+a) \left(\frac{u_0^2}{2} \right)_x \\ + \frac{1}{2\epsilon k_1(a-1)}(\rho^2)_x + \mathcal{O}(\epsilon^2, \epsilon\delta^2) = 0.$$

Furthermore applying the Galilean transformation introduced in Section 3.1 to (3.31) gives

$$(3.32) \quad m_t - \alpha u_{0,x} + \epsilon k_1(1+a) \left(\frac{u_0^2}{2} \right)_x \\ + \frac{1}{2\epsilon k_1(a-1)}(\rho^2)_x + \mathcal{O}(\epsilon^2, \epsilon\delta^2) = 0.$$

We also note that to leading order we have $u_0 = m + \mathcal{O}(\delta^2)$, and so we may write

$$(3.33) \quad \left(\frac{u_0^2}{2} \right)_x = \frac{1}{1+a}(\alpha u_{0,x}m + u_0m_x) + \mathcal{O}(\delta^2),$$

and accordingly (3.31) may be written as

$$(3.34) \quad m_t - \alpha u_{0,x} + \epsilon a u_{0,x} m + \epsilon u_0 m_x + \frac{1}{2\epsilon k_1(a-1)}(\rho^2)_x + \mathcal{O}(\epsilon^2, \epsilon\delta^2) = 0.$$

This is the second member (3.36b) of our two component system (3.36), up to the following rescaling:

$$(3.35) \quad x \rightarrow \beta_0 x, \quad t \rightarrow \beta_0 t, \quad u \rightarrow \frac{1}{2\epsilon k_1} u, \quad \rho \rightarrow \sqrt{\frac{\kappa}{2}(a-1)} \rho,$$

where $\kappa > 0$ is an arbitrary parameter. With regard to this rescaling, we note that it is important here that u, m be of order $O(\epsilon)$, otherwise following the rescaling u, m would be $O(1/\epsilon)$ and all subsequent considerations would fail. Under this rescaling (3.15) and (3.34) become,

$$(3.36a) \quad \rho_t + u\rho_x + (a-1)u_x\rho = 0,$$

$$(3.36b) \quad m_t - \alpha u_{0,x} + a u_{0,x} m + u_0 m_x + \kappa \rho \rho_x = 0,$$

where terms of order $\mathcal{O}(\epsilon, \delta^2)$ and higher have been neglected. The coefficients k_1, k_2 and k_3 are completely determined by the constraints (3.8), (3.10) and (3.30) in terms of a, c , and α as follows:

$$\begin{aligned} k_1 &= \frac{1}{(1+c^2)(a+1)} + \frac{c^2}{a+1}, \\ k_2 &= \left(\frac{1}{(a+1)(1+c^2)} + \frac{c^2(1-a)}{2(a+1)} - \frac{1}{2} \right) k_1, \\ k_3 &= \frac{k_1}{6(c-\alpha)}. \end{aligned}$$

Following minor relabelling, equations (3.36) are transformed to (1.1), which constitutes the two component system that will be investigated in the remainder of this article.

Remark 1. In the particular case $a = 2$ we reduce to the integrable two-component equation which was derived in [30].

Remark 2. When $a = \frac{c^4+1}{(c^2+1)^2}$ we have $k_2 = 0$, and as a result of this a number of transformations become linear, in particular (3.1). Accordingly, in this setting it is convenient to express the physical variable η in terms of the auxiliary variables θ or ρ , which should be useful in practical applications.

4. GEOMETRIC REFORMULATION AS A GEODESIC FLOW

In this section, we recast the system (1.1) as a geodesic flow on the tangent bundle of a suitable infinite-dimensional Lie group. For simplicity, we will focus on the periodic case (i.e on the circle \mathbb{S}^1), but there is no obstacle to working on the non-periodic case (i.e on the real line), provided we correctly impose some conditions at infinity. When $\alpha = 0, a = 2$ and $\kappa = 1$, the system (3.36) reduces to the *two component Camassa–Holm equation* (2CH)

$$(4.1) \quad \begin{cases} m_t + u m_x = -2u_x m - \rho \rho_x, \\ \rho_t + u \rho_x = -u_x \rho, \end{cases}$$

where $m = u - u_{xx}$ and which was first derived in [38]. In [16], the system (4.1) was recast as the geodesic equations for a right-invariant Riemannian metric on a semi-direct product $\text{Diff}^\infty(\mathbb{S}^1) \circledast C^\infty(\mathbb{S}^1)$, where $\text{Diff}^\infty(\mathbb{S}^1)$ is the group of orientation-preserving, smooth diffeomorphisms of the circle and $C^\infty(\mathbb{S}^1)$ is the space of real, smooth functions on \mathbb{S}^1 . The full system (1.1) (with $\alpha = 0$) has been studied in the short-wave limit $m = -u_{xx}$ in [15]. It is shown there that geometrically the two-component system (1.1) corresponds to a geodesic flow with respect to a linear, right-invariant, symmetric connection on a suitable semi-direct product. In addition, if $a = 2$ the connection is compatible with the metric induced by the norm

$$\|u_{xx}\|_{L^2(\mathbb{S}^1)} + \|\rho\|_{L^2(\mathbb{S}^1)},$$

where $u \in C^\infty(\mathbb{S}^1)/\mathbb{R}$ and $\rho \in C^\infty(\mathbb{S}^1)$. We find here for (1.1) a similar situation: if $a = 2$ the geodesic flow is metric (cf. Theorem 4.1) and if $a \neq 2$, the system (1.1) cannot be realized as a *metric* geodesic flow, at least for a large class of inertia operators.

The above described geometric picture was in fact developed in [17], where it had been shown that *every quadratic evolution equation* defined on the Lie algebra \mathfrak{g} of a Lie group G can be interpreted as the *geodesic equations* for a *right-invariant linear connection* on G , although in this case the connection does not derive necessarily from an invariant metric. To be able to use this framework, we rewrite the system (3.36) as

$$(4.2) \quad \begin{cases} m_t = \alpha u_x - a u_x m - u m_x - \kappa \rho \rho_x, \\ \rho_t = -u \rho_x - (a - 1) u_x \rho, \\ \alpha_t = 0, \end{cases}$$

which is a quadratic evolution equation in the variables

$$(u, \rho, \alpha) \in C^\infty(\mathbb{S}^1) \times C^\infty(\mathbb{S}^1) \times \mathbb{R}.$$

We will consider the Fréchet Lie group

$$C^\infty G := (\text{Diff}^\infty(\mathbb{S}^1) \circledast C^\infty(\mathbb{S}^1)) \times \mathbb{R},$$

where the group product is given by

$$(\varphi_1, f_1, s_1) * (\varphi_2, f_2, s_2) := (\varphi_1 \circ \varphi_2, f_2 + f_1 \circ \varphi_2, s_1 + s_2)$$

for $(\varphi_1, f_1, s_1), (\varphi_2, f_2, s_2) \in \text{Diff}^\infty(\mathbb{S}^1) \times C^\infty(\mathbb{S}^1) \times \mathbb{R}$, and where \circ denotes composition of mappings. The neutral element is $(\text{id}, 0, 0)$ and the inverse of an element (φ, f, s) is given by

$$(\varphi, f, s)^{-1} = (\varphi^{-1}, -f \circ \varphi^{-1}, -s).$$

The Lie algebra of $C^\infty G$ is the vector space $C^\infty \mathfrak{g} := C^\infty(\mathbb{S}^1) \oplus C^\infty(\mathbb{S}^1) \oplus \mathbb{R}$ endowed with the Lie bracket

$$[(u_1, \rho_1, \alpha_1), (u_2, \rho_2, \alpha_2)] := (u_{1,x} u_2 - u_{2,x} u_1, \rho_{1,x} u_2 - \rho_{2,x} u_1, 0),$$

for $(u_1, \rho_1, \alpha_1), (u_2, \rho_2, \alpha_2) \in C^\infty \mathfrak{g}$.

Let R_g denote the right translation on $C^\infty G$ by the element $g = (\varphi_2, f_2, s_2)$, and $T_h R_g$ be its tangent map at $h = (\varphi_1, f_1, s_1)$. Then, we have

$$(4.3) \quad T_h R_g \cdot V = (v_1 \circ \varphi_2, \sigma_1 \circ \varphi_2, \alpha_1),$$

where $V = (v_1, \sigma_1, \alpha_1)$. Note that its expression does not depend on h . In particular, we will have a similar expression for the second order tangent map. More precisely, if $X = (\delta\varphi_1, \delta f_1, \delta s_1, \delta v, \delta\sigma, \delta\alpha)$, we get

$$(4.4) \quad T_V(TR_g).X = (\delta\varphi_1 \circ \varphi_2, \delta f_1 \circ \varphi_2, \delta s_1, \delta v_1 \circ \varphi_2, \delta\sigma_1 \circ \varphi_2, \delta\alpha_1).$$

Given a smooth path $g(t) := (\varphi(t), f(t), s(t)) \in C^\infty G$, we define its *Eulerian velocity*, which lies in the Lie algebra $C^\infty \mathfrak{g}$, by

$$U(t) = TR_{g^{-1}(t)}.g_t(t).$$

Let (u, ρ, α) be the three components of U . Then, using (4.3), we get

$$u = \varphi_t \circ \varphi^{-1}, \quad \rho = f_t \circ \varphi^{-1}, \quad \alpha = s_t.$$

Introducing the *Lagrangian velocity* $V := (v, \sigma, \alpha)$, where

$$v := \varphi_t = u \circ \varphi, \quad \sigma := f_t = \rho \circ \varphi, \quad \alpha := s_t,$$

we have

$$v_t = (u_t + uu_x) \circ \varphi, \quad \sigma_t = (\rho_t + u\rho_x) \circ \varphi,$$

and we may rewrite equation (4.2) as

$$\begin{cases} v_t = (A^{-1}([A, u]u_x + \alpha u_x - au_x Au - \kappa\rho\rho_x)) \circ \varphi, \\ \sigma_t = ((1-a)u_x\rho) \circ \varphi, \\ \alpha_t = 0, \end{cases}$$

where $A = 1 - D^2$ and $[A, u]v := A(uv) - uA(v)$. Therefore, the system (4.2) is equivalent to

$$(4.5) \quad \begin{cases} \partial_t(\varphi, f, s) = (v, \sigma, \alpha), \\ \partial_t(v, \sigma, \alpha) = S_{(\varphi, f, s)}(v, \sigma, \alpha), \end{cases}$$

where

$$(4.6) \quad S_{(\varphi, f, s)} := TTR_{(\varphi, f, s)} \circ S \circ TR_{(\varphi, f, s)}^{-1},$$

and

$$(4.7) \quad S(u, \rho, \alpha) := S_{(\text{id}, 0, 0)}(u, \rho, \alpha) \\ \left(A^{-1}([A, u]u_x + \alpha u_x - au_x Au - \kappa\rho\rho_x), (1-a)u_x\rho, 0 \right).$$

The *second order vector field*

$$(4.8) \quad F(g, V) = (g, V, V, S_g(V))$$

defined on

$$TC^\infty G \simeq (\text{Diff}^\infty(\mathbb{S}^1) \times C^\infty(\mathbb{S}^1) \times \mathbb{R}) \times (C^\infty(\mathbb{S}^1) \oplus C^\infty(\mathbb{S}^1) \oplus \mathbb{R})$$

is quadratic on V and is called a *spray*. It corresponds to the *geodesic flow of a symmetric linear connection* on $C^\infty G$ (see [36, Section IV.3]).

Remark 3. Note that the relation (4.6) is just the fourth component of the equation

$$F(TR_g.U) = TTR_g.F(U),$$

where $g = (\varphi, f, s)$, $U = (u, \rho, \alpha)$ and which traduces the right-invariance of the spray $F : TC^\infty G \rightarrow TTC^\infty G$.

When the parameter $a = 2$, the spray F derives from a right-invariant metric on $C^\infty G$ and the system of equations (4.2) corresponds to the *Arnold-Euler equations* of this metric. Let us briefly recall this formalism. Given a Lie group G and its Lie algebra \mathfrak{g} , any inner product on \mathfrak{g} induces a *right-invariant* Riemannian metric on G by extending it by right-translation. If this inner product on \mathfrak{g} is represented by an invertible operator $\mathbb{A} : \mathfrak{g} \rightarrow \mathfrak{g}^*$, for historical reasons, going back to the work of Euler on the motion of the rigid body, this operator \mathbb{A} is called the *inertia operator* of the system. Then a path $g(t)$ on G is a geodesic for this right-invariant Riemannian metric, if and only if, its Eulerian velocity $u(t) := TR_{g(t)^{-1}}.\dot{g}(t)$ is solution of the so-called Arnold-Euler equation

$$u_t = -\mathbb{B}(u, u),$$

where

$$\mathbb{B}(u_1, u_2) := \frac{1}{2} \left(\text{ad}_{u_1}^\top u_2 + \text{ad}_{u_2}^\top u_1 \right),$$

where $u_1, u_2 \in \mathfrak{g}$ and ad_u^\top is the adjoint of the operator ad_u , with respect to \mathbb{A} (see [1] for instance).

Theorem 4.1. *Given $\kappa > 0$, the system (4.2) is, for $a = 2$, the Arnold-Euler equation on the Lie algebra $C^\infty \mathfrak{g}$ of the Fréchet Lie group $C^\infty G$ associated to the inner product*

$$(4.9) \quad \langle (u_1, \rho_1, \alpha_1), (u_2, \rho_2, \alpha_2) \rangle = \int_{\mathbb{S}^1} u_1 u_2 dx + \int_{\mathbb{S}^1} u_{1,x} u_{2,x} dx \\ + \kappa \int_{\mathbb{S}^1} \rho_1 \rho_2 dx - \frac{1}{2} \int_{\mathbb{S}^1} (u_1 \alpha_2 + u_2 \alpha_1) dx + \frac{1}{2} \alpha_1 \alpha_2,$$

where $u_i, \rho_i \in C^\infty(\mathbb{S}^1)$ and $\alpha_i \in \mathbb{R}$, for $i = 1, 2$.

Remark 4. Note that the norm induced by (4.9) on $C^\infty \mathfrak{g}$

$$\|(u, \rho, \alpha)\|^2 = \int_{\mathbb{S}^1} u_x^2 dx + \int_{\mathbb{S}^1} \left(u - \frac{\alpha}{2} \right)^2 dx + \frac{\alpha^2}{2} + \int_{\mathbb{S}^1} \kappa \rho^2 dx$$

is equivalent to the Hilbert-norm

$$\|(u, \rho, \alpha)\|^2 = \|u\|_{H^1}^2 + \|\rho\|_{L^2}^2 + |\alpha|^2.$$

Proof. The inner product (4.9) can be rewritten as

$$\langle (u_1, \rho_1, \alpha_1), (u_2, \rho_2, \alpha_2) \rangle = \int_{\mathbb{S}^1} (u_1, \rho_1, \alpha_1) \cdot \mathbb{A}(u_2, \rho_2, \alpha_2) dx$$

where the inertia operator \mathbb{A} is defined by

$$\mathbb{A}(u, \rho, \alpha) = \left(Au - \frac{\alpha}{2}, \kappa \rho, \frac{1}{2} \left(\alpha - \int_{\mathbb{S}^1} u \right) \right), \quad (u, \alpha, \rho) \in C^\infty \mathfrak{g},$$

$Au := u - u_{xx}$, and \cdot stands for the usual inner product in \mathbb{R}^3 . Set $U_i = (u_i, \rho_i, \alpha_i)$, for $i = 1, 2, 3$. We have

$$\text{ad}_{U_1} U_2 = (u_{1,x} u_2 - u_{2,x} u_1, \rho_{1,x} u_2 - \rho_{2,x} u_1, 0),$$

and therefore, after some integrations by parts, we get

$$\langle \text{ad}_{U_1} U_2, U_3 \rangle = \int_{\mathbb{S}^1} (u_2, \rho_2, \alpha_2) \cdot (f, g, 0) dx,$$

where

$$f = 2u_{1,x}Au_3 + u_1Au_{3,x} - \alpha_3u_{1,x} + \kappa\rho_{1,x}\rho_3, \quad g = \kappa(u_1\rho_3)_x.$$

Note now that the equation

$$\mathbb{A}(\tilde{u}, \tilde{\rho}, \tilde{\alpha}) = (f, g, 0)$$

has a unique solution which is given by

$$\tilde{u} = A^{-1}f + \int_{\mathbb{S}^1} f dx, \quad \tilde{\rho} = \frac{1}{\kappa}g, \quad \tilde{\alpha} = 2 \int_{\mathbb{S}^1} f dx.$$

Thus we have $\text{ad}_{U_1}^\top U_3 = (\tilde{u}, \tilde{\rho}, \tilde{\alpha})$, where

$$\begin{aligned} \tilde{u} = A^{-1} \left(2u_{1,x}Au_3 + u_1Au_{3,x} - \alpha_3u_{1,x} + \kappa\rho_{1,x}\rho_3 \right) \\ + \int_{\mathbb{S}^1} (u_{1,x}Au_3 + \kappa\rho_{1,x}\rho_3) dx, \end{aligned}$$

and

$$\tilde{\rho} = (u_1\rho_3)_x, \quad \tilde{\alpha} = 2 \int_{\mathbb{S}^1} (u_{1,x}Au_3 + \kappa\rho_{1,x}\rho_3) dx.$$

We conclude therefore that

$$\mathbb{B}(U, U) = \left(A^{-1}(2u_xAu + uAu_x - \alpha u_x + \kappa\rho_x\rho), (u\rho)_x, 0 \right),$$

where $U = (u, \rho, \alpha)$ and that the equation

$$U_t = -\mathbb{B}(U, U)$$

is equivalent to (4.2), when $a = 2$. \square

5. WELL-POSEDNESS OF THE EQUATION

Our strategy will be to study the Cauchy problem for the geodesic equations (4.5). Following Ebin & Marsden's approach [14], if we can prove *local existence and uniqueness of geodesics* of the ODE (4.5) on TG , then the PDE (4.2) is *well-posed*. To do so, we need to introduce an approximation of the Fréchet–Lie group $\text{Diff}^\infty(\mathbb{S}^1)$ by *Hilbert manifolds*. Let $H^q(\mathbb{S}^1)$ be the completion of $C^\infty(\mathbb{S}^1)$ for the norm

$$\|u\|_{H^q} := \left(\sum_{k \in \mathbb{Z}} (1 + k^2)^q |\hat{u}_k|^2 \right)^{1/2},$$

where $q \in \mathbb{R}, q \geq 0$. We recall that $H^q(\mathbb{S}^1)$ is a multiplicative algebra for $q > 1/2$ (cf. [40, Theorem 2.8.3]). This means that

$$\|uv\|_{H^q(\mathbb{S}^1)} \lesssim \|u\|_{H^q(\mathbb{S}^1)} \|v\|_{H^q(\mathbb{S}^1)}, \quad u, v \in H^q(\mathbb{S}^1).$$

A C^1 diffeomorphism φ of \mathbb{S}^1 is of class H^q if for any of its lifts to \mathbb{R} , $\tilde{\varphi}$, we have

$$\tilde{\varphi} - \text{id} \in H^q(\mathbb{S}^1).$$

For $q > 3/2$, the set $\mathcal{D}^q(\mathbb{S}^1)$ of C^1 -diffeomorphisms of the circle which are of class H^q has the structure of a *Hilbert manifold*, modelled on $H^q(\mathbb{S}^1)$ (see [14] or [29]). Contrary to $\text{Diff}^\infty(\mathbb{S}^1)$, the manifold $\mathcal{D}^q(\mathbb{S}^1)$ is only a *topological group* and *not a Lie group* (composition and inversion in $\mathcal{D}^q(\mathbb{S}^1)$

are continuous but *not differentiable*). More precisely, the following regularity properties for $q > 3/2$ are well-known (see [29] for instance).

(1) The mapping

$$u \mapsto R_\varphi(u) := u \circ \varphi, \quad H^q(\mathbb{S}^1) \rightarrow H^q(\mathbb{S}^1)$$

is smooth, for any $\varphi \in \mathcal{D}^q(\mathbb{S}^1)$.

(2) The mapping

$$(u, \varphi) \mapsto u \circ \varphi, \quad H^{q+k}(\mathbb{S}^1) \times \mathcal{D}^q(\mathbb{S}^1) \rightarrow H^q(\mathbb{S}^1)$$

is of class C^k .

(3) The mapping

$$\varphi \mapsto \varphi^{-1}, \quad \mathcal{D}^{q+k}(\mathbb{S}^1) \rightarrow \mathcal{D}^q(\mathbb{S}^1)$$

is of class C^k .

Moreover the following slightly sharpen result was established in [18, Corollary B.6].

Lemma 5.1. *Let $q > 3/2$. Then, the mappings*

$$(\varphi, v) \mapsto v \circ \varphi, \quad \mathcal{D}^q(\mathbb{S}^1) \times H^q(\mathbb{S}^1) \rightarrow H^{q-1}(\mathbb{S}^1)$$

and

$$(\varphi, v) \mapsto v \circ \varphi^{-1}, \quad \mathcal{D}^q(\mathbb{S}^1) \times H^q(\mathbb{S}^1) \rightarrow H^{q-1}(\mathbb{S}^1)$$

are C^1 .

Remark 5. The tangent bundle of the Hilbert manifold $\mathcal{D}^q(\mathbb{S}^1)$ is trivial because if $\mathfrak{t} : T\mathbb{S}^1 \rightarrow \mathbb{S}^1 \times \mathbb{R}$ is a smooth trivialisation of the tangent bundle of the circle, then

$$\Psi : T\mathcal{D}^q(\mathbb{S}^1) \rightarrow \mathcal{D}^q(\mathbb{S}^1) \times H^q(\mathbb{S}^1), \quad \xi \mapsto \mathfrak{t} \circ \xi$$

defines a smooth vector bundle isomorphism (see [14, Page 107]).

As suggested in [16], we shall use the following choice for the Hilbert approximation of $C^\infty G$:

$$H^q G := \mathcal{D}^q(\mathbb{S}^1) \times H^{q-1}(\mathbb{S}^1) \times \mathbb{R},$$

which is defined for $q > 3/2$.

Theorem 5.2. *The spray F defined on $T\text{Diff}^\infty(\mathbb{S}^1)$ by (4.8) extends smoothly to a spray*

$$F_q : TH^q G \rightarrow TTH^q G,$$

for each $q > 3/2$.

Proof. We first observe that the quadratic operator S defined in (4.7) can be rewritten as

$$S(u, \rho, \alpha) = \left(\frac{1}{2} A^{-1} D (2\alpha u - \kappa \rho^2 + (a-3)u_x^2 - au^2), (1-a)\rho u_x, 0 \right),$$

where D stands for d/dx . Therefore, the two non-vanishing components S^1 , S^2 of $S_{(\varphi, f, s)}(v, \sigma, \alpha)$ are

$$S^1 = \frac{1}{2} (A^{-1} D)_\varphi \left(2\alpha v - \kappa \sigma^2 + (a-3) \frac{1}{\varphi_x^2} v_x^2 - av^2 \right)$$

and

$$S^2 = (1 - a) \frac{1}{\varphi_x} \sigma v_x,$$

where

$$(A^{-1}D)_\varphi := R_\varphi (A^{-1}D) R_{\varphi^{-1}}.$$

Let $q > 3/2$, $\varphi \in \mathcal{D}^q(\mathbb{S}^1)$, $v \in H^q(\mathbb{S}^1)$ and $\sigma \in H^{q-1}(\mathbb{S}^1)$. Then, the expression S^2 belongs to $H^{q-1}(\mathbb{S}^1)$ because $H^{q-1}(\mathbb{S}^1)$ is a multiplicative algebra and the expression S^1 belongs to $H^q(\mathbb{S}^1)$ because, furthermore,

$$R_{\varphi^{-1}} \in \mathcal{L}(H^{q-1}(\mathbb{S}^1), H^{q-1}(\mathbb{S}^1)), \quad R_\varphi \in \mathcal{L}(H^q(\mathbb{S}^1), H^q(\mathbb{S}^1))$$

by [18, Lemma B.2], and

$$A^{-1}D = DA^{-1} \in \mathcal{L}(H^{q-1}(\mathbb{S}^1), H^q(\mathbb{S}^1)).$$

It remains to be shown that S^1 and S^2 depend smoothly on φ, v, σ , which will achieve the proof. First, it is clear that the mapping

$$(\varphi, v, \sigma) \mapsto (1 - a) \frac{1}{\varphi_x} \sigma v_x, \quad \mathcal{D}^q(\mathbb{S}^1) \times H^q(\mathbb{S}^1) \times H^{q-1}(\mathbb{S}^1) \rightarrow H^{q-1}(\mathbb{S}^1)$$

is smooth because $H^{q-1}(\mathbb{S}^1)$ is a multiplicative algebra. For the same reason, it is clear that

$$(\varphi, v, \sigma) \mapsto 2\alpha v - \kappa \sigma^2 + (a - 3) \frac{1}{\varphi_x^2} v_x^2 - av^2,$$

from $\mathcal{D}^q(\mathbb{S}^1) \times H^q(\mathbb{S}^1) \times H^{q-1}(\mathbb{S}^1)$ to $H^{q-1}(\mathbb{S}^1)$ is smooth. Therefore, if we can prove that

$$(\varphi, w) \mapsto (A^{-1}D)_\varphi w, \quad \mathcal{D}^q(\mathbb{S}^1) \times H^{q-1}(\mathbb{S}^1) \rightarrow H^q(\mathbb{S}^1)$$

is smooth, we are done, using the chain rule. Now, observe that $(1 - D)$ and $(1 + D)$ belong to $\text{Isom}(H^q(\mathbb{S}^1), H^{q-1}(\mathbb{S}^1))$, for $q \geq 1$, and that

$$A^{-1}D = \frac{1}{2} ((1 - D)^{-1} - (1 + D)^{-1}),$$

because $A = (1 - D)(1 + D)$. Therefore

$$(A^{-1}D)_\varphi = \frac{1}{2} ((1 - D)_\varphi^{-1} - (1 + D)_\varphi^{-1}).$$

But

$$D_\varphi = \frac{1}{\varphi_x} D$$

and hence

$$\varphi \mapsto (1 \pm D)_\varphi$$

from $\mathcal{D}^q(\mathbb{S}^1)$ to $\text{Isom}(H^q(\mathbb{S}^1), H^{q-1}(\mathbb{S}^1))$ is smooth. Moreover, since the set $\text{Isom}(H^q(\mathbb{S}^1), H^{q-1}(\mathbb{S}^1))$ is open in $\mathcal{L}(H^q(\mathbb{S}^1), H^{q-1}(\mathbb{S}^1))$ and the mapping

$$P \mapsto P^{-1}, \quad \text{Isom}(H^q(\mathbb{S}^1), H^{q-1}(\mathbb{S}^1)) \rightarrow \text{Isom}(H^{q-1}(\mathbb{S}^1), H^q(\mathbb{S}^1))$$

is smooth, we conclude that $(1 - D)_\varphi^{-1} w$ and $(1 + D)_\varphi^{-1} w$ both depend smoothly on (φ, w) . \square

Applying the Cauchy–Lipschitz (or Picard–Lindelöf) theorem in the Hilbert manifold TH^qG , we obtain the following local-existence result for the geodesics.

Corollary 5.3. *Let $q > 3/2$. For each initial data $(g_0, V_0) \in TH^qG$, there exists a unique non-extendable solution*

$$(g, V) \in C^\infty(J_q, TH^qG),$$

of the Cauchy problem (4.5) for the geodesic equations, with $g(0) = g_0$ and $V(0) = V_0$, defined on some maximal interval of existence $J_q = J_q(g_0, V_0)$, which is open and contains 0. Moreover, the solution depends smoothly on the initial data.

Finally, taking account of lemma 5.1, we obtain the following.

Corollary 5.4. *Let $q > 5/2$ and $\alpha \in \mathbb{R}$. For each initial data*

$$(u_0, \rho_0) \in H^q(\mathbb{S}^1) \oplus H^{q-1}(\mathbb{S}^1),$$

there exists a unique non-extendable solution

$$(u, \rho) \in C^0(J_q, H^q(\mathbb{S}^1) \oplus H^{q-1}(\mathbb{S}^1)) \cap C^1(J_q, H^{q-1}(\mathbb{S}^1) \oplus H^{q-2}(\mathbb{S}^1)),$$

of the evolution equation (4.2), with $(u(0), \rho(0)) = (u_0, \rho_0)$, defined on some maximal interval of existence $J_q = J_q(u_0, \rho_0)$, which is open and contains 0. Moreover, the solution depends continuously on the initial data.

Furthermore, using the right-invariance of the spray, an argument due to Ebin & Marsden [14] implies that the interval of existence J_q in theorem 5.3 is independent of the Sobolev index q and that for any smooth initial condition the solution exists in the smooth category. A proof of the following result follows along the lines of the proof of [18, Theorem 1.1]. We will only sketch it. Let Φ_q be the flow of the spray F_q and R_τ be the (right) action of the rotation group \mathbb{S}^1 on H^qG , defined by

$$(R_\tau \cdot (\varphi, f, s))(x) := (\varphi(x + \tau), f(x + \tau), s),$$

where $(\varphi, f, s) \in H^qG$. Note that if $V = (\varphi, f, s, v, \sigma, \alpha) \in T\mathcal{D}^{q+1}(\mathbb{S}^1)$, then

$$s \mapsto TR_s \cdot V, \quad \mathbb{S}^1 \rightarrow TH^qG$$

is a C^1 map, and that

$$\frac{d}{ds} R_\tau \cdot V = (\varphi_x, f_x, 0, v_x, \sigma_x, 0).$$

Moreover, since the spray F_q and its flow Φ_q are invariant under this action, we get

$$T_{V_0} \Phi_q(t) \cdot (\varphi_{0,x}, f_{0,x}, 0, v_{0,x}, \sigma_{0,x}, 0) = (\varphi_x(t), f_x(t), 0, v_x(t), \sigma_x(t), 0),$$

from which we deduce that if $V_0 \in TH^{q+1}G$ then so is $V(t)$, and conversely (by reversing time). We therefore also have a well-posedness result for (4.2) in the smooth category.

Corollary 5.5. *Let $\alpha \in \mathbb{R}$ and $(u_0, \rho_0) \in C^\infty(\mathbb{S}^1) \times C^\infty(\mathbb{S}^1)$ be given. Then, there exists an open interval J with $0 \in J$ and a unique non-extendable solution*

$$(u, \rho) \in C^\infty(J, C^\infty(\mathbb{S}^1) \times C^\infty(\mathbb{S}^1))$$

of (4.2) with initial datum (u_0, ρ_0) .

6. GLOBAL SOLUTIONS

Given the fact that the Camassa-Holm equation is embedded in the system (1.1), classical results, cf. [6, 9, 8] make clear that not all solutions to (1.1) can exist globally. Global in time solutions to (1.1) in the case $a = 2$, $\alpha = 0$, $\kappa = 1$ have been constructed in [10]. In this section, we will not construct global solutions, but we provide a criterion for a strong solution to the full system (1.1) to be defined for all time. Most of the proofs are similar to the ones given in [19] and will only be sketched. Throughout this section, we work in the smooth category, and we let

$$(g, V) \in C^\infty(J, TC^\infty G)$$

be the unique solution of the Cauchy problem (4.5), emanating from

$$(g_0, V_0) \in TC^\infty G,$$

and defined on the *maximal time interval* J . The corresponding solution (u, ρ) of the Arnold-Euler equation (4.2) is a path

$$(u, \rho) \in C^\infty(J, C^\infty(\mathbb{S}^1) \oplus C^\infty(\mathbb{S}^1)),$$

with $(u(0), \rho(0)) = (u_0, \rho_0)$. The *momentum* $m(t) := Au(t)$ is defined as a path

$$m \in C^\infty(J, C^\infty(\mathbb{S}^1)).$$

We will now establish several *a priori* estimates that will lead to a global existence result.

Lemma 6.1. *We have*

$$(\rho \circ \varphi)(t) \cdot \varphi_x^{a-1}(t) = (\rho \circ \varphi)(0) \cdot \varphi_x^{a-1}(0),$$

for all $t \in J$.

Proof. Let φ be the flow of the time dependent vector field u . We have

$$\frac{d}{dt} (\rho \circ \varphi \cdot \varphi_x^{a-1}) = (\rho_t + \rho_x u + (a-1)\rho u_x) \circ \varphi \cdot \varphi_x^{a-1},$$

which vanishes because (u, ρ) solves (4.2). \square

Corollary 6.2. *Suppose that $\rho_0 > 0$. Then $\rho(t) > 0$ for all $t \in J$.*

Proof. Due to lemma 6.1, we have

$$(\rho \circ \varphi)(t) = (\rho \circ \varphi)(0) \frac{\varphi_x^{a-1}(0)}{\varphi_x^{a-1}(t)} > 0,$$

which achieves the proof because φ is an orientation-preserving diffeomorphism. \square

Corollary 6.3. *Suppose that $\|u_x(t)\|_\infty$ is bounded on any bounded subinterval of J . Then $\|\rho(t)\|_\infty$ is bounded on any bounded subinterval of J .*

Proof. Let φ be the flow of the time dependent vector field u and set

$$\alpha(t) = \max_{x \in \mathbb{S}^1} 1/\varphi_x(t).$$

Note that α is a continuous function. Let I denote any bounded subinterval of J , and set

$$K = \sup_{t \in I} \|u_x(t)\|_\infty.$$

From equation $\varphi_t = u \circ \varphi$, we deduce that

$$(1/\varphi_x)_t = -(u_x \circ \varphi)/\varphi_x,$$

and therefore, we have

$$\alpha(t) \leq \alpha(0) + K \int_0^t \alpha(s) ds.$$

Thus $\|1/\varphi_x(t)\|_\infty$ is bounded on I due to Gronwall's lemma and the conclusion follows from lemma 6.1. \square

Lemma 6.4. *Suppose that $\|u_x(t)\|_\infty$ is bounded on any bounded subinterval of J . Then*

$$\|m(t)\|_{H^k}^2 + \|\rho(t)\|_{H^{k+1}}^2$$

is bounded on any bounded subinterval of J , for all non-negative integer k .

Proof. We have

$$\begin{aligned} \frac{d}{dt} \left(\|m\|_{L^2}^2 + \|\rho\|_{H^1}^2 \right) &= \alpha \int u_x m - a \int u_x m^2 - \int u m m_x \\ &\quad - \kappa \int \rho \rho_x m - \int u \rho \rho_x - (a-1) \int u_x \rho^2 - \int u_x \rho_x^2 \\ &\quad - \int u \rho_x \rho_{xx} - (a-1) \int u_{xx} \rho \rho_x - (a-1) \int u_x \rho_x^2. \end{aligned}$$

After integrating by parts the terms

$$- \int u m m_x = \frac{1}{2} \int u_x m^2, \quad \text{and} \quad - \int u \rho_x \rho_{xx} = \frac{1}{2} \int u_x \rho_x^2,$$

using the Cauchy-Schwarz inequality, and the fact that $\|u\|_{H^2} \lesssim \|m\|_{L^2}$, we obtain

$$\frac{d}{dt} \left(\|m\|_{L^2}^2 + \|\rho\|_{H^1}^2 \right) \lesssim (1 + \|u_x\|_\infty + \|\rho\|_\infty) \left(\|m\|_{L^2}^2 + \|\rho\|_{H^1}^2 \right)$$

and the conclusion holds from corollary 6.3 and by Gronwall's lemma. Suppose now that $k \geq 1$. We will show by induction on k that

$$(6.1) \quad \frac{d}{dt} \left(\|m\|_{H^k}^2 + \|\rho\|_{H^{k+1}}^2 \right) \lesssim (1 + \|m\|_{H^{k-1}} + \|\rho\|_{H^k}) \left(\|m\|_{H^k}^2 + \|\rho\|_{H^{k+1}}^2 \right),$$

which will achieve the proof, using Gronwall's lemma. For $k = 1$, we only need to estimate the term

$$\int m_x m_{tx} + \int \rho_{xx} \rho_{txx},$$

because

$$\|u_x\|_\infty + \|\rho\|_\infty \lesssim \|m\|_{L^2} + \|\rho\|_{H^1}.$$

We have first

$$\begin{aligned} \int m_x m_{tx} &= \alpha \int u_{xx} m_x - a \int u_{xx} m m_x - a \int u_x m_x^2 \\ &\quad - \int u_x m_x^2 - \int u m_{xx} m_x - \kappa \int \rho_x^2 m_x - \kappa \int \rho \rho_{xx} m_x. \end{aligned}$$

Integrating by parts the term

$$-\int um_{xx}m_x = \frac{1}{2} \int u_x m_x^2,$$

using the Cauchy–Schwarz inequality and the fact that $H^1(\mathbb{S}^1)$ is a multiplicative algebra, we get

$$\begin{aligned} \int m_x m_{tx} &\lesssim \|u\|_{H^2} \|m\|_{H^1} + \|u\|_{H^2} \|m\|_{H^1}^2 + \|u_x\|_{\infty} \|m\|_{H^1}^2 \\ &\quad + \|\rho_x^2\|_{L^2} \|m\|_{H^1} + \|\rho\|_{\infty} \|\rho\|_{H^2} \|m\|_{H^1}. \end{aligned}$$

Now, using the following estimates

$$\|u\|_{H^2} \lesssim \|m\|_{L^2}, \quad \|u_x\|_{\infty} \lesssim \|m\|_{L^2}, \quad \|\rho\|_{\infty} \lesssim \|\rho\|_{H^1},$$

together with

$$\|\rho_x^2\|_{L^2} \lesssim \|\rho\|_{H^1} \|\rho\|_{H^2},$$

we obtain

$$\int m_x m_{tx} \lesssim (1 + \|m\|_{L^2} + \|\rho\|_{H^1}) \left(\|m\|_{H^1}^2 + \|\rho\|_{H^2}^2 \right).$$

We have next

$$\begin{aligned} \int \rho_{xx} \rho_{txx} &= - \int u_{xx} \rho_x \rho_{xx} - 2 \int u_x \rho_{xx}^2 - \int u \rho_{xxx} \rho_{xx} \\ &\quad - (a-1) \int u_{xxx} \rho \rho_{xx} - 2(a-1) \int u_{xx} \rho_x \rho_{xx} - (a-1) \int u_x \rho_{xx}^2, \end{aligned}$$

from which we deduce

$$\begin{aligned} \int \rho_{xx} \rho_{txx} &\lesssim \|u\|_{H^2} \|\rho_x^2\|_{H^1} + \|u_x\|_{\infty} \|\rho\|_{H^2}^2 + \|u_x\|_{\infty} \|\rho\|_{H^2}^2 \\ &\quad + \|\rho\|_{\infty} \|u\|_{H^3} \|\rho\|_{H^2} + \|u\|_{H^2} \|\rho_x^2\|_{H^1} + \|u_x\|_{\infty} \|\rho\|_{H^2}^2. \end{aligned}$$

Using the fact that $H^1(\mathbb{S}^1)$ is a multiplicative algebra and the following estimates

$$\|u\|_{H^2} \lesssim \|m\|_{L^2}, \quad \|u_x\|_{\infty} \lesssim \|m\|_{L^2}, \quad \|u\|_{H^3} \lesssim \|m\|_{H^1}, \quad \|\rho\|_{\infty} \lesssim \|\rho\|_{H^1},$$

we thus obtain

$$\int \rho_{xx} \rho_{txx} \lesssim (\|m\|_{L^2} + \|\rho\|_{H^1}) \left(\|m\|_{H^1}^2 + \|\rho\|_{H^2}^2 \right),$$

which achieves the proof for $k = 1$, using again Gronwall's lemma and the first *a priori* estimate obtained on $\|m\|_{L^2} + \|\rho\|_{H^1}$. Assuming now that (6.1) is true for $k \geq 1$, we will show that the same is true for $k + 1$. We have

$$\begin{aligned} \frac{d}{dt} \left(\|m\|_{H^{k+1}}^2 + \|\rho\|_{H^{k+2}}^2 \right) &= \frac{d}{dt} \left(\|m\|_{H^k}^2 + \|\rho\|_{H^{k+1}}^2 \right) \\ &\quad + 2 \int m_t^{(k+1)} m^{(k+1)} + 2 \int \rho_t^{(k+2)} \rho^{(k+2)}. \end{aligned}$$

Thus, we need to estimate

$$(6.2) \quad \alpha \int (u_x)^{(k+1)} m^{(k+1)} - a \int (u_x m)^{(k+1)} m^{(k+1)} - \int (u m_x) m^{(k+1)} \\ - \kappa \int (\rho \rho_x)^{(k+1)} m^{(k+1)} - \int (u \rho_x)^{(k+2)} \rho^{(k+2)} - (a-1) \int (u_x \rho)^{(k+2)} \rho^{(k+2)}.$$

Using the following estimate derived from the Leibnitz formula:

$$\left| \int (fg)^{(n+1)} h^{(n+1)} \right| \lesssim \|f\|_{H^n} \|g\|_{H^{n+1}} \|h\|_{H^{n+1}} \\ + \left| \int f g^{(n+1)} h^{(n+1)} \right| + \left| \int f^{(n+1)} g h^{(n+1)} \right|,$$

where $f, g, h \in C^\infty(\mathbb{S}^1)$ and $n \geq 1$, we deduce, using some integration by parts, that each term in (6.2) is bounded, up to a multiplicative constant by

$$(1 + \|m\|_{H^k} + \|\rho\|_{H^{k+1}}) \left(\|m\|_{H^{k+1}}^2 + \|\rho\|_{H^{k+2}}^2 \right),$$

which achieves the proof. \square

In [19] the following distance on $\mathcal{D}^q(\mathbb{S}^1)$ was introduced,

$$d_q(\varphi_1, \varphi_2) := d_{C^0}(\varphi_1, \varphi_2) + \|\varphi_{1x} - \varphi_{2x}\|_{H^{q-1}} + \|1/\varphi_{1x} - 1/\varphi_{2x}\|_\infty,$$

which makes $(\mathcal{D}^q(\mathbb{S}^1), d_q)$ a complete metric space. Following [19, Section 5], we can check that the spray F_q defined in Theorem 5.2 is bounded on bounded sets of TH^qG (since the proof is similar, we redirect to this reference for the details). As a consequence of the *a priori* estimates obtained in Corollary 6.3 and Lemma 6.4, we obtain therefore the following theorem.

Theorem 6.5. *Let $(u_0, \rho_0) \in C^\infty(\mathbb{S}^1) \times C^\infty(\mathbb{S}^1)$ and*

$$(u, \rho) \in C^\infty(J, C^\infty(\mathbb{S}^1) \oplus C^\infty(\mathbb{S}^1)),$$

be the solution of (4.2) with $(u(0), \rho(0)) = (u_0, \rho_0)$. Suppose that $\|u_x(t)\|_\infty$ is bounded on any bounded subinterval of J . Then, the solution $(u(t), \rho(t))$ is defined for $t \in \mathbb{R}$.

Remark 6. Note that theorem 6.5 is still true for a solution

$$(u, \rho) \in C^0(J_q, H^q(\mathbb{S}^1) \oplus H^{q-1}(\mathbb{S}^1)) \cap C^1(J_q, H^{q-1}(\mathbb{S}^1) \oplus H^{q-2}(\mathbb{S}^1)),$$

for q big enough but the proof requires the use of a Friedrich's mollifier and Kato-Ponce estimate (see [19, Theorem 5.1]), to establish the *a priori* estimate given in lemma 6.4.

Remark 7. It is well-known that the Camassa-Holm equation (CH) has solutions which blow-up in finite time, cf. [9]. One way to exhibit such solutions uses the fact that a solution which is spatially odd initially will remain spatially odd at each time. This is a consequence of the following symmetry of the equation

$$u(x) \mapsto -u(-x).$$

It could be interesting to note that equation (4.2) is invariant under the transformation

$$u(x) \mapsto -u(-x), \quad \rho(x) \mapsto -\rho(-x), \quad \alpha \mapsto -\alpha,$$

but unfortunately not by the transformation

$$u(x) \mapsto -u(-x), \quad \rho(x) \mapsto -\rho(-x),$$

unless $\alpha = 0$. In particular, there is no reason, contrary to (CH), that a solution (u, ρ) of (4.2), which is spatially odd for $t = 0$ will remain odd at time t . There are classes other than odd initial data which lead to a finite time blow-up for the Camassa-Holm equation. The specification of compatible initial conditions for system (1.1) with $\alpha \neq 0$ remains open.

ACKNOWLEDGEMENTS

D. Henry, B. Kolev and T. Lyons were supported by the Irish Research Council–Campus France PHC “Ulysses” programme. All the authors would like to thank Rossen Ivanov for his stimulating discussions, and Cathy and Michel for their kind hospitality at *Les Grandes Molières* during the preparation of this work. The authors would like to thank the anonymous referee for helpful suggestions and comments.

REFERENCES

- [1] V. I. Arnold. Sur la géométrie différentielle des groupes de Lie de dimension infinie et ses applications à l’hydrodynamique des fluides parfaits. *Ann. Inst. Fourier (Grenoble)*, 16(fasc. 1):319–361, 1966.
- [2] J. C. Burns. Long waves on running water. *Proc. Cambridge Phil. Soc.*, 49:695–706, 1953.
- [3] R. Camassa, D. Holm, and J. Hyman. A new integrable shallow water equation. *Adv. Appl. Mech.*, 31:1–33, 1994.
- [4] R. Camassa and D. D. Holm. An integrable shallow water equation with peaked solitons. *Phys. Rev. Lett.*, 71(11):1661–1664, 1993.
- [5] M. Chen, S.-Q. Liu, and Y. Zhang. A two-component generalization of the Camassa-Holm equation and its solutions. *Lett. Math. Phys.*, 75(1):1–15, 2006.
- [6] A. Constantin. Existence of permanent and breaking waves for a shallow water equation: a geometric approach. *Ann. Inst. Fourier (Grenoble)*, 50(2):321–362, 2000.
- [7] A. Constantin. *Nonlinear water waves with applications to wave-current interactions and tsunamis*, volume 81 of *CBMS-NSF Regional Conference Series in Applied Mathematics*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2011.
- [8] A. Constantin and J. Escher. Global weak solutions for a shallow water equation. *Ind. Univ. Math. J.*, 47:1527–1545, 1998.
- [9] A. Constantin and J. Escher. Wave breaking for nonlinear nonlocal shallow water equations. *Acta Math.*, 181(2):229–243, 1998.
- [10] A. Constantin and R. I. Ivanov. On an integrable two-component Camassa-Holm shallow water system. *Phys. Lett. A*, 372(48):7129–7132, 2008.
- [11] A. Constantin and B. Kolev. Geodesic flow on the diffeomorphism group of the circle. *Comment. Math. Helv.*, 78(4):787–804, 2003.
- [12] A. Constantin and D. Lannes. The hydrodynamical relevance of the Camassa-Holm and Degasperis-Procesi equations. *Arch. Ration. Mech. Anal.*, 192(1):165–186, 2009.
- [13] A. Degasperis and M. Procesi. Asymptotic integrability. In *Symmetry and perturbation theory (Rome, 1998)*, pages 23–37. World Sci. Publ., River Edge, NJ, 1999.
- [14] D. G. Ebin and J. E. Marsden. Groups of diffeomorphisms and the motion of an incompressible fluid. *Ann. of Math. (2)*, 92:102–163, 1970.
- [15] J. Escher. Non-metric two-component Euler equations on the circle. *Monatsh. Math.*, 167(3-4):449–459, 2012.
- [16] J. Escher, M. Kohlmann, and J. Lenells. The geometry of the two-component Camassa-Holm and Degasperis-Procesi equations. *J. Geom. Phys.*, 61(2):436–452, 2011.

- [17] J. Escher and B. Kolev. The Degasperis-Procesi equation as a non-metric Euler equation. *Math. Z.*, 269(3-4):1137–1153, 2011.
- [18] J. Escher and B. Kolev. Right-invariant Sobolev metrics of fractional order on the diffeomorphism group of the circle. *Journal of Geometric Mechanics*, 6(3):335–372, 2014.
- [19] J. Escher and B. Kolev. Geodesic Completeness for Sobolev H^s -metrics on the Diffeomorphism Group of the Circle. *Journal of Evolution Equations*, 2014. To appear.
- [20] J. Escher, O. Lechtenfeld, and Z. Yin. Well-posedness and blow-up phenomena for the 2-component Camassa-Holm equation. *Discrete Contin. Dyn. Syst.*, 19(3):493–513, 2007.
- [21] J. Escher, Y. Liu, and Z. Yin. Global weak solutions and blow-up structure for the Degasperis-Procesi equation. *J. Funct. Anal.*, 241(2):457–485, 2006.
- [22] J. Escher, Y. Liu, and Z. Yin. Shock waves and blow-up phenomena for the periodic Degasperis-Procesi equation. *Indiana Univ. Math. J.*, 56(1):87–117, 2007.
- [23] J. Escher and Z. Yin. Well-posedness, blow-up phenomena, and global solutions for the b -equation. *J. Reine Angew. Math.*, 624:51–80, 2008.
- [24] L. P. Euler. Du mouvement de rotation des corps solides autour d'un axe variable. *Mémoires de l'académie des sciences de Berlin*, 14:154–193, 1765.
- [25] N. C. Freeman and R. S. Johnson. Shallow water waves on shear flows. *J. Fluid Mech.*, 42:401–409, 1970.
- [26] D. Henry. Compactly supported solutions of a family of nonlinear partial differential equations. *Dyn. Contin. Discrete Impuls. Syst. Ser. A Math. Anal.*, 15(1):145–150, 2008.
- [27] D. Henry. Infinite propagation speed for a two component Camassa-Holm equation. *Discrete Contin. Dyn. Syst. Ser. B*, 12(3):597–606, 2009.
- [28] D. Henry. Persistence properties for a family of nonlinear partial differential equations. *Nonlinear Anal.*, 70(4):1565–1573, 2009.
- [29] H. Inci, T. Kappeler, and P. Topalov. *On the Regularity of the Composition of Diffeomorphisms*, volume 226 of *Memoirs of the American Mathematical Society*. American Mathematical Society, first edition, mar 2013.
- [30] R. Ivanov. Two-component integrable systems modelling shallow water waves: the constant vorticity case. *Wave Motion*, 46(6):389–396, 2009.
- [31] R. I. Ivanov. Extended Camassa-Holm hierarchy and conserved quantities. *Z. Naturforsch.*, 61(4):133–138, 2006.
- [32] R. I. Ivanov. Water waves and integrability. *Philos. Trans. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci.*, 365(1858):2267–2280, 2007.
- [33] R. S. Johnson. Camassa-Holm, Korteweg-de Vries and related models for water waves. *J. Fluid Mech.*, 455:63–82, 2002.
- [34] R. S. Johnson. The Camassa-Holm equation for water waves moving over a shear flow. *Fluid Dynam. Res.*, 33(1-2):97–111, 2003. In memoriam Prof. Philip Gerald Drazin 1934–2002.
- [35] R. S. Johnson. *Singular perturbation theory*. Mathematical and Analytical Techniques with Applications to Engineering. Springer, New York, 2005. Mathematical and analytical techniques with applications to engineering, With a foreword by Alan Jeffrey.
- [36] S. Lang. *Fundamentals of Differential Geometry*, volume 191 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1999.
- [37] G. Misiołek. Classical solutions of the periodic Camassa-Holm equation. *Geom. Funct. Anal.*, 12(5):1080–1104, 2002.
- [38] P. J. Olver and P. Rosenau. Tri-Hamiltonian duality between solitons and solitary-wave solutions having compact support. *Phys. Rev. E (3)*, 53(2):1900–1906, 1996.
- [39] G. Thomas and G. Klopman. Wave-current interactions in the nearshore region. In *Gravity Waves in Water of Finite Depth.*, volume 40 of *Advances in Fluid Mechanics.*, pages 215–319. WIT, Southampton, England, 1997.
- [40] H. Triebel. *Theory of Function Spaces*. Birkhäuser Boston, 1983.

INSTITUTE FOR APPLIED MATHEMATICS, UNIVERSITY OF HANOVER, D-30167 HANOVER,
GERMANY

E-mail address: `escher@ifam.uni-hannover.de`

DEPARTMENT OF APPLIED MATHEMATICS, UNIVERSITY COLLEGE CORK, WESTERN
ROAD, CORK, IRELAND

E-mail address: `d.henry@ucc.ie`

AIX MARSEILLE UNIVERSITÉ, CNRS, CENTRALE MARSEILLE, I2M, UMR 7373, 13453
MARSEILLE, FRANCE

E-mail address: `boris.kolev@math.cnrs.fr`

SCHOOL OF MATHEMATICAL SCIENCES, DUBLIN INSTITUTE OF TECHNOLOGY, KEVIN
STREET, DUBLIN 8, IRELAND

E-mail address: `tony.lyons@mydit.ie`