Post-Industrial Revolution changes in large-scale atmospheric pollution of the northern hemisphere by heavy metals as documented in central Greenland snow and ice

Jean-Pierre Candelone and Sungmin Hong

Laboratoire de Glaciologie et Géophysique de l'Environnement du Centre National de la Recherche Scientifique
Saint Martin d'Hères, France

Christian Pellone

Centre de Recherche et d'Essais des Machines Hydrauliques de Grenoble, Saint Martin d'Hères, France

Claude F. Boutron¹

Laboratoire de Glaciologie et Géophysique de l'Environnement du Centre National de la Recherche Scientifique
Saint Martin d'Hères, France

Abstract. Pb, Zn, Cd and Cu have been measured using ultraclean procedures in various sections of a 70.3-m snow/ice core covering the past 220 years (including the Industrial Revolution) drilled at Summit, central Greenland. These time series are the first reliable ones ever published for Zn, Cd, and Cu; for Pb they are the first verification of the pioneering data published more than two decades ago by C. Patterson and his coworkers [Murozumi *et al.*, 1969]. For all four heavy metals, concentrations are found to have markedly increased up until the 1960s and 1970s before decreasing significantly during the following few decades. The timing and the amplitude of the observed changes differ significantly however from one metal to another. Comparison with concentration values obtained by analyzing ancient Holocene ice dated 7760 years B.P., that is, before humans started to impact on the atmosphere, show that no detectable increase occurred for Zn, Cd, and Cu before the Industrial Revolution. On the other hand, Pb concentrations were already one order of magnitude above natural values in late 18th century ice. Cumulative deposition of heavy metals to the whole Greenland ice cap since the Industrial Revolution ranges from 3200 t for Pb to 60 t for Cd.

1. Introduction

Investigating the occurrence of heavy metals in the well-preserved dated snow and ice layers deposited in the large Greenland ice sheet during the last few centuries offers a unique way to reconstruct the history of the large-scale pollution of the atmosphere of the northern hemisphere for these toxic metals especially since the Industrial Revolution [Wolff and Peel, 1985; Peel, 1989; Wolff, 1990; Boutron *et al.*, 1994, 1995].

This was first illustrated by C. Patterson and his coworkers in a pioneering paper published twenty five years ago [Murozumi *et al.*, 1969]. Using revolutionary ultraclean sampling and laboratory techniques [Patterson and Settle, 1976], they obtained the first reliable time series for Pb concentration in Greenland snow and ice for the 1753–1965 time period from the analysis of large-size snow blocks collected from the walls of shafts or trenches at two locations in north west Greenland: Camp Century (77°10'N, 61°08'W, elevation 1866 m above sea level (a.s.l.)) and the "virgin trench site" (76°46'N, 58°52'W, elevation 1800 m (a.s.l.)) ~ 80 km

from Camp Century. In addition, they also obtained the first estimate of natural Pb concentrations in Greenland Holocene ice by analyzing a single block of blue ice dated ~ 2800 years B.P. collected at Camp Tuto (76°25'N, 68°20'W). They found that the Pb concentration was exceedingly low, ~ 0.8 pg/g, 2800 years B.P., before man started to impact significantly on the atmosphere, and that this natural Pb could be explained simply by a soil and rock dust contribution. In the 1750s, Pb concentration was found to be ~ 10 pg/g, that is one order of magnitude higher than the value for 2800 years B.P.: it clearly evidenced for the first time that the Arctic atmosphere was already strongly contaminated in Pb before the onset of the Industrial Revolution. From the 1750s to 1965, a further continuous ~ 20 fold increase was observed, the increase being especially rapid after the 1930s as a response to the introduction and subsequent massive use of Pb alkyl additives [Nriagu, 1990] in automobile gasoline. On the whole, Pb concentrations appeared to have massively increased by about 200 fold from several thousand years ago to the 1960s in Greenland snow and ice, providing the first clear evidence of massive large-scale pollution of the atmosphere of the northern hemisphere for this toxic metal. And it is at least partially because of this landmark work that the United States and various other countries severely limited their use of Pb alkyl additives in gasoline from the early 1970s.

During the following two and half decades, various other investigators tried to emulate this pioneering work, not only for Pb but also for several other heavy metals such as Cd, Hg, Cu,

¹Also at UFR de Mécanique, Université Joseph Fourier de Grenoble, Domaine Universitaire, Grenoble, France.

and Zn [e.g. Weiss *et al.*, 1971 a,b, 1975; Carr and Wilkiss, 1973; Cragin *et al.*, 1975; Herron *et al.*, 1977; Appelquist *et al.*, 1978]. None of them succeeded in producing reliable time series for the past few centuries, mainly because they were unable to fully control contamination problems during field sampling, laboratory analysis or both.

The only reliable data which became available during these 25 years were either for ancient ice or for recent (post-1965) snow. C. Patterson and his coworkers were able to successfully meet the very difficult requirements of decontaminating deep Greenland ice cores for heavy metals analysis [Ng and Patterson, 1981]. It allowed them to analyze, for Pb, a few sections of the Camp Century 1387-m ice core whose age ranged from 2700 to 5500 years B.P., and to confirm that the natural Pb concentration in Greenland Holocene ice was really at the picogram-per-gram level as previously claimed by Murozumi *et al.* [1969] from the analysis of their single block of blue ice. For recent snow, various reliable data were published from the analysis of shallow samples collected in southern or central Greenland either from the walls of hand dug pits [Wolff and Peel, 1988; Boyle *et al.*, 1994; Savarino *et al.*, 1994; Candelone *et al.*, 1995] or using specially designed acid cleaned all-plastic augers [Boutron *et al.*, 1991]. These last data show that Pb concentrations have decreased by a factor of 7.5 since the mid-1960s in central Greenland as a consequence of the reduction in the use of Pb alkyl additives especially in the United States. During the same period, Cd and Zn concentrations were found to have decreased by a factor of 2.5 in response to the increasing efforts made in various countries of the northern hemisphere to reduce the emissions of these metals to the atmosphere from the combustion of fossil fuels, metal processing activities, and refuse incineration. They also revealed that there are pronounced intra annual (seasonal) variations in heavy metals concentrations in recent Greenland snow.

Confronted with the lack of confirmation of the Pb time series of Murozumi *et al.* [1969] and with the lack of time series for other metals for the past few centuries, we decided to collect and analyze for Pb, Zn, Cd, and Cu new samples covering the post-Industrial Revolution period, that is the past ~ 220 years, using improved ultraclean procedures. The results are interpreted and discussed in the light of available data on anthropogenic emissions of these metals to the atmosphere.

2. Experiment

2.1. Field Sampling

The samples were collected in July 1989 at Summit (72°34'N, 37°37'N, elevation 3238 m (a.s.l.), mean annual temperature - 32°C, mean annual snow accumulation rate 23 g cm⁻² yr⁻¹) as part of the European "Eurocore" program. They consisted of a snow/ice core 9.8 cm in diameter and 70.3 m deep, which was electromechanically drilled only for our heavy metals measurements. To minimize local contamination problems, the exact sampling site was selected about 500 m from the Eurocore camp. In order to minimize contamination brought to the outside of the core during drilling operations, we used a stainless steel drill [Gillet *et al.*, 1984] which was coated with polytetrafluoroethylene (PTFE).

Core handling was kept to a strict minimum. Each core section was handled by operators wearing full clean room

clothing and shoulder length polyethylene gloves, using specially designed acid cleaned low density polyethylene (LDPE) pistons (to push the core section out of the drill if necessary), brushes (to remove the cutting chips from the outside of the core), gutters (to keep the core sections before packing), and saws.

Each core section (about 1 m long) was packed inside double-sealed LDPE bags and transported frozen to Grenoble, France.

2.2. Decontamination of the Core Sections

Despite the great precautions which were taken in the field to minimize contamination brought to the outside of the core, significant contamination was found to be present on the outside of the core sections. This contamination was especially pronounced for Zn: for this metal, measured concentrations in the first centimeter or so from the outside ranged indeed from about 1000 to 5000 pg/g, that is, concentrations higher by up to 2 orders of magnitude than the original Zn concentration in the snow. This Zn contamination was thought to originate at least partially from the galvanized steel cable of the electromechanical drill.

Each of the analyzed core sections was therefore mechanically decontaminated before analysis, using ultraclean procedures, in order to remove the contaminated outside veneer layers and get the uncontaminated innermost part of the core. The decontamination took place inside a laminar flow clean bench located in a - 15°C cold room.

For low and medium-density snow sections (depth down to ~20 m) the decontamination procedure was similar to that described by Boutron *et al.* [1991] and Görlach and Boutron [1992]. It involved breaking the core sections with a sharp-edged LDPE splitting wedge in order to obtain 20 or 15 cm long sections with fresh untouched surfaces at both extremities, then extracting the central part (diameter: 4 cm) of these 20 or 15 cm sections by hammering a home made cylindrical LDPE beaker into the center of one of the untouched surfaces parallel to the core axis. Additional subsamples were also obtained at increasing distances from the axis using smaller diameter LDPE beakers.

Below ~ 20 m, with the higher density of the snow, it was no longer possible to use this procedure, and a new improved procedure, described in detail by Candelone *et al.* [1994], was employed. It involved supporting the core section horizontally in a LDPE home made lathe [Candelone *et al.*, 1994] then chiselling successive veneers of snow (or ice for the deepest sections) in progression from the outside towards the center, using the same stainless steel chisels used by C. Patterson and his coworkers in their Antarctic ice cores work [Boutron and Patterson, 1986]. The two extremities of the inner core (diameter: 4 cm), obtained when the procedure was completed, were then broken using chisels while firmly held with all LDPE homemade tongs. A distinct advantage of this improved procedure is that it is possible to handle sections of variable length and obtain inner cores which integrate a given time period, for instance an exact full year.

The procedural blank for the chiselling procedure was accurately determined by processing an artificial ice core [Candelone *et al.*, 1994] made by freezing ultrapure water [Boutron, 1990] whose heavy metals content was known beforehand. It ranged from 0.015 pg/g for Cd up to 0.25 pg/g

for Cu [Candelone *et al.*, 1994], that is contributions which are extremely small compared to heavy metals concentrations in post-Industrial Revolution Greenland snow and ice.

2.3. Checking the Efficiency of the Decontamination Procedure

As previously emphasised by various authors [Ng and Patterson, 1981; Boutron and Patterson, 1986; Peel, 1989; Wolff, 1990; Boutron *et al.*, 1994, 1995], it is essential to determine changes in the concentrations of each heavy metal from the outside to the inside of each analyzed core section to make certain that concentrations measured in the central part, obtained after the decontamination is completed, do represent the original concentration in the snow or ice. If a clear plateau of concentration is obtained in the central part, it indicates that external contamination has not penetrated to this central part: the plateau concentration will then represent the original concentration in the snow or ice, providing an appropriate blank correction is made for the whole analytical procedure. If, on the other hand, a continuous decrease of concentration is observed to the most central parts, this indicates that external contamination has probably reached the center of the core: the corresponding concentration value then gives only an upper limit of the original concentration in the snow or ice.

Outside-inside concentration profiles have been obtained for all four heavy metals for each of the investigated core sections. In most cases, good plateaus of concentrations were observed, as illustrated in Figure 1 for Cu and Zn in the 61.56-61.85-m section. A continuous decrease of concentrations from the outside to the center was observed in a few cases only, as shown in Figure 2 for Cu in the 69.93-70.17-m section and Zn in the 51.76-52.23-m section.

2.4. Analytical Procedures

Each veneer layer and inner core was analyzed separately using ultraclean procedures [Boutron, 1990]. Pb, Cd, and Cu

Figure 1. Measured Zn and Cu concentrations as a function of radius in the 61.56-61.85 section of the 70.3m snow/ice core electromechanically drilled at Summit, central Greenland. Good plateaus of concentrations are obtained in the central part of the core section.

Figure 2. Measured heavy metals concentrations as a function of radius in two sections of the 70.3-m snow/ice core electromechanically drilled at Summit, central Greenland. (A) Cu in the 69.93-70.17-m section (year 1773); (B) Zn in the 51.76-52.23-m section (years 1841-1842). No satisfying plateau of concentrations is obtained in the central part of the core section for these two cases.

were determined by graphite furnace atomic absorption spectrometry (GFAAS) after preconcentration ($\sim \times 50$) by non boiling evaporation [Görlach and Boutron, 1990]. Zn, Al, and Na were measured by GFAAS without preconcentration. The GFAAS measurements were performed with a Perkin Elmer 2380 spectrometer and HGA 500 graphite furnace, using electrodeless discharge lamps for Pb, Cd, and Zn. Pyrolytically coated graphite tubes from Applied Optics were used. The precision of the data is estimated to be $\pm 10\%$.

Various 0.1% HNO_3 (ultrapure twice-distilled HNO_3 from US National Institute of Standards and Technology [Paulsen *et al.*, 1988]) aliquots were also taken for subsequent determination of Pb by isotope dilution mass spectrometry (IDMS) [Rosman *et al.*, 1993, 1994], Pb isotopes by thermal ionization mass spectrometry (TIMS) [Rosman *et al.*, 1993, 1994], ethyl and methyl Pb compounds by capillary gas chromatography with microwave-induced plasma atomic emission spectrometry (CGC-MIP-AES) [Lobinski *et al.*, 1993, 1994], and Hg by cold vapor atomic fluorescence spectrometry (CVAFS) [Vandal *et al.*, 1993].

Comparative determination of Pb by GFAAS after preconcentration and IDMS indicated that the IDMS values were systematically slightly lower, by $\sim 20\%$, than the GFAAS values when these last values were corrected using the slope (0.82) of the calibration line of the preconcentration procedure given by Görlach and Boutron [1990]. They were, on the other hand, in excellent agreement if this correction was not made. Taking into account the fact that IDMS is essentially an absolute technique, we have therefore decided to no longer apply a correction for the preconcentration procedure not only for Pb, but also for Cd, Cu, and Zn (for these last metals, this results in decreasing the values by 24%, 11%, and 15%, respectively). It must however be emphasised that these relatively small differences do not significantly influence the geochemical interpretation of the snow and ice data.

2.5. Dating of the Samples

The dating of the core was based on detailed continuous oxygen isotopes data obtained in a nearby parallel core [H. Clausen, personal communication, 1993]. It indicates that the bottom of the core (70.3 m) corresponds to the year 1772, that is, slightly before the beginning of the first Industrial Revolution. The whole core then covers continuously a 217-year time period. The precision of the dating is estimated to be about ± 2 years at the bottom of the core.

3. Results and Discussion

3.1. Presentation of the Data

Since comprehensive heavy metals data had already been obtained for the 1967-1989 period from the analysis of a 10.7-m snow core drilled at a nearby location [Boutron *et al.*, 1991], we mainly analyzed the part of the 70.3-m core corresponding to the pre-1966 period. Special attention was moreover given to the 1945-1966 period (depth interval 20-11 m), which was

expected to correspond to a rapid increase of Pb concentrations in response to the massive use of Pb alkyl additives in gasoline after World War II.

Table 1 displays Pb, Cd, Zn, and Cu concentrations measured in the central part of each of the 26 investigated core sections. For the core sections which were decontaminated using the stainless steel chisels, (see section 2.2), that is, the sections whose depth was greater than 20 m, the chiselling procedure was conducted so that the length of the inner core obtained after the decontamination was completed integrated exactly 1 or 2 years of snow or ice accumulation at the corresponding depth (estimated from the detailed oxygen isotopes profiles). For these sections, each data point given in Table 1 then exactly corresponds to a 1 or 2 year time interval. It is then not influenced by intra annual (seasonal) variations of the concentrations [Wolff and Peel, 1988; Boyle *et al.*, 1994; Savarino *et al.*, 1994; Candelone *et al.*, 1995].

For the core sections which were decontaminated using the earlier procedure (see section 2.2) of Boutron *et al.* [1991] and Görlach and Boutron [1992], that is, the sections whose depth was less than 20 m, the situation is more complicated. The depth interval associated with each data point often does not

Table 1. Summit, Central Greenland: Heavy Metals Concentrations Measured in the Central Part of Various Sections of a 70.3-m Snow/Ice Core

Depth Interval*, m	Age†, Calendar Year	Measured Concentration, pg/g			
		Pb	Zn	Cd	Cu
7.48-7.93	1974.5	84	60	1.7	7.6
9.79-10.11	1969.6	109	46	1.5	3.2
10.92-11.25	1965.4	99	60	1.3	8.3
11.44-11.77	1964.1	88	129	1.5	9
11.96-12.25	1963	90	75	1.3	7.9
12.54-12.86	1961.6	65	72	0.85	13
12.98-13.27	1960.4	60	49	0.78	4
13.37-13.67	1959.2	126	155	2.2	15
13.74-14.17	1958.2	59	54	0.82	3.2
14.38-14.69	1956.9	90	74	1.2	6.4
15.17-15.62	1954.7	67	75	1.2	7.5
15.80-16.17	1953.2	81	134	3.3	17
16.69-16.97	1951.2	67	67	1.4	7.3
17.30-17.58	1949.7	88	71	2.9	7.1
18.08-18.54	1948	49	79	2.1	6.8
18.66-19.11	1946.6	31	42	1.2	4.9
21.10-21.45	1940	41	75	1.1	8.6
23.64-24.08	1933	18	90	1.1	9.3
25.48-25.92	1928	45	65	2.0	6.9
29.77-30.13	1915	34	34	1.2	5.2
37.27-37.60	1891	63	30	1.1	4.9
43.90-44.45	1869-1870	18	44	0.68	2.7
51.76-52.23	1841-1842	17	22	0.15	2.4
58.20-58.50	1817	4.4	29	0.23	3.5
61.56-61.85	1805	12	20	0.21	6.1
69.93-70.17	1773	7.6	16	0.27	2.8

*Surface: July 1989.

†Estimated age for the middle of the corresponding core section.

correspond exactly to 1 year of snow accumulation, since it was determined by the cumulative length of the LDPE beakers hammered into the core. It can be seen in Table 1 that the corresponding depth interval was often ~ 30 to 35 cm, which represents slightly less than 1 year of snow accumulation at these depths (at 15 m, for instance, 1 year of snow accumulation is approximately equal to 40 cm). Because these samples do not represent exactly 1 year of accumulation, the corresponding data points are probably influenced by seasonal variations in concentrations: this will result in a larger scatter in the data points, as it will be seen in the time series discussed in sections 3.2 and 3.3

3.2. Changes in Zn, Cd, and Cu Concentrations From 1773 to 1992

Changes in the concentrations of these three metals from 1773 to 1992 are shown in Figure 3 as documented by combining the data obtained in this work from the analysis of the 70.3-m core with the data given by *Boutron et al.* [1991] for the 1967-1989 period (averaged so that each data point corresponds approximately to 1 year of snow accumulation)

Figure 3. Changes in Zn, Cd, and Cu concentrations in snow/ice deposited at Summit, central Greenland, from 1773 to 1992. Solid circles: this work (70.3-m core); solid triangles: averaged values from *Boutron et al.* [1991] (10.7-m core); open triangles: mean yearly values for 1990, 1991, and 1992 from *Candelone et al.* [1995] (1.6-m snow pit). The general time trends are shown with spline-smoothed curves.

and with the mean yearly values for the years 1990, 1991, and 1992 given by *Candelone et al.* [1995]. The data of *Boutron et al.* [1991] were obtained from the analysis of a 10.7-m snow core drilled using a specially designed all-plastic acid cleaned mechanical auger. The data of *Candelone et al.* [1995] were obtained from the analysis of a continuous sequence of 30 samples collected by pushing ultraclean wide mouth LDPE bottles into the walls of a 1.6-m deep hand-dug snow pit. This is the first reliable time series ever published for these three metals, since previously published data [*Weiss et al.*, 1975; *Herron et al.*, 1977] were plagued by major contamination problems.

A common feature of the three time series shown in Figure 3 is the existence of a clear increasing trend until the 1960s and 1970s, followed by a significant decreasing trend. But a detailed examination of the profiles reveals pronounced differences for the three metals.

First of all, the timing of the observed increase is not the same. For Zn, our data indicate that the increase started as early as the 1770s. It was, however, fairly slow until the ~ 1900s, then became more rapid during the first half of the 20th century. For Cd and Cu, on the other hand, the increase did not start until the 1850s (the maximum shown by the spline function around the 1800s for Cu is probably not significant since it is due only to a single data point). Its rate was, moreover, fairly stable during the following 1 century or so.

Another difference is with the amplitude of the observed increases. Cd shows the greatest increase (~ x 8), followed by Zn (~ x 5) and Cu (~ x 4).

Also, the time when the maximum is observed is not the same. The spline-smoothed curves shown in Figure 3 suggest that the maximum was reached earlier for Zn (around ~ 1961) than for Cd and Cu (around ~ 1972 and 1969, respectively).

Finally, the amplitude of the recent decrease differs from one metal to another. Cd is found to have decreased by a factor of ~ 2.3 during the past 2 decades down to 1992 values, which are similar to those around the 1870s. For Zn, the decrease appears to be smaller (factor of ~ 1.7), with the 1992 values similar to those in the 1920s. The decrease for Cu appears to be hardly significant (factor of ~ 1.2), with the 1992 values close to those observed in the 1930s.

These changes cannot be linked with changes in emissions from natural sources to the atmosphere, such as rock and soil dust, sea salt spray, and volcanoes [*Nriagu*, 1989]. The corresponding emissions have indeed not significantly changed since the 18th century, as evidenced, for instance, for the rock and soil dust and sea-salt spray contributions by the fact that no time trend is observed since the 1770s for Al and sea salt Na (Table 2). They are then, without doubt, the result of major changes in the large scale anthropogenic emissions of these metals to the atmosphere of the northern hemisphere.

There is a wealth of data available on the world production of Cd, Cu, and Zn during the past century or so, for instance, in extremely detailed publications of the United Kingdom government [*Imperial Mineral Resources Bureau*, 1913-1920; *Imperial Institute*, 1920-1944; *Colonial Geological Survey* 1945-1951; *National Environmental Research Council*, 1952-1970; *National Environmental Research Council*, 1970-1990] and the U.S. Bureau of Mines [1930-1990] or in specialized books devoted to given metals [e.g. *Nriagu*, 1978, 1979a, b, c]. Although the past trends in metals production have often been used as surrogates for trends in past metals emissions to the

Table 2. Measured Al and Sea-Salt Na (Na_{ss}) Concentrations in Typical Samples Whose Age Ranges from 1773 to 1992

Year	Measured Concentration, ng/g	
	Al	Na_{ss}
1773	4.3	2.4
1817	5.6	4.4
1870	7.6	4.1
1915	5.0	2.3
1940	8.6	0.7
1968	3.7	0.9
1992*	5.6	3.2

Concentrations of Na_{ss} were calculated by correcting measured Na concentrations for the crustal component evaluated from Al concentrations using the Na/Al abundance ratio in the mean crustal material [Taylor and McLennan, 1985].

* From Candelone et al. [1995].

atmosphere [e.g., Wolff and Suttie, 1994], they however do not satisfyingly represent these trends.

On the other hand, the available data on world emissions (which originate mainly from non-ferrous metals production and also from coal combustion, steel and iron manufacturing, and refuse incineration) are much sparser. Even for recent years the best available estimates [Nriagu and Pacyna, 1988; Pacyna, 1994] remain rather uncertain. For instance, the ranges for the estimates of worldwide emissions for Zn, Cd, and Cu for the year 1983, given by Nriagu and Pacyna [1988], are (in metric tons per year) 70,000–194,000, 3,100–12,000, and 19,900–50,900 respectively, that is, uncertainties by up to a factor of 4. The few data on historical changes in these emission fluxes [e.g., Nriagu 1979d] are even sparser.

Table 3 compares the mean ratios of total emissions (anthropogenic plus natural, 1983 values)/natural emissions given by Nriagu [1989] with the enhancement factors which we observe in Summit snow and ice from 1773 to 1983 (that is, not the enhancement factors discussed above, which were for the time period from 1773 to the time when the maximum is reached). It must be noted that this comparison relies upon the fact that concentrations measured in the ice in 1773 are entirely natural, which is indeed the case, as will be discussed in section 3.4. It can be seen in Table 3 that the increase factors are in remarkable agreement with the mean ratios, which shows that the historical changes in Zn, Cd, and Cu deposition in Greenland are consistent with the estimates of emissions.

Finally, the decreasing trend which we observe from the early 1960s for Zn and from the early 1970s for Cd and possibly Cu appears to be consistent with the very significant reported decrease in the emissions in the northern hemisphere as a consequence of the abatement measures which were taken in various countries of the northern hemisphere during the past decades [Pacyna, 1989, 1991; Pacyna et al., 1993; Thomas and Spiro, 1994]. For instance, Pacyna [1991] estimated that emissions of Cd and Zn from Europe, which are likely to be a

significant contribution to Cd and Zn deposited in Greenland, peaked in the 1970s, and that the 1980 emissions were about three times lower than the pre-mid-1970s ones. His estimates were obtained from a comprehensive review of changes in emission factors for these two metals in Europe from the 1950s up to 1985 for major source categories such as non ferrous metal production and combustion of fuels [Pacyna, 1991]. In addition to the impact of these abatement measures it is also likely that a significant reduction of the emissions also resulted from the very strong economic recession which has affected the states of the former Soviet Union during the past few years. These reductions are thought to have been especially important for the highly polluting non ferrous metal production activities in the Russian Arctic, for instance, in the Kola Peninsula [Jaffe et al., 1995].

3.3. Changes in Pb Concentrations From 1773 to 1992

Changes in Pb concentrations from 1773 to 1992 are shown in Figure 4. This time series is the first verification and extension of the pioneering data published 25 years ago by Murozumi et al. [1969] for the years 1753–1965, which are also shown in Figure 4.

For the time period covered by both time series, the general trends obtained from both sets of data are similar: our data confirm the very large increase in concentration from the middle of the 18th century to the 1960s seen in the earlier work. However, our data appear to be systematically lower than those of Murozumi et al. [1969]. A possible explanation for this discrepancy is that it would be linked with the different geographical characteristics of the sampling sites. The samples of Murozumi et al. [1969] were indeed collected at a much lower altitude than our samples (1800 instead of 3238 m a.s.l.): the high altitude-free troposphere of Summit is likely to be less contaminated. Moreover, the sampling sites of Murozumi et al. [1969] were located in north west Greenland, that is, an area which is more affected by the Arctic haze phenomenon [Barrie, 1986] than central Greenland. An alternative explanation could, however, be that the data of Murozumi et al. [1969], although of outstanding quality at the time when they were obtained, suffered slight but, however, significant contamination. They were indeed not meeting all

Table 3. Comparison of Zn, Cd, and Cu Levels

	Zn	Cd	Cu
Total/Natural world emissions*	3.9	6.9	2.3
Enhancement factor in Greenland snow and ice from 1773 to 1983	3.4	6.3	2.6

Comparison of the Mean Ratio of Total World Emissions (Anthropogenic Plus Natural, 1983 values) / Natural World Emissions With the Enhancement Factor in Summit, Central Greenland, Snow and Ice from 1773 to 1983.

* Nriagu and Pacyna [1988].

Figure 4. Changes in Pb concentrations in snow/ice deposited at Summit, central Greenland, from 1773 to 1992. Solid circles: this work (70.3-m core); solid triangles: averaged values from *Boutron et al.* [1991] (10.7-m core); open triangles: mean yearly values for 1990, 1991 and 1992 from *Candelone et al.* [1995] (1.6-m snow pit). The general time trend is shown with a spline-smoothed curve. Also shown, with open circles, are the original data of *Murozumi et al.* [1969] for the Camp Century area in north west Greenland (the dotted line shown on the figure is the original one given by *Murozumi et al.* [1969]).

standards of quality by which heavy metals work is now judged, especially regarding the necessity of taking outside-inside profiles across the samples. Such profiles, which were first proposed by *Ng and Patterson* [1981], were not studied for the snow blocks analyzed by *Murozumi et al.* [1969].

A more detailed examination of the data in Figure 4 indicates that there are different growth regimes between 1773 and 1992. From 1773 to the ~ 1890s, first, a rather slow increase (~ 6.3 fold in ~ 130 years) is observed. It is attributed to the progressive increase in Pb emissions to the atmosphere of the northern hemisphere, linked with the anthropogenic sources which were dominant at that time when Pb alkyl additives in gasoline were still not used. The most important of these sources were non ferrous metals production (Pb, Cu-Ni, and Zn-Cd), iron and steel manufacturing, and coal and wood combustion [*Nriagu*, 1978, 1979d; *Patterson*, 1980; *Nriagu and Pacyna*, 1988; *Wolff and Suttie*, 1994].

The spline-smoothed curve shown in Figure 4 suggests that this ~ 130 year period was followed by a few decades around the 1920s-1930s during which Pb concentrations seem to have slightly decreased. From our data points, it is difficult to determine whether this decrease is significant or not: more data points would obviously be needed, especially for the 1880s-1920s. It is however interesting to see that such a decline around the 1920s-1930s was also suggested by the data of *Murozumi et al.* [1969] (see Figure 4). If real, it could indicate that Pb emissions to the atmosphere declined during the big economic recession, as already documented for other pollutants such as SO_2 [e.g. *Möller*, 1984].

Starting from the ~ 1930s, a large and rapid increase is then observed until ~ 1970 (~ 3.4 fold in ~ 40 years). Although the contribution from other sources such as non ferrous metals production certainly remained very significant during that

period, and probably played a role in the observed increase, there is little doubt that this rapid increase was largely caused by the introduction (in 1923) and massive increase in the use of Pb alkyl additives in automobile and aviation gasoline [*Nriagu*, 1990]. The consumption of additives massively increased from 1923 to 1970, reaching a peak value of ~ 400,000 t/yr in the early 1970s. It is estimated that ~ 70% of the lead content in the additives was emitted to the atmosphere via the exhaust pipes of the engines. The largest part of these emissions occurred in the northern hemisphere, where there are the most cars (95% in 1948; 93% now [*Wolff and Suttie* 1994]).

Further unambiguous evidence of the key role played by Pb alkyl additives in the rapid increase of concentrations from the 1930s to ~ 1970 was recently obtained thanks to the direct measurement of organo-Pb compounds (Et_3Pb^+ and $\text{Et}_2\text{Pb}^{2+}$) in aliquots of part of the post-1923 samples analyzed in this work [*Lobinski et al.*, 1994]. These compounds, which have no natural origin and are unique tracers of gasoline-related pollution [*Granjean*, 1983; *Van Cleuvenbergen and Adams*, 1990] were indeed found to be present above detection limit (10 fg/g *Lobinski et al.*, 1993]) in our samples from the 1930s, with a marked increasing trend to the 1970s. Additional evidence was also provided by the determination of the isotopic pattern of Pb (Pb^{204} , Pb^{206} , Pb^{207} , and Pb^{208}) in other aliquots [*Rosman et al.*, 1994, 1995].

After this final increase a clear maximum of Pb concentrations is observed in ~ 1970, Figure 4, followed by a pronounced decrease from ~ 1970 to 1992 (~ 6.5 fold in 2 decades) which brings the 1992 values back down to the concentrations observed in the mid-19th century (Figure 4). Part of this decrease is probably due to the reduction in emissions from non ferrous metal production and steel and iron manufacturing, resulting from abatement measures

Figure 5. (a) Changes in Pb concentration in snow/ice at Summit, central Greenland, from 1955 to 1986 (spline-smoothed curve). (b) Changes in Pb (in metric tons) used to manufacture Pb alkyl antiknock additives for gasoline during the same time period [Larbey and Bain, 1988]. It is estimated that ~ 70% of the amount of the lead content in the additive is emitted to the atmosphere through the exhaust pipe.

[Pacyna, 1989, 1991; Pacyna *et al.*, 1993], although the largest part undoubtedly originates from the massive fall in the use of Pb alkyl additives in gasoline from the early 1970s [Nriagu, 1990]. This is illustrated in Figure 5, which shows the striking parallelism between the observed changes in Pb concentration in Summit snow from 1955 to 1986 and the reported world (in fact mostly northern hemisphere) consumption of additives

during the same period [Larbey and Bain, 1988]. Both curves do peak nearly at the same time (~ 1970). Moreover, the relative changes are very similar (decrease by a factor of ~ 4 from 1970 to 1986).

This fall was due both to the pioneering work of Murozumi *et al.* [1969] and to the generalization of catalytic exhaust pipes (these pipes cannot work when using leaded gasoline). It included two aspects. First, the increasing use of unleaded gasoline, which now represents virtually 100% of the gasoline sold in the United States and in Canada [Loranger and Zayed, 1994] and an ever increasing percentage of gasoline sold in European countries (48% in France in 1994). Secondly, the very pronounced decline in the Pb content of leaded gasoline: in France, for instance, it is now almost 4 fold lower (0.15 g/L since June 1991) than it was 15 years ago (0.55 g/L in April 1976), with intermediate steps in January 1979 (0.50 g/L), January 1981 (0.40 g/L), and March 1989 (0.25 g/L)).

3.4. Comparison With Concentrations in Prehuman-Activities Ancient Holocene Ice

Table 4 compares Pb, Zn, Cd, and Cu concentrations observed in our oldest sample, dated on 1773, with those measured in Greenland ice 7760 years old, that is, Holocene ice deposited before man started to impact significantly on the atmosphere. These last data were obtained from the analysis of a 1286.5-m deep section of the 3028.8-m GRIP deep ice core drilled in 1990-1992 at Summit [Hong *et al.*, 1994; S. Hong, personal communication, 1994].

For Zn, Cd, and Cu, concentrations in 1773 appear to be very similar to those 7760 years ago, which shows that in the second half of the 18th century the troposphere of the remote Arctic areas of the northern hemisphere was still uncontaminated for these three metals. At that time, Zn, Cd, and Cu were still entirely derived from natural sources, mainly rock and soil dust [Hong *et al.*, 1994; S. Hong, personal communication, 1994]. When combined with the time series shown in Figure 3, the overall picture is, then, that the large-scale pollution of the troposphere of the northern hemisphere started from ~ the 1770s for Zn and from ~ the 1850s for Cd and Cu.

For Pb, on the other hand, the situation is very different: concentrations measured in 1773 are found to be more than 1 order of magnitude higher than in ancient Holocene ice (Table 4). This clearly confirms for the first time the claim made long ago by C. Patterson and his coworkers [Murozumi

Table 4. Summit, Central Greenland: Comparison of Heavy Metals Concentrations Measured in Ice Dated for 1773 (Second Half of the 18th Century, This Work) and in Prehuman-Activities Holocene Ice Dated for 7760 Years B.P.

Age of Ice, Years		Measured Concentrations, pg/g			
Calendar	B.P.	Pb	Zn	Cd	Cu
1773 A.D.	222	7.6	16	0.27	≤2.8
5766 B.C.	7760	0.55	23	0.25	1.7

(from Hong *et al.* [1994] and S. Hong, personal communication, 1994).

Figure 6. Changes in Pb concentrations in ice/snow at Summit, central Greenland, from 3000 years B.P. to 1992. Years 3000–470 B.P.: data from *Hong et al.* [1994]; years 1773–1992: this work. Note the logarithmic scale on the vertical axis.

et al., 1969; *Patterson*, 1980] that the troposphere of the remote areas of the northern hemisphere was already heavily contaminated with this toxic metal long before the onset of the Industrial Revolution. Actually, this large-scale pollution started as early as ~ 3000 years ago. This was recently evidenced by *Hong et al.* [1994] from the analysis of various sections of the 3028.8-m GRIP deep ice core whose age ranged from 3000 to 470 years B.P. (depths: 619.3 to 129.3 m). It was evident from ~ 2500 to 1700 years B.P., that is, during the flourishing of the Greco-Roman civilization, as a consequence of anthropogenic Pb emissions to the atmosphere linked with the extensive Greek

and Roman Pb/Ag mining and smelting activities. During that period, Pb concentrations in Greenland ice reached values of ~ 3 pg/g [*Hong et al.*, 1994], which is ~ 6 fold above natural values and only 2.5 fold lower than the 1773 concentration value. After the fall of the Roman Empire, it declined down to values close to the natural ones. Finally, it started to increase again progressively from 1500 to 470 years ago, that is, during the Medieval and Renaissance times [*Hong et al.*, 1994]. Pb concentration 470 years ago (i.e., in 1524) was close to 4 pg/g, that is ~ 2 times lower than in 1773.

The overall picture is, then, that the history of the large-scale pollution of the troposphere of the northern hemisphere during the past 3 millennia has been rather lively, with a succession of increases and decreases (Figure 6). Two maxima are observed in Figure 6: the first one around 2000 years ago, at the peak of Roman power and influence; the second one around 1970, at the peak of the use of Pb alkyl additives in gasoline. The amplitude of these changes is much larger than that for any other pollutant species: the Pb concentration measured in Greenland snow in 1970 (~ 110 pg/g, Figure 4) is about 200 times higher than in pre human activities Holocene ice (0.55 pg/g, Table 4).

3.5. Past and Present-Day Atmospheric Deposition Fluxes to the Greenland Ice Sheet

The past and present day atmospheric deposition fluxes of Pb, Zn, Cd, and Cu at Summit can be estimated by combining the heavy metals concentrations measured in the snow or ice with the snow/ice accumulation rate of $23 \text{ g H}_2\text{O cm}^{-2} \text{ yr}^{-1}$ at Summit (a value which has remained essentially unchanged during the past few thousand years). We have made the calculations for different times (Table 5): 7760 years B.P., which gives the pre human activity Holocene fluxes; 1773, which corresponds to the onset of the first Industrial Revolution; 1850, which is the time when Cd and Cu concentrations started to increase; and 1992. In addition, Table 5 also presents the deposition fluxes for the 1960s and 1970s

Table 5. Heavy Metals Deposition Fluxes at Summit, Central Greenland

Year	Flux, $10^{-10} \text{ g cm}^{-2} \text{ yr}^{-1}$			
	Pb	Zn	Cd	Cu
7760 B.P.*	0.13	5.3	0.06	0.39
1773	1.8	3.7	0.06	$\leq 0.64^\dagger$
1850‡	3.5	7.0	0.06	0.53
1992‡	3.9	12	0.18	1.7
1960s and 1970s (maximum)	25	20	0.41	2.2

*Concentration values given by *Hong et al.* [1994] and S. Hong, personal communication, 1994.

[†]No satisfying plateau of concentration was obtained for Cu in the 1773 sample.

[‡]For 1850 and 1992 the concentration values used for the calculations are those given by the spline-smoothed curves shown in Figures 3 and 4.

Table 6. Past and Present-Day Deposition Fluxes of Heavy Metals to the Greenland Ice Cap

Year	Flux, t yr ⁻¹			
	Pb	Zn	Cd	Cu
7760 B.P.	0.24	9.8	0.11	0.72
1773	3.3	6.8	0.11	≤1.2
1850	6.5	13	0.11	1.0
1992	7.2	22	0.33	3.1
1960s and 1970s (maximum)	46	37	0.75	4.1

when the concentrations reached their maximum. The fluxes so calculated range from $0.06 \times 10^{-10} \text{ g cm}^{-2} \text{ yr}^{-1}$ (Cd until 1850) to $25 \times 10^{-10} \text{ g cm}^{-2} \text{ yr}^{-1}$ (Pb in the early 1970s).

Extrapolation to the whole Greenland ice cap (surface of $1.4 \times 10^6 \text{ km}^2$; mean accumulation rate of $34.5 \text{ g H}_2\text{O cm}^{-2} \text{ yr}^{-1}$) would require that the deposition patterns are known for the different areas of the ice cap, which is not the case. Besides the Summit data, there are indeed only very few reliable heavy metals data for other locations in Greenland, which, moreover, are not for the same time periods [Murozumi *et al.*, 1969; Wolff and Peel, 1988]. We have then assumed that the deposition patterns for heavy metals to the Greenland ice cap are similar to those for non sea-salt sulfate (nss sulfate) [Legrand, 1995]. For this compound, extensive data are now available for various areas of Greenland [Legrand, 1995]. They show that the nss sulfate deposition fluxes are correlated with the snow/ice accumulation rates. The mean annual deposition flux of nss sulfate to the Greenland ice cap was found to be $\sim 1.04 \times 10^{-6} \text{ g cm}^{-2} \text{ yr}^{-1}$, that is, ~ 1.3 times the value for Summit ($0.8 \times 10^{-6} \text{ g cm}^{-2} \text{ yr}^{-1}$). The heavy metals deposition fluxes so obtained are given in Table 6. They range from 0.11 t yr^{-1} (Cd before 1850) up to 46 t yr^{-1} (Pb in the early 1970s).

Finally, it is interesting to evaluate the cumulative anthropogenic deposition flux of heavy metals to the Greenland ice cap from 1773 to 1992, that is, from the onset of the first Industrial Revolution to present. The calculations were made from the spline-smoothed curves shown in Figures 3 and 4 after subtracting the natural contribution estimated from the 7760 years B.P. sample. The cumulative deposition flux during this 220-year period is found to be $\sim 3200 \text{ t}$ for Pb, which is 1 order of magnitude higher than the deposition (400 t) documented for the Greco-Roman times [Hong *et al.*, 1994]. For Zn, Cd, and Cu the fluxes are ~ 2500 , 60 , and 200 t respectively.

Acknowledgments. This work is a contribution of the Eurocore program funded by the Commission of European Communities and Switzerland within the framework of Cost 611. It was supported by the French Ministry of the Environment (grant 92295), the Institut National des Sciences de l'Univers, and the University Joseph Fourier of Grenoble. We thank C. Rado, R.J. Delmas, and F. Maupetit for their participation in field sampling.

References

- Appelquist, H., K. O. Jensen, T. Sevel, and C. Hammer, Mercury in the Greenland ice sheet, *Nature*, **273**, 657-659, 1978.
- Barrie, L. A., Arctic air pollution: an overview of current knowledge, *Atmos. Environ.*, **20**, 643-663, 1986.
- Boutron, C. F., and C. C. Patterson, Lead concentration changes in Antarctic ice during the Wisconsin/Holocene transition, *Nature*, **323**, 222-225, 1986.
- Boutron, C.F., A clean laboratory for ultralow concentration heavy metal analysis, *Fresenius Z. Anal. Chem.* **337**, 482-491, 1990.
- Boutron, C. F., U. Görlach, J. P. Candelone, M. A. Bolshov, and R. J. Delmas, Decrease in anthropogenic lead, cadmium and zinc in Greenland snows since the late 1960s, *Nature*, **353**, 153-156, 1991.
- Boutron, C. F., J. P. Candelone, and S. Hong, Past and recent changes in the large scale tropospheric cycles of lead and other heavy metals as documented in Antarctic and Greenland snow and ice: a review, *Geochim. Cosmochim. Acta*, **58**, 3217-3225, 1994.
- Boutron, C. F., J. P. Candelone, and S. Hong, Greenland snow and ice cores: unique archives of the large scale pollution of the troposphere of the Northern Hemisphere for lead and other heavy metals, *Sci. Total Environ.*, **160/161**, 233-241, 1995.
- Boyle, E. A., R. M. Sherrell, and M. P. Bacon, Lead variability in the western North Atlantic Ocean and central Greenland ice: implications for the search of decadal trends in anthropogenic emissions, *Geochim. Cosmochim. Acta*, **58**, 3227-3238, 1994.
- Candelone, J. P., S. Hong, and C. F. Boutron, An improved method for decontaminating polar snow or ice cores for heavy metal analysis, *Anal. Chim. Acta*, **299**, 9-16, 1994.
- Candelone, J. P., J. L. Jaffrezo, S. Hong, C. Pellone, C. I. Davidson and C. F. Boutron, Seasonal variations in heavy metal concentrations in present day Greenland snow, *Earth Planet. Sci. Lett.*, in press, 1995.
- Carr, R. A., and P. E. Wilkiss, Mercury in the Greenland ice sheet: further data, *Science*, **181**, 843-844, 1973.
- Colonial Geological Survey, *Statistical Summary of the Mineral Industry. (Production, Imports and Exports)*, vols 1945-1951, Miner. Resour. Div., Her Majesty's Stationery Off., London, 1945-1951.
- Cragin, J. H., M. M. Herron, and C. C. Langway Jr., The chemistry of 700 years of precipitation at Dye 3, Greenland, *Res. Rep. 341*, U.S. Army Cold Regions Res. Eng. Lab., Hanover, New Hampshire, 1975.

- Gillet, F., D. Donnou, C. Girard, A. Manouvrier, C. Rado, and G. Ricou, Ice core quality in electro-mechanical drilling, in *Ice Drilling Technology, Rep. 84-34*, edited by G. Holdsworth, K. C. Kuivinen, and J. H. Rand, U.S. Army Cold Regions Res. Eng. Lab., Hanover, New Hampshire, Rep. 84-34, 1984.
- Görlach, U., and C. F. Boutron, Preconcentration of lead, cadmium, copper and zinc in water at the pg/g level by non-boiling evaporation, *Anal. Chim. Acta*, 236, 391-388, 1990.
- Görlach, U. and C. F. Boutron, Changes in heavy metals concentrations in Antarctic snows from 1940 to 1980, *J. Atmos. Chem.* 14, 205-222, 1992.
- Granjean, P., Health significance of organolead compounds, in *Lead versus Health*, edited by M. Rutter and R. Russell Jones, pp 179-189, John Wiley, New York, 1983.
- Herron, M. M., C. C. Langway Jr., H. V. Weiss and J. H. Cragin, Atmospheric trace metals and sulfate in the Greenland ice sheet, *Geochim. Cosmochim. Acta*, 41, 915-920, 1977.
- Hong, S., J. P. Candelone, C. C. Patterson and C. F. Boutron, Greenland ice evidence of hemispheric pollution for lead two millennia ago by Greek and Roman civilizations, *Science*, 265, 1841-1843, 1994.
- Imperial Mineral Resources Bureau, *The Mineral Industry of the British Empire and Foreign Countries. Statistical Summary (Production, Imports and Exports)*, vols 1913-1920, His Majesty's Stationery Off. London, 1913-1920.
- Imperial Institute, *The Mineral Industry of the British Empire and Foreign Countries. Statistical Summary (Production, Imports and Exports)*, vols 1920-1944, His Majesty's Stationery Off., London, 1920-1944.
- Jaffe, D., Cerundolo B., Rickers J., Stolzberg R. and A. Baklanov, Deposition of sulfate and heavy metals in the Kola peninsula, *Sci. Total Environ.*, 160/161, 127-134, 1995.
- Larbey R. J. and J. R. Bain, Trends in the use of lead as a gasoline antiknock additive, paper presented at International Lead and Zinc Conference, The Associated Octel Company Limited, London, October 5-7, 1988.
- Legrand, M., Sulphur-derived species in polar ice: a review, in *Ice Core Studies of Global Biogeochemical Cycles*, edited by R. J. Delmas, NATO-ARW Ser. I, Vol. 30, Kluwer, Dordrecht, pp. 91-120, 1995.
- Lobinski, R., C. F. Boutron, J. P. Candelone, S. Hong, J. Szpunar-Lobinska, and F. C. Adams, Speciation analysis of organolead compounds in Greenland snow at the femtogram-per-gram level by capillary gas chromatography/atomic emission spectrometry, *Anal. Chem.*, 65, 2510-2515, 1993.
- Lobinski, R., C. F. Boutron, J. P. Candelone, S. Hong, J. Szpunar-Lobinska, and F. C. Adams, Present century snow core record of organolead pollution in Greenland, *Environ. Sci. Technol.*, 28, 1467-1471, 1994.
- Loranger, S., and J. Zayed, Manganese and lead concentrations in ambient air and emission rates from unleaded and leaded gasoline between 1981 and 1992 in Canada: a comparative study, *Atmos. Environ.*, 28, 1645-1651, 1994.
- Möller, D., Estimation of the global man-made sulphur emission, *Atmos. Environ.*, 18, 19-27, 1984.
- Murozumi, M., T. J. Chow and C. C. Patterson, Chemical concentrations of pollutant lead aerosols, terrestrial dusts and sea salts in Greenland and Antarctic snow strata, *Geochim. Cosmochim. Acta*, 33, 1247-1294, 1969.
- National Environmental Research Council, *Statistical Summary of the Mineral Industry. World Production, Imports and Exports*, vols 1952-1970, Inst. of Geol. Sci. Miner. Resour. Div., Her Majesty's Stationery Off., London, 1952-1970.
- National Environmental Research Council, *Statistics. Production: Exports, Imports*, vols 1970-1990, Inst. of Geol. Sci. Miner. Stat. and Econ. Unit, Her Majesty's Stationery Off., London, 1970-1990.
- Ng, A. and C. C. Patterson, Natural concentrations of lead in ancient Arctic and Antarctic ice, *Geochim. Cosmochim. Acta*, 45, 2109-2121, 1981.
- Nriagu, J. O. (Ed.), *Biogeochemistry of Lead in the Environment*, Elsevier, New York, 1978.
- Nriagu, J. O. (Ed.), *Zinc in the Environment*, John Wiley, New York, 1979a.
- Nriagu, J. O. (Ed.), *Cadmium in the Environment*, Wiley, New York, 1979b.
- Nriagu, J. O. (Ed.), *Copper in the Environment*, Wiley, New York, 1979c.
- Nriagu, J. O., Global inventory of natural and anthropogenic emissions of trace metals to the atmosphere, *Nature*, 279, 409-411, 1979d.
- Nriagu, J. O., A global assessment of natural sources of atmospheric trace metals, *Nature*, 338, 47-49, 1989.
- Nriagu, J. O., The rise and fall of leaded gasoline, *Sci. Total Environ.*, 92, 13-28, 1990.
- Nriagu, J. O. and J. M. Pacyna, Quantitative assessment of worldwide contamination of air, water and soils by trace metals, *Nature*, 323, 134-139, 1988.
- Pacyna, J. M., Technological parameters affecting atmospheric emissions of trace elements from major anthropogenic sources, in *Control and Fate of Atmospheric Trace Metals*, edited by J. M. Pacyna and B. Ottar, pp. 15-31, Kluwer Academic, Norwell, Mass., 1989.
- Pacyna, J. M., Emission Factors of Atmospheric Cd, Pb and Zn for Major Source Categories in Europe in 1950 through 1985, *Rep. OR 30-91, Norw. Inst. for Air Res.*, Lillestrom, Norway, 1991.
- Pacyna, J. M., Emissions of pollutants and their control, in *Topics in Atmospheric and Interstellar Physics and Chemistry*, edited by C. F. Boutron, pp. 135-159, Les Editions de Physique, Paris, 1994.
- Pacyna, J. M., E. Voldner, T.F. Bidleman, G. Evans and G.J. Keeler, Emissions, atmospheric transport and deposition of heavy metals and persistent organic pollutants, in *Proceedings of the 1st Workshop on Emissions and Modelling of Atmospheric Transport of Persistent Organic Pollutants and Heavy Metals*, edited by J.M. Pacyna, E. Voldner, G.J. Keeler and G. Evans, pp. 45-137, EMEP-CCC Rep. 7/93, Norw. Inst. for Air Res., Lillestrom, Norway, 1993.
- Patterson, C. C., An alternative perspective - Lead pollution in the human environment: origin, extent and significance, in *Lead in the Human Environment*, pp. 265-349, Natl. Acad. of Sci., Washington, D. C., 1980.
- Patterson, C. C. and D. M. Settle, The reduction of orders of magnitude errors in lead analysis of biological materials and natural waters by evaluating and controlling the extent and sources of industrial lead contamination introduced during sample collection and analysis, in *Accuracy in Trace Analysis*, edited by P. La Fleur, Natl. Bur. Stand. Spec. Publ. 422, 321-351, 1976.
- Paulsen, P. J., E. S. Beary, D. S. Bushee and J. R. Moody, Inductively coupled plasma mass spectrometric analysis of ultrapure acids, *Anal. Chem.*, 60, 971-975, 1988.
- Peel, D. A., Trace metals and soot in ice cores, in *The Environmental Record in Glaciers and Ice Sheets (Dahlem Konferenzen)*, edited by H. Oeschger and C. C. Langway Jr., pp. 207-223, John Wiley, New York, 1989.
- Rosman, K. J. R., W. Chisholm, C. F. Boutron, J. P. Candelone and U. Görlach, Isotopic evidence for the sources of lead in Greenland snows since the late 1960s, *Nature*, 362, 333-335, 1993.

- Rosman, K. J. R., W. Chisholm, C. F. Boutron, J. P. Candelone and S. Hong, Isotopic evidence to account for changes in the concentration of lead in Greenland snow between 1960 and 1988, *Geochim. Cosmochim. Acta*, **58**, 3265-3269, 1994.
- Rosman, K. J. R., W. Chisholm, C. F. Boutron, J. P. Candelone, and S. Hong, The changing isotopic composition of lead in snow deposited in central Greenland since the 18th century, *Earth Planet. Sci. Lett.*, in press, 1995.
- Savarino, J., C. F. Boutron and J. L. Jaffrezo, Short-term variations of Pb, Cd, Zn and Cu in recent Greenland snow, *Atmos. Environ.* **28**, 1731-1737, 1994.
- Taylor, S. R. and S. M. McLennan, *The Continental Crust: its composition and evolution*, Blackwell, Cambridge, Mass., 1985.
- Thomas, V. and T. G. Spiro, Emissions and exposure to metals: cadmium and lead, in *Industrial Ecology and Global Change*, edited by R. Socolow, C. Andrew, F. Berkhout, and V. Thomas, pp. 297-317, Cambridge Univ. Press, New York, 1994.
- U.S. Bureau of Mines, *Mineral Yearbooks*, vols 1930-1990, Washington, D. C., 1930-1990.
- Van Cleuvenbergen, R. J. A. and F. C. Adams, Organolead compounds, in *Handbook of Environmental Chemistry*, edited by O. Hutzinger, Springer-Verlag, New York, 1990.
- Vandal, G. M., W. F. Fitzgerald, C. F. Boutron, and J. P. Candelone, Variations in mercury deposition to Antarctica over the past 34,000 years, *Nature*, **362**, 621-623, 1993.
- Weiss, H. V., M. Koide, and E. D. Goldberg, Mercury in the Greenland ice sheet: evidence of recent input by man, *Science*, **174**, 692-694, 1971a.
- Weiss, H. V., M. Koide, and E. D. Goldberg, Selenium and sulfur in a Greenland ice sheet: relation to fossil fuel combustion, *Science*, **172**, 261-263, 1971b.
- Weiss, H. V., K. Bertine, M. Koide, and E. D. Goldberg, The chemical composition of a Greenland glacier, *Geochim. Cosmochim. Acta*, **39**, 1-10, 1975.
- Wolff, E. W., Signals of atmospheric pollution in polar snow and ice, *Antarct. Sci.*, **2**, 189-205, 1990.
- Wolff, E. W., and D. A. Peel, The record of global pollution in polar snow and ice, *Nature*, **313**, 535-540, 1985.
- Wolff, E. W., and D.A. Peel, Concentrations of cadmium, copper, lead and zinc in snow from near Dye 3 in south Greenland, *Ann. Glaciol.*, **10**, 193-197, 1988.
- Wolff, E. W. and E. D. Suttie, Antarctic snow record of southern hemisphere lead pollution, *Geophys. Res. Lett.*, **21**, 781-784, 1994.

C.F. Boutron, J.P. Candelone and S. Hong, Laboratoire de Glaciologie et Géophysique de l'Environnement du CNRS, 54, rue Molière, B.P. 96, 38402 Saint Martin d'Hères, France.

C. Pellone, Centre de Recherche et d'Essais des Machines Hydrauliques de Grenoble, Domaine Universitaire, B.P. 95 X, 38402 Saint Martin d'Hères, France.

(Received September 21, 1994; revised February 9, 1995; accepted February 24, 1995.)