



Crustal and upper mantle structure beneath south-western margin of the Arabian Peninsula from teleseismic tomography

Félicie Korostelev, Clémence Basuyau, Sylvie Leroy, Christel Tiberi, Abdulhakim Ahmed, Graham Stuart, Derek Keir, Frédérique Rolandone, Ismail Al Ganad, Khaled Khanbari, et al.

► To cite this version:

Félicie Korostelev, Clémence Basuyau, Sylvie Leroy, Christel Tiberi, Abdulhakim Ahmed, et al.. Crustal and upper mantle structure beneath south-western margin of the Arabian Peninsula from teleseismic tomography. *Geochemistry, Geophysics, Geosystems*, 2014, pp.1-40. 10.1002/2014GC005316 . hal-01015496

HAL Id: hal-01015496

<https://hal.science/hal-01015496>

Submitted on 26 Jun 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**¹ Crustal and upper mantle structure beneath
² south-western margin of the Arabian Peninsula from
³ teleseismic tomography**

Félicie Korostelev,^{1,2} Clémence Basuyau,³ Sylvie Leroy,^{1,2} Christel Tiberi,⁴

Abdulhakim Ahmed,^{1,2,5} Graham W. Stuart,⁶ Derek Keir,⁷ Frédérique

Rolandone,^{1,2} Ismail Al Ganad,⁸ Khaled Khanbari⁹ and Lapo Boschi¹

Corresponding author: F. Korostelev, Univ. Paris 06 CNRS ISTEP-UPMC, Paris, France.

(felicie.korostelev@upmc.fr)

¹Sorbonne Universités, UPMC Univ Paris

4 **Abstract.** We image the lithospheric and upper asthenospheric struc-
5 ture of western continental Yemen with 24 broadband stations to evaluate

06, UMR 7193, Institut des Sciences de la
Terre Paris (iSTeP), F-75005 Paris, France

²CNRS, UMR 7193, Institut des Sciences
de la Terre Paris (iSTeP), F-75005 Paris,
France

³Univ. Paris Diderot, Institut de
Physique du Globe de Paris, Paris, France.

⁴CNRS Géosciences Montpellier, France.

⁵Seismological and Volcanological
Observatory Center, Dhamar, Yemen.

⁶School of Earth and Environment,
University of Leeds, Leeds, UK.

⁷National Oceanography Centre
Southampton, University of Southampton,
Southampton, U.K.

⁸Yemen Geological Survey and mineral
Resources Board, Sana'a, Yemen.

⁹Sana'a University, Yemen Remote
Sensing and GIS Center, Sana'a, Yemen.

6 the role of the Afar plume on the evolution of the continental margin and
7 its extent eastwards along the Gulf of Aden. We use teleseismic tomography
8 to compute relative P-wave velocity variations in southwestern Yemen down
9 to 300 km depth. Published receiver function analysis suggest a dramatic and
10 localized thinning of the crust in the vicinity of the Red Sea and the Gulf
11 of Aden, consistent with the velocity structure that we retrieve in our model.
12 The mantle part of the model is dominated by the presence of a low veloc-
13 ity anomaly in which we infer partial melting just below thick Oligocene flood
14 basalts and recent off-axis volcanic events (from 15 Ma to present). This low
15 velocity anomaly could correspond to an abnormally hot mantle and could
16 be responsible for dynamic topography and recent magmatism in western
17 Yemen. Our new P-wave velocity model beneath western Yemen suggests the
18 young rift flank volcanoes beneath margins and on the flanks of the Red Sea
19 rift are caused by focused small-scale diapiric upwelling from a broad region
20 of hot mantle beneath the area. Our work shows that relatively hot mantle,
21 along with partial melting of the mantle, can persist beneath rifted margins
22 after breakup has occurred.

1. Introduction

The Afar triple junction, where the Red Sea, East African and Gulf of Aden rifts intersect (Fig.1), is a key region to understand how continental breakup occurred under the influence of a plume and abnormally hot mantle [e.g. *Bastow et al.*, 2011]. Many seismological studies have been carried out in north-east Africa and Arabia to determine the depths of the Moho [e.g. *Ahmed et al.*, 2013; *Mechie et al.*, 2013] and to image upper mantle structure and understand regional geodynamics [e.g. *Bastow et al.*, 2005; *Benoit et al.*, 2003, 2006; *Montagner et al.*, 2007; *Zhao*, 2007; *Sicilia et al.*, 2008; *Chang and Van der Lee*, 2011; *Koulakov*, 2011] and how it is connected with global mantle flow [e.g. *Montelli et al.*, 2006; *Boschi et al.*, 2007, 2008; *Forte et al.*, 2010; *Moucha and Forte*, 2011]. However no previous study has the resolution in continental Yemen on the Gulf of Aden margin due to the lack of seismic stations. Surface wave studies [e.g. *Debayle et al.*, 2001; *Sebai et al.*, 2006; *Montagner et al.*, 2007; *Sicilia et al.*, 2008; *Chang et al.*, 2011; *Chang and Van der Lee*, 2011] lack sufficient lateral resolution to image the detail of lithospheric and uppermost mantle structures.

The YOCMAL project (YOung Conjugate MArgins Laboratory) deployed 23 broadband stations in a network running from the Red Sea margin to Aden city, passing through the Yemeni highlands and Sana'a city (Fig.2). Using classical teleseismic tomography [*Aki et al.*, 1977] on recordings from these stations together with a permanent Geofon station (DAMY), we image the relative velocity variations of P-waves in the crust and upper mantle down to 300 km depth. We thus: (1) characterize the lithospheric structure of the rifted margins of the Gulf of Aden and Red Sea system of western Yemen; (2) locate

the presence of asthenospheric upwellings in the region and their interaction with the lithosphere.

2. Geodynamic setting

The Red Sea and Gulf of Aden rifts are connected to the East African rift at the Afar triple junction, in the 'Horn of Africa'. In the Afar triple junction region, the presence of flood basalts [e.g. *Baker et al.*, 1996; *Hofmann et al.*, 1997; *George et al.*, 1998] and an abnormally low velocity upper mantle anomaly [e.g. *Debayle et al.*, 2001; *Bastow et al.*, 2005] could be due to the presence of a mantle plume (Fig.1). Around 35 Ma ago, the current rift system began to open, under the influence of the Afar plume [e.g. *Leroy et al.*, 2012]. The flood basalts of Ethiopia and Yemen are the signatures of voluminous eruptions produced during the Paleogene [*Ebinger and Sleep*, 1998] with highest eruption rates at 31 Ma [*Baker et al.*, 1996; *Hofmann et al.*, 1997; *George et al.*, 1998]. A renewed phase of volcanism took place 24 Ma ago that corresponds with the synchronous appearance of basaltic dikes and gabbros along the Red Sea, from Afar and Yemen to northern Egypt. From 25 to 16 Ma, a series of basaltic, trachytic and rhyolitic dikes were emplaced along the Red Sea margins [*Zumbo et al.*, 1995], at the same time as emplacement of large granitic batholiths (Jabal Hufash and Jabal Bura, see Fig.2). These granites are oriented north-south [*George et al.*, 1998] and located at the border of the Great Escarpment, a 1000 km long sudden change in altitude, from 200 m on the west in the Tihama Plain, to more than 1000 m in the Yemeni highlands. In addition, syn-rift (30 to 16 Ma) basaltic flows dipping towards the sea (seaward dipping reflectors - SDRs) have been imaged under the Tihama Plain [*Davison et al.*, 1994]. In the Red Sea, a second phase of opening began 14 Ma ago synchronous with the formation of the Dead Sea transform fault to the north.

This phase is accompanied by increased extension rates and rift-flank uplift [*Courtillot et al.*, 1999].

From 12 Ma to present, magmatic provinces developed within a 2500 km radius of Afar and southwestern Yemen as far away as western Saudi Arabia, Jordan and northern Syria, [*Zumbo et al.*, 1995; *Bertrand et al.*, 2003; *Coulié et al.*, 2003], see Fig.1. Recent magmatism has occurred offshore in the ocean-continent transition of the Eastern Gulf of Aden margin [*Lucazeau et al.*, 2009; *Autin et al.*, 2010; *Watremez et al.*, 2011], off-axis Sheba ridge [*d'Acremont et al.*, 2010] and below the southern Oman continental margin [*Basuyau et al.*, 2010] (Fig.1). The increased magmatism caused by extension above a plume was first thought to stop at the Shukra al Sheik fracture zone [e.g. *Hébert et al.*, 2001], but these recent results indicate that the limit could be further east (Fig.1), as proposed by *Leroy et al.* [2010a].

The Gulf of Aden is characterized by two stretched continental margins systems: non volcanic margins in the east and volcanic margins in the west near the Afar triple junction. The volcanic margins are associated with syn-rift SDRs up to 5 km thick [*Tard et al.*, 1991; *Leroy et al.*, 2012]. SDRs are also sparsely found in the east especially at the ocean-continent transition [*Autin et al.*, 2010; *Leroy et al.*, 2010b] but no syn-rift volcanism has been found east of the longitude 46°E, see Fig.1 [*Leroy et al.*, 2012]. The study region is near the edge of the African superplume, as shown by large scale global tomographic models [e.g. *Debaille et al.*, 2001; *Sebai et al.*, 2006; *Boschi et al.*, 2007, 2008] and by petrologically derived temperature estimates of the mantle [*Rooney et al.*, 2012; *Rolandone et al.*, 2013].

3. Data

Data have been collected from 23 temporary broadband stations deployed from March 2009 to March 2010 (YOCMAL project), and from one permanent station (DAMY, Geofon). The network extends from the Red Sea margin to the Gulf of Aden margin, passing through the Yemeni highlands (Fig.2). The sensors deployed were Guralp 40T (sampling rate 50 sps, 30 s natural period), 6TD (sampling rate 40 sps, 30 s natural period) and ESPD (sampling rate 40 sps, 60 s natural period). This network configuration allows imaging structures with a high resolution, and down to 300 km depth (the surface extent of our network). We selected 200 teleseismic events with clear first P-wave arrivals from earthquakes with magnitude ≥ 5.5 . Among them, 142 events arrived as P, 35 events as PP and 23 events as PKP (Fig.3), and were picked on consistent peaks in the first cycle. We used 2456 delay times calculated relative to the IASP91 Earth model [Kennett and Engdahl, 1991]. For each residual, a picking error was assigned within the range ± 0.05 to ± 0.2 s. As the events registered come mostly from east / northeast and have a range of epicentral distances, the rays are crossing, and we expect our best resolution in this direction.

4. Method

Our delay time residuals were inverted using the ACH teleseismic tomography method first used by Aki *et al.* [1974], then developed by Weiland *et al.* [1995], Zeyen and Achauer [1997] and Jordan and Achauer [1999]. As the inversion uses a 3D iterative ray tracing at each step, the initial location of the parameterization nodes affects the inversion result [e.g. Calò *et al.*, 2008]. To reduce this problem, we averaged the results of four inversions with the same parameters of inversion, but with a meshing shifted by half a node (10 km)

towards the east, north, and northeast compared with our initial reference model. This 'Average Smooth model' technique smooths the local effects due to the meshing [Evans and Achauer, 1993]. The resulting velocity model is presented in Fig.4 as depth slices and in Fig.5 as depth cross-sections.

Our network's dimensions are 360 km from North to South, 260 km from West to East, so our investigation depth (i.e. the depth until which we have resolution) is estimated to be ~ 300 km [Evans and Achauer, 1993]. In our initial model, velocities are organized in successive horizontal layers of nodes, with an interpolation gradient between each of them [Thurber, 1983]. The minimum distance between two nodes horizontally is 20 km (i.e. the minimum distance between two stations). The node spacing is 40 to 100 km at the edges of the model. The initial model was the IASP91 reference Earth model. Ten levels of nodes were placed in depth between 0 and 500 km, so we have nine layers (nodes at 0, 30, 45, 70, 100, 150, 200, 250, 300 and 500 km depth). The thickness of the crust is important in the inversion process [e.g. Zhao *et al.*, 1994, 2006]. In our study we use the receiver functions results of Ahmed *et al.* [2013] to compute a time correction for each station (Fig.6). This correction accounts for lateral variations in the crustal thickness, which are difficult to constrain by our teleseismic data alone; seismic velocity within the crust is treated as a free parameter in our inversion. The correction for each station is computed on the absolute residuals and is in the range -0.611 s (for the station located on the thinnest crust) to 0.160 s (for the station located above the thicker crust). This allows a reduction in the propagation of crustal structure into the deeper layers of the velocity model.

The smoothing factor, which limits the short wavelength velocity variations, and the ini-

132 tial standard deviation associated with each node of the initial model for each of the nine
 133 iterations were chosen after a series of tests and are respectively 0.001 and 0.007 km.s⁻¹.
 134 The smoothing value chosen is the same as *Tiberi et al.* [2008] and *Basuyau et al.* [2010].
 135 Tests showed that the results are not significantly different for a standard deviation be-
 136 tween 0.005 and 0.01 km.s⁻¹. We tested these inversion parameters in order to get a
 137 stable model, choosing parameters that decrease the root mean square of residuals (RMS)
 138 through the inversion's nine iterations. The overall decrease of the RMS is 55% in the
 139 final model.

5. Results

5.1. Checkerboard test

140 We use a checkerboard test to assess the resolution of our inversion. The synthetic
 141 checkerboard model consisted of rectangular velocity anomalies of +5% and -5% velocity
 142 variations at depths of 70, 200 and 300km depth (Fig.7a). The size of the anomalies in-
 143 creases around the edges as the method requires the nodes to be further apart at the edges
 144 of the model. We use the same inversion parameters (smoothing, standard deviation) as
 145 in our actual inversion.

146 The results of the checkerboard test (Fig.7b) show that at 70 km depth, due to the con-
 147 centration of crossing rays under the stations, we have the best resolution, with ~ 20 %
 148 recovered amplitude. At 200 km depth, the crossing rays cover a larger area, so the
 149 anomalies are well retrieved. At 300 km depth, the eastern retrieved anomalies are much
 150 better constrained east of 44°E, due to the concentration of ray paths coming from the
 151 east. The Fig.7b shows the piercing points under our network at 300 km depth, at our

maximum investigation depth. Below 300 km depth, the rays are too dispersed to give good resolution.

5.2. Crustal-scale structures

The thickness of the crust in our study area is estimated from receiver function analysis to be between 14 km at the coast and 35 km inland [Ahmed *et al.*, 2013]. The crust is represented by the first two layers of our model, and is characterized by the highest anomaly contrasts in P-wave velocity, which have a range of $\pm 4\%$ (Fig.4 and 5). These anomalies are related to geological structures observed at the surface. At 30 km depth, the strong low velocity anomaly (-4%) beneath the center of our network (stations MAWI to DALA) correlates with the high topography of the plateau (>2000 m above sea level). We interpret this pattern as due to the thicker crust (>30 km) beneath the Yemeni highlands [Ahmed *et al.*, 2013].

Under stations located near the Red Sea (north-westernmost part of the network) and Gulf of Aden (southernmost part of the network) margins there are high velocity anomalies above 30 km due to the thinner crust (<30 km). Ahmed *et al.* [2013] estimated the thickness of the crust to be ~ 22 km in the coastal areas, and less than 14 km for the Red Sea margin. These high velocity anomalies are located beneath SDRs, which is consistent with the emplacement of sub-aerial volcanic material during rifting [Tard *et al.*, 1991; Davison *et al.*, 1994; Bastow and Keir, 2011; Leroy *et al.*, 2012].

The high velocity anomalies under the stations ANID and SUGH could be due to a thin crust as they are in the Red Sea coastal area, or to Tertiary granitic intrusions of Jabal Hufash and Jabal Bura respectively [Geoffroy *et al.*, 2002], see Fig.2. The high velocity anomaly corresponding to Jabal Hufash is imaged down to 45 km depth, which is slightly

deeper than that of Jabal Bura (30 km depth). This can be explained by significant smearing due to the higher amplitude of the Jabal Hufash anomaly. There are lower velocities under the stations UAYA and ZUWA, on the Tihama plain (Fig.5b), probably due to ~ 4000 m of low velocity sediments [El-Anbaawy *et al.*, 1992; Davison *et al.*, 1994].

5.3. Upper mantle structures

The resulting upper-mantle P-wave anomalies are in the range of $\pm 2\%$. The most striking pattern is a low velocity anomaly located under the Yemeni highlands at 45 km depth, apparently dipping northeastward down to 300 km. It reaches its maximum amplitude at 70 km depth (east of the DAMY station) beneath the volcanic field of Dhamar (Fig.2), where there are two active volcanoes [Manetti *et al.*, 1991]. The northern part of this anomaly is located, at 70 km depth, under the volcanic field of Sanaa, which is also still active [Manetti *et al.*, 1991], see Fig.5.

There is a second low-velocity anomaly located under the southwestern corner of Yemen and the stations MOKA and HAKI (Fig.5c and d). This low velocity anomaly is nearly vertical and is recognized from the surface to 300 km. It is located just beneath the volcanic area of Jabal An Nar (Fig.5d), which was active during late Miocene, around 10 Ma, [Manetti *et al.*, 1991].

Even if we corrected the residuals from the crustal portion, our results are likely to include effects from Moho variations. This is because the corrections are based on receiver functions which display a strong trade-off between Moho depth and crustal velocity model. In addition, we took a 1D velocity model to be consistent with the receiver functions study of [Ahmed *et al.*, 2013] and this can generate errors [Martin and Ritter, 2005]. The lack of detailed 3D crustal information precludes us from going further with the crustal

corrections than we have done. To estimate the resolution of our models in both the crust and the mantle, we proceed to synthetic tests. We test whether the low velocity anomaly beneath the high plateau is dipping towards the northeast because of smearing along ray paths (Fig.3) in the next section. We investigate by means of synthetic tests whether: (1) the velocity variations observed in our resulting model are smearing downwards into the mantle, (2) the low velocity anomalies under the southwestern corner of Yemen and under the high plateaus are artefacts, and whether we can determine at what depth they are located, (3) the dipping low-velocity anomaly is related to the presence of partial melt or not.

6. Synthetic tests and presence of melt

6.1. Propagation of crustal signal

We test the smearing with depth of a crustal anomaly (0-30 km) without a crustal correction by introducing a -5% anomaly around stations in the Yemeni Highlands and a +5% anomaly under the two continental margins (Fig.8a). The rest of the input model is laterally uniform. The results of the inversion of this synthetic model (Fig.8b and c) show that the velocity variations are well recovered in location but with 40% lower amplitude. Between 45 and 100 km depth, we can discern very low amplitude anomalies ($<0.5\%$), corresponding to a small amount of vertical smearing. At 100 km depth, we have a -1% anomaly under the Yemeni highlands. Approximately half of this signal could be due to a smearing of the crust signal. No significant anomaly can be seen beneath 100 km depth. We conclude that crustal velocity anomalies do not propagate to deeper layers of our model and that there is an authentic low velocity anomaly in the upper mantle. This

test shows that the use of a crustal correction is relevant and is necessary to limit the extent of the propagation of crustal velocity structure into deeper layers of the model.

6.2. Low velocity anomalies beneath Southwestern Yemen

Using a synthetic input model, we simulate the resulting geometry of our final P wave model (see auxiliary material). We computed a series of tests (available in auxiliary material), and here we present the most relevant example. We first place a -5% anomaly under the three Yemeni volcanic fields of Sana'a, Dhamar and Marib from the base of the crust down to 200 km depth. The other -5% anomaly is placed from the base of the crust to the base of our model, under the southwestern corner of Yemen and southern Red Sea. Figure 9 shows the retrieved inversion image for a SW-NE profile (compare with the observed results in Fig.9c). Although the anomaly amplitudes within the crust are not retrieved, the dipping anomaly is quite well retrieved in the synthetic output model, as well as the low velocity anomaly beneath Jabal An Nar volcanic field (Fig.9). This tests shows that the dipping low velocity anomaly under the Yemeni volcanic fields can be explained by a 220x260 km mantle upwelling between the base of the crust and 200 km depth. Moreover, the low velocity anomaly beneath Jabal An Nar can be explained by a large zone of hotter mantle.

6.3. Presence of partial melt

Traveltime tomography gives the present state of the upper mantle in terms of velocity variations but it precludes any direct interpretation concerning their origin. Indeed, several factors, such as temperature, chemical composition or anisotropy can affect the velocity of seismic waves [e.g. *Karato*, 1993; *Sobolev et al.*, 1996]. *Karato* [1993] demon-

strates that a purely thermal origin leads to a linear relationship between P and S residuals with a slope of 2.9. *Gao et al.* [2004] proposed that a slope higher than 2.9 between P and S residuals for the same events highlights the presence of partial melting and several authors used the relationship to explain the presence of negative velocity anomalies in the upper mantle [e.g. *Bastow et al.*, 2005; *Basuyau et al.*, 2010]. This method considers relative delay times, so that problems associated with amplitude recovery (e.g. due to differing numbers of traveltime observations and regularization levels) and other artefacts associated with the inversion procedure (i.e. parameterization and ray-path accuracy) do not complicate the comparison of the data.

We thus selected the events coming from northeast (Russia, Japan, China), and picked the S arrival on the transverse component for stations located above the Yemeni Highlands, so that the rays chosen are passing through the northeastward dipping low-velocity anomaly. In Fig.10, following *Gao et al.* [2004], *Bastow et al.* [2005] and *Basuyau et al.* [2010], we plot S versus P relative travel-time residuals, and find out that our data are consistent with a slope >2.9 , thus implying the presence of partial melt along the rays coming from northeast. The presence of Holocene volcanoes (in Sana'a, Dhamar and Marib volcanic fields) on the surface helps support the idea that there is partial melt being created in the asthenosphere and intruding the lithosphere. Moreover, the isotopic signatures of the melts from the three volcanic fields suggest a strong asthenospheric component [*Manetti et al.*, 1991], which is consistent with our results.

7. Discussion

7.1. Crustal-scale structures

We produce a relative velocity model for the propagation of P waves down to 300 km beneath western Yemen. The low velocity anomaly ($\sim -4\%$) located below the Yemeni highlands between 0 and ~ 35 km corresponds to a region of 30-35 km thick crust [*Ahmed et al.*, 2013]. At 30 km depth, we observe a dramatic transition from very low to high velocities (-4% to $+4\%$) under the coast of the Red Sea and Gulf of Aden. We interpret this sudden short length scale (<40 km for the Red Sea margin, <100 km for the Gulf of Aden margin) variation as a lateral transition between a thick crust and a thinned, intruded and stretched crust. In addition, our new seismic images showing lower mantle velocities under the southwestern corner of Yemen is consistent with ongoing rifting above a thermal anomaly in the underlying mantle, due to the Afar plume. Under these conditions, melt generated by adiabatic decompression of the asthenosphere beneath thinned and stretched lithosphere migrates upwards to intrude or underplate continental crust and extrude as mostly basalt flows (Oligocene flood basalts) [*White and McKenzie*, 1989]. Our positive V_p anomaly near the base of the crust under the Red Sea and Gulf of Aden margin are consistent with melt produced from an abnormally hot mantle which enriches the melt in MgO [*White and McKenzie*, 1989], capable of producing intrusions/underplate of up to 7.2 km/s rather than 6.8 km/s from melting normal mantle. These high velocity anomalies are focused mainly into narrow zones of denser material, away from the most stretched areas. That is because the mantle temperature was highest at the start of continental break-up [*White et al.*, 2008]. Such lower crustal intrusions/underplating are a common feature of volcanic margins such as the north Atlantic [*Geoffroy*, 2005; *Mjelde et al.*, 2005;

277 *White et al.*, 2008].

278 Placing constraints on the timing of the underplating is difficult. If emplaced before the
 279 eruption of SDRs and continental break-up as has been proposed for the north Atlantic (at
 280 least 10 to 15 Ma off Norway, *Gernigon et al.* [2006]) then it would have an influence on the
 281 structural development of the margin and partly consist of high pressure granulite/eclogite
 282 lower crustal rocks [*Gernigon et al.*, 2006].

283 At shallower depths of 0-30 km in our model, seismic anomalies are directly related to
 284 the geological units observed at the surface. The granitic intrusions of Jabal Hufash and
 285 Jabal Bura are associated with high velocity anomalies (up to +4%, see Fig.2). The depth
 286 extent of the anomaly below the granitic intrusions of Jabal Hufash may indicate a deeper
 287 root, as hypothesized by *Denèle et al.* [2012]. It could also be explained by a stronger
 288 smearing effect due to the high amplitude of the Jabal Hufash anomaly.

7.2. Deep structure of the margins

289 Our synthetic tests show that deep anomalies cannot be explained by the smearing
 290 in depth of crustal anomalies. We interpret the low velocity anomaly (-2%) between
 291 ~35 km and 300 km depth under the highest topography as abnormally hot mantle up-
 292 welling. This low velocity hotter mantle is located below the stations DAMY, RUSA and
 293 YSLE (Fig.5), which are located on the thick Oligocene flood basalts and three more
 294 recent volcanic fields (15 Ma to present) volcanic areas, e.g. Dhamar and Sana'a [*Davison*
 295 *et al.*, 1994; *Pik et al.*, 2008; *Leroy et al.*, 2010b], see Fig.2. Additionally, we infer presence
 296 of partial melt in the crust or uppermost mantle to be responsible for this low velocity
 297 anomaly (Fig.10). A similar pattern of upper mantle off-axis upwelling has also been
 298 found in the southern Red Sea rift of Afar and explained by small diapiric upwellings

(<100 km) [*Hammond et al.*, 2013]. We surmise that a similar mechanism is the best explanation for our observations. This idea is supported by the large asthenospheric component in the magma from the three Yemeni volcanic fields inferred from trace element and isotope geochemistry [*Manetti et al.*, 1991]. Observations of recent dike intrusions at Harrat Lunayirr in Saudi Arabia [*Pallister et al.*, 2010] show rifted margin magmatism may be important in accomodating extension after breakup along the full length of the Red Sea margin [*Ebinger and Belachew*, 2010].

Our new relative P-wave velocity model beneath western Yemen suggests the young rift flank volcanoes on the margin of the Red Sea rift are caused by focused small-scale diapiric upwelling from a broad region of hot mantle beneath the area. Our work shows that relatively hot mantle, along with partial melting of the mantle, can persist beneath rifted margins after breakup has occurred. These findings have important implications for interpreting the thermal history and deformation of volcanic rifted margins after breakup is achieved since most breakup models assume rift margin volcanism ceases after seafloor spreading starts.

Buoyant hot mantle may contribute to a dynamic topography of the Yemeni high plateau. Almost all the topography in this region could indeed have a dynamic origin, because the rift-flank uplift from flexure [*Daradich et al.*, 2003] is not sufficient to produce high topographies over such a large area. Numerical modeling of the plume/lithosphere interaction predict an uplift of about 500 to 1500 m in less than 0.3 to 0.5 Ma after the plume initiation [*d'Acremont et al.*, 2003]. Moreover, *White and McKenzie* [1989] explained that an increase of about 150°C in the mantle leads to a dynamic uplift, but that the addition of dense igneous material under a thinned crust produces an immediate subsidence of

more than 2 km in order to maintain isostatic equilibrium. We observe a similar pattern on the Red Sea and Gulf of Aden margins, with high Yemeni plateau dynamically supported by a hot mantle (Fig.5a, b and d), and a subsiding area, for example the Tihama Plain, underlain by seaward dipping reflectors, and ultra-mafic bodies accreted under the crust. This hypothesis should however be tested for Yemen by gravity and isostatic modeling.

The weak low velocity anomaly imaged under the southwestern corner of Yemen, beneath the MOKA station, is underneath the Miocene volcanic area of Jabal An Nar. Our synthetic test (Fig.9) shows that this low velocity anomaly may be explained by a large zone of hotter mantle. This feature could be due to the Afar plume signal, located only ~ 300 km away, in the Afar depression.

8. Conclusions

We performed an inversion of P-wave teleseismic data to image lithospheric structure beneath the SW of the Yemen, the southern Red Sea and western Gulf of Aden margins. The crustal part of the model is dominated by a possible ultra-mafic underplating beneath the Red Sea and Gulf of Aden margins, a sudden thinning of the crust for this volcanic margin, and Tertiary granitic intrusions (Jabal Hufash and Jabal Bura). In the mantle, we image a low velocity anomaly in which we infer partial melting just below the highest topography, the thick Oligocene flood basalts and other off-axis volcanic regions (from 15 Ma to present). This low velocity anomaly could correspond to an abnormally hot mantle and could be responsible for dynamic topography and recent magmatism in western Yemen. Some hot material has also been inferred under the southwestern corner of Yemen and may be due to the Afar plume signal.

9. Appendix : Other synthetic tests

Acknowledgments. This project was funded by the ANR YOCMAL (Agence Na-
tionale de la Recherche), CNRS-INSU-PICS Yemen, GSMRB Yemen and is in the frame-
work of the Actions Marges program. Seismic equipment from SEIS-UK is funded by
NERC under loan number 873. We thank David Hawthorn, Alex Brisbane and Vic-
toria Lane for their efforts during the deployment and servicing of network, the French
Embassy in Yemen (J. G. Sarkis, J. Dechezlepretre and C. Bousquet), local governors and
the people of the Yemen governerates for their help during the field work.

References

- Ahmed, A., C. Tiberi, S. Leroy, G. W. Stuart, D. Keir, J. Sholan, K. Khanbari, I. Al-
Ganad, and C. Basuyau (2013), Crustal structure of the rifted volcanic margins and
uplifted plateau of Western Yemen from receiver function analysis, *Geophysical Journal
International*, *193*(3), 1673–1690.
- Aki, K., A. Christofferson, E. Husebye, and C. Powell (1974), Three-dimensional seismic
velocity anomalies in the crust and upper-mantle under the USGS, California seismic
array, *Eos Trans. AGU*, *56*, 1145.
- Aki, K., A. Christoffersson, and E. Husebye (1977), Determination of the three-
dimensional seismic structure of the lithosphere, *Journal of Geophysical Research*, *82*(2),
277–296.
- Autin, J., S. Leroy, M. Beslier, E. d’Acremont, P. Razin, A. Ribodetti, N. Bellahsen,
C. Robin, and K. Al Toubi (2010), Continental break-up history of a deep magma-poor
margin based on seismic reflection data (northeastern Gulf of Aden margin, offshore

Oman), *Geophysical Journal International*, 180(2), 501–519.

Baker, J., L. Snee, and M. Menzies (1996), A brief Oligocene period of flood volcanism in Yemen: implications for the duration and rate of continental flood volcanism at the Afro-Arabian triple junction, *Earth and Planetary Science Letters*, 138(1), 39–55.

Bastow, I., and D. Keir (2011), The protracted development of the continent-ocean transition in Afar, *Nature Geoscience*, 4(4), 248–250.

Bastow, I., G. Stuart, J. Kendall, C. Ebinger, et al. (2005), Upper-mantle seismic structure in a region of incipient continental breakup: Northern Ethiopian Rift, *Geophysical Journal International*, 162(2), 479–493.

Bastow, I., D. Keir, and E. Daly (2011), The Ethiopia Afar Geoscientific Lithospheric Experiment (EAGLE): Probing the transition from continental rifting to incipient seafloor spreading, *Volcanism and Evolution of the African Lithosphere*, 478, 51–76.

Basuyau, C., C. Tiberi, S. Leroy, G. Stuart, A. Al-Lazki, K. Al-Toubi, and C. Ebinger (2010), Evidence of partial melting beneath a continental margin: case of Dhofar, in the Northeast Gulf of Aden (Sultanate of Oman), *Geophysical Journal International*, 180(2), 520–534.

Benoit, M., A. Nyblade, J. VanDecar, and H. Gurrola (2003), Upper mantle P wave velocity structure and transition zone thickness beneath the Arabian Shield, *Geophys. Res. Lett*, 30(10), 1531.

Benoit, M., A. Nyblade, and J. VanDecar (2006), Upper mantle P-wave speed variations beneath Ethiopia and the origin of the Afar hotspot, *Geology*, 34(5), 329–332.

Bertrand, H., G. Chazot, J. Blichert-Toft, and S. Thorvaldsson (2003), Implications of widespread high- μ volcanism on the Arabian Plate for Afar mantle plume and litho-

sphere composition, *Chemical Geology*, 198(1), 47–61.

Boschi, L., T. Becker, and B. Steinberger (2007), Mantle plumes: Dynamic models and seismic images, *Geochemistry, Geophysics, Geosystems*, 8(10).

Boschi, L., T. W. Becker, and B. Steinberger (2008), On the statistical significance of correlations between synthetic mantle plumes and tomographic models, *Physics of the Earth and Planetary Interiors*, 167(3), 230–238.

Calò, M., C. Dorbath, D. Luzio, S. Rotolo, and G. d’Anna (2008), WAM tomography in the southern Tyrrhenian region. Petrological inferences and hypotheses on the fluid circulation in the subducting Ionian slab and adjoining mantle domain, *Bollettino di Geofisica Teorica ed Applicata*, 42(2), 136–141.

Chang, S., and S. Van der Lee (2011), Mantle plumes and associated flow beneath Arabia and East Africa, *Earth and Planetary Science Letters*, 302(3-4), 448–454.

Chang, S., M. Merino, S. Van der Lee, S. Stein, and C. Stein (2011), Mantle flow beneath Arabia offset from the opening Red Sea, *Geophysical Research Letters*, 38(4), L04,301.

Coulié, E., X. Quidelleur, P. Gillot, V. Courtillot, J. Lefèvre, and S. Chiesa (2003), Comparative K–Ar and Ar/Ar dating of Ethiopian and Yemenite Oligocene volcanism: implications for timing and duration of the Ethiopian traps, *Earth and Planetary Science Letters*, 206(3), 477–492.

Courtillot, V., C. Jaupart, I. Manighetti, P. Tapponnier, and J. Besse (1999), On causal links between flood basalts and continental breakup, *Earth and Planetary Science Letters*, 166(3), 177–195.

d’Acremont, E., S. Leroy, and E. Burov (2003), Numerical modelling of a mantle plume: the plume head-lithosphere interaction in the formation of an oceanic large igneous

- 410 province, *Earth and Planetary Science Letters*, 206(3-4), 379–396.
- 411 d’Acremont, E., S. Leroy, M. Maia, P. Gente, and J. Autin (2010), Volcanism, jump and
412 propagation on the Sheba ridge, eastern Gulf of Aden: segmentation evolution and
413 implications for oceanic accretion processes, *Geophysical Journal International*, 180(2),
414 535–551.
- 415 Daradich, A., J. Mitrovica, R. Pysklywec, S. Willett, and A. Forte (2003), Mantle flow,
416 dynamic topography, and rift-flank uplift of Arabia, *Geology*, 31(10), 901–904.
- 417 Davison, I., M. Al-Kadasi, S. Al-Khirbash, S., A. Al-Subbary, J. Baker, S. Blakey,
418 D. Bosence, C. Dart, R. Heaton, K. McClay, et al. (1994), Geological evolution of the
419 southeastern Red Sea Rift margin, Republic of Yemen, *Geological Society of America*
420 *Bulletin*, 106(11), 1474–1493.
- 421 Debayle, E., J. L  v  que, and M. Cara (2001), Seismic evidence for a deeply rooted low-
422 velocity anomaly in the upper mantle beneath the northeastern Afro/Arabian continent,
423 *Earth and Planetary Science Letters*, 193(3), 423–436.
- 424 Den  le, Y., R. Pik, and S. Leroy (2012), Thermal and tectonic histories of the Jabel-Bura
425 per-alkaline granite on the south-eastern margin of the Red Sea (Yemen), *Magmatic*
426 *Rifting and Active Volcanism Conference, Addis Abeba, Ethiopia.*, pp. 107–108.
- 427 Ebinger, C., and M. Belachew (2010), Geodynamics: Active passive margins, *Nature*
428 *Geoscience*, 3, 670–671.
- 429 Ebinger, C., and N. Sleep (1998), Cenozoic magmatism throughout east Africa resulting
430 from impact of a single plume, *Nature*, 395(6704), 788–791.
- 431 El-Anbaawy, M. I. H., M. A. H. Al-Aawah, K. A. Al-Thour, and M. Tucker (1992),
432 Miocene evaporites of the Red Sea Rift, Yemen Republic: Sedimentology of the Salif

halite: *Sedimentary Geology*, *81*, 61–71.

Evans, J., and U. Achauer (1993), Teleseismic velocity tomography using the ACH method: theory and application to continental-scale studies, in *Seismic Tomography Theory and Practice*, pp. 319–360, Chapman and Hall, London.

Forte, A. M., S. Quéré, R. Moucha, N. A. Simmons, S. P. Grand, J. X. Mitrovica, and D. B. Rowley (2010), Joint seismic–geodynamic–mineral physical modelling of African geodynamics: A reconciliation of deep-mantle convection with surface geophysical constraints, *Earth and Planetary Science Letters*, *295*(3), 329–341.

Gao, W., S. Grand, W. Baldrige, D. Wilson, M. West, J. Ni, and R. Aster (2004), Upper mantle convection beneath the Central Rio Grande rift imaged by P and S wave tomography, *Journal of Geophysical Research*, *109*, B03305.

Geoffroy, L. (2005), Volcanic passive margins, *C R Geoscience*, *337*, 1395–1408.

Geoffroy, L., P. Huchon, and K. Khanbari (2002), Did Yemeni Tertiary granites intrude neck zones of a stretched continental upper crust?, *Terra Nova*, *10*(4), 196–200.

George, R., N. Rogers, and S. Kelley (1998), Earliest magmatism in Ethiopia: evidence for two mantle plumes in one flood basalt province, *Geology*, *26*(10), 923.

Gernigon, L., F. Lucazeau, F. Brigaud, J. Ringenbach, S. Planke, and B. Le Gall (2006), A moderate melting model for the Vøring margin (Norway) based on structural observations and a thermo-kinematical modelling: Implication for the meaning of the lower crustal bodies, *Tectonophysics*, *412*(3), 255–278.

Hammond, J., J.-M. Kendall, G. Stuart, C. Ebinger, I. Bastow, D. Keir, A. Ayele, M. Belachew, B. Goitom, G. Ogubazghi, and T. Wright (2013), Mantle upwelling and initiation of rift segmentation beneath the Afar Depression, *Geology*, *41*(6), 635–638.

- Hébert, H., C. Deplus, P. Huchon, K. Khanbari, and L. Audin (2001), Lithospheric structure of the nascent west Gulf of Aden spreading center inferred from gravity data, *Journal of Geophysical Research*, *106*, 26–345.
- Hofmann, C., V. Courtillot, G. Féraud, P. Rochette, G. Yirgu, E. Ketefo, and P. R. (1997), Timing of the Ethiopian flood basalt event and implications for plume birth and global change, *Nature*, *389*, 838–841.
- Jordan, M., and U. Achauer (1999), A new method for the 3-D joint inversion of teleseismic delaytimes and Bouguer gravity data with application to the French Massif Central, in *EOS Trans.*, AGU Fall Meeting.
- Karato, S. (1993), Importance of anelasticity in the interpretation of seismic tomography, *Geophysical Research Letters*, *20*, 1623–1626.
- Kennett, B., and E. Engdahl (1991), Traveltimes from global earthquake location and phase identification, *Geophysical Journal International*, *105*(429-465).
- Koulakov, I. (2011), High-frequency P and S velocity anomalies in the upper mantle beneath Asia from inversion of worldwide traveltime data, *Journal of Geophysical Research: Solid Earth (1978–2012)*, *116*(B4).
- Leroy, S., E. d’Acremont, C. Tiberi, C. Basuyau, J. Autin, F. Lucazeau, and H. Sloan (2010a), Recent off-axis volcanism in the eastern Gulf of Aden: Implications for plume–ridge interaction, *Earth and Planetary Science Letters*, *293*(1), 140–153.
- Leroy, S., F. Lucazeau, E. d’Acremont, L. Watremez, J. Autin, S. Rouzo, N. Bellahsen, C. Tiberi, C. Ebinger, M. Beslier, et al. (2010b), Contrasted styles of rifting in the eastern Gulf of Aden: A combined wide-angle, multichannel seismic, and heat flow survey, *Geochemistry Geophysics Geosystems*, *11*(7), Q07,004.

- 479 Leroy, S., P. Razin, J. Autin, F. Bache, E. dAcremont, L. Watremez, J. Robinet, C. Bau-
480 rion, Y. Denèle, N. Bellahsen, et al. (2012), From rifting to oceanic spreading in the
481 Gulf of Aden: a synthesis, *Arabian Journal of Geosciences*, pp. 1–43.
- 482 Lucazeau, F., S. Leroy, J. Autin, A. Bonneville, B. Goutorbe, L. Watremez, E. DAcre-
483 mont, D. Düsünür, F. Rolandone, P. Huchon, et al. (2009), Post-rift volcanism and
484 high heat-flow at the ocean-continent transition of the eastern Gulf of Aden, *Terra*
485 *nova*, *21*(4), 285–292.
- 486 Manetti, P., G. Capaldi, S. Chiesa, L. Civetta, S. Conticelli, M. Gasparon, L. Volpe,
487 and G. Orsi (1991), Magmatism of the eastern Red Sea margin in the northern part of
488 Yemen from Oligocene to present, *Tectonophysics*, *198*(2), 181–202.
- 489 Martin, M., and J. Ritter (2005), High-resolution teleseismic body-wave tomography be-
490 neath SE Romania–I. Implications for three-dimensional versus one-dimensional crustal
491 correction strategies with a new crustal velocity model, *Geophysical Journal Interna-*
492 *tional*, *162*(2), 448–460.
- 493 Mechie, J., Z. Ben-Avraham, M. Weber, H.-J. Götze, I. Koulakov, A. Mohsen, and
494 M. Stiller (2013), The distribution of Moho depths beneath the Arabian plate and
495 margins, *Tectonophysics*, ([http:// dx.doi.org/10.1016/j.tecto.2012.11.015](http://dx.doi.org/10.1016/j.tecto.2012.11.015)).
- 496 Mjelde, R., T. Raum, B. Myhren, H. Shimamura, Y. Murai, T. Takanami, R. Karpuz,
497 and U. Næss (2005), Continent-ocean transition on the Vøring Plateau, NE Atlantic,
498 derived from densely sampled ocean bottom seismometer data, *Journal of geophysical*
499 *research*, *110*(B5), B05,101.
- 500 Montagner, J., B. Marty, E. Stutzmann, D. Sicilia, M. Cara, R. Pik, J. Lévêque, G. Roult,
501 E. Beucler, E. Debayle, et al. (2007), Mantle upwellings and convective instabilities re-

vealed by seismic tomography and helium isotope geochemistry beneath eastern Africa,
Geophys. Res. Lett., *34*, L21,303.

Montelli, R., G. Nolet, F. Dahlen, and G. Masters (2006), A catalogue of deep mantle
plumes: new results from finite-frequency tomography, *Geochem. Geophys. Geosyst.*,
7(11), 007.

Moucha, R., and A. M. Forte (2011), Changes in African topography driven by mantle
convection, *Nature Geoscience*, *4*(10), 707–712.

Pallister, J., W. McCausland, S. Jónsson, Z. Lu, H. Zahran, S. El Hadidy, A. Aburukbah,
I. Stewart, P. Lundgren, R. White, et al. (2010), Broad accommodation of rift-related
extension recorded by dyke intrusion in Saudi Arabia, *Nature Geoscience*, *3*(10), 705–
712.

Pik, R., B. Marty, J. Carignan, G. Yirgu, and T. Ayalew (2008), Timing of East African
Rift development in southern Ethiopia: Implication for mantle plume activity and evo-
lution of topography, *Geology*, *36*(2), 167.

Rolandone, F., F. Lucazeau, S. Leroy, J.-C. Mareschal, R. Jorand, B. Goutorbe, and
H. Bouquerel (2013), New heat flow measurements in Oman and the thermal state of
the Arabian Shield and Platform, *Tectonophysics*, *589*, 77–89.

Rooney, T., C. Herzberg, and I. Bastow (2012), Elevated mantle temperature beneath
East Africa, *Geology*, *40*(1), 27–30.

Sebai, A., E. Stutzmann, J. Montagner, D. Sicilia, and E. Beucler (2006), Anisotropic
structure of the African upper mantle from Rayleigh and Love wave tomography, *Physics
of the Earth and Planetary Interiors*, *155*(1), 48–62.

- 524 Sicilia, D., J. Montagner, M. Cara, E. Stutzmann, E. Debayle, J. Lépine, J. Lévêque,
525 E. Beucler, A. Sebai, G. Roult, et al. (2008), Upper mantle structure of shear-waves
526 velocities and stratification of anisotropy in the Afar Hotspot region, *Tectonophysics*,
527 *462*(1-4), 164–177.
- 528 Sobolev, S. V., H. Zeyen, G. Stoll, F. Werling, R. Altherr, and K. Fuchs (1996), Upper
529 mantle temperatures from teleseismic tomography of French Massif Central including
530 effects of composition, mineral reactions, anharmonicity, anelasticity and partial melt,
531 *Earth and Planetary Science Letters*, *139*(1), 147–163.
- 532 Tard, F., P. Masse, F. Walgenwitz, and P. Gruneisen (1991), The volcanic passive margin
533 in the vicinity of Aden, Yemen, *Bulletin Centres Recherche Exploration–Production*
534 *Elf-Aquitaine*, *15*, 1–9.
- 535 Thurber, C. H. (1983), Earthquake locations and three-dimensional crustal
536 structure in the Coyote Lake Area, central California, *J. Geophys. Res.*,
537 *88*(B10)(doi:10.1029/JB088iB10p08226), 8226–8236.
- 538 Tiberi, C., A. Deschamps, J. Déverchère, C. Petit, J. Perrot, D. Appriou, V. Mordvinova,
539 T. Dugaarma, M. Ulzibaat, and A. Artemiev (2008), Asthenospheric imprints on the
540 lithosphere in Central Mongolia and Southern Siberia from a joint inversion of grav-
541 ity and seismology (MOBAL experiment), *Geophysical Journal International*, *175*(3),
542 1283–1297.
- 543 Watremez, L., S. Leroy, S. Rouzo, E. d’Acremont, P. Unternehr, C. Ebinger, F. Lu-
544 cazeau, and A. Al-Lazki (2011), The crustal structure of the north-eastern Gulf of
545 Aden continental margin: insights from wide-angle seismic data, *Geophysical Journal*
546 *International*, *184*(2), 575–594.

Weiland, C., L. Steck, P. Dawson, and V. Korneev (1995), Nonlinear teleseismic tomography at Long Valley caldera, using three-dimensional minimum travel time ray tracing, *Journal of geophysical research*, *100*(B10), 20,379–20.

White, R., and D. McKenzie (1989), Magmatism at rift zones: the generation of volcanic continental margins and flood basalts, *Journal of Geophysical Research*, *94*(B6), 7685–7729.

White, R., L. Smith, A. Roberts, P. Christie, N. Kusznir, A. Roberts, D. Healy, R. Spitzer, A. Chappell, J. Eccles, et al. (2008), Lower-crustal intrusion on the North Atlantic continental margin, *Nature*, *452*(7186), 460–464.

Zeyen, H., and U. Achauer (1997), Joint Inversion of Teleseismic Delay Times and Gravity Anomaly Data for Regional Structures. Theory and Synthetic Examples, *Nato ASI Series 1 Disarmament Technologies*, *17*, 155–168.

Zhao, D. (2007), Seismic images under 60 hotspots: search for mantle plumes, *Gondwana Research*, *12*(4), 335–355.

Zhao, D., A. Hasegawa, and H. Kanamori (1994), Deep structure of Japan subduction zone as derived from local, regional, and teleseismic events, *Journal of Geophysical Research: Solid Earth (1978–2012)*, *99*(B11), 22,313–22,329.

Zhao, D., J. Lei, T. Inoue, A. Yamada, and S. S. Gao (2006), Deep structure and origin of the Baikal rift zone, *Earth and Planetary Science Letters*, *243*(3), 681–691.

Zumbo, V., G. Féraud, H. Bertrand, and G. Chazot (1995), $^{40}\text{Ar}/^{39}\text{Ar}$ chronology of Tertiary magmatic activity in Southern Yemen during the early Red Sea-Aden Rifting, *Journal of volcanology and geothermal research*, *65*(3-4), 265–279.

FIGURES

Figure 1. Geodynamic map of Arabia. Yellow triangles are for YOCMAL Network stations in southwestern Yemen. The magmatism older than 20 Ma is represented in purple, whereas the younger magmatism is in pink (modified from *Davison et al.* [1994]). J.: Jordan, UAE: United Arab Emirates, SS.: Shukra el Sheik Fracture Zone.

Figure 2. (a) Topographic map of Southwestern Yemen. The Yemeni highlands, above 1000 m high, are mainly constituted by basaltic trapps. These basalts are 3 km thick. The volcanic Pliocene to present day volcanic fields of Sanaa and Dhamar are represented in red. Jabal Hufash and Jabal Bura Tertiary batholiths are in blue (modified from *Davison et al.* [1994]). Black dots are for YOCMAL seismological stations. (b) Geological sketch of the northern part of our study area. Red stars are for Pliocene-Quaternary volcanoes. The batholithes are located along the Great Escarpment, which runs parallel to the Red Sea margin and the Tihama Plain. Left of the Great Escarpment, altitudes are below 200 m.

Figure 3. Azimuthal distribution of the events used on our study.

Figure 4. Final P-wave velocity model obtained from inversion. Seismic stations are located with the black triangles. Note the color bar change between the first two layers and the other slices of the model, and the scale is saturated for the first layer.

Figure 5. Cross sections in the final P-wave velocity model obtained from inversion. (a) Cross section along the Red Sea margin, (b) Cross section along the Gulf of Aden margin, (c) Cross section through the granitic intrusions of Jabal Hufash and Jabal Bura, (d) Southwest-Northeast cross section of the corner of Yemen.

Figure 6. Crustal correction applied for the stations. These corrections were computed from the crustal thickness obtained by receiver function analysis [*Ahmed et al.*, 2013].

Figure 7. Checkerboard test for the inversion of seismological data. (a) Synthetic input model for P-wave velocity, (b) depth slices at 70, 200 and 300 km through the retrieved velocity model, piercing points: impact points of the rays with the layer located at 300 km depth (c) North-South cross section in the retrieved velocity model.

Figure 8. Synthetic test for the propagation of crustal signal. (a) Synthetic output model, (b) depth slices through the retrieved velocity model. (c) Cross sections of the Aden and Red Sea margins in the synthetic model.

Figure 9. Synthetic test for two low velocity anomalies under the Southwestern Yemen. (a) SW-NE cross section in the input model. (b) SW-NE cross section in the output model. (c) SW-NE cross-section of the corner of Yemen from our final P-wave velocity model.

Figure 10. Plot of P-wave versus S-wave relative arrival-time residuals for all the stations and events coming from Northeast (Japan, China, Russia). The solid red line is the least square fit to our data (with the slope of the line), and the dashed line is a slope of 2.9 (thermal effect only). Standard deviation is 0.31 for the best-fit line and 0.36 for the 2.9 gradient.

Figure 11. Auxiliary material. Series of synthetic tests for constraining the low velocity anomalies under the Southwestern Yemen. (a) Location of the cross section. (b) SW-NE cross-section in the final P-wave tomography model presented Fig 5d of the paper (c to j) SW-NE cross-sections of a series of tests with low-velocity anomalies of distinct widths and depths.