A short proof of the zero-two law for cosine functions

Jean Esterle

► To cite this version:

Jean Esterle. A short proof of the zero-two law for cosine functions. Archiv der Mathematik, 2015, 105 (4), pp.381-387. hal-01147799

HAL Id: hal-01147799

<https://hal.science/hal-01147799>

Submitted on 1 May 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A short proof of the zero-two law for cosine functions

Jean Esterle

Abstract : Let $(C(t))_{t \in \mathbb{R}}$ be a cosine function in a unital Banach algebra. We give a simple proof of the fact that if $\limsup_{t \rightarrow 0} \|C(t) - 1_A\| < 2$, then $\limsup_{t \rightarrow 0} \|C(t) - 1_A\| = 0$.

Keywords : Cosine function, scalar cosine function, commutative local Banach algebra.

AMS classification : Primary 46J45, 47D09, Secondary 26A99

1 Introduction

Recall that a cosine function taking values in a unital normed algebra A with unit element 1_A is a family $C = (C(t))_{t \in \mathbb{R}}$ of elements of A satisfying the so-called d'Alembert equation

$$C(0) = 1_A, C(s+t) + C(s-t) = 2C(s)C(t) \quad (s \in \mathbb{R}, t \in \mathbb{R}). \quad (1)$$

One can define in a similar way cosine sequences $(C(n))_{n \in \mathbb{Z}}$. A cosine sequence depends only on the value of $C(1)$, since we have, for $n \geq 1$,

$$C(-n) = C(n) = T_n(C(1)),$$

where $T_n(x) = \sum_{k=0}^{[n/2]} C_n^k x^{n-2k} (x^2 - 1)^k$ is the n^{th} -Tchebyshev polynomial.

Strongly continuous operator valued cosine functions play an important role in the study of abstract nonlinear second order differential equations, see for example [11]. In a paper to appear in the Journal of Evolution Equations [9], Schwenninger and Zwart showed that if a strongly continuous cosine family $(C(t))_{t \in \mathbb{R}}$ of bounded operators on a Banach space X satisfies $\limsup_{t \rightarrow 0} \|C(t) - I_X\| < 2$, then the generator a of this cosine function is a bounded operator, so that $\limsup_{t \rightarrow 0} \|C(t) - I_X\| = \limsup_{t \rightarrow 0} \|\cos(ta)I_X - I_X\| = 0$, and they asked whether a similar zero-two law holds for general cosine functions $(C(t))_{t \in \mathbb{R}}$. This

question was answered positively by Chojnacki in [5]. Using a sophisticated argument based on ultrapowers, Chonajcki deduced this zero-two law from the fact that if a cosine sequence $C(t)$ satisfies $\sup_{t \in \mathbb{R}} \|C(t) - 1_A\| < 2$, then $C(t) = 1_A$ for $t \in \mathbb{R}$. This second result, which was obtained independently by the author in [7], is proved by Chojnacki in [5] by adapting methods used by Bobrowski, Chojnacki and Gregosiewicz in [3] to show that if a cosine sequence $(C(t))_{t \in \mathbb{R}}$ satisfies $\sup_{t \in \mathbb{R}} \|C(t) - \cos(at)1_A\| < \frac{8}{3\sqrt{3}}$ for some $a \in \mathbb{R}$, then $C(t) = \cos(at)1_A$ for $t \in \mathbb{R}$, a result also obtained independently by the author in [7], which improves previous results of [2], [4] and [10].

The purpose of this paper is to give a short direct proof of the zero-two law. The zero-two law for complex-valued cosine functions is a folklore result, which easily implies that if $\limsup_{t \rightarrow 0} \rho(C(t) - 1_A) < 2$ then $\limsup_{t \rightarrow 0} \rho(C(t) - 1_A) = 0$, where $\rho(x)$ denotes the spectral radius of an element x of a Banach algebra, see section 2. Our proof of the zero-two law is then based on the fact that if $\|C(t) - 1_A\| \leq 2$, and if $\rho\left(C\left(\frac{t}{2}\right) - 1_A\right) < 1$, then we have

$$C\left(\frac{t}{2}\right) = \sqrt{1_A - \frac{C(t) - 1_A}{2}},$$

where $\sqrt{1_A - u}$ is defined by the usual series for $\|u\| \leq 1$. It follows from this identity and from the fact that the coefficients of the Taylor series at the origin of the function $t \rightarrow 1 - \sqrt{1 - t}$ are positive that in this situation we have

$$\left\| C\left(\frac{t}{2}\right) - 1_A \right\| \leq 1 - \sqrt{1 - \left\| \frac{C(t) - 1_A}{2} \right\|},$$

and the zero-two law follows.

Notice that if we replace the constant 2 by $\frac{3}{2}$ a "three line argument" due to Arendt [1] shows that if $\limsup_{t \rightarrow 0} \|C(t) - 1_A\| < \frac{3}{2}$ then $\limsup_{t \rightarrow 0} \|C(t) - 1_A\| = 0$. The proof presented here has some analogy with Arendt's proof, and the difficulty to estimate $\|(1_A + C(\frac{t}{2}))^{-1}\|$ is circumvented in the present paper by using the formula above.

The author wishes to thank W. Chonajcki for giving information about the reference [5]. He also thanks F. Schwenninger for valuable discussions about the content of the paper.

2 The zero-two law for the spectral radius

The zero-two law for scalar cosine functions pertains to folklore, but we could not find a reference in the literature for the following certainly well-known lemma, which is a variant of proposition 3.1 of [7].

Lemma 2.1. *Let $c = (c(t))_{t \in \mathbb{R}}$ be a complex-valued cosine function. Then c satisfies one of the following conditions*

- (i) $\limsup_{t \rightarrow 0} |c(t) - 1| = +\infty$,
- (ii) $\limsup_{t \rightarrow 0} |c(t) - 1| = 2$,
- (iii) $\limsup_{t \rightarrow 0} |c(t) - 1| = 0$.

First assume that $M := \limsup_{t \rightarrow 0} |c(t)| < +\infty$, and denote by S the set of all complex numbers α for which there exists a sequence $(t_m)_{m \geq 1}$ of positive real numbers such that $\lim_{m \rightarrow +\infty} t_m = 0$ and $\lim_{m \rightarrow +\infty} c(t_m) = \alpha$. Then $|\alpha| \leq M$ for every $\alpha \in S$. Notice that if $\alpha \in S$, and if a sequence $(t_m)_{m \in \mathbb{Z}}$ satisfies the above conditions with respect to α , then $T_n(\alpha) = \lim_{m \rightarrow +\infty} T_n(C(t_m)) = \lim_{m \rightarrow +\infty} C(nt_m)$, and so $T_n(\alpha) \in S$, and $|T_n(\alpha)| \leq M$ for $n \geq 1$. Now write $\alpha = \cos(z) = \sum_{k=0}^{+\infty} \frac{z^{2k}}{(2k)!}$, and set $u = \operatorname{Re}(z)$, $v = \operatorname{Im}(z)$. We have, for $n \geq 1$,

$$T_n(\alpha) = \cos(nz) = \frac{e^{inu} e^{-nv} + e^{-inu} e^{nv}}{2}.$$

Since $\sup_{n \geq 1} |T_n(\alpha)| \leq M$, we have $v = 0$, $S \subset [-1, 1]$, and $\limsup_{t \rightarrow 0} |c(t) - 1| \leq 2$.

Now assume that $S \neq \{1\}$, and let $\alpha \in S \setminus \{1\}$. We have $\alpha = \cos(u)$ for some $u \in \mathbb{R}$. We see as above that $\cos(nu) \in S$ for every $n \geq 1$. If u/π is irrational, then the set $\{e^{inu}\}_n \geq 1$ is dense in the unit circle $\mathbb{T}$, and so $S = [-1, 1]$ since S is closed, and in this situation $\limsup_{t \rightarrow 0} |c(t) - 1| = 2$.

Now assume that u/π is rational, and let $s \geq 2$ be the smallest positive integer such that $e^{ius} = 1$. Then $e^{\frac{2i\pi}{s}} = e^{ipu}$ for some positive integer p , and so $\cos(\frac{2\pi}{s}) \in S$. Let $(t_m)_{m \geq 1}$ be a sequence of positive reals such that $\lim_{m \rightarrow +\infty} t_m = 0$ and $\lim_{m \rightarrow +\infty} c(t_m) = \cos(\frac{2\pi}{s})$, let $q \geq 2$, and let β be a limit point of the sequence $c(\frac{t_m}{s^{q-1}})_{m \geq 1}$. There exists $y \in \mathbb{R}$ such that $\cos(y) = \beta$ and such that $s^{q-1}y = \frac{2\pi}{s} + 2k\pi$, with $k \in \mathbb{Z}$. Then $y = (1 + ks)\frac{2\pi}{s^q}$. Since $\gcd(1 + ks, s^q) = 1$, there exists a positive integer r such that $ry - \frac{2\pi}{s^q} \in 2\pi\mathbb{Z}$, so that $\cos(\frac{2\pi}{s^q}) \in S$. This implies that $\cos(\frac{2p\pi}{s^q}) \in S$ for $p \geq 1, q \geq 1$, and S is dense in $[-1, 1]$. Since S is closed, we obtain again $S = [-1, 1]$, which implies that $\limsup_{t \rightarrow 0} |c(t) - 1| = 2$. So if neither (i) nor (ii) holds, we have $S = \{1\}$, which implies (iii). $\square$

Notice that if a cosine function $C = (C(t))_{t \in \mathbb{R}}$ in a Banach algebra A satisfies $\sup_{|t| \leq \eta} \|C(t)\| \leq M < +\infty$ for some $\eta > 0$, then $\sup_{|t| \leq L} \|C(t)\| < +\infty$ for every $L > 0$, since $\sup_{|t| \leq n\eta} \|C(t)\| \leq \sup_{\|y\| \leq M} \|T_n(y)\|$ for every $n \geq 1$, where T_n denotes the n -th Tchebyshev polynomial. In particular if a complex-valued cosine function $c = (c(t))_{t \in \mathbb{R}}$ satisfies (iii), then the identity

$$(1 - c(s - t))(1 - c(s + t)) = (c(s) - c(t))^2$$

shows as is well-known that the cosine function c is continuous on $\mathbb{R}$, which implies that $c(t) = \cos(ta)$ for some $a \in \mathbb{C}$.

If A is commutative and unital, we will denote 1_A the unit element of A , and we will denote by $\hat{A}$ the space of all characters on A , equipped with the Gelfand topology, i.e. the compact topology induced by the weak* topology on the unit ball of the dual space of A .

Proposition 2.2. *Let $C = (C(t))$ be a cosine function in a unital Banach algebra A . If $\limsup_{t \rightarrow 0} \rho(C(t) - 1_A) < 2$, then $\limsup_{t \rightarrow 0} \rho(C(t) - 1_A) = 0$.*

Proof: We may assume that unital Banach algebra A is generated by $(C(t))_{t \in \mathbb{R}}$. Let $\chi \in \hat{A}$. Then the cosine complex-valued function $(\chi(C(t)))_{t \in \mathbb{R}}$ satisfies condition (iii) of the lemma, and so there exists $a_\chi \in \mathbb{C}$ such that we have

$$\chi(C(t)) = \cos(ta_\chi) \quad (t \in \mathbb{R}).$$

Set $u_\chi = \operatorname{Re}(a_\chi)$, $v_\chi = \operatorname{Im}(a_\chi)$. We have

$$\rho(C(t) - 1) \geq |1 - \cos(tu_\chi) \operatorname{ch}(tv_\chi)|.$$

If the family $(u_\chi)_{\chi \in \hat{A}}$ were unbounded, there would exist a sequence $(t_n)_{n \geq 1}$ of real numbers converging to zero and a sequence $(\chi_n)_{n \geq 1}$ of characters of A such that $\cos(t_n u_{\chi_n}) = -1$, and we would have $\rho(C(t_n) - 1) \geq 2$ for $n \geq 1$. So the family $(u_\chi)_{\chi \in \hat{A}}$ is bounded. If the family $(v_\chi)_{\chi \in \hat{A}}$ were unbounded, there would exist a sequence $(t'_n)_{n \geq 1}$ of real numbers converging to zero and a sequence $(\chi'_n)_{n \geq 1}$ of characters of A such that $\lim_{n \rightarrow +\infty} \operatorname{ch}(t'_n v_{\chi'_n}) = +\infty$. But this would imply that $\limsup_{t \rightarrow 0} \rho(C(t)) = +\infty$. Hence the family $(a_\chi)_{\chi \in \hat{A}}$ is bounded, and we have

$$\limsup_{t \rightarrow 0} \rho(C(t) - 1_A) = \lim_{t \rightarrow 0} \sup_{\chi \in \hat{A}} |\cos(ta_\chi) - 1| = 0.$$

□

3 The zero-two law for cosine functions

Set $\alpha_n = \frac{1}{n!} \frac{1}{2} \left(\frac{1}{2} - 1\right) \dots \left(\frac{1}{2} - n + 1\right)$ for $n \geq 1$, with the convention $\alpha_0 = 0$, and for $|z| < 1$, set

$$\sqrt{1-z} = \sum_{n=0}^{+\infty} (-1)^n \alpha_n z^n,$$

so that $(\sqrt{1-z})^2 = 1-z$, and $\sqrt{1-t}$ is the positive square root of $1-t$ for $t \in (-1, 1)$. Also $\operatorname{Re}(\sqrt{1-z}) > 0$ for $|z| < 1$.

We have, for $t \in [0, 1)$,

$$\sum_{n=1}^{+\infty} (-1)^{n-1} \alpha_n t^n = 1 - \sqrt{1-t}.$$

Since $(-1)^{n-1} \alpha_n \geq 0$ for $n \geq 1$, the series $\sum_{n=1}^{+\infty} |\alpha_n| = \sum_{n=1}^{+\infty} (-1)^{n-1} \alpha_n$ is convergent, and we have

$$\sum_{n=1}^{+\infty} |\alpha_n| t^n = 1 - \sqrt{1-t} \quad (0 \leq t \leq 1).$$

Now let A be a commutative unital Banach algebra, and let $x \in A$ such that $\|x\| \leq 1$. Set

$$\sqrt{1_A - x} = \sum_{n=0}^{+\infty} (-1)^n \alpha_n x^n.$$

Then $(\sqrt{1_A - x})^2 = 1_A - x$, and we have

$$\left\| 1_A - \sqrt{1_A - x} \right\| = \left\| \sum_{n=1}^{+\infty} (-1)^n \alpha_n x^n \right\| \leq \sum_{n=1}^{+\infty} |\alpha_n| \|x\|^n = 1 - \sqrt{1 - \|x\|} \quad (2)$$

Notice also that if A is commutative, then we have

$$\operatorname{Re} \left(\chi \left(\sqrt{1_A - x} \right) \right) = \operatorname{Re} \left(\sqrt{1 - \chi(x)} \right) \geq 0 \quad (\chi \in \hat{A}). \quad (3)$$

We obtain the following formula

Lemma 3.1. *Let $(C(t))_{t \in \mathbb{R}}$ be a cosine function in a unital Banach algebra A . Assume that $\|C(t) - 1_A\| \leq 2$ and that $\rho((C(\frac{t}{2}) - 1) < 1$. Then we have*

$$C\left(\frac{t}{2}\right) = \sqrt{1_A - \frac{1_A - C(t)}{2}}.$$

Proof : The abstract version of the formula $\sin^2\left(\frac{u}{2}\right) = \frac{1 - \cos(u)}{2}$ gives

$$1_A - C\left(\frac{t}{2}\right)^2 = \frac{1_A - C(t)}{2}, \quad C\left(\frac{t}{2}\right)^2 = 1_A - \frac{1_A - C(t)}{2} = \left(\sqrt{1_A - \frac{1_A - C(t)}{2}} \right)^2,$$

$$\left(C\left(\frac{t}{2}\right) - \sqrt{1_A - \frac{1_A - C(t)}{2}} \right) \left(C\left(\frac{t}{2}\right) + \sqrt{1_A - \frac{1_A - C(t)}{2}} \right) = 0.$$

We may assume that A is commutative. Let $\chi \in \hat{A}$. Since $\rho\left(C\left(\frac{t}{2}\right) - 1_A\right) < 1$, we have $\operatorname{Re}\left(\chi\left(C\left(\frac{t}{2}\right)\right)\right) > 0$. Since $\operatorname{Re}\left(\chi\left(\sqrt{1_A - \frac{1_A - C(t)}{2}}\right)\right) \geq 0$, $C\left(\frac{t}{2}\right) + \sqrt{1_A - \frac{1_A - C(t)}{2}}$ is invertible in A , and $C\left(\frac{t}{2}\right) - \sqrt{1_A - \frac{1_A - C(t)}{2}} = 0$. $\square$

Theorem 3.2. *Let $(C(t))_{t \in \mathbb{R}}$ be a cosine sequence in a Banach algebra. If $\limsup_{t \rightarrow 0} \|C(t) - 1_A\| < 2$, then $\limsup_{t \rightarrow 0} \|C(t) - 1_A\| = 0$.*

Proof : It follows from proposition 2.2 and lemma 3.1 that there exists $\eta > 0$ such that we have, for $|t| \leq \eta$,

$$\|C(t) - 1_A\| < 2, C\left(\frac{t}{2}\right) = \sqrt{1_A - \frac{1_A - C(t)}{2}}.$$

Using (1), we obtain, for $|t| \leq \eta$,

$$\left\|C\left(\frac{t}{2}\right) - 1_A\right\| \leq 1 - \sqrt{1 - \left\|\frac{C(t) - 1_A}{2}\right\|}.$$

Set $l = \limsup_{t \rightarrow 0} \|C(t) - 1_A\|$. We obtain

$$l \leq 1 - \sqrt{1 - \frac{l}{2}} \leq 1,$$

and so $l = 0$. $\square$

Notice that the proof above gives a little bit more than the zero-two law : if $\|1 - C(t)\| \leq 2$ and $\rho\left(1 - C\left(\frac{t}{2}\right)\right) < 1$ for $|t| \leq \eta$, then we have, for $n \geq 1$,

$$\sup_{|t| \leq 2^{-n}\eta} \|C(t) - 1_A\| \leq u_n,$$

where the sequence u_n satisfies $u_0 = 2, u_{n+1} = 1 - \sqrt{1 - \frac{u_n}{2}}$ for $n \geq 1$, and $\lim_{n \rightarrow +\infty} u_n = 0$, which gives an explicit control on the convergence to 0 of $\|C(t) - 1_A\|$ as $t \rightarrow 0$.

Notice also that the fact that the coefficients of the Taylor expansion at the origin of the function $t \rightarrow 1 - \sqrt{1 - t}$ are positive was used in [6] to show that $\|x^2 - x\| \geq 1/4$ for every quasinilpotent element x of a Banach algebra such that $|x| \geq 1/2$. Similar argument were used in [8] to show that if a semigroup $(T(t))_{t \geq 0}$ in a Banach algebra A satisfies $\limsup_{t \rightarrow 0^+} \|T(t) - T((n+1)t)\| < \frac{n}{(n+1)^{1+\frac{1}{n}}}$ for some $n \geq 1$, then there exists an idempotent J of A such that $\lim_{t \rightarrow 0} \|T(t) - J\| = 0$, so that $\limsup_{t \rightarrow 0^+} \|T(t) - T((n+1)t)\| = 0$.

Références

- [1] W. Arendt, *A $0-3/2$ - Law for Cosine Functions*. Ulmer Seminare, Funktionalanalysis und Evolutionsgleichungen, 17 (2012), 349-350.
- [2] A. Bobrowski and W. Chojnacki, *Isolated points of some sets of bounded cosine families, bounded semigroups, and bounded groups on a Banach space*, Studia Math. 217 (2013), 219-241.
- [3] A. Bobrowski, W. Chojnacki, and A. Gregosiewicz, *On close-to-scalar one-parameter cosine families*, J. Math. Anal. Appl. (2015), <http://dx.doi.org/10.1016/j.jmaa.2015.04.012>.
- [4] W. Chojnacki, *On cosine families close to scalar cosine families*, to appear in J. Aust. Math. Soc., arXiv:1411.0854.
- [5] W. Chojnacki, *Around Schwenninger and Zwart's zero-two law for cosine families*, submitted.
- [6] J. Esterle, *Quasimultipliers, representations of H^∞ , and the closed ideal problem for commutative Banach algebras*, Springer Lect. Notes 975 (1983), 66-162.
- [7] J. Esterle, *Bounded cosine functions close to continuous bounded scalar cosine functions*, submitted. <https://hal.archives-ouvertes.fr/hal-01111839>
- [8] J. Esterle and A. Mokhtari, *Distance entre éléments d'un semi-groupe dans une algèbre de Banach*, J. Func. An 195 (2002), 167-189.
- [9] F. Schwenninger and H. Zwart, *Zero-two laws for cosine families*, J. Evolution Eq., to appear, arXiv:1402.1304v3.
- [10] F. Schwenninger and H. Zwart, *Less than one, implies 0*, submitted.
- [11] C. Travis and G. Webb, *Cosine families and abstract nonlinear second order differential equation*, Acta Math. Acad. Sci. Hungar 32(1978), 75-96.

Jean Esterle
 IMB, UMR 5251
 Université de Bordeaux
 351, cours de la Libération
 33405 - Talence (France)
esterle@math.u-bordeaux1.fr