



HAL
open science

Ideotype construction from an architectural model of pea

Romain Barillot, Didier Combes, Pierre Huynh, Abraham Escobar Gutiérrez

► **To cite this version:**

Romain Barillot, Didier Combes, Pierre Huynh, Abraham Escobar Gutiérrez. Ideotype construction from an architectural model of pea. ECA international Conference, Jul 2012, Rennes, France. 64 p., 2012, Plant and canopy architecture impact on disease epidemiology and pest development. hal-01187874

HAL Id: hal-01187874

<https://hal.science/hal-01187874>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

IDEOTYPE CONSTRUCTION FROM AN ARCHITECTURAL MODEL OF PEA.

Romain Barillot¹, D. Combes², P. Huynh¹ and A.J. Escobar-Gutiérrez².

¹LUNAM Université, Groupe Ecole Supérieure d'Agriculture, UPSP Laboratoire d'Ecophysiologie Végétale & Agroécologie, BP 30748, F-49007 Angers cedex 01, France.

²INRA, UR4 P3F, Equipe Ecophysiologie des plantes fourragères, Le Chêne - RD 150, BP 6, F-86600 Lusignan, France.

There is an increasing need for growing legume species in Europe in order to reduce dependency on imported vegetable proteins. Currently, pea (*Pisum sativum*) is the principle source of vegetable proteins. However, the productivity rate of pea crop is still under its potential mainly due to fungal foliar diseases. The most damaging ones are caused by *Ascochyta pisi*, *Mycosphaerella pinodes* and *Phoma medicaginis*.

Dynamics of such diseases have been shown to depend on plant architecture *e.g.* Leaf Area Index (LAI), plant height, internode length (Le May *et al.*, 2009). Virtual models of canopy architecture therefore appear as suitable tools for studying and modelling plant–pathogen interactions and diseases dispersal.

We have developed a 3D architectural model of pea growth based on the L-system formalism and developed on the L-Py platform (Boudon *et al.*, 2010). Initial parameters were based on data from a field-grown crop of winter pea *cv* Lucy. The above ground architecture is represented as a succession of phytomers emitted by main stems and branches. Phytomers are considered as a collection of organs encoded as modules that support their state *i.e.* age, length, topology and geometry. The rate of phytomer emission was set for main stems and branches. The number of branches and their time of emergence are attributes of each node of stems.

Coupled with epidemiologic models, this simulator can be used as a conceptual framework for studying the effects of pea architectural parameters on disease development and dispersal. Indeed, the present model is able to build contrasted canopies of ideotypes with regard to LAI and its spatial distribution (plant density, phytomer number, branching ability, axis orientation, leaf growth), plant height (internode growth), foliage orientation (leaf geometry) and dynamics of growth (rate of phytomers and branch emission, organ growth kinetics). Moreover, the formalism chosen in the present study also allows to assess the benefits of intercropping systems, comprising pea-wheat mixtures, towards disease pressure such as the wheat/*Septoria tritici* pathosystem (Robert *et al.*, 2008).

References:

- Boudon F, Cokelaer T, Pradal C, Godin C. (2010). L-Py, an open L-systems framework in Python. In: Dejong T, Da Silva D eds. *6th International Workshop on Functional-Structural Plant Models*. Davis, CA.
- Le May C, Ney B, Lemarchand E, Schoeny A, Tivoli B. (2009). Effect of pea plant architecture on spatiotemporal epidemic development of ascochyta blight (*Mycosphaerella pinodes*) in the field. *Plant Pathology*, **58**: 332-343.
- Robert C, Fournier C, Andrieu B, Ney B. (2008). Coupling a 3D virtual wheat (*Triticum aestivum*) plant model with a *Septoria tritici* epidemic model (Septo3D): a new approach to investigate plant–pathogen interactions linked to canopy architecture. *Functional Plant Biology*, **35**: 997-1013.

Keywords:

Architectural model, Ideotype, L-System, LPy, *Pisum sativum*