

Computing the Chow variety of quadratic space curves

Peter Bürgisser, Kathlén Kohn, Pierre Lairez, Bernd Sturmfels

► **To cite this version:**

Peter Bürgisser, Kathlén Kohn, Pierre Lairez, Bernd Sturmfels. Computing the Chow variety of quadratic space curves. Sixth International Conference on Mathematical Aspects of Computer and Information Sciences (MACIS 2015), Nov 2015, Berlin, Germany. hal-01188213v2

HAL Id: hal-01188213

<https://hal.archives-ouvertes.fr/hal-01188213v2>

Submitted on 27 Nov 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Computing the Chow Variety of Quadratic Space Curves

Peter Bürgisser^{1*}, Kathlén Kohn^{1**}, Pierre Lairez^{1*}, and Bernd Sturmfels^{1,2**}

¹ Institute of Mathematics, Technische Universität Berlin, Germany

² Department of Mathematics, University of California, Berkeley, USA

Abstract. Quadrics in the Grassmannian of lines in 3-space form a 19-dimensional projective space. We study the subvariety of coisotropic hypersurfaces. Following Gel'fand, Kapranov and Zelevinsky, it decomposes into Chow forms of plane conics, Chow forms of pairs of lines, and Hurwitz forms of quadric surfaces. We compute the ideals of these loci.

Keywords: Chow variety, coisotropic hypersurface, Grassmannian, space curve, computation

Introduction

The Chow variety, introduced in 1937 by Chow and van der Waerden [4], parameterizes algebraic cycles of any fixed dimension and degree in a projective space, each given by its Chow form. The case of curves in \mathbb{P}^3 goes back to an 1848 paper by Cayley [3]. A fundamental problem, addressed by Green and Morrison [8] as well as Gel'fand, Kapranov and Zelevinsky [6, §4.3], is to describe the equations defining Chow varieties. We present a definitive computational solution for the smallest non-trivial case, namely for cycles of dimension 1 and degree 2 in \mathbb{P}^3 .

The *Chow form* of a cycle of degree 2 is a quadratic form in the Plücker coordinates of the Grassmannian $G(2, 4)$ of lines in \mathbb{P}^3 . Such a quadric in $G(2, 4)$ represents the set of all lines that intersect the given cycle. Quadratic forms in Plücker coordinates form a projective space \mathbb{P}^{19} . The Chow variety we are interested in, denoted $G(2, 2, 4)$, is the set of all Chow forms in that \mathbb{P}^{19} . The aim of this note is to make the concepts in [3, 4, 8] and [6, §4.3] completely explicit.

We start with the 9-dimensional subvariety of \mathbb{P}^{19} whose points are the *coisotropic quadrics* in $G(2, 4)$. By [6, §4.3, Theorem 3.14], this decomposes as the Chow variety and the variety of Hurwitz forms [9], representing lines that are tangent to a quadric surface in \mathbb{P}^3 . Section 1 studies the ideal generated by the coisotropy conditions. We work in a polynomial ring in 20 variables, one for each quadratic Plücker monomial on $G(2, 4)$ minus one for the Plücker relation. We derive the coisotropic ideal from the differential characterization of coisotropy. Proposition 1 exhibits the decomposition of this ideal into three minimal primes. In particular, this shows that the coisotropic ideal is radical, and it hence resolves

* This author was partially supported by DFG grant BU 1371/2-2.

** This author was supported by a Fellowship from the Einstein Foundation Berlin.

the degree 2 case of a problem posed in 1986 by Green and Morrison [8]. They wrote: ‘We do not know whether [the differential characterization of coisotropy] generates the full ideal of these Chow variables.’

Section 2 derives the radical ideal of the Chow variety $G(2, 2, 4)$ in \mathbb{P}^{19} . Its two minimal primes represent Chow forms of plane conics and Chow forms of pairs of lines. We also study the characterization of Chow forms among all coisotropic quadrics by the vanishing of certain differential forms. These represent the integrability of the α -distribution in [6, §4.3, Theorem 3.22]. After saturation by the irrelevant ideal, the integrability ideal is found to be radical.

1 Coisotropic Quadrics

The Grassmannian $G(2, 4)$ is a quadric in \mathbb{P}^5 . Its points are lines in \mathbb{P}^3 . We represent these lines using dual Plücker coordinates $\mathbf{p} = (p_{01}, p_{02}, p_{03}, p_{12}, p_{13}, p_{23})$ subject to the Plücker relation $p_{01}p_{23} - p_{02}p_{13} + p_{03}p_{12}$. Following [9, §2], by *dual* coordinates we mean that p_{ij} is the ij -minor of a 2×4 -matrix whose rows span the line. The generic quadric in $G(2, 4)$ is written as a generic quadratic form

$$Q(\mathbf{p}) = \mathbf{p} \cdot \begin{pmatrix} c_0 & c_1 & c_2 & c_3 & c_4 & c_5 \\ c_1 & c_6 & c_7 & c_8 & c_9 & c_{10} \\ c_2 & c_7 & c_{11} & c_{12} & c_{13} & c_{14} \\ c_3 & c_8 & c_{12} & c_{15} & c_{16} & c_{17} \\ c_4 & c_9 & c_{13} & c_{16} & c_{18} & c_{19} \\ c_5 & c_{10} & c_{14} & c_{17} & c_{19} & c_{20} \end{pmatrix} \cdot \mathbf{p}^T. \quad (1)$$

The quadric $Q(\mathbf{p})$ is an element in $V := \mathbb{C}[\mathbf{p}]_2 / \mathbb{C}\{p_{01}p_{23} - p_{02}p_{13} + p_{03}p_{12}\} \simeq \mathbb{C}^{21} / \mathbb{C}$. Hence, $\mathbf{c} = (c_0, c_1, \dots, c_{20})$ serves as homogeneous coordinates on $\mathbb{P}^{19} = \mathbb{P}(V)$, which – due to the Plücker relation – need to be understood modulo

$$c_5 \mapsto c_5 + \lambda, \quad c_9 \mapsto c_9 - \lambda, \quad c_{12} \mapsto c_{12} + \lambda. \quad (2)$$

The coordinate ring $\mathbb{Q}[V]$ is a subring of $\mathbb{Q}[c_0, c_1, \dots, c_{20}]$, namely it is the invariant ring of the additive group action (2). Hence $\mathbb{Q}[V]$ is the polynomial ring in 20 variables $c_0, c_1, c_2, c_3, c_4, c_5 - c_{12}, c_6, c_7, c_8, c_9 + c_{12}, c_{10}, c_{11}, c_{13}, \dots, c_{20}$.

We are interested in the \mathbf{c} 's that lead to *coisotropic* hypersurfaces of $G(2, 4)$. For these, the tangent space at any point ℓ , considered as a subspace of $T_\ell G(2, 4) = \text{Hom}(\ell, \mathbb{C}^4/\ell)$, has the form $\{\varphi \mid \varphi(a) = 0\} + \{\varphi \mid \text{im}(\varphi) \subset M\}$, for some $a \in \ell \setminus \{0\}$ and some plane M in \mathbb{C}^4/ℓ . By [6, §4.3, (3.24)], the quadric hypersurface $\{Q(\mathbf{p}) = 0\}$ in $G(2, 4)$ is coisotropic if and only if there exist $s, t \in \mathbb{C}$ such that

$$\frac{\partial Q}{\partial p_{01}} \cdot \frac{\partial Q}{\partial p_{23}} - \frac{\partial Q}{\partial p_{02}} \cdot \frac{\partial Q}{\partial p_{13}} + \frac{\partial Q}{\partial p_{03}} \cdot \frac{\partial Q}{\partial p_{12}} = s \cdot Q + t \cdot (p_{01}p_{23} - p_{02}p_{13} + p_{03}p_{12}). \quad (3)$$

Equivalently, the vector $(t, s, -1)^T$ is in the kernel of the 21×3 matrix in Figure 1. The 3×3 minors of this matrix are all in the subring $\mathbb{Q}[V]$. The *coisotropic ideal* I is the ideal of $\mathbb{Q}[V]$ generated by these minors. The subscheme $V(I)$ of $\mathbb{P}^{19} =$

each expressing the relevant geometric condition. This construction ensures that the ideals are prime. We then verified that their intersection equals I . For details, check our computations, using the link given at the end of this article.

From the geometric perspective of [6], the third prime P_{Squares} is extraneous, because nonreduced hypersurfaces in $G(2, 4)$ are excluded by Gel'fand, Kapranov and Zelevinsky. Theorem 3.14 in [6, §4.3] concerns irreducible hypersurfaces, and the identification of Chow forms within the coisotropic hypersurfaces [6, §4.3, Theorem 3.22] assumes the corresponding polynomial to be squarefree. With this, the following would be the correct ideal for the coisotropic variety in \mathbb{P}^{19} :

$$P_{\text{Hurwitz}} \cap P_{\text{ChowLines}} = (I : P_{\text{Squares}}) . \quad (5)$$

This means that the reduced coisotropic quadrics in $G(2, 4)$ are either Chow forms of curves or Hurwitz forms of surfaces. The ideal in (5) has codimension 10, degree 92, and is minimally generated by 175 cubics and 20 quartics in $\mathbb{Q}[V]$.

A slightly different point of view on the coisotropic ideal is presented in a recent paper of Catanese [2]. He derives a variety in $\mathbb{P}^{20} = \mathbb{P}(\mathbb{C}[\mathbf{p}]_2)$ which projects isomorphically onto our variety $V(I) \subset \mathbb{P}^{19}$. The center of projection is the Plücker quadric. To be precise, Proposition 4.1 in [2] states the following: For every $Q \in \mathbb{C}[\mathbf{p}]_2 \setminus \mathbb{C}(p_{01}p_{23} - p_{02}p_{13} + p_{03}p_{12})$ satisfying (3) there is a unique $\lambda \in \mathbb{C}$ such that the quadric $Q_\lambda := Q + \lambda \cdot (p_{01}p_{23} - p_{02}p_{13} + p_{03}p_{12})$ satisfies

$$\frac{\partial Q_\lambda}{\partial p_{01}} \cdot \frac{\partial Q_\lambda}{\partial p_{23}} - \frac{\partial Q_\lambda}{\partial p_{02}} \cdot \frac{\partial Q_\lambda}{\partial p_{13}} + \frac{\partial Q_\lambda}{\partial p_{03}} \cdot \frac{\partial Q_\lambda}{\partial p_{12}} = t \cdot (p_{01}p_{23} - p_{02}p_{13} + p_{03}p_{12}) \quad (6)$$

for some $t \in \mathbb{C}$. This implies that $V(I)$ is isomorphic to the variety of all $Q \in \mathbb{P}(\mathbb{C}[\mathbf{p}]_2) \setminus \{p_{01}p_{23} - p_{02}p_{13} + p_{03}p_{12}\}$ satisfying (6). Let I_2 be generated by the 2×2 minors of the 21×2 matrix that is obtained by deleting the middle column of the matrix in Figure 1. Then $V(I_2)$ contains exactly those $Q \in \mathbb{P}(\mathbb{C}[\mathbf{p}]_2)$ satisfying (6), and $V(I)$ is the projection of $V(I_2)$ from the center $(p_{01}p_{23} - p_{02}p_{13} + p_{03}p_{12})$. The ideal I_2 has codimension 11, degree 92, and is minimally generated by 20 quadrics. Interestingly, Catanese shows furthermore in [2, Theorem 3.3] that a hypersurface in $G(2, 4)$ is coisotropic if and only if it is selfdual in \mathbb{P}^5 with respect to the inner product given by the Plücker quadric.

2 The Chow Variety

In this section we study the Chow variety $G(2, 2, 4)$ of one-dimensional algebraic cycles of degree two in \mathbb{P}^3 . By [6, §4.1, Ex. 1.3], the Chow variety $G(2, 2, 4)$ is the union of two irreducible components of dimension eight in \mathbb{P}^{19} , one corresponding to planar quadrics and the other to pairs of lines. Formally, this means that $G(2, 2, 4) = V(P_{\text{ChowConic}}) \cup V(P_{\text{ChowLines}})$, where $P_{\text{ChowConic}}$ is the homogeneous prime ideal in $\mathbb{Q}[V]$ whose variety comprises the Chow forms of irreducible curves of degree two in \mathbb{P}^3 . The ideal $P_{\text{ChowConic}}$ has codimension 11 and degree 92, and it is minimally generated by 21 quadrics and 35 cubics. The radical ideal $P_{\text{ChowConic}} \cap P_{\text{ChowLines}}$ has codimension 11, degree $232 = 92 + 140$, and it is minimally generated by 230 cubics.

Since $G(2, 2, 4)$ should be contained in the coisotropic variety $V(I)$, it seems that $P_{\text{ChowConic}}$ is missing from the decomposition (4). Here is the explanation:

Proposition 2. *Every Chow form of a plane conic in \mathbb{P}^3 is also a Hurwitz form. In symbols, $P_{\text{Hurwitz}} \subset P_{\text{ChowConic}}$ and thus $V(P_{\text{ChowConic}}) \subset V(P_{\text{Hurwitz}})$.*

Our first proof is by computer: just check the inclusion of ideals in Macaulay2. For a conceptual proof, we consider a 4×4 -symmetric matrix $M = M_0 + \epsilon M_1$, where $\text{rank}(M_0) = 1$. By [9, eqn. (1)], the Hurwitz form of the corresponding quadric surface in \mathbb{P}^3 is $Q(\mathbf{p}) = \mathbf{p}(\wedge_2 M)\mathbf{p}^T$. Divide by ϵ and let $\epsilon \rightarrow 0$. The limit is the Chow form of the plane conic defined by restricting M_1 to $\ker(M_0) \simeq \mathbb{P}^2$. This type of degeneration is familiar from the study of complete quadrics [5]. Proposition 2 explains why the locus of irreducible curves is not visible in (4).

Gel'fand, Kapranov and Zelevinsky [6, §4.3] introduce a class of differential forms in order to discriminate Chow forms among all coisotropic hypersurfaces. In their setup, these forms represent the integrability of the α -distribution $\mathcal{E}_{\alpha, Z}$. We shall apply the tools of computational commutative algebra to shed some light on the characterization of Chow forms via integrability of α -distributions.

For this, we use local affine coordinates instead of Plücker coordinates. A point in the Grassmannian $G(2, 4)$ is represented as the row space of the matrix

$$\begin{pmatrix} 1 & 0 & a_2 & a_3 \\ 0 & 1 & b_2 & b_3 \end{pmatrix}. \quad (7)$$

We express the quadrics Q in (1) in terms of the local coordinates a_2, a_3, b_2, b_3 , by substituting the Plücker coordinates with the minors of the matrix (7), i.e.,

$$p_{01} = 1, \quad p_{02} = b_2, \quad p_{03} = b_3, \quad p_{12} = -a_2, \quad p_{13} = -a_3, \quad p_{23} = a_2b_3 - b_2a_3. \quad (8)$$

We consider the following differential 1-forms on affine 4-space:

$$\begin{aligned} \alpha_1^1 &:= \frac{\partial Q}{\partial a_2} da_2 + \frac{\partial Q}{\partial a_3} da_3, & \alpha_2^1 &:= \frac{\partial Q}{\partial a_2} db_2 + \frac{\partial Q}{\partial a_3} db_3, \\ \alpha_1^2 &:= \frac{\partial Q}{\partial b_2} da_2 + \frac{\partial Q}{\partial b_3} da_3, & \alpha_2^2 &:= \frac{\partial Q}{\partial b_2} db_2 + \frac{\partial Q}{\partial b_3} db_3. \end{aligned}$$

By taking wedge products, we derive the 16 differential 4-forms

$$dQ \wedge d\alpha_j^i \wedge \alpha_l^k = q_{ijkl} \cdot da_2 \wedge da_3 \wedge db_2 \wedge db_3 \quad \text{for } i, j, k, l \in \{1, 2\}. \quad (9)$$

Here the expressions q_{ijkl} are certain polynomials in $\mathbb{Q}[V][a_2, a_3, b_2, b_3]$.

Theorems 3.19 and 3.22 in [6, §4.3] state that a squarefree coisotropic quadric Q is a Chow form if and only if all 16 coefficients q_{ijkl} are multiples of Q . By taking normal forms of the polynomials q_{ijkl} modulo the principal ideal $\langle Q \rangle$, we obtain a collection of 720 homogeneous polynomials in \mathfrak{c} . Among these, 58 have degree three, 340 have degree four, and 322 have degree five. The aforementioned result implies that these 720 polynomials cut out $G(2, 2, 4)$ as a subset of \mathbb{P}^{19} .

The *integrability ideal* $J \subset \mathbb{Q}[V]$ is generated by these 720 polynomials and their analogues from other affine charts of the Grassmannian, obtained by permuting columns in (7). We know that $V(J)$ equals the union of $G(2, 2, 4)$ with all double hyperplanes in $G(2, 4)$ (corresponding to P_{Squares}) set-theoretically. Maple, Macaulay2 and Magma verified for us that it holds scheme-theoretically:

Proposition 3. *The integrability ideal J is minimally generated by 210 cubics. Writing \mathfrak{m} for the irrelevant ideal $\langle c_0, c_1, \dots, c_{20} \rangle$ of $\mathbb{Q}[V]$, we have*

$$\sqrt{J} = (J : \mathfrak{m}) = P_{\text{ChowConic}} \cap P_{\text{ChowLines}} \cap P_{\text{Squares}}. \quad (10)$$

Conclusion

We reported on computational experiments with hypersurfaces in the Grassmannian $G(2, 4)$ that are associated to curves and surfaces in \mathbb{P}^3 . For degree 2, all relevant parameter spaces were described by explicit polynomials in 20 variables. All ideals and computations discussed in this note can be obtained at

www3.math.tu-berlin.de/algebra/static/pluecker/

Many possibilities exist for future work. Obvious next milestones are the ideals for the Chow varieties of degree 3 cycles in \mathbb{P}^3 , and degree 2 cycles in \mathbb{P}^4 . Methods from representation theory promise a compact encoding of their generators, in terms of irreducible $\text{GL}(4)$ -modules. Another question we aim to pursue is motivated by the geometry of condition numbers [1]: express the volume of a tubular neighborhood of a coisotropic quadric in $G(2, 4)$ as a function of \mathbf{c} .

References

1. Bürgisser, P., Cucker, F.: Condition: The Geometry of Numerical Algorithms. Springer, Heidelberg (2013)
2. Catanese, F.: Cayley forms and self-dual varieties. Proceedings of the Edinburgh Mathematical Society 57, 89–109 (2014)
3. Cayley, A.: On the theory of elimination. Cambridge and Dublin Math. J. 3, 116–120 (1848)
4. Chow, W.-L., van der Waerden, B.L.: Zur algebraischen Geometrie. IX. Über zugeordnete Formen und algebraische Systeme von algebraischen Mannigfaltigkeiten. Mathematische Annalen 113, 696–708 (1937)
5. DeConcini, C., Goresky, M., MacPherson, R., Procesi, C.: On the geometry of quadrics and their degenerations. Comment. Math. Helvetici 63, 337–413 (1988)
6. Gel'fand, I.M., Kapranov, M.M., Zelevinsky, A.V.: Discriminants, Resultants and Multidimensional Determinants. Birkhäuser, Boston (1994)
7. Grayson, D., Stillman, M.: Macaulay2, a software system for research in algebraic geometry. Available at www.math.uiuc.edu/Macaulay2/
8. Green, M., Morrison, I.: The equations defining Chow varieties. Duke Mathematical Journal 53, 733–747 (1986)
9. Sturmfels, B.: The Hurwitz form of a projective variety. [arXiv:1410.6703](https://arxiv.org/abs/1410.6703).